A mycobacterial DivIVA domain-containing protein involved in cell length and septation

Pickford, Hayleah; Alcock, Emily; Singh, Albel; Kelemen, Gabriella; Bhatt, Apoorva

DOI:

[10.1099/mic.0.000952](https://doi.org/10.1099/mic.0.000952)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Pickford, H, Alcock, E, Singh, A, Kelemen, G & Bhatt, A 2020, 'A mycobacterial DivIVA domain-containing protein involved in cell length and septation', *Microbiology*, vol. 2020, no. 9, 000952, pp. 817-825.
<https://doi.org/10.1099/mic.0.000952>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

A mycobacterial DivIVA domain-containing protein involved in cell length and septation

Hayleah Pickford¹, Emily Alcock², Albel Singh¹, Gabriella Kelemen² and Apoorva Bhatt^{1,*}

Abstract

Mycobacterial cells elongate via polar deposition of cell wall material, similar to the filamentous *Streptomyces* species, which contain a tip-organizing centre. Coiled-coiled proteins such as DivIVA play an important role in this process. The genome of *Mycobacterium tuberculosis*, the causative agent of tuberculosis, encodes many coiled-coil proteins that are homologous to DivIVA with a potential role in mycobacterial cell elongation. Here we describe studies on *Mycobacterium smegmatis* MSMEG_2416, a homologue of *M. tuberculosis* Rv2927c. Two previous independent studies showed that MSMEG_2416 was involved in septation (subsequently referred to as *sepIVA*). Contrary to these previous reports, we found *sepIVA* to be dispensable for growth in laboratory media by generating a viable null mutant. The mutant strain did, however, show a number of differences, including a change in colony morphology and biofilm formation that could be reversed on complementation with *sepIVA* as well as Rv2927c, the *sepIVA* homologue from *M. tuberculosis*. However, analysis of cell wall lipids did not reveal any alterations in lipid profiles of the mutant strain. Microscopic examination of the mutant revealed longer cells with more septa, which occurred at irregular intervals, often generating mini-compartments, a profile similar to that observed in the previous studies following conditional depletion, highlighting a role for *sepIVA* in mycobacterial growth.

INTRODUCTION

A majority of widely studied rod-shaped bacteria such as *Escherichia coli* and *Bacillus subtilis* elongate by the lateral deposition of new cell wall material along the whole length of the bacterium. It is the highly regulated septal positioning and formation that results in the formation of morphologically identical daughter cells. The essential bacterial tubulin homologue FtsZ [1] is crucial in the initiation of a contractile protofilament ring, the Z ring, at the site of septum formation. The Min system, active and highly studied in both *E. coli* and *B. subtilis*, is composed of a number of interacting proteins that inhibit Z ring formation at sites distant from the mid cell, resulting in septum formation at a defined range at the centre of the cell [2, 3]. *B. subtilis* possesses a septum-determining protein termed DivIVA [4], which recognizes membranes of negative curvature [5], hence its localization at the cell poles, and the septum during the initiation of septation [4]. DivIVA is responsible for the maintenance of a high concentration of the FtsZ inhibitor, MinC, at the cell poles,

ensuring correct positioning of septum formation at the mid cell [3].

Members of the genus *Mycobacterium*, including the tuberculosis-causing *Mycobacterium tuberculosis*, exhibit polar growth with newly synthesized cell wall deposited at the poles of the rod-shaped bacterium in contrast to the lateral deposition seen in *Bacillus* and other bacteria. Mycobacteria do not possess *min* homologues, yet they do possess a *divIVA* homologue, termed *wag31*. Depletion of Wag31 in the fast-growing saprophyte *Mycobacterium smegmatis* produced cells that were rounded at one pole, progressively becoming more coccoid in shape, and eventually, lysing [6]. Depletion of Wag31 resulted in reduced Wag31 localization at the cell poles, with the subsequent inability to direct nascent peptidoglycan synthesis at the poles [6]. The outer section of the mycobacterial cell wall consists of a lipid-rich layer that contain unique lipids, including mycolic acids [7], and Wag31 is also known to recruit enzymes involved in the biosynthesis of these lipidic components to poles of growing cells [8–11].

Received 26 November 2019; Accepted 29 June 2020; Published 17 July 2020

Author affiliations: ¹School of Biosciences and Institute of Microbiology and Infection, University of Birmingham, UK; ²School of Biological Sciences, University of East Anglia, Norwich, UK.

***Correspondence:** Apoorva Bhatt, a.bhatt@bham.ac.uk

Keywords: *Mycobacterium*; septation; division; cytoskeleton; polar growth; tuberculosis.

Abbreviations: BLAST, basic local alignment search tool; c.f.u., colony forming units; 2D-TLC, 2 dimension thin layer chromatography; IMD, intracellular membrane domain; PBP, penicillin binding protein; TIPOC, Tip Organising Centre; TSB, tryptic soy broth.

Three supplementary figures are available with the online version of this article.

000952 © 2020 The Authors

This is an open-access article distributed under the terms of the Creative Commons Attribution License. This article was made open access via a Publish and Read agreement between the Microbiology Society and the corresponding author's institution.

and at the time were guided solely by the predicted essentiality of the *M. tuberculosis* *sepIVA* gene (*Rv2927c*) [17], anticipating *sepIVA* to be an essential gene in *M. smegmatis*. However, prior to using a gene essentiality testing tool [18] to study cells conditionally depleted of *sepIVA*, we first attempted to generate a knockout of *sepIVA* in *M. smegmatis* to validate a potential inability to delete *sepIVA* in a wild-type (WT) strain. We transduced *M. smegmatis* mc²155 with phΔMSMEG2416, a recombinant phage designed to replace *sepIVA* with a hygromycin resistance cassette (*hyg*) by specialized transduction [19]. Surprisingly, we were able to generate hygromycin-resistant transductants, indicating a replacement of *sepIVA* with *hyg* in the WT strain. One such transductant was analysed by whole-genome sequencing, confirming the replacement of *sepIVA* with *hyg* (Fig. S1, available in the online version of this article). The transductant was selected for further analysis and is referred to as Δ*sepIVA*. The ability to generate a viable *sepIVA* mutant in WT *M. smegmatis* demonstrated that the gene was not essential for the viability and growth of *M. smegmatis* in laboratory media, which in this case was tryptic soy broth (TSB) agar. We further investigated possible factors that may have affected our ability to obtain a viable *sepIVA* mutant, contrary to the earlier reports of essentiality. The two previous studies used 7H9 and 7H10 for growth [14, 15], while we used TSB agar to select for transductants. To probe potential effects of media, we tested the ability of the Δ*sepIVA* strain to grow on plates of three Middlebrook media, 7H9+agar, 7H10 and 7H11, compared to TSB agar. While the mutant strain formed smaller colonies on the Middlebrook media plates, we did not observe any significant changes in the efficiency of obtaining colony-forming units (c.f.u.) between the different agar media, suggesting that the media used for the generation of recombinant strains did not play a role in the differing outcome of our study (Fig. S2). Next, we explored the possibility that a second, pre-existing mutation in a fraction of the transduced *M. smegmatis* cells may have enabled us to isolate a viable *sepIVA* mutant. Synthetic lethality can lead to an inability to generate knockouts of otherwise non-essential genes unless the lethality is abrogated by the concurrent loss of a second gene function. This has been observed in genes involved in the α-glucan pathway of *M. tuberculosis* [20]. Using the whole-genome sequence, we

performed a variant call analysis to compare single-nucleotide polymorphisms (SNPs) of the parental WT *M. smegmatis* strain and the Δ*sepIVA* mutant strain, comparing both to the reference genome sequence of *M. smegmatis* mc²155 (NC_008596.1). One particular SNP, in a gene required for septation, stood out from the list of SNPs found only in the Δ*sepIVA* strain: a single nucleotide change in the gene *ftsW* that resulted in the change of an aspartate residue at position 91 to an asparagine. FtsW is required for septation and for the mid-cell positioning of penicillin-binding protein 3 (PBP3) [21]. In an interaction unique to mycobacteria, FtsW also forms a ternary complex that includes FtsZ and PBP3 with a potential role in septal peptidoglycan synthesis [21]. Given its association with septation, we reckoned that if a pre-existing mutated *ftsW* gene rendered the resident cell amenable to the generation of a viable *sepIVA* mutant, expression of a WT copy of *ftsW* in the mutant strain would be lethal. However, we were able to obtain viable transformants when an integrating plasmid carrying *M. smegmatis* *ftsW* with its native promoter was electroporated into the Δ*sepIVA* strain, suggesting that the mutation in *ftsW* was unlikely to have influenced our ability to generate a viable *sepIVA* knockout in *M. smegmatis* (one such transformant is shown in Fig. S3). Our recombinant phage was designed to replace residues 466–602 of the 738 bp *sepIVA* gene with the *hyg-sacB* cassette from the allelic exchange vector, thus retaining a substantial section of the 5' end of *sepIVA* open reading frame after allelic exchange that resulted in the knockout strain. It is possible that if there was a readthrough into the replacement cassette after allelic exchange, a shorter peptide containing the first 155 amino acids of the N-terminus of SepIVA (246 aa long) would be produced, and subsequently aided the formation of a viable mutant strain. Using web-based translation tools [22] we were able to test the sequences obtained from the allelic exchange vector and predicted a 174 aa long peptide retaining 155 aa from the N-terminus of SepIVA to be formed. Due to a lack of antibodies against SepIVA, we do not have evidence that this peptide is produced in our Δ*sepIVA* strain, but the phenotypes displayed by our mutant strain (described below) suggest that while this putative shorter SepIVA may have enabled us to generate a viable knockout strain, it did not retain full functionality.

Fig. 2. Colonies of *M. smegmatis* WT, Δ*sepIVA*, Δ*sepIVA*-C and Δ*sepIVA*-CRv strains grown on TSB agar. Scale bar, 5 mm.

Deletion of *sepIVA* alters colony morphology and biofilm formation

While we were able to generate a null mutant of *sepIVA* in *M. smegmatis*, there was a striking difference in the appearance of colonies of Δ*sepIVA* as compared to the parental (wild-type WT) strain, *M. smegmatis* mc²155 (Fig. 2). Colonies of the Δ*sepIVA* strain appeared 'drier' and more convoluted compared to the WT strain, with the phenotype more apparent when cultured on tryptic soy agar supplemented with 0.05% Tween-80 (Fig. 2). Introduction of an integrating plasmid-borne copy of *sepIVA* with its native promoter into Δ*sepIVA* restored colony morphology to that of the WT strain, indicating that the observed changes were solely due to the loss of *sepIVA*. Colony morphology could also be restored

Fig. 3. Pellicles of *M. smegmatis* wild-type (WT), $\Delta sepIVA$ and $\Delta sepIVA$ -C strains cultured in 24 well plates containing Sauton's media (each well has a diameter of 15 mm). The pictures represent one of two repeats.

in $\Delta sepIVA$ transformed with a plasmid-borne copy of the *M. tuberculosis* *sepIVA* (Rv2927c), indicating that the *M. tuberculosis* homologue could functionally complement the $\Delta sepIVA$ mutant. The mutant also demonstrated an impaired ability to form air-water interface biofilms (pellicles) (Fig. 3). In spite of these differences of growth morphologies, we surprisingly observed no changes in the growth rates of the $\Delta sepIVA$ strain when compared to the WT and complemented strains (Fig. 4).

Cell wall lipid profiles of the $\Delta sepIVA$ mutant

Alterations in the outer cell wall lipids in mycobacteria are often responsible for changing colony morphology [23, 24]. Also, cell wall lipid biosynthesis enzymes are known to interact with the cytoskeletal machinery to co-ordinate cell wall biogenesis with polar growth [6, 25]. To probe changes in cell wall lipids in the mutants, we grew colonies of the WT, mutant and complemented strains on agar plates with [14 C]-acetic acid to label cell wall lipids. Apolar and polar fractions of the lipids were extracted from scraped colonies and analysed by two-dimensional thin-layer chromatography (2D TLC) (Figs 5 and 6). Surprisingly, no alterations in lipid profiles were seen in the mutant strain, indicating that the

Fig. 4. Growth curve of *M. smegmatis* WT, $\Delta sepIVA$ and $\Delta sepIVA$ -C strains in TSB supplemented with 0.05% Tween-80, and grown at 37 °C with agitation.

altered colony morphology was not caused by a change in cell wall lipid composition.

Loss of *sepIVA* affects average cell length

To further probe the effects of loss of *sepIVA*, we observed mid-log phase cultures of the mutant strain using light microscopy, comparing them to those of the WT strain. Cells of the $\Delta sepIVA$ strain were significantly longer ($P < 0.05$), when compared to the WT and complemented strains (Fig. 7). The average length of cells of the mutant strain was 8.8 μ m compared to 5.4 μ m for the WT strain and 6.5 μ m for the complemented strain, suggesting that loss of *sepIVA* affected average cell length, possibly due to defects in division.

Loss of *sepIVA* leads to altered septation patterns

To further query the long cell phenotype we observed in the $\Delta sepIVA$ strain, we investigated the formation of septa in the mutant strain using fluorescent vancomycin staining. The mutant strain showed an irregular, more frequent septation pattern when compared to the WT strain; the longer cells of $\Delta sepIVA$ had more cross walls and often generated mini-compartments (Fig. 8). Septation phenotypes were restored to WT patterns in the complemented strain. These findings showed that loss of *sepIVA* dysregulated patterns of septum formation in *M. smegmatis*.

DISCUSSION

Mycobacteria require both common and bespoke cytoskeletal proteins that drive the distinct polar growth pattern seen in this genus and other related bacterial genera, such as *Streptomyces*. At the start of the study described here we were interested in probing the function of *MSMEG_2416* (termed *sepIVA*, and a homologue of *M. tuberculosis* Rv2927c), a gene encoding a coiled-coiled protein that shares a domain with the septum forming protein DivIVA. We were able to generate a viable null mutant of *M. smegmatis* *sepIVA* to help us address the role of this gene in mycobacterial growth. During the course of this study, two independent reports on mycobacterial septal factors were published [14, 15], outlining a role for *sepIVA* in septum formation in mycobacterial cells. Contrary to our studies, both studies reported *sepIVA* as an essential gene in *M. smegmatis*. We thus continued our studies and characterized the *M. smegmatis* *sepIVA* mutant we had generated. Deletion of *sepIVA* also seemed to produce longer cells and affected the culture characteristics of the strain, including the formation of altered colony morphology. It also affected septation patterns in the mutant strain, which produced irregular septation.

Similar to our approach, Jain *et al.* used specialized transduction for allelic exchange of the native copy of *sepIVA*, but using a FLAS-tagged, Tet-regulated second copy of *sepIVA* in a merodiploid strain [15]. Wu *et al.*, on the other hand, were unable to generate a conditional mutant using an approach that involved generating a merodiploid with *sepIVA* under the control of a non-native promoter. Instead they engineered a strain containing a DAS-tag at the C-terminus of *SepIVA*. The

Fig. 5. 2D TLC of apolar lipids extracted from *M. smegmatis* WT, $\Delta sepIVA$ and $\Delta sepIVA-C$ strains grown on TSB agar containing [^{14}C]-acetic acid at 37 °C for 7 days. The presence of Tween-80 on the plates is indicated by a '+'. Solvent systems A-D are as described [28]. TAG, triacylated glycerol; DAG, diacylated glycerol; TDM, trehalose monomycolate; GMM, glucose monomycolate; FFA, free fatty acids, FMA, free mycolic acids.

DAS-tag made SepIVA amenable to conditional depletion by controlled degradation, which led to the formation of filamentous cells [14]. It is unlikely that downstream effects on *MSMEG_2417* might have led to a different outcome of essentiality in these two studies, as *MSMEG_2417* is not essential in *M. smegmatis* [26]. Moreover, the conditional mutants

described in these studies showed similar phenotypes to WT *M. smegmatis* under conditions that allowed for expression of SepIVA function [14, 15]. *Rv2927c* is predicted to be essential in *M. tuberculosis*, but *M. smegmatis* can tolerate the loss of genes known to be essential in *M. tuberculosis*. Moreover, a study by Dragset *et al.* [26] identified an *in vitro* gene

Fig. 6. 2D TLC of polar lipids extracted from *M. smegmatis* WT, $\Delta sepIVA$ and $\Delta sepIVA-C$ strains grown on TSB agar containing [^{14}C]-acetic acid at 37 °C for 7 days. The presence of Tween-80 on the plates is indicated by a '+'. Solvent systems D and E are as described [28]. DAT, diacylated trehalose; PI, Ac_2PIM_2 and Ac_2PIM_6 indicate phospholipids.

Fig. 7. Microscopy and cell length analysis of WT *M. smegmatis*, $\Delta sepIVA$ and $\Delta sepIVA-C$. Strains were cultured in TSB supplemented with 0.05% Tween-80, and visualized at 100 $\times$ magnification. (a) A random sample of 250 cells from mid-exponential phase cultures (two biological repeats) were measured to determine average cell lengths. The deletion of *sepIVA* resulted in the production of significantly longer cells (b), when compared to WT *M. smegmatis* and $\Delta sepIVA-C$. ****, $P < 0.0001$. Scale bar, 10 μ M.

essentiality set for *M. smegmatis* that indicated that *sepIVA* is a non-essential gene.

We queried various reasons likely to explain the contrast between our report of non-essentiality of *sepIVA* and those of *sepIVA* essentiality in the two previous studies [14, 15]. We ruled out the role of growth media, as the mutant generated in this study was able to grow on the media used in the previous reports. We did identify a SNP in another cell division-associated gene, *ftsW*, and queried its role in the viability of our $\Delta sepIVA$ mutant. We postulated that if the WT copy of *ftsW* was dominant, its introduction into the $\Delta sepIVA$ strain would cause lethality. However, the transformants obtained after electroporation of $\Delta sepIVA$ were viable after plating on selective media. It was also likely that the mutated *ftsW* allele was the dominant allele. However, the ability to restore all phenotypes of $\Delta sepIVA$ to those of a WT strain solely by complementation with a recombinant copy of *sepIVA* suggest that the mutated *ftsW* allele had no role to play in the observed phenotypes, and was unlikely to have affected the ability to generate a viable *sepIVA* mutant. The complete open reading frame for *sepIVA* was not deleted in our $\Delta sepIVA$ mutant and it was possible that a putative 174 aa long peptide retaining 155 aa from the N-terminus of SepIVA was formed due to a readthrough into the *hyg-sacB* replacement cassette. As we

Fig. 8. Fluorescence microscopy and analysis of septation in WT *M. smegmatis*, $\Delta sepIVA$ and $\Delta sepIVA-C$. Strains were cultured to mid-exponential phase in TSB supplemented with 0.05% Tween-80, and the cell membrane and chromosomal matter was stained using fluorescent vancomycin (green) and propidium iodide (red), respectively, and one representative image is shown for each strain (a). The distance between septa of a population of cells was recorded and analysed statistically (b). A total of 116, 232 and 32 septal distance measurements (septum to septum, or septum to cell end) were made for WT, $\Delta sepIVA$ and $\Delta sepIVA-C$, respectively (two biological replicates). *SepIVA* deletion resulted in aberrant septation, with the distance between septa to septa or cell end being statistically shorter than for WT *M. smegmatis*. ****, $P < 0.0001$.

do see a septation defect in our mutant $\Delta sepIVA$ strain, this shorter hypothetical peptide was unlikely to have retained functions directly related to septation. However, we cannot rule out a second role for the N-terminal of SepIVA in distinct interactions with the cell division apparatus, which dictate the essentiality of *sepIVA* in mycobacteria. Thus, after considering various possibilities, including the potential role of growth media, suppressor mutations and a potential truncated SepIVA produced in the $\Delta sepIVA$ strain, we were unable to conclusively identify a sole defining reason for our ability to obtain a viable null mutant of *sepIVA* in contrast to the two previous reports of essentiality. However, despite the differences in reports of essentiality, the rest of our studies report similar phenotypes to the conditional mutants characterized in these previous studies, particularly the elongated cells and altered septation patterns. Thus, this work, along with the previous findings of Wu *et al.* [14] and Jain *et al.* [15], does identify a key role for *sepIVA* in septation and warrants further studies to decipher its precise role.

METHODS

Bacterial strains and culture conditions

M. smegmatis mc²155 was cultured in tryptic soy broth (TSB) supplemented with Tween-80 (0.05%) to prevent clumping. *E. coli* strains (TOP10 and HB101) were cultured in Luria-Bertani broth. All bacterial strains were cultured at 37°C with agitation. Where required, the following antibiotics were

Table 1. List of bacterial strains, phages and plasmids used in this work

Description		Source
Bacterial strains		
WT	<i>M. smegmatis</i> mc ² 155	[29]
$\Delta sepIVA$	<i>M. smegmatis</i> mc ² 155 in which <i>sepIVA</i> (<i>MSMEG2416</i>) is replaced with <i>hyg</i>	This work
$\Delta sepIVA-C$	Complemented strain of $\Delta sepIVA$, containing pMV306- <i>MSMEG2416</i>	This work
$\Delta sepIVA-CRv$	Complemented strain of $\Delta sepIVA$, containing pMV306- <i>Rv2927c</i>	This work
Plasmids		
p0004s	Allelic exchange substrate vector containing <i>hyg</i>	[19]
p $\Delta MSMEG_{2416}$	p0004s derivative constructed for the allelic exchange of <i>sepIVA</i> (<i>MSMEG_{2416}</i>)	This work
pMV306	Integrative <i>E. coli</i> - <i>Mycobacterium</i> shuttle vector; kan ^r	[27]
pMV306- <i>MSMEG2416</i>	pMV306 containing <i>MSMEG2416</i> with its native promoter	This work
pMV306- <i>Rv2927c</i>	pMV306 containing <i>Rv2927c</i> with its native promoter	This work
Phages		
phAE159	Temperature-sensitive derivative of mycobacteriophage TM4	[30]
ph $\Delta MSMEG2416$	phAE159 derivative designed for replacing <i>sepIVA</i> (<i>MSMEG2416</i>) with <i>hyg</i> .	This work

used for selection: hygromycin (150 $\mu\text{g ml}^{-1}$ for *E. coli* and 100 $\mu\text{g ml}^{-1}$ for *M. smegmatis*) and kanamycin (50 $\mu\text{g ml}^{-1}$ for *E. coli* and 25 $\mu\text{g ml}^{-1}$ for *M. smegmatis*). When examining pellicle formation of *M. smegmatis* strains, cultures were diluted in Sauton's media to OD₆₀₀ 0.05, and were cultured in 24-well plates. Cultures were incubated at 37°C, 5% CO₂, with pellicles examined following 4 days of incubation. For testing survival on Middlebrook media, 10 ml of serial 10-fold dilutions of the *M. smegmatis* strains were plated on 7H9 broth+1.5% agar, 7H10 or 7H11 plates (performed in triplicate).

Construction of mutant strains

All of the plasmids and phages utilized in this work are outlined in Table 1. Approximately 1 kb of the left and right flanking regions of *MSMEG2416* were PCR-amplified from *M. smegmatis* mc²155 genomic DNA using the primer pairs MS2416_LL (5'-TTTTTTTTTCCATAAATTGGCCTCG AAGAGGAACACAAG-3') and MS2416_LR (5'-TTTT TTTTCCATTCTTGGGATGTTGCCGTCTCGATG-3'), and MS2416_RL (5'-TTTTTTTTTCCATAGATTGGT TCGTGGCGAGTGCGACATC-3') and MS2416_RR (5'-TTTTTTTTTCCATCTTTTGGGATGGCGGTGGTACAGC

TCTTC-3'). Van9II restriction sites were incorporated into the 5' end of each primer. The resultant PCR products were digested with Van9II and cloned into Van9II-digested p0004s. Recombinant plasmids were verified by Van9II digestion and sequencing. The sequence confirmed plasmid was linearized with PacI and cloned into PacI-digested pAE159. The resultant *phΔMSMEG2416* phasmid DNA was used to generate high-titre phage particles, and specialized transduction was performed as described previously [19]. *M. smegmatis ΔsepIVA* was confirmed by Southern blot analysis and whole-genome sequencing.

MSMEG2416 and *Rv2927c*, and their native promoters, were PCR-amplified from *M. smegmatis* mc²155 and *M. tuberculosis* H37Rv genomic DNA, respectively, using the primer pairs MS2416_F (5'-GCGTCTAGAGATCGCGCCGACGCGTCCGC-3') and MS2416_R (5'-GCGAAGCTTGCGGTGCTTCGCGCGTTCG-3'), and *Rv2927c_F* (5'-GCGTCTAGACACGCTCGGGCAAGATCGCC-3') and *Rv2927c_R* (5'-GCGAAGCTTGCCATAAGAGAAATCCTAC-3'). Digested PCR products were cloned into the digested integrative vector, pMV306 [27]. The resultant plasmids were verified by digestion and sequencing (Table 1). Sequence-confirmed plasmids were electroporated into *M. smegmatis* mc²155 *ΔsepIVA*. Transformants were screened by selection on kanamycin and hygromycin plates, and confirmed as complemented strains by PCR. One such transformant complemented with *MSMEG_2416 (sepIVA)* was termed *ΔsepIVA-C* and one with *Rv2927c* was termed *ΔsepIVA-CRv*. Similarly, to test the 'poisoning' effect of WT *ftsW* on the *sepIVA* mutant, *M. smegmatis ftsW* was PCR-amplified with its native promoter sequence and cloned into pMV306. The plasmid was transformed into both WT *M. smegmatis* mc²155 and *ΔsepIVA*.

Microscopic examination of *M. smegmatis* strains

When measuring cell lengths, cells from mid-exponential phase cultures were visualized using a Nikon A1R inverted confocal microscope. A random sample of 250 individual cells were measured (visualized from randomly selected regions of the slides). Statistical analysis of cell lengths was performed with a two-way analysis of variance (ANOVA), using Bonferroni for multiple comparisons. *, $P < 0.05$. For fluorescence microscopy, chromosomal material was visualized by staining with propidium iodide, and newly synthesized peptidoglycan in septa was stained using fluorescent vancomycin in the same way as described for *Streptomyces* [13], using mid-exponential phase cultures. Staining of nascent peptidoglycan was performed by incubating growing cells using $2 \mu\text{g ml}^{-1}$ BODIPY FL vancomycin (Molecular Probes) and $2 \mu\text{g ml}^{-1}$ unlabelled vancomycin (Sigma), together with propidium iodide (Sigma, $10 \mu\text{g ml}^{-1}$) as described for *Streptomyces* [13]. Samples stained using fluorescent vancomycin were not fixed; and while propidium iodide is not expected to stain live, non-fixed samples, we found that $10 \mu\text{g ml}^{-1}$ propidium iodide routinely stained chromosomes of live *M. smegmatis* cells as well as cells of *Streptomyces*. Samples were viewed using a

Zeiss Axioplan 2 microscope with an AxioCamMR camera and 100×1.4 NA Ph3 objective.

Funding

The authors received no specific grant from any funding agency.

Acknowledgements

H. P. is funded by a BBSRC MIBTP (Midlands Integrative Biosciences Training Partnership) PhD studentship. We would also like to thank Alessandro Di Maio of the University of Birmingham, Birmingham Advanced Light Microscopy Facility for advice on fluorescence microscopy. Genome sequencing was provided by MicrobesNG (<http://www.microbesng.uk>), which was supported by the BBSRC (grant number BB/L024209/1).

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Erickson HP. Ftsz, a tubulin homologue in prokaryote cell division. *Trends Cell Biol* 1997;7:362–367.
2. de Boer PA, Crossley RE, Rothfield LI. A division inhibitor and a topological specificity factor coded for by the minicell locus determine proper placement of the division septum in *E. coli*. *Cell* 1989;56:641–649.
3. Marston AL, Thomaides HB, Edwards DH, Sharpe ME, Errington J. Polar localization of the mind protein of *Bacillus subtilis* and its role in selection of the mid-cell division site. *Genes Dev* 1998;12:3419–3430.
4. Edwards DH, Errington J. The *Bacillus subtilis* DivIVA protein targets to the division septum and controls the site specificity of cell division. *Mol Microbiol* 1997;24:905–915.
5. Lenarcic R, Halbedel S, Visser L, Shaw M, Wu LJ et al. Localisation of DivIVA by targeting to negatively curved membranes. *Embo J* 2009;28:2272–2282.
6. Kang C-M, Nyayapathy S, Lee J-Y, Suh J-W, Husson RN. Wag31, a homologue of the cell division protein DivIVA, regulates growth, morphology and polar cell wall synthesis in mycobacteria. *Microbiology* 2008;154:725–735.
7. Daffé M, Draper P. The envelope layers of mycobacteria with reference to their pathogenicity. *Adv Microb Physiol* 1998;39:131–203.
8. Mukherjee P, Sureka K, Datta P, Hossain T, Barik S et al. Novel role of Wag31 in protection of mycobacteria under oxidative stress. *Mol Microbiol* 2009;73:103–119.
9. Plocinski P, Arora N, Sarva K, Blaszczyk E, Qin H et al. *Mycobacterium tuberculosis* CwsA interacts with CrgA and Wag31, and the CrgA-CwsA complex is involved in peptidoglycan synthesis and cell shape determination. *J Bacteriol* 2012;194:6398–6409.
10. Xu W-xi, Zhang L, Mai J-tao, Peng R-chao, Yang E-zhuo et al. The Wag31 protein interacts with AccA3 and coordinates cell wall lipid permeability and lipophilic drug resistance in *Mycobacterium smegmatis*. *Biochem Biophys Res Commun* 2014;448:255–260.
11. Meniche X, Otten R, Siegrist MS, Baer CE, Murphy KC et al. Subpolar addition of new cell wall is directed by DivIVA in mycobacteria. *Proc Natl Acad Sci U S A* 2014;111:E3243–E3251.
12. Bagchi S, Tomenius H, Belova LM, Ausmees N. Intermediate filament-like proteins in bacteria and a cytoskeletal function in *Streptomyces*. *Mol Microbiol* 2008;70:1037–1050.
13. Holmes NA, Walshaw J, Leggett RM, Thibessard A, Dalton KA et al. Coiled-Coil protein Scy is a key component of a multiprotein assembly controlling polarized growth in *Streptomyces*. *Proc Natl Acad Sci U S A* 2013;110:E397–E406.
14. Wu KJ, Zhang J, Baranowski C, Leung V, Rego EH et al. Characterization of conserved and novel septal factors in *Mycobacterium smegmatis*. *J Bacteriol* 2018;200.

15. Jain P, Malakar B, Khan MZ, Lochab S, Singh A *et al.* Delin-eating FtsQ-mediated regulation of cell division. *J Biol Chem* 2018;293:12331–12349.
16. Hayashi JM, Luo C-Y, Mayfield JA, Hsu T, Fukuda T *et al.* Spatially distinct and metabolically active membrane domain in mycobac-teria. *Proc Natl Acad Sci U S A* 2016;113:5400–5405.
17. Griffin JE, Gawronski JD, Dejesus MA, Iøerger TR, Akerley BJ *et al.* High-resolution phenotypic profiling defines genes essential for mycobacterial growth and cholesterol catabolism. *PLoS Pathog* 2011;7:e1002251.
18. Bhatt A, Jacobs WR. Gene essentiality testing in mycobacterium smegmatis using specialized transduction. *Methods Mol Biol* 2009;465:325–336.
19. Bardarov S, Bardarov S, Pavelka MS, Sambandamurthy V, Larsen M, Tufariello J *et al.* Specialized transduction: an efficient method for generating marked and unmarked targeted gene disruptions in Mycobacterium tuberculosis, M. bovis BCG and M. smegmatis. *Microbiology* 2002;148:3007–3017.
20. Kalscheuer R, Syson K, Veeraraghavan U, Weinrick B, Biermann KE *et al.* Self-Poisoning of Mycobacterium tuberculosis by targeting GlgE in an alpha-glucan pathway. *Nat Chem Biol* 2010;6:376–384.
21. Datta P, Dasgupta A, Singh AK, Mukherjee P, Kundu M *et al.* Interaction between FtsW and penicillin-binding protein 3 (PBP3) directs pbp3 to mid-cell, controls cell septation and mediates the formation of a trimeric complex involving FtsZ, FtsW and PBP3 in mycobacteria. *Mol Microbiol* 2006;62:1655–1673.
22. Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD *et al.* ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res* 2003;31:3784–3788.
23. Hunter RL, Venkataprasad N, Olsen MR. The role of trehalose dimycolate (cord factor) on morphology of virulent M. tuberculosis in vitro. *Tuberculosis* 2006;86:349–356.
24. Rastogi S, Singh AK, Pant G, Mitra K, Sashidhara KV *et al.* Down-Regulation of PE11, a cell wall associated esterase, enhances the biofilm growth of Mycobacterium tuberculosis and reduces cell wall virulence lipid levels. *Microbiology* 2017;163:52–61.
25. Jani C, Eoh H, Lee JJ, Hamasha K, Sahana MB *et al.* Regulation of polar peptidoglycan biosynthesis by Wag31 phosphorylation in mycobacteria. *BMC Microbiol* 2010;10:327.
26. Dragset MS, Iøerger TR, Zhang YJ, Mærk M, Ginbot Z *et al.* Genome-Wide phenotypic profiling identifies and Categorizes genes required for mycobacterial low iron fitness. *Sci Rep* 2019;9:11394.
27. Stover CK, de la Cruz VF, Fuerst TR, Burlein JE, Benson LA *et al.* New use of BCG for recombinant vaccines. *Nature* 1991;351:456–460.
28. Dobson G, Minnikin DE, Minnikin SM, Parlett JH, Goodfellow M. *Systematic Analysis of Complex Mycobacterial Lipids: In Chemical Methods in Bacterial Systematics*; 1985. pp. 237–266.
29. Snapper SB, Melton RE, Mustafa S, Kieser T, Jacobs WR. Isolation and characterization of efficient plasmid transfor-mation mutants of Mycobacterium smegmatis. *Mol Microbiol* 1990;4:1911–1919.
30. Kriakov J, Lee S, Jacobs WR. Identification of a regulated alkaline phosphatase, a cell surface-associated lipoprotein, in *Mycobacte-rium smegmatis*. *J Bacteriol* 2003;185:4983–4991.

Edited by: R. Manganelli and P. Brodin

Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at microbiologyresearch.org.