Indomethacin markedly blunts cerebral perfusion and reactivity, with little cognitive consequence in healthy young and older adults

Shoemaker, Leena; Wilson, Luke; Lucas, Sam; Machado, Liana; Walker, Robert; Cotter, J.D.

DOI:

[10.1113/JP280118](https://doi.org/10.1113/JP280118)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Shoemaker, L, Wilson, L, Lucas, S, Machado, L, Walker, R & Cotter, JD 2021, 'Indomethacin markedly blunts cerebral perfusion and reactivity, with little cognitive consequence in healthy young and older adults', *The Journal of Physiology*, vol. 599, no. 4, pp. 1097-1113. <https://doi.org/10.1113/JP280118>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Shoemaker, L.N., Wilson, L.C., Lucas, S.J.E., Machado, L., Walker, R.J. and Cotter, J.D. (2021), Indomethacin markedly blunts cerebral perfusion and reactivity, with little cognitive consequence in healthy young and older adults. *J Physiol*, 599: 1097-1113., which has been published in final form at: <https://doi.org/10.1113/JP280118>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The Journal of Physiology

<https://jp.msubmit.net>

JP-RP-2020-280118R1

Title: Indomethacin markedly blunts cerebral perfusion and reactivity, with little cognitive consequence in healthy young and older adults.

Authors: Leena Shoemaker
Luke Wilson
Samuel Lucas
Liana Machado
Robert Walker
James Cotter

Author Conflict: No competing interests declared

Author Contribution: Leena Shoemaker: Conception or design of the work; Acquisition or analysis or interpretation of data for the work; Drafting the work or revising it critically for important intellectual content; Final approval of the version to be published; Agreement to be accountable for all aspects of the work Luke Wilson: Conception or design of the work; Acquisition or analysis or interpretation of data for the work; Drafting the work or revising it critically for important intellectual content; Final approval of the version to be published; Agreement to be accountable for all aspects of the work Samuel Lucas: Conception or design of the work; Acquisition or analysis or interpretation of data for the work; Drafting the work or revising it critically for important intellectual content; Final approval of the version to be published; Agreement to be accountable for all aspects of the work Liana Machado: Conception or design of the work; Acquisition or analysis or interpretation of data for the work; Drafting the work or revising it critically for important intellectual content; Final approval of the

Disclaimer: This is a confidential document.

version to be published; Agreement to be accountable for all aspects of the work Robert Walker: Acquisition or analysis or interpretation of data for the work; Drafting the work or revising it critically for important intellectual content; Final approval of the version to be published; Agreement to be accountable for all aspects of the work James Cotter: Conception or design of the work; Acquisition or analysis or interpretation of data for the work; Drafting the work or revising it critically for important intellectual content; Final approval of the version to be published; Agreement to be accountable for all aspects of the work

Running Title: CEREBRAL BLOOD FLOW, COGNITION, AND AGING

Dual Publication: No

Funding: University of Otago - School of Physical Education, Sport and Exercise Sciences: Leena N Shoemaker, N/A; University of Otago - Department of Medicine: Leena N Shoemaker, N/A We acknowledge funding support from the Lottery Health Research grant, Department of Internal Affairs to LCW, which assisted with the ethics application and protocol design.

Indomethacin markedly blunts cerebral perfusion and reactivity, with little cognitive consequence in healthy young and older adults.

Shoemaker, L.N.¹, Wilson, L.C.², Lucas, S.J.E.^{3,5,6}, Machado, L.⁴, Walker, R.J.², Cotter, J.D.¹

¹ School of Physical Education, Sport and Exercise Sciences, University of Otago, Dunedin, NZ

² Dunedin School of Medicine, University of Otago, Dunedin, NZ

³ Department of Physiology, University of Otago, Dunedin, NZ

⁴ Department of Psychology, University of Otago, Dunedin, NZ

⁵ School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, UK

⁶ Centre for Human Brain Health, University of Birmingham, Birmingham, UK

ORCID ID

Shoemaker (0000-0003-3924-0717)

Wilson (0000-0003-2636-098X)

Lucas (0000-0002-8713-2457)

Machado (0000-0002-0856-3831)

Walker (0000-0003-3366-09560)

Cotter (0000-0002-6014-9865)

Correspondence: James D. Cotter

School of Physical Education, Sport and Exercise Sciences,

University of Otago, Dunedin, NZ

+64 3 479 9109

jim.cotter@otago.ac.nz

Running Title: Cerebral blood flow, cognition, and aging

Key Points Summary

- Cognitive function depends on adequate cerebrovascular perfusion and control. However, it is unknown if acutely-reduced cerebral blood flow (CBF) impairs cognition in healthy adults.
- In this study we used a placebo-controlled, single-blinded, randomised cross-over design to test the hypothesis that acutely-reduced CBF (using a pharmacological aid; indomethacin) would impair cognition in young and older healthy adults.
- At baseline, older adults had lower cognitive performance and CBF, but similar cerebrovascular reactivity to CO₂ and dynamic cerebral autoregulation compared to young adults.
- In both young and older adults, cognitive performance on a mental switching task was slightly (7%) reduced after indomethacin, but not significantly associated with reductions in CBF (~31%).
- These results indicate that cognitive performance is broadly resilient against a ~31% reduction in CBF *per se* in healthy young and older adults.

Abstract

Cognitive function depends on adequate cerebrovascular perfusion and control. However, it is unknown if acutely-reduced cerebral blood flow (CBF) impairs cognition in healthy adults. Using a placebo-controlled, single-blinded, randomised cross-over design, we tested the hypothesis that acutely-reduced CBF (using indomethacin [1.2 mg/kg oral dose]) would impair cognition in young (n=13; 25±4 y) and older (n=12; 58±6 y) healthy adults. CBF and cerebrovascular control were measured using middle cerebral artery blood velocity (MCAv_{mean}) and its reactivity to hypercapnia (CVR_{HYPER}) and hypocapnia (CVR_{HYPO}), respectively. Cognitive function was assessed using a computerised battery including response time tasks. Baseline comparisons revealed that older adults had 14% lower MCAv_{mean} and 15% lower cognitive performance (all p≤0.048) but not lower CVR_{HYPER/HYPO} (p≥0.26). Linear and rank-based mixed models revealed that indomethacin decreased MCAv_{mean} by 31% [95% CI:-35,-26], CVR_{HYPER} by 68% [IQR:-94,-44] and CVR_{HYPO} by 50% [IQR:-83,-33] (treatment-effect; all p<0.01), regardless of age. Baseline CVR_{HYPER/HYPO} were strongly associated with their indomethacin-induced reductions (r=0.70 to 0.89, p<0.01). Mental switching performance was impaired 7% [IQR:0,19] after indomethacin (p=0.04), but not significantly associated with reductions in MCAv_{mean} (Young: rho=-0.31, p=0.30; Older: rho=0.06, p=0.86). Conclusion: indomethacin reduced MCAv_{mean} and impaired cognition slightly, however no

- 1 clear association was evident in younger or older adults. Older adults had poorer cognition and
- 2 lower $\text{MCAv}_{\text{mean}}$ but similar $\text{CVR}_{\text{HYPER/HYPO}}$.
- 3 **Keywords:** aging, cognition, cerebral blood flow, hypercapnia, indomethacin

1 Introduction

2 Cognitive function has an immediate and critical reliance on adequate cerebrovascular control,
 3 perfusion, and metabolism (Barnes *et al.*, 2013). While an acute increase in perfusion does not
 4 measurably improve cognitive performance (Shoemaker *et al.*, 2019b; Shoemaker *et al.*, 2020), the
 5 extent to which cognition is resilient against *lower* cerebral perfusion is unknown. To our
 6 knowledge, no one has imposed a (reversible) reduction of cerebral blood flow (CBF) in healthy
 7 individuals to test if an *acute* reduction in CBF or cerebrovascular responsiveness impairs
 8 cognition.

9 It is possible to acutely reduce CBF and cerebrovascular reactivity to CO₂ (CVR_{CO2}; i.e.,
 10 modulation of vascular tone in response to increases and/or decreases in CO₂) in young and older
 11 healthy adults pharmacologically. Indomethacin, a non-steroidal anti-inflammatory drug, inhibits
 12 the enzyme cyclooxygenase (COX) and thus prostaglandin synthesis. While similar to other
 13 COXinhibitors (e.g., naproxen, ibuprofen), only indomethacin reduces CBF in healthy humans,
 14 without changing cerebral metabolic rate (Hohimer *et al.*, 1985; Kraaier *et al.*, 1992) or plasma
 15 catecholamine concentrations (Wennmalm *et al.*, 1983; Staessen *et al.*, 1984; Green *et al.*, 1987).
 16 Indomethacin reduces basal CBF by 19 - 42% and CVR_{CO2} by 50 - 65% in both young and older
 17 adults (Eriksson *et al.*, 1983; Wennmalm *et al.*, 1983; Jensen *et al.*, 1993; Markus *et al.*, 1994;
 18 Kastrup *et al.*, 1999; Bruhn *et al.*, 2001; St. Lawrence *et al.*, 2002; Xie *et al.*, 2006; Ivancev *et al.*,
 19 2009; Xie *et al.*, 2009; Barnes *et al.*, 2012a; Hoiland *et al.*, 2015; Peltonen *et al.*, 2015; Hoiland *et al.*,
 20 2016; Peltonen *et al.*, 2016), and alters dynamic cerebral autoregulation (in young males; (Smirl
 21 *et al.*, 2014)).

22 Aging is associated with independent reductions in cerebral perfusion and some aspects of
 23 cognition, so older adults may be more cognitively susceptible to acute reductions in CBF (i.e.,
 24 lower reserve). A brief reduction in perfusion impairs cognition in patients with cardiovascular
 25 disease (Marshall *et al.*, 2001) and with end-stage kidney disease during haemodialysis (Findlay *et al.*,
 26 2019). Yet, the impact of acute reductions in cerebral perfusion on cognition has not been
 27 addressed in healthy adults. Young adults, with higher cognition, CBF and perhaps CVR_{CO2} (Lucas
 28 *et al.*, 2012; Bailey *et al.*, 2013), may be able to cognitively tolerate acute reductions in cerebral
 29 perfusion better than older counterparts (i.e., have a higher reserve).

30 Indomethacin reduces cerebral perfusion to a similar extent as occurs with healthy aging [i.e., ~30%
 31 (Ainslie *et al.*, 2008)]. Indomethacin therefore provides a means to acutely eliminate age-related

differences in resting CBF and the possible differences in CVR_{CO_2} , and their potential impact on cognitive function. Therefore, the primary aim of this study was to test whether and to what extent an acute reduction in CBF and cerebrovascular control would impair cognition in young and older adults. A secondary aim was to elucidate the extent to which age-related reductions in cognition at baseline are modulated by the usually-observed impairment in CBF and cerebrovascular control. Expecting that older adults would show lower baseline CBF (~30%) and lower cognitive performance in both response time and working memory tasks, we hypothesised that an acute reduction of CBF (i.e., with indomethacin) *per se* would impair cognition, but more so in older adults.

Methods

This study was approved by the New Zealand Central Health and Disability Ethics Committee (18/CEN/142) in accordance with the standards set by the Declaration of Helsinki. This study was prospectively registered in the Australian New Zealand Clinical Trials Registry on 27/09/2018 (ACTRN12618001603202). All participants gave written informed consent prior to data collection.

Participants

Prospective participants were invited to take part if they were aged 18-35 or 50-75 y, with no sign of cognitive impairment and were not smokers. Exclusion criteria were known cardiovascular, cerebrovascular, neurological, metabolic, respiratory, renal, or haematological disease or condition, or current usage of medication such as cardiac glycosides, aminoglycosides, diuretics, anticoagulants, antihypertensive, aspirin or corticosteroids. The use of nonsteroidal anti-inflammatory drugs was restricted for a minimum of 7 d prior to the initial study visit and continued through the final experimental visit. Older females were recruited only if they reported being postmenopausal or having experienced amenorrhea for a minimum of 12 mo. Young females were in luteal menstrual phase or on oral contraception (active pill phase) for all familiarisation and experimental visits. All participants were screened for cognitive impairment using the Montreal Cognitive Assessment (MoCA©). A score of 25 or higher was required for participation, as a score below 25 represents abnormally low cognitive performance (Nasreddine *et al.*, 2005).

Twenty-nine participants were recruited and undertook familiarisation. One subsequently moved to a different city, two were excluded due to insonation difficulties of the MCA and PCA from poor temporal windows, and one was excluded due to a low MoCA© score. Therefore, 25 participants

were subsequently randomised after familiarisation, using a computer-generated and counter-balanced allocation. The allocation sequence was concealed until the moment of assignment.

Participants reported to the laboratory having abstained from caffeine and food for a minimum of 2 h, and from strenuous exercise for 12 h. Diet and activity were kept consistent for the 24 h prior to each visit.

Experimental Procedures

This study used a placebo-controlled, single-blinded, randomised cross-over design, in which two groups (young vs. older) each completed two conditions in cross-over fashion (indomethacin vs. placebo). Seven of thirteen young adults completed the indomethacin condition first, while the twelve older adults were counter-balanced. Participant information is in Table 1. Participants were blinded to the treatment. Measures of cognitive function and cerebrovascular control were assessed before and after ingestion of indomethacin and a placebo (Figure 1). Conditions were undertaken at least 10 d after a familiarisation visit and separated by at least 72 h. Testing was completed between 7:00 AM and 9:00 PM, with time-of-day consistent within participants (within 30 min).

Familiarisation

During the familiarisation visit participants completed the MoCA®. After full instrumentation, participants completed familiarisation rounds of cerebrovascular reactivity to hypercapnia (CVR_{HYPER}) and hypocapnia (CVR_{HYPO}). In addition, they sufficiently practiced (as evidenced by a plateau in performance) the cognitive battery (i.e., Pro, Anti, Pro/Anti and Backward Digit Span) to minimise future concern of practice effects. Pro, Anti, Pro/Anti was practiced three times, while the Backward Digit Span was practiced once.

After the experimental conditions, below, participants undertook a fourth visit for estimating their peak rate of oxygen consumption ($\dot{V}O_{2peak}$) on the treadmill. A standard submaximal paradigm was used; 3 stages of four minutes, each with mildly increasing intensities (+10-15% of calculated heart rate reserve (Karvonen, 1957)) (Golding *et al.*, 1989; Akalan *et al.*, 2008). Heart rate (HR) and $\dot{V}O_2$ were averaged across the fourth minute of each stage, and $\dot{V}O_{2peak}$ subsequently estimated from a linear regression using predicted maximal HR (i.e., 220 minus age (Fox, 1973)).

Experimental Conditions

Participants attended the laboratory for the placebo and indomethacin conditions, where they sat comfortably in a semi-recumbent chair throughout (i.e., ~3.5 h), except when undertaking dynamic cerebral autoregulation (CA) tests (~8 min). After instrumentation and collection of baseline data, participants completed the cognitive battery twice and tests of dynamic CA, CVR_{HYPER} and CVR_{HYPO} . Measures of CVR_{CO_2} (CVR_{HYPER} followed by CVR_{HYPO}) were completed after dynamic CA as hypocapnia can profoundly affect cerebrovascular control (Ide *et al.*, 2003). Dynamic CA and $CVR_{HYPER/HYPO}$ were measured to characterise cerebrovascular control mechanisms alongside changes in cerebral blood velocity in anterior and posterior regions of the brain, in response to indomethacin in young and older adults. Participants then received 100 mg of an anti-nausea medication, Simethicone (De-Gas®, Pfizer Australia Pty Limited) and 1.2 mg/kg (rounded to the nearest 25 mg) of either placebo or indomethacin. Simethicone has previously been used (Barnes *et al.*, 2012b) to prevent GI upset, a potential side-effect of indomethacin.

After a minimum of 60 min of rest (during which participants read or watched Netflix, consistent between conditions), cognitive function and cerebrovascular control measures were repeated in the same sequence as at baseline.

Measurements

Cerebrovascular and Cardiovascular measures

Cerebral blood velocity was measured in both the middle and posterior cerebral arteries using Transcranial Doppler Ultrasound (Spencer Doppler, Sterling VA, USA). Depth was controlled between 45-60 mm for the middle cerebral artery (MCA) and 60-70 mm for the posterior cerebral artery (PCA). Carotid compression was used to differentiate the MCA from the PCA. The probe was then secured in place with a headband device (Marc 600 Headframe) to maintain insonation angle and position.

Heart rate and its rhythm were measured continuously using a standard lead-II electrocardiogram (ADInstruments, Dunedin, NZ). Blood pressure was measured on a beat-by-beat basis using finger photoplethysmography (Finometer®Pro, Finapres Medical Systems, Enschede, The Netherlands), calibrated against manual blood pressure measurements. Mean MCA velocity ($MCAv_{mean}$) and mean arterial pressure (MAP) were first calculated as the mean time integrals and then divided by cardiac period.

Participants breathed through a leak-free respiratory mask (Hans-Rudolf 8980, Kansas City, MO) attached to a turbine and two-way valve, as described in Shoemaker *et al.* (2019a). Partial pressure of end-tidal CO₂ (PETCO₂) was measured continuously using an online gas analyser (Model CD-3A Carbon Dioxide Analyser, AEI Technologies, Pittsburgh, USA). The PETCO₂ was controlled using a customised dynamic end-tidal clamping system (C.E.T. Gas Clamp, School of Physical Education, Sport and Exercise Sciences, University of Otago, Dunedin, NZ). Hypercapnia was achieved by titrating CO₂ into room air using solenoid-valve control, resulting in +5 and +10 mm Hg steps in PETCO₂ from baseline. A single step for hypocapnia (-10 mm Hg) was achieved by increasing ventilation with verbal coaching and visual feedback.

All data were sampled continuously (1 kHz) using an analogue-to-digital converter (Powerlab /16SP ML795; ADInstruments, Dunedin, NZ) and stored for later analysis on Labchart software (version 7, ADInstruments).

Cerebrovascular Control

For both CVR_{HYPER} and CVR_{HYPO}, data were averaged across 30 s after a minimum of 2 min from the onset of PETCO₂ manipulation, in agreement with the CVR_{HYPER} recommendations for lower error and individual variability (Burley *et al.*, 2020). The CVR_{HYPER/HYPO} were calculated using linear regression as the change in MCAv_{mean} divided by the change in mean PETCO₂ (cm/s/mm Hg). The CVR_{HYPER} was determined as the slope of two +5 mm Hg hypercapnia steps. Likewise, CVR_{HYPO} was determined as the slope of the change from baseline to -10 mm Hg. Exceptions to these procedures were as follows: 3 of 25 participants had one CVR_{HYPER} discarded and 8 of 25 had one CVR_{HYPO} discarded due to validity problems in baseline or elevated/reduced PETCO₂, and were thus excluded from inferential analyses of this measure; 6 of 25 participants had CVR_{HYPER} calculated from one CO₂ step, due to insufficient clamping of PETCO₂; 9 participants had one CO₂ epoch beginning 75 - 120 s after hypercapnia onset, for reasons of data integrity (e.g., prioritising signal measurement quality or stability), and in all cases these were time matched as closely as possible with their corresponding intra-trial CVR_{HYPER}.

Dynamic CA was determined using a sit-to-stand protocol, performed for 5 min at a frequency of 0.05 Hz. Transfer function analysis for pressure-flow relations was computed using commercially-available software (Ensemble version 1.0.0.14, Elucimed Ltd, Wellington, NZ) as described in Tzeng *et al.* (2012). Briefly, Ensemble uses the R-R interval from the electrocardiogram to obtain precise beat-to-beat BP and MCAv signals, which are spline interpolated and resampled at 4 Hz for

spectral analysis and TFA based on the Welch algorithm. Recordings are sub-divided into overlapping (50%) windows which are then linearly detrended and passed through a Hanning window prior to fast Fourier transform analysis. The cross-spectrum between the two signals (BP and MCAv) was divided by the auto spectrum of the input signal (BP) to derive transfer function values of coherence, absolute gain, and phase. These values were selectively analysed at the point estimate of the 0.05 Hz driven frequency. All coherence values included in analysis were above the statistically-calculated threshold (0.63). The Ensemble algorithms for TFA have been cross-validated in Meel-van den Abeelen *et al.* (2014) and Tzeng *et al.* (2012).

Cognitive Function

Choice response time, inhibition, and mental switching were tested using a response time battery consisting of Pro, Anti, and Pro/Anti tasks, respectively (described in Shoemaker *et al.* (2019b) and based on (Guiney *et al.*, 2015). Participants were instructed to respond to a green (Pro) or red (Anti) visual stimulus by pressing a button on the corresponding (Pro) or opposite (Anti) side to which the stimulus appeared. Response time (ms) and accuracy were recorded for all trials. Accuracy-adjusted response time (aRT) was used to account for any speed-accuracy trade-off, and was calculated as:

$$\text{aRT} = \text{Median response time of correct responses} / (1 - \text{error rate}).$$

This measure can be interpreted like unadjusted response time and has been used previously in peer-reviewed research (Guiney *et al.*, 2019) with high within-day reliability following multiple practise trials (Shoemaker *et al.*, 2020).

Working memory was assessed using the Backward Digit Span task. This task involves participants recalling a list of numbers in *reverse* order. The task started with 3 digits and continued up to 9. Each level had two trials, whereby the same number of digits was presented verbally (e.g., 4-8-1 and 5-3-8). An unsuccessful trial was where the digits were not recalled in perfect reverse order. The task was stopped when two unsuccessful sequential trials of the same length occurred. The highest recalled digit span was recorded.

Lastly, participants were asked to report their “feeling” (overall feeling state) pre- and post- placebo and indomethacin, using the 11-point Affect Feeling Scale (Giblin, 2011). Participants also reported any gastrointestinal upset and tiredness on a 7-point Likert scale.

Statistical Analysis

A priori power calculations with an expected moderate ($\eta p^2 = 0.06$) effect size revealed that a sample of 24 participants was needed to obtain 80% power with 2 groups and 4 measurements ($\alpha = 0.05$). Data were analysed using R (R Development Core Team, 2008) and graphed with GraphPad Prism (Prism Version 8, GraphPad Software, CA, USA). An alpha of 0.05 was used for each analysis.

Independent two-tailed t-tests were performed to compare baseline measures of cerebrovascular and cognitive characteristics between groups. Primary and secondary outcome variables were assessed for homogeneity of variances with Levene's test. Linearity and approximate normal distribution of model and individual residuals were assessed qualitatively using visual inspection of histograms and Q-Q plots and formally tested with Shapiro-Wilk's Test. If the assumptions of a parametric test were met (i.e., homogeneity of variance, linearity, normality, and independence), then the data were analysed using a linear mixed-effect model for repeated measures. Only $MCA_{v_{mean}}$, $PCA_{v_{mean}}$, MAP, CVC, and Phase (from CA measures) met these assumptions and were tested using the parametric approach. Variance-covariance structure, model inclusion of random and fixed factors, and weighting of model errors was assessed using Akaike's Information Criteria (AIC). At minimum, condition (2 levels: indomethacin, placebo), time (2 levels: pre, post), age (continuous variable), and their interactions were treated as fixed factors. Sex and fitness were included according to AIC values. Fitness did not have a significant fit to any model. Participant was included as a random effect. Post-hoc testing for significant interactions of parametric testing was completed with Tukey's HSD. All results from the linear mixed-effect models are reported using mean $\pm$ SD or 95% confidence intervals [CI: lower limit, upper limit].

Non-parametric rank-based analysis method was used for $CVR_{HYPER/HYPO}$, Gain and Coherence [from CA measures], heart rate, $PETCO_2$, and cognitive data using the R package developed by Noguchi *et al.* (2012), to assess the three-way interaction of age-by-condition-by-time. Due to the package allowing for only three factors, effects of sex were tested using an additional analysis, of condition-by-time-by-sex. P-values are reported from the ANOVA-type statistic. Multiple comparisons for significant interactions were completed with Bonferroni's post-hoc adjustment. All results from the nonparametric rank-based methods are reported as median and interquartile ranges [IQR: quartile 1, quartile 3]. Lastly, correlation data involving cognition were analysed using Spearman's rho correlation coefficient (ρ) while $CVR_{HYPER/HYPO}$ were analysed using Pearson correlation coefficients (r).

Reliability was assessed using intraclass correlation (ICC) estimates and their 95% confidence intervals, calculated based on a mean-rating, absolute agreement, 2-way mixed-effects (test-retest) model. Consecutive pairwise comparisons of reliability from 2 trials for both between-day (i.e., placebo and indomethacin) and within-day (e.g., pre- and post-placebo), as calculated by ICC, show excellent (>0.9), good ($0.76-0.9$), moderate ($0.5-0.75$), and poor (<0.5) reliability (Table 2, as described by Koo and Li (2016)). Reliability is presented also as the coefficient of variation ($SD/mean \times 100$).

Cognitive, cerebro- and cardio-vascular data are reported in the results from all 25 participants. However, $PCAv_{mean}$ was accessible for the entire protocol in only 20 participants ($n = 13$ young and $n = 7$ older). Measures of CA are reported from 24 participants due to a poor beat-to-beat blood pressure recording from one older adult ($n = 13$ young and $n = 11$ older). The CVR_{CO_2} measures are reported as described above.

In text and Figure 6, all means with 95% confidence intervals and medians with interquartile ranges are calculated from the difference between the change from indomethacin (i.e., ‘post’ minus ‘pre’) and the change from placebo, and thus represent the treatment effect.

Results

Reliability of measures

Resting measures of $MCAv_{mean}$ and $PCAv_{mean}$ (Table 2) showed good reliability across days (i.e., baseline measures) and excellent reliability within one day (placebo condition). aRT measures show good-to-excellent reliability within a single day and good reliability between days. The functional CBF measures of $CVR_{HYPER/HYPO}$ show good-to-excellent reliability within days and poor reliability between days.

Cerebro- and Cardio-vascular Responses

Baseline comparisons between groups revealed that $MCAv_{mean}$ was 14% lower in older adults ($p = 0.048$), whereas $PCAv_{mean}$, CVR_{HYPER} and CVR_{HYPO} were not significantly lower ($p \geq 0.263$). Older adults had lower heart rate (13%; $p = 0.049$) and CVC (17%; $p = 0.043$), whereas MAP ($p = 0.405$) and $PETCO_2$ ($p = 0.179$) were similar between groups. Older adults had lower aerobic fitness (estimated $\dot{V}O_{2peak}$) than young adults. However, according to a new $\dot{V}O_{2peak}$ calculation that accounts for age, sex, height, weight and exercise mode (de Souza e Silva *et al.*, 2018), both groups

were equivalently (Table 1; two-tailed T-test, $p = 0.308$) *more* fit than would be expected for their respective wider populations.

Older adults did not have greater cerebro- or cardio-vascular sensitivity to indomethacin, as all age-by-condition-by-time interactions were non-significant ($p \geq 0.249$). Furthermore, there were no significant main effects or interactions with sex for any cerebro- or cardio-vascular variable (all $p \geq 0.174$; Figure 2).

Indomethacin reduced $MCAv_{mean}$, CVR_{HYPER} , and CVR_{HYPO} (all interaction effects and subsequent pairwise comparisons $p < 0.001$ vs. pre-indomethacin and post-placebo). Specifically, $MCAv_{mean}$ declined by 31% [CI: -35, -26], CVR_{HYPER} by 68% [IQR: -94, -44] and CVR_{HYPO} by 50% [IQR: -83, -33], irrespective of age. Baseline CVR_{HYPER} (young: $r = 0.81$, older: $r = 0.89$) and CVR_{HYPO} (young: $r = 0.70$, older: $r = 0.89$) were strongly associated (all $p \leq 0.02$) with their indomethacin-induced change (Figure 3). Specifically, participants who had the largest CVR_{HYPER} and CVR_{HYPO} at baseline had the largest decrease post-indomethacin (Figure 3), in both young and older groups. Regardless of age, indomethacin increased MAP ($8 \pm 12\%$) and decreased HR ($7 \pm 8\%$), CVC ($34 \pm 17\%$) and $PCAv_{mean}$ ($26 \pm 14\%$; all time-by-condition interaction: $p \leq 0.001$).

Cerebral Autoregulation (Figure 4)

In all conditions, coherence was above the statistically-calculated threshold (i.e., > 0.63 , Figure 4A), allowing for the interpretation of gain and phase. Age groups were not different for coherence, gain and phase (all $p \geq 0.08$) at 0.05 Hz. Indomethacin dampened both BP and $MCAv_{mean}$ power ($p < 0.046$ vs. pre-indomethacin, $p < 0.007$ vs. post-placebo). It reduced gain by 30% ([IQR: -46, 1]; Figure 4B) and increased phase by 57% ([CI: 4, 110]; all $p < 0.001$ vs. pre-indomethacin and post-placebo). Coherence was lowered by 4% ([IQR: -11, 4]; condition-by-time interaction $p = 0.006$; $p < 0.001$ vs. pre-indomethacin and post-placebo). Coherence was 3% higher in males than females (main effect of sex: $p = 0.001$).

Cognitive Performance (Figure 5)

At baseline, older adults had 14-15% lower performance than young adults (all $p < 0.001$) in Pro (14%), Anti (15%) and Pro/Anti (15%) tasks, but not lower working memory ($p = 0.663$; Figure 5D). Furthermore, there were no significant main effects or interactions involving sex for any cognitive outcome measures (all $p \geq 0.377$).

Older adults' cognitive performance was not more affected by indomethacin than that of young adults' (age-by-condition-by-time interaction: all $p \geq 0.377$). Pro/Anti performance was 7% [IQR: 0, 19] worse post-indomethacin ($p \leq 0.042$ vs. pre-indomethacin and post-placebo), however Pro (4% [IQR: 0,11]) and Anti (4% [IQR: -4,17]) performance were not measurably affected (time-by-condition interaction; $p = 0.061$ and $p = 0.181$ respectively) but showed the same pattern as Pro/Anti.

Working memory improved during the placebo condition (1 AU [IQR: 0,1]; $p = 0.001$ vs pre-placebo and post-indomethacin) but not during the indomethacin condition (0 AU [IQR: -1,0]; $p = 0.219$ vs. pre-indomethacin). Participants reported slightly but significantly worse overall feelings after indomethacin than after placebo (-1 ± 1 vs 0 ± 1 AU, respectively; $p = 0.038$), but not more gastrointestinal upset or tiredness (both $p \geq 0.120$).

Associations between Changes in Cerebrovascular and Cognitive Function (Figure 6)

The indomethacin-induced reduction in $MCAv_{mean}$ was not associated with an acute change in aRT within either age group (all $\rho \leq 0.34$; $p \geq 0.249$; Figure 6). Additionally, changes in CVR_{HYPER} and CVR_{HYPO} were not reliably associated with Pro ($\rho = 0.22$ and $\rho = -0.18$, respectively), Anti ($\rho = -0.11$ and $\rho = -0.06$, respectively), or Pro/Anti performance ($\rho = -0.29$ and $\rho = 0.22$, respectively; all $p \geq 0.167$).

Discussion

The novel findings of the current study were that executive cognitive function (mental switching and short-term memory) was broadly resilient to a moderate (31%) acute reduction in $MCAv_{mean}$ in healthy young and older adults. Other findings are valuable because they have seldom been shown and/or are equivocal in the literature. Specifically, the current results show that: (i) those who had the largest cerebrovascular reactivities to CO_2 at baseline had the largest decrease post-indomethacin; (ii) healthy older adults had lower $MCAv_{mean}$ and cognitive performance than healthy younger adults did, which is consistently reported, but (iii) they did not have lower CVR_{HYPER} , CVR_{HYPO} , or CA, which is equivocal in the literature.

The effects of indomethacin on cognitive function and cerebral perfusion.

The current findings only partially supported our hypothesis that an acute reduction in $MCAv_{mean}$ (i.e., 31% reduction with indomethacin) *per se* would impair cognition function and would be more evident in older adults. The cognitive function domain of mental switching ability (i.e., Pro/Anti)

was impaired by ~7% in young and older adults post-indomethacin, and showed a small to moderate effect size (Cohen's $d_z = 0.42$) that was beyond the within-day coefficient of variation (3.8%). There was no evidence that the acute reductions in mental switching ability and $MCAv_{mean}$ were associated, but this is difficult to identify and characterise within a single-dose study.

Prior to the current study, the impact of acute reductions in CBF on cognitive performance had been addressed only in clinical cohorts. Marshall *et al.* (2001) concluded that a 23 - 54% reduction in perfusion from an acute internal carotid artery balloon test occlusion (30 min) resulted in transient (and reversible) reductions of sustained attention (response time) in patients with inoperable pericavernous aneurysms or head and neck tumours, despite markedly varied cognitive responses between patients. Findlay *et al.* (2019) showed in end-stage kidney disease patients that $MCAv_{mean}$ decreased significantly by 10% during at least 2 h of haemodialysis (a pro-inflammatory state), which moderately (Spearman's $\rho -0.32$) correlated with the intradialytic decline in executive function (trail making tasks; 13.5 s slower). Clinical populations with particular susceptibility to cerebral hypoperfusion during head-up tilt (i.e., postural tachycardia and chronic fatigue syndromes) have also shown simultaneous cognitive impairment on working memory and attention tasks (n-back) (Stewart *et al.*, 2012; Medow *et al.*, 2014). Interestingly, phenylephrine restored both the head-up tilt-induced cerebral hypoperfusion and the impaired cognitive performance in chronic fatigue syndrome patients (Medow *et al.*, 2014). Although we acutely reduced cerebral perfusion substantively (~31%) in *healthy* young and older adults, we did not find evidence to support a similar cognitive impairment related to the decreased flow. Healthy individuals with greater fitness, such as those recruited for this study, can buffer physiological strain - such as inflammation (Hamer & Steptoe, 2007) and oxidative stress (Radak *et al.*, 2005; Radak *et al.*, 2008) - to a higher degree. Higher fitness additionally provides higher cerebral perfusion chronically (Ainslie *et al.*, 2008). Collectively, such attributes of fitness may act as a buffer against functional and cognitive impairments caused by short-term reductions in perfusion. Although seemingly unrelated to concurrent reductions in cerebral perfusion, cognition was impaired after ingestion of indomethacin. In an acute sense, indomethacin is an *anti*-inflammatory drug that does not appear to alter cerebral metabolism (Pickard & MacKenzie, 1973; Sakabe & Siesjö, 1979; Dahlgren *et al.*, 1981; Wennmalm *et al.*, 1981; Jensen *et al.*, 1991). However, indomethacin is reported to promote oxidative stress in the small intestine and kidney of rodents by virtue of drug-induced generation of reactive oxygen species and decreased level of anti-oxidants and oxygen uptake (Basivireddy *et al.*, 2002; Varghese *et al.*, 2009; Tomita *et al.*, 2014). The brain is particularly vulnerable to oxidative

stress, but the effect of indomethacin on acute cerebral oxidative stress (and cognition) has not been addressed. Therefore, the mechanism by which cognition is acutely impaired with ingestion of indomethacin remains unclear. One possibility is that participants were distracted by overall feelings of (mild) discomfort, leading to worse performance. But if this occurred, a reduction in cognitive performance might be expected across all measures. Future research should consider implementing a fatigue and gastrointestinal upset control such as a visually draining or vertigo-inducing stimulus.

Indomethacin inhibits vasodilating prostaglandin synthesis, leading to cerebral vasoconstriction increasing resistance and reducing cerebral perfusion. As expected, and in agreement with previous literature (Eriksson *et al.*, 1983; Wennmalm *et al.*, 1983; Jensen *et al.*, 1993; Markus *et al.*, 1994; Kastrup *et al.*, 1999; Bruhn *et al.*, 2001; St. Lawrence *et al.*, 2002; Xie *et al.*, 2006; Ivancev *et al.*, 2009; Xie *et al.*, 2009; Barnes *et al.*, 2012a; Hoiland *et al.*, 2015; Peltonen *et al.*, 2015; Hoiland *et al.*, 2016; Peltonen *et al.*, 2016), we also observed decreased CVR_{HYPER} and CVR_{HYP0} across young and older groups. These reductions (68% and 50%, respectively) were beyond the day-to-day and within-day variability of CVR measures (Table 2: 11 – 24%) and may appear greater than what is typically observed following indomethacin ingestion (i.e., ~30-65%). One reason may be that these reactivities represent the indomethacin treatment-effect, a reduction greater than the effect of time (i.e., controlling for within-day variability). Our mean $\pm$ standard deviation values for indomethacin-related reductions (pre vs. post indomethacin) are similar to those typically reported for both CVR_{HYP0} (Young: $51 \pm 24\%$; Older: $59 \pm 21\%$) and CVR_{HYPER} (Young: $67 \pm 20\%$; Older: $74 \pm 13\%$). Although our data (Figure 3) appear to support a role of prostaglandin synthesis in CVR_{CO2}, it is important to consider that indomethacin is the only COX-inhibitor to reduce both basal cerebral blood flow and reactivity to CO₂ (Eriksson *et al.*, 1983; Wennmalm *et al.*, 1984; Markus *et al.*, 1994; Hoiland *et al.*, 2016), despite other potent COX-inhibitors (e.g., aspirin and naproxen) having similar inhibition of the cerebrovascular production of prostaglandins (Chemtob *et al.*, 1991). Therefore, it is possible that indomethacin reduces CBF via a mechanism(s) independent of prostaglandin synthesis inhibition. Indeed, indomethacin has numerous inhibitory and rapid-acting enzyme and cellular actions (Flower, 1974; Chemtob *et al.*, 1991) that are likely to cause systemic vasoconstriction independent of COX inhibition. One such action is via cyclic AMP-dependent protein kinase inhibition, as discussed by Hoiland *et al.* (2016) and Hoiland and Ainslie (2017). Briefly, cyclic AMP is involved in the regulation of vascular smooth muscle tone (Adelstein *et al.*, 1978) and indomethacin has been shown to inhibit cAMP-dependent protein

kinase (Kantor & Hampton, 1978; Goueli & Ahmed, 1980). However, our finding that individuals with heightened CVR_{CO_2} at rest experience greater indomethacin-related reductions (Figure 3) is in line with previous literature. Specifically, Kastrup *et al.* (1997) showed that the indomethacin-induced decrease in CVR_{HYPER} is linearly correlated with initial baseline CVR_{HYPER} ($r = 0.74$), and we have extended this finding to show the same relation with CVR_{HYPO} , as well as for older adults ($r = 0.87$). Given the dominant role of CVR_{CO_2} in cerebrovascular control, this has numerous implications for lifestyle and pharmacological interventions that impact cerebrovascular tone, and on assessment of cerebrovascular health (Burley *et al.*, 2016).

Although $CVR_{HYPER/HYPO}$ was *reduced* after indomethacin, it is likely by the same drug-induced vasoconstrictive-effect that CA appears to be *enhanced*. Dynamic CA offers mechanistic insight into the indomethacin-related reductions in cerebral perfusion, and how indomethacin may alter the pressure-flow relation in young and older adults. Decreases in perfusion and increases in vascular resistance intensify the signal power for blood pressure and weaken signal power for blood flow. These power changes result in significant decreases in gain and increases in phase after indomethacin, in agreement with existing literature in young men (Smirl *et al.*, 2014), new born and fetal lambs (Van Bel *et al.*, 1993; Van Bel *et al.*, 1995), and head-injured humans (Puppo *et al.*, 2007). A decrease in gain indicates that less flow is transmitted per unit (i.e. mm Hg) of pressure, i.e., blood pressure is having less influence on $MCAv_{mean}$. Changes in blood pressure normally trigger a rapid downstream vasoconstrictive response. However, in this scenario indomethacin has pre-emptively caused systemic vasoconstriction, facilitating the CA response. Similarly, an increase in phase indicates the flow response to a pressure-pulse is lengthened. This likely results from increased vascular resistance. Therefore, we are careful to interpret “enhanced” CA as anything but increased vascular tone – which is typically a sign of vascular dysfunction. Importantly, our data extend the findings of Smirl *et al.* (2014), such that indomethacin alters the cerebral pressure-flow relation not only in young men, but also in young women and healthy older adults, with no apparent sex-differences.

Baseline Group and Cerebrovascular Characteristics

The CBF in the middle cerebral artery declines by ~5% every ten years (Grolimund & Seiler, 1988), or 28 - 50% between ages 30 and 70 y (Heo *et al.*, 2010; Ogoh *et al.*, 2014). The prefrontal cortex – which is often associated with cognitive function - also has volumetric declines of ~5% per decade after age 20 y (Raz *et al.*, 2004). In addition to other structural changes (Bhogal *et al.*, 2016), the

age-related decrease in CBF may be partly attributed to a loss of prostaglandin function (Barnes *et al.*, 2012b). The current study shows a 14% difference in $MCAv_{mean}$ between young and older adults (Figure 2A), despite the older group having high levels of fitness and being sampled from an academic population. The reduction in anterior cerebral perfusion is likely due to a degree of vascular dysfunction, as evident from reduced CVC in older adults (Figure 2C). Lower CVC reflects less vasculature, higher vascular resistance (Tarumi & Zhang, 2018), or both. However, older adults did *not* show a significant difference in posterior cerebral blood velocity compared to young adults (Figure 2B). Although $PCAv_{mean}$ has been reported to decline ~3.7% every 10 years (Grolmund & Seiler, 1988), more recent literature reports that older (age-range: 40-73 y) adults have similar posterior perfusion to young (20-30 y) adults (Krejza *et al.*, 1999; Sorond *et al.*, 2005; Sorond *et al.*, 2008). The current study has limited power ($n=20$) and age range (25 vs 58 y) to address this issue but provides data for future meta-analytic study.

Older adults also did not have a measurable impairment in cerebrovascular control at rest, as determined by CVR_{HYPO} , CVR_{HYPER} , and dynamic CA. The finding that CVR_{CO_2} was not impacted by age agrees with Braz *et al.* (2017), who also found no difference between young and older groups of trained and sedentary adults. At this time, there is no clear evidence whether CVR_{CO_2} is impacted by healthy aging, or not (see review by Hoiland *et al.* (2019)). It is possible that any differences between age groups were missed due to CVR_{CO_2} being a variable measure, despite using a reliable measure of CO_2 control (i.e., computerised clamping) and following the current recommendations for lower intra-individual variability (Burley *et al.*, 2020). That being said, we are confident our findings are accurate as the reliability is similar to or better than what has been reported elsewhere, particularly in studies using TCD (Wilson *et al.*, 2010; McDonnell *et al.*, 2013). Cerebrovascular reactivity to CO_2 is a strong perturbation used to assess cerebrovascular control. The cerebrovascular response to CO_2 is highly integrated and affects many interrelated physiological systems that have direct and indirect impacts on cerebrovascular control. Thus, the consistently reported variability in this measure might be attributable to the numerous physiological systems involved. Furthermore, environmental influences (e.g., stress, prolonged sitting, fasting, circadian rhythms) affect these systems and may cause the appearance of changed cerebrovascular control by virtue of CVR_{CO_2} . Cognitive functions, such as executive functioning, also decline with age (Li *et al.*, 2001; Colcombe & Kramer, 2003; Kramer *et al.*, 2003; Brown *et al.*, 2010). The older adults in the current study were, on average, only 15% worse with response time tasks than their younger counterparts were (Figure 5). This may be because the older adults recruited for this study

were exceptionally healthy, aerobically fit, and cognitively active. For instance, 75% of the older group were current or recently-retired academic staff of the University. Thus, this older group may underestimate the cognitive and cerebrovascular change expected for an “older” population, even if healthy.

One consideration is that an acute reduction in CBF was used to partially inform chronic age-related effects. In an acute sense, cognition was not evidently impaired by virtue of decreased CBF *per se*. This does not mean that a *chronic* loss of, or reductions in, CBF or cerebrovascular control would not lead to cognitive impairment. In fact, meta-analyses reveal that early-stage cognitive decline is associated with abnormal cerebral haemodynamics, including reductions in perfusion and CO₂-reactivity (Beishon *et al.*, 2017). Wolters *et al.* (2017) have also demonstrated that cerebral hypoperfusion was a risk factor for cognitive impairment after a 6-y follow-up with 4759 adults. Additionally, a decrease in CBF over 3 y correlated to a decrease in global cognition (mini-mental state exam; $r = 0.59$) of 27 hypertensive and cognitively-sound older adults (Kitagawa *et al.*, 2009). Thus, there may be a case for chronic reductions in cerebral perfusion impacting cognition.

Limitations

Due to the nature of a correlation analysis, it is possible that the true effect of reduced perfusion on cognition was missed due to the consistent reduction in $MCAv_{mean}$ between participants, limiting the spread of data. Future research could administer graded doses of indomethacin across multiple days to measure any cognitive reactivity to reductions in perfusion within participants. It is also possible that chronic reductions in perfusion impair cognition (as occurs with aging). One avenue to investigate this notion may be via long-term use of indomethacin. However, Eriksson *et al.* (1983) found that cerebral perfusion normalises after one week of oral indomethacin intake (1.5 mg/kg). Moreover, this study was designed to measure any change in cognition that occurs with an *acute* reduction in perfusion, and not to simulate the changes that occur with aging.

As normal for studies using a cross-sectional design, the young group cannot be interpreted as younger versions of the older group. Although both groups were healthy and had above normal fitness, it is not certain that each younger adult will maintain this status across their lifespan. The results may therefore underestimate any age-related effects on cerebrovascular or cognitive function, due to the above-normal fitness and cognitive status in our older cohort. This is encouraging to an aging population, such that a healthy lifestyle may protect against functional (cognitive) effects of acutely reduced cerebral perfusion, which may occur during surgery,

dehydration, orthostasis, and heat stress. As such, indomethacin is only one model to acutely reduce CBF. Importantly, indomethacin reduces cerebral perfusion without manipulating the local tissue metabolism. Maintaining cerebral metabolism was paramount to addressing the question of whether acute reductions in blood flow *per se* negatively affect cognitive performance. Therefore, although using indomethacin to reduce CBF may not be generalisable to a “real-world” context of acute CBF reductions, it was the best option to limit physiological and psychological confounding factors.

The attending researchers were not blinded due to the obvious nature of CBF decline with indomethacin. For example, quality recordings and participant welfare were ensured by keeping a close watch on all physiological variables during the testing protocol; thus, the researcher would be immediately aware of a drop in $MCAv_{mean}$ caused by indomethacin. However, the research team took great care to ensure participants were blinded to each condition and to ensure the same monitoring of participants, and minimal verbal engagement, regardless of treatment, to reduce potential confounding conditions.

The current study also relied on the usual assumption that changes in CO_2 increase flow by dilation of downstream vessels without a meaningful change in MCA (or PCA) vessel diameter. This has been extensively disproven with MRI, wherein the MCA may dilate up to 7% (Coverdale *et al.*, 2014; Verbree *et al.*, 2014; Al-Khazraji *et al.*, 2018). However, any vessel dilation would result in an overall underestimation of changes in flow. Furthermore, by using a drug to purposefully reduce flow and increase vascular resistance, we may be underestimating the total reduction in flow.

Indeed, Kellawan *et al.* (2020) demonstrated that COX inhibition via indomethacin reduces MCA cross-sectional area by 0.2 mm. Lastly, although sex was included as a potential confounding factor within the statistical design, we did not power the study to test for sex differences.

Conclusion

Cognitive performance on a mental switching task was slightly (~7%) worse after an oral dose of indomethacin. However, we did not find evidence of an association between the reduction in performance and the 31% reduction in cerebral perfusion, even in older adults. Although older adults had lower $MCAv_{mean}$ and worse cognitive performance at baseline, both groups experienced a similar reduction in cerebral perfusion and cognition after indomethacin. Therefore, cognitive performance may not be influenced by a ~31% reduction in CBF *per se* in healthy young and older adults. These findings are encouraging to a healthy aging population, as both young and older individuals appear broadly resilient to acute reductions in cerebral perfusion up to ~31%.

Acknowledgements

We thank our participants for their time and effort; Travis Gibbons for his assistance with early data acquisition; Nigel Barrett and Gavin Kennedy for their technical support and expertise; and the School of Physical Education, Sport and Exercise Sciences and Department of Medicine for funding support. Additionally, we acknowledge funding support from the Lottery Health Research grant, Department of Internal Affairs to LCW, which assisted with the ethics application and protocol design.

Author Contribution

All authors contributed to the study conception and design and interpretation of results. Data collection and analysis were performed by LNS, LCW and JDC. The first draft of the manuscript was written by LNS and all authors commented on previous versions of the manuscript. RW provided clinical oversight related to the safety of indomethacin in the study design and medical clearance for older participants. All authors read and approved the final manuscript.

Disclosures

The authors declare that there is no conflict of interest.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Figure Legends

Figure 1. Timeline summary of experimental protocol, which took ~3.5 h (including instrumentation) and was undertaken twice. Time was allocated after cognitive tasks and before CA to allow for recovery of blood pressure and heart rate. *Abbreviations:* BL, baseline; CA, cerebrovascular autoregulation; CVR_{CO_2} , cerebral blood velocity hypercapnic (CVR_{HYPER}) and hypocapnic (CVR_{HYPO}) reactivity. Green/Red boxes represent a cognitive battery measuring visuomotor processing speed (Pro), inhibitory control (Anti) and mental switching (Pro/Anti) performance.

Figure 2. Cardio- and cerebro-vascular outcomes pre- and post- placebo and indomethacin for young (black bars, $n = 13$) and older (white bars; $n = 12$) adults were analysed using linear (Panels A, B, C, D) and nonparametric (Panels E, F, G, H) mixed-models and are therefore represented with mean or median (*) bars, respectively. *Abbreviations:* $MCAv_{mean}$, mean middle cerebral artery blood velocity; $PCAv_{mean}$, mean posterior cerebral artery blood velocity (Young: $n = 13$; Older: $n = 7$ CVC, cerebrovascular conductance; MAP, mean arterial pressure; HR, heart rate; $PETCO_2$, pressures of end-tidal carbon dioxide; CVR_{HYPO} (Young: $n = 11$; Older: $n = 6$), cerebrovascular reactivity to hypocapnia; CVR_{HYPER} (Young: $n = 13$; Older: $n = 9$), cerebrovascular reactivity to hypercapnia. ^a $p \leq 0.010$ vs young (main effect of age); ^b $p < 0.001$ vs pre-indomethacin, regardless of age (condition-by-time interaction); ^c $p < 0.001$ vs post-placebo, regardless of age (condition-by-time interaction).

Figure 3. Cerebrovascular reactivity (CVR_{CO_2}) to hypercapnia (grey; $n = 22$) and hypocapnia (black; $n = 17$) at rest (x-axis) is strongly associated with the indomethacin-induced reduction (y-axis) in both young (closed circles) and older (open circles) adults. i.e., those with the greatest CVR_{CO_2} at rest experienced the greatest reduction in CVR_{CO_2} after a dose of indomethacin.

Figure 4. Transfer function analysis of dynamic cerebral autoregulation during a sit-stand protocol (0.05 Hz) in young (black bars; $n = 13$) and older (white bars; $n = 7$) adults pre- and post-indomethacin. Both pre- and post-placebo (not shown) and indomethacin data were included in each mixed-model analysis. Bars represent median (*, Panels A and B) and mean (Panel C). ^b $p \leq 0.033$ vs pre-indomethacin, regardless of group; ^c $p \leq 0.038$ vs post-placebo, regardless of group.

Figure 5. Young (black bars; $n = 13$) and older (white bars; $n = 12$) adults' cognitive performance as determined by accuracy-adjusted response time (aRT) for Pro, Anti, and Pro/Anti batteries

(panels A-C) and working memory score (panel D) pre- and post-placebo and indomethacin using nonparametric mixed-models (bars represent median). ^a $p < 0.001$ vs young (main effect of age); ^b $p \leq 0.040$ vs pre-indomethacin, regardless of age; ^c $p \leq 0.042$ vs post-placebo, regardless of age.

Figure 6. Spearman's rho correlations between absolute (Panels A and B) and relative (Panels C and D) changes in accuracy-adjusted response time (aRT) and mean middle cerebral artery blood velocity (MCAv_{mean}). All change scores are calculated as the difference between the change from baseline with indomethacin and the change from baseline with placebo for young adults (panel A and C; closed circles, $n = 13$) and older adults (panel B and D; open circles, $n = 12$) during Pro (Green), Anti (Red), and Pro/Anti (Black) tasks.

1 **Tables**2 *Table 1. Participant Characteristics*

	Young n = 13 (6 female)	Older n = 12 (6 female)
Age (y)	25 ± 4	58 ± 6*
Mass (kg)	78 ± 18	73 ± 13
Height (cm)	175 ± 10	169 ± 10
Estimated $\dot{V}O_{2peak}$ (mL/min/kg)	52 ± 8	42 ± 11*
% of Predicted $\dot{V}O_{2peak}$	131 ± 22	130 ± 26
MoCA© Score (/30)	29 ± 1	28 ± 1
Resting Systolic BP (mm Hg)	107 ± 10	115 ± 8
Resting Diastolic BP (mm Hg)	68 ± 7	69 ± 6

3 *Note.* Baseline measures are reported as mean ± standard deviation from familiarisation and pre-
4 placebo conditions. Abbreviations: $\dot{V}O_{2peak}$, peak rate of oxygen consumption (mL/min/kg);
5 MoCA©, Montreal Cognitive Assessment – any score above 25 is considered “normal”. % of
6 Predicted $\dot{V}O_{2peak}$ was calculated as the percent difference between estimated $\dot{V}O_{2peak}$ (treadmill)
7 and the predicted $\dot{V}O_{2peak}$ using the equation from de Souza e Silva *et al.* (2018) *p ≤ 0.048 vs.
8 Young, using Student’s independent samples T-test.

Table 2. Reliability of dependent variables, shown as coefficients of variation (CV) and intraclass correlations (ICC).

		CV (%)	ICC	[95% CI]
A. Across 2 Days (Baseline)	MCAV _{mean}	5.4%	0.89	[0.75 - 0.95]
	PCAV _{mean}	9.4%	0.85	[0.62 - 0.94]
	CVR _{HYPER}	15.5%	0.48	[-0.21 - 0.78]
	CVR _{HYP0}	23.6%	0.49	[-0.15 - 0.78]

	Pro aRT	5.4%	0.79	[0.53 - 0.91]
	Anti aRT	6.1%	0.75	[0.44 - 0.89]
	Pro/Anti aRT	4.5%	0.84	[0.63 - 0.93]
B.	MCA _{v_{mean}}	4.3%	0.96	[0.91 - 0.98]
Within 1 Day	PCA _{v_{mean}}	5.9%	0.96	[0.90 - 0.98]
(Two measures, 2	CVR _{HYPER}	12.9%	0.88	[0.71 - 0.95]
h apart during the	CVR _{HYPO}	10.9%	0.91	[0.77 - 0.97]
placebo	Pro aRT	3.9%	0.92	[0.81 - 0.96]
condition)	Anti aRT	5.2%	0.86	[0.67 - 0.94]
	Pro/Anti aRT	3.8%	0.92	[0.82 - 0.96]

- 1 *Abbreviations:* CI, confidence interval; MCA_{v_{mean}}, mean middle cerebral artery blood velocity;
- 2 PCA_{v_{mean}}, mean posterior cerebral artery blood velocity; CVR_{HYPER}, cerebrovascular hypercapnic
- 3 reactivity; CVR_{HYPO}, cerebrovascular hypocapnic reactivity; aRT, accuracy-adjusted response time.
- 4 Reliabilities represent data from 25 participants for MCA_{v_{mean}}, PCA_{v_{mean}}, Pro, Anti, and Pro/Anti
- 5 (A and B), 23 participants for CVR_{HYPER} (A and B), 21 for CVR_{HYPO} (A), and 18 for CVR_{HYPO} (B).

References

- Adelstein RS, Conti MA, Hathaway DR & Klee CB. (1978). Phosphorylation of smooth muscle myosin light chain kinase by the catalytic subunit of adenosine 3': 5'-monophosphate-dependent protein kinase. *J Biol Chem* **253**, 8347-8350.
- Ainslie PN, Cotter JD, George KP, Lucas S, Murrell C, Shave R, Thomas KN, Williams MJ & Atkinson G. (2008). Elevation in cerebral blood flow velocity with aerobic fitness throughout healthy human ageing. *J Physiol* **586**, 4005-4010.
- Akalan C, Robergs RA & Kravitz L. (2008). Prediction of VO₂max from an individualized submaximal cycle ergometer protocol. *J Exerc Physiol Online* **11**, 1-17.
- Al-Khazraji BK, Shoemaker LN, Gati JS, Szekeres T & Shoemaker JK. (2018). Reactivity of larger intracranial arteries using 7 T MRI in young adults. *J Cereb Blood Flow Metab*, 271678x18762880.
- Bailey DM, Marley CJ, Brugniaux JV, Hodson D, New KJ, Ogoh S & Ainslie PN. (2013). Elevated aerobic fitness sustained throughout the adult lifespan is associated with improved cerebral hemodynamics. *Stroke* **44**, 3235-3238.
- Barnes JN, Schmidt JE, Nicholson WT & Joyner MJ. (2012a). Cyclooxygenase inhibition abolishes age-related differences in cerebral vasodilator responses to hypercapnia. *J Appl Physiol* **112**, 1884-1890.
- Barnes JN, Taylor JL, Kluck BN, Johnson CP & Joyner MJ. (2013). Cerebrovascular reactivity is associated with maximal aerobic capacity in healthy older adults. *J Appl Physiol* **114**, 1383-1387.

- 1 Basivireddy J, Vasudevan A, Jacob M & Balasubramanian KA. (2002). Indomethacin-induced
2 mitochondrial dysfunction and oxidative stress in villus enterocytes. *Biochem Pharmacol*
3 **64**, 339-349.
- 4
- 5 Beishon L, Haunton VJ, Panerai RB & Robinson TG. (2017). Cerebral Hemodynamics in Mild
6 Cognitive Impairment: A Systematic Review. *J Alzheimers Dis* **59**, 369-385.
- 7
- 8 Bertsch K, Hagemann D, Hermes M, Walter C, Khan R & Naumann E. (2009). Resting cerebral
9 blood flow, attention, and aging. *Brain Res Bull* **1267**, 77-88.
- 10
- 11 Bhogal AA, De Vis JB, Siero JC, Petersen ET, Luijten PR, Hendrikse J, Philippens ME &
12 Hoogduin H. (2016). The BOLD cerebrovascular reactivity response to progressive
13 hypercapnia in young and elderly. *Neuroimage* **139**, 94-102.
- 14
- 15 Braz I, Flück D, Lip G, Lundby C & Fisher J. (2017). Impact of aerobic fitness on cerebral blood
16 flow and cerebral vascular responsiveness to CO₂ in young and older men. *Scand J Med Sci*
17 *Sports* **27**, 634-642.
- 18
- 19 Brown AD, McMorris CA, Longman RS, Leigh R, Hill MD, Friedenreich CM & Poulin MJ.
20 (2010). Effects of cardiorespiratory fitness and cerebral blood flow on cognitive outcomes
21 in older women. *Neurobiol Aging* **31**, 2047-2057.
- 22
- 23 Bruhn H, Fransson P & Frahm J. (2001). Modulation of cerebral blood oxygenation by
24 indomethacin: MRI at rest and functional brain activation. *J Magn Reson Imaging* **13**, 325-
25 334.
- 26
- 27 Burley CV, Bailey DM, Marley CJ & Lucas SJ. (2016). Brain train to combat brain drain; focus on
28 exercise strategies that optimize neuroprotection. *Exp Physiol* **101**, 1178-1184.
- 29

- 1 Burley CV, Lucas RA, Whittaker AC, Mullinger K & Lucas SJ. (2020). The CO₂-stimulus duration
2 and steady-state time-point used for data extraction alters the cerebrovascular reactivity
3 outcome measure. *Exp Physiol*. **105**, 893-903
- 4
- 5 Chambers CD, Stokes MG & Mattingley JB. (2004). Modality-specific control of strategic spatial
6 attention in parietal cortex. *Neuron* **44**, 925-930.
- 7
- 8 Chemtob S, Beharry K, Barna T, Varma DR & Aranda JV. (1991). Differences in the effects in the
9 newborn piglet of various nonsteroidal antiinflammatory drugs on cerebral blood flow but
10 not on cerebrovascular prostaglandins. *Pediatr Res* **30**, 106-111.
- 11
- 12 Colcombe S & Kramer AF. (2003). Fitness effects on the cognitive function of older adults: a meta-
13 analytic study. *Psychol Sci* **14**, 125-130.
- 14
- 15 Coverdale NS, Badrov MB & Shoemaker JK. (2017). Impact of age on cerebrovascular dilation
16 versus reactivity to hypercapnia. *J Cereb Blood Flow Metab* **37**, 344-355.
- 17
- 18 Coverdale NS, Gati JS, Opalevych O, Perrotta A & Shoemaker JK. (2014). Cerebral blood flow
19 velocity underestimates cerebral blood flow during modest hypercapnia and hypocapnia. *J*
20 *Appl Physiol* **117**, 1090-1096.
- 21
- 22 Dahlgren N, Nilsson B, Sakabe T & Siesjö Bk. (1981). The effect of indomethacin on cerebral
23 blood flow and oxygen consumption in the rat at normal and increased carbon dioxide
24 tensions. *Acta Physiol Scand* **111**, 475-485.
- 25
- 26 de Souza e Silva CG, Kaminsky LA, Arena R, Christle JW, Araújo CGS, Lima RM, Ashley EA &
27 Myers J. (2018). A reference equation for maximal aerobic power for treadmill and cycle
28 ergometer exercise testing: Analysis from the FRIEND registry. *Eur J Prev Cardiol* **25**, 742-
29 750.

- 1
- 2 Eriksson S, Hagenfeldt L, Law D, Patrono C, Pinca E & Wennmalm A. (1983). Effect of
- 3 prostaglandin synthesis inhibitors on basal and carbon dioxide stimulated cerebral blood
- 4 flow in man. *Acta Physiol Scand* **117**, 203-211.
- 5
- 6 Findlay MD, Dawson J, Dickie DA, Forbes KP, McGlynn D, Quinn T & Mark PB. (2019).
- 7 Investigating the relationship between cerebral blood flow and cognitive function in
- 8 hemodialysis patients. *J Am Soc Nephrol* **30**, 147-158.
- 9
- 10 Flower RJ. (1974). Drugs which inhibit prostaglandin biosynthesis. *Pharmacol Rev* **26**, 33-67.
- 11
- 12 Fox EL. (1973). A simple, accurate technique for predicting maximal aerobic power. *J Appl Physiol*
- 13 **35**, 914-916.
- 14
- 15 Giblin LM. (2011). The Effect of Unexpected Exercise Duration on Rating of Perceived Exertion in
- 16 an Untrained, Sedentary Population.
- 17
- 18 Golding LA, Myers CR, SINNING K & Sinning S. (1989). Y's way to physical fitness: the
- 19 complete guide to fitness testing and instruction.
- 20
- 21 Goueli SA & Ahmed K. (1980). Indomethacin and inhibition of protein kinase reactions. *Nature*
- 22 **287**, 171-172.
- 23
- 24 Green RS, Leffler CW, Busija DW, Fletcher AM & Beasley DG. (1987). Indomethacin does not
- 25 alter the circulating catecholamine response to asphyxia in the neonatal piglet. *Pediatr Res*
- 26 **21**, 534.
- 27
- 28 Grolimund P & Seiler R. (1988). Age dependence of the flow velocity in the basal cerebral
- 29 arteries—a transcranial Doppler ultrasound study. *Ultrasound Med Biol* **14**, 191-198.

- 1
- 2 Guiney H, Lucas SJ, Cotter JD & Machado L. (2015). Evidence cerebral blood-flow regulation
- 3 mediates exercise-cognition links in healthy young adults. *Neuropsychology* **29**, 1-9.
- 4
- 5 Guiney H, Lucas SJE, Cotter JD & Machado L. (2019). Investigating links between habitual
- 6 physical activity, cerebrovascular function, and cognitive control in healthy older adults.
- 7 *Neuropsychologia*.
- 8
- 9 Hamer M & Steptoe A. (2007). Association between physical fitness, parasympathetic control, and
- 10 proinflammatory responses to mental stress. *Psychosom Med* **69**, 660-666.
- 11
- 12 Heo S, Prakash RS, Voss MW, Erickson KI, Ouyang C, Sutton BP & Kramer AF. (2010). Resting
- 13 hippocampal blood flow, spatial memory and aging. *Brain Res* **1315**, 119-127.
- 14
- 15 Hohimer AR, Richardson B, Bissonnette J & Machida C. (1985). The effect of indomethacin on
- 16 breathing movements and cerebral blood flow and metabolism in the fetal sheep. *J Dev*
- 17 *Physiol* **7**, 217-228.
- 18
- 19 Hoiland RL & Ainslie PN. (2017). Reply from Ryan L. Hoiland and Philip N. Ainslie. *J Physiol*
- 20 **595**, 3673-3675.
- 21
- 22 Hoiland RL, Ainslie PN, Wildfong KW, Smith KJ, Bain AR, Willie CK, Foster G, Monteleone B &
- 23 Day TA. (2015). Indomethacin-induced impairment of regional cerebrovascular reactivity:
- 24 implications for respiratory control. *J Physiol* **593**, 1291-1306.
- 25
- 26 Hoiland RL, Fisher JA & Ainslie PN. (2019). Regulation of the Cerebral Circulation by Arterial
- 27 Carbon Dioxide. *Compr Physiol* **9**, 1101-1154.
- 28

- 1 Hoiland RL, Tymko MM, Bain AR, Wildfong KW, Monteleone B & Ainslie PN. (2016). Carbon
2 dioxide-mediated vasomotion of extra-cranial cerebral arteries in humans: a role for
3 prostaglandins? *J Physiol* **594**, 3463-3481.
- 4
- 5 Ide K, Eliasziw M & Poulin MJ. (2003). Relationship between middle cerebral artery blood velocity
6 and end-tidal PCO₂ in the hypocapnic-hypercapnic range in humans. *J Appl Physiol* **95**,
7 129-137.
- 8
- 9 Ivancev V, Bakovic D, Obad A, Breskovic T, Palada I, Joyner MJ & Dujic Z. (2009). Effects of
10 indomethacin on cerebrovascular response to hypercapnea and hypocapnea in breath-hold
11 diving and obstructive sleep apnea. *Respir Physiol Neurobiol* **166**, 152-158.
- 12
- 13 Jensen K, Freundlich M, Bünemann L, Therkelsen K, Hansen H & Cold G. (1993). The effect of
14 indomethacin upon cerebral blood flow in healthy volunteers. *Acta Neurochir (Wien)* **124**,
15 114-119.
- 16
- 17 Jensen K, Öhrström J, Cold G & Astrup J. (1991). The effects of indomethacin on intracranial
18 pressure, cerebral blood flow and cerebral metabolism in patients with severe head injury
19 and intracranial hypertension. *Acta Neurochir (Wien)* **108**, 116-121.
- 20
- 21 Kantor HS & Hampton M. (1978). Indomethacin in submicromolar concentrations inhibits cyclic
22 AMP-dependent protein kinase. *Nature* **276**, 841-842.
- 23
- 24 Karvonen MJ. (1957). The effects of training on heart rate: A longitudinal study. *Ann Med Exp Biol*
25 *Fenn* **35**, 307-315.
- 26
- 27 Kastrup A, Happe V, Hartmann C & Schabet M. (1999). Gender-related effects of indomethacin on
28 cerebrovascular CO₂ reactivity. *J Neurol Sci* **162**, 127-132.
- 29

- 1 Kastrup A, Thomas C, Hartmann C & Schabet M. (1997). Sex dependency of cerebrovascular CO₂
2 reactivity in normal subjects. *Stroke* **28**, 2353-2356.
- 3
- 4 Kellawan JM, Peltonen GL, Harrell JW, Roldan-Alzate A, Wieben O & Schrage WG. (2020).
5 Differential contribution of cyclooxygenase to basal cerebral blood flow and hypoxic
6 cerebral vasodilation. *Am J Physiol Regul Integr Comp Physiol* **318**, R468-R479.
- 7
- 8 Kitagawa K, Oku N, Kimura Y, Yagita Y, Sakaguchi M, Hatazawa J & Sakoda S. (2009).
9 Relationship between cerebral blood flow and later cognitive decline in hypertensive
10 patients with cerebral small vessel disease. *Hypertens Res* **32**, 816-820.
- 11
- 12 Koo TK & Li MY. (2016). A guideline of selecting and reporting intraclass correlation coefficients
13 for reliability research. *J Chiropr Med* **15**, 155-163.
- 14
- 15 Kraaier V, Van Huffelen A, Wieneke G, Van der Worp H & Bär P. (1992). Quantitative EEG
16 changes due to cerebral vasoconstriction. Indomethacin versus hyperventilation-induced
17 reduction in cerebral blood flow in normal subjects. *Electroencephalogr Clin Neurophysiol*
18 **82**, 208-212.
- 19
- 20 Kramer AF, Colcombe SJ, McAuley E, Eriksen KI, Scalf P, Jerome GJ, Marquez DX, Elavsky S &
21 Webb AG. (2003). Enhancing brain and cognitive function of older adults through fitness
22 training. *J Mol Neurosci* **20**, 213-221.
- 23
- 24 Krejza J, Mariak Z, Walecki J, Szydlak P, Lewko J & Ustymowicz A. (1999). Transcranial color
25 Doppler sonography of basal cerebral arteries in 182 healthy subjects: age and sex
26 variability and normal reference values for blood flow parameters. *Am JRoentgenol* **172**,
27 213-218.
- 28
- 29 Li S-C, Lindenberger U & Sikström S. (2001). Aging cognition: from neuromodulation to
30 representation. *Trends Cogn Sci* **5**, 479-486.

- 1
- 2 Lucas SJ, Ainslie PN, Murrell CJ, Thomas KN, Franz EA & Cotter JD. (2012). Effect of age on
- 3 exercise-induced alterations in cognitive executive function: relationship to cerebral
- 4 perfusion. *Exp Gerontol* **47**, 541-551.
- 5
- 6 Markus HS, Vallance P & Brown MM. (1994). Differential effect of three cyclooxygenase
- 7 inhibitors on human cerebral blood flow velocity and carbon dioxide reactivity. *Stroke* **25**,
- 8 1760-1764.
- 9
- 10 Marshall RS, Lazar RM, Pile-Spellman J, Young WL, Duong DH, Joshi S & Ostapovich N.
- 11 (2001). Recovery of brain function during induced cerebral hypoperfusion. *Brain* **124**,
- 12 1208-1217.
- 13
- 14 McDonnell MN, Berry NM, Cutting MA, Keage HA, Buckley JD & Howe PR. (2013). Transcranial
- 15 Doppler ultrasound to assess cerebrovascular reactivity: reliability, reproducibility and
- 16 effect of posture. *PeerJ* **1**, e65.
- 17
- 18 Medow MS, Sood S, Messer Z, Dzogbeta S, Terilli C & Stewart JM. (2014). Phenylephrine
- 19 alteration of cerebral blood flow during orthostasis: effect on n-back performance in
- 20 chronic fatigue syndrome. *J Appl Physiol* **117**, 1157-1164.
- 21
- 22 Meel-van den Abeelen AS, Simpson DM, Wang LJ, Slump CH, Zhang R, Tarumi T, Rickards CA,
- 23 Payne S, Mitsis GD & Kostoglou K. (2014). Between-centre variability in transfer function
- 24 analysis, a widely used method for linear quantification of the dynamic pressure-flow
- 25 relation: The CARNet study. *Med Engineer Physics* **36**, 620-627.
- 26
- 27 Myerson J, Emery L, White DA & Hale S. (2003). Effects of age, domain, and processing demands
- 28 on memory span: Evidence for differential decline. *Aging Neuropsychol Cog* **10**, 20-27.
- 29

- 1 Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL &
2 Chertkow H. (2005). The Montreal Cognitive Assessment, MoCA: a brief screening tool
3 for mild cognitive impairment. *J Am Geriatr Soc* **53**, 695-699.
- 4
- 5 Noguchi K, Gel YR, Brunner E & Konietzschke F. (2012). nparLD: an R software package for the
6 nonparametric analysis of longitudinal data in factorial experiments. *JStat Softw* **50**.
- 7
- 8 Ogoh S, Tsukamoto H, Hirasawa A, Hasegawa H, Hirose N & Hashimoto T. (2014). The effect of
9 changes in cerebral blood flow on cognitive function during exercise. *Physiol Rep* **2**.
- 10
- 11 Peltonen GL, Harrell JW, Aleckson BP, LaPlante KM, Crain MK & Schrage WG. (2016). Cerebral
12 blood flow regulation in women across menstrual phase: differential contribution of
13 cyclooxygenase to basal, hypoxic, and hypercapnic vascular tone. *Am J Physiol Regul*
14 *Integr Comp Physiol* **311**, R222-231.
- 15
- 16 Peltonen GL, Harrell JW, Rousseau CL, Ernst BS, Marino ML, Crain MK & Schrage WG. (2015).
17 Cerebrovascular regulation in men and women: stimulus-specific role of cyclooxygenase.
18 *Physiol Rep* **3**.
- 19
- 20 Phillips AA, Chan FH, Zheng MMZ, Krassioukov AV & Ainslie PN. (2016). Neurovascular
21 coupling in humans: Physiology, methodological advances and clinical implications. *J*
22 *Cereb Blood Flow Metab* **36**, 647-664.
- 23
- 24 Pickard J & MacKenzie E. (1973). Inhibition of prostaglandin synthesis and the response of baboon
25 cerebral circulation to carbon dioxide. *Nature New Biol* **245**, 187.
- 26
- 27 Puppo C, Lopez L, Farina G, Caragna E, Moraes L, Iturralde A & Biestro A. (2007). Indomethacin
28 and cerebral autoregulation in severe head injured patients: a transcranial Doppler study.
29 *Acta Neurochir (Wien)* **149**, 139-149.

- 1
- 2 Radak Z, Chung HY & Goto S. (2005). Exercise and hormesis: oxidative stress-related adaptation
- 3 for successful aging. *Biogerontology* **6**, 71-75.
- 4
- 5 Radak Z, Chung HY, Koltai E, Taylor AW & Goto S. (2008). Exercise, oxidative stress and
- 6 hormesis. *Ageing Res Rev* **7**, 34-42.
- 7
- 8 Raz N, Gunning-Dixon F, Head D, Rodrigue KM, Williamson A & Acker JD. (2004). Aging,
- 9 sexual dimorphism, and hemispheric asymmetry of the cerebral cortex: replicability of
- 10 regional differences in volume. *Neurobiol Aging* **25**, 377-396.
- 11
- 12 Sakabe T & Siesjö BK. (1979). The effect of indomethacin on the blood flow—metabolism couple
- 13 in the brain under normal, hypercapnic and hypoxic conditions. *Acta Physiol Scand* **107**,
- 14 283-284.
- 15
- 16 Salthouse TA. (2009). When does age-related cognitive decline begin? *Neurobiol Aging* **30**, 507-
- 17 514.
- 18
- 19 Shoemaker LN, Wilson LC, Lucas SJ, Machado L & Cotter JD. (2019a). Cerebrovascular
- 20 regulation is not blunted during mental stress. *Exp Physiol* **104**, 1678-1687.
- 21
- 22 Shoemaker LN, Wilson LC, Lucas SJ, Machado L & Cotter JD. (2020). Acute exercise-related
- 23 cognitive effects are not attributable to changes in end-tidal CO₂ or cerebral blood
- 24 velocity. *Eur J Appl Physiol*, 1-13.
- 25
- 26 Shoemaker LN, Wilson LC, Lucas SJ, Machado L, Thomas KN & Cotter JD. (2019b). Swimming-
- 27 related effects on cerebrovascular and cognitive function. *Physiol Rep* **7**.
- 28

- 1 Smirl JD, Tzeng YC, Monteleone BJ & Ainslie PN. (2014). Influence of cerebrovascular resistance
2 on the dynamic relationship between blood pressure and cerebral blood flow in humans.
3 *JAppl Physiol* **116**, 1614-1622.
- 4
- 5 Sorond FA, Khavari R, Serrador JM & Lipsitz LA. (2005). Regional cerebral autoregulation during
6 orthostatic stress: age-related differences. *J Gerontol* **60**, 1484-1487.
- 7
- 8 Sorond FA, Schnyer DM, Serrador JM, Milberg WP & Lipsitz LA. (2008). Cerebral blood flow
9 regulation during cognitive tasks: effects of healthy aging. *Cortex* **44**, 179-184.
- 10
- 11 St. Lawrence KS, Ye FQ, Lewis BK, Weinberger DR, Frank JA & McLaughlin AC. (2002). Effects
12 of indomethacin on cerebral blood flow at rest and during hypercapnia: an arterial spin
13 tagging study in humans. *J MRI* **15**, 628-635.
- 14
- 15 Staessen J, Cattaert A, Fagard R, Lijnen P, Moerman E, De Schaepdryver A & Amery A. (1984).
16 Hemodynamic and humoral effects of prostaglandin inhibition in exercising humans. *J Appl*
17 *Physiol* **56**, 39-45.
- 18
- 19 Stewart JM, Medow MS, Messer ZR, Baugham IL, Terilli C & Ocon AJ. (2012). Postural
20 neurocognitive and neuronal activated cerebral blood flow deficits in young chronic fatigue
21 syndrome patients with postural tachycardia syndrome. *Am J Physiol-Heart C* **302**, H1185-
22 H1194.
- 23
- 24 Tarumi T & Zhang R. (2018). Cerebral blood flow in normal aging adults: cardiovascular
25 determinants, clinical implications, and aerobic fitness. *J Neurochem* **144**, 595-608.
- 26
- 27 Tomita T, Sadakata H, Tamura M & Matsui H. (2014). Indomethacin-induced generation of
28 reactive oxygen species leads to epithelial cell injury before the formation of intestinal
29 lesions in mice. *J Physiol Pharmacol* **65**, 435-440.

- 1
- 2 Townsend JT & Ashby FG. (1983). *Stochastic modeling of elementary psychological processes*.
- 3 CUP Archive.
- 4
- 5 Tzeng Y-C, Ainslie PN, Cooke WH, Peebles KC, Willie CK, MacRae BA, Smirl JD, Horsman HM
- 6 & Rickards CA. (2012). Assessment of cerebral autoregulation: the quandary of
- 7 quantification. *Am J Physiol-Heart* **303**, H658-H671.
- 8
- 9 Van Bel F, Bartelds B, Teitel DF & Rudolph AM. (1995). Effect of indomethacin on cerebral blood
- 10 flow and oxygenation in the normal and ventilated fetal lamb. *Pediatr Res* **38**, 243.
- 11
- 12 Van Bel F, Klautz RJ, Steendijk P, Schipper IB, Teitel DF & Baan J. (1993). The influence of
- 13 indomethacin on the autoregulatory ability of the cerebral vascular bed in the newborn
- 14 lamb. *Pediatr Res* **34**, 178.
- 15
- 16 Varghese J, Faith M & Jacob M. (2009). Zinc prevents indomethacin-induced renal damage in rats
- 17 by ameliorating oxidative stress and mitochondrial dysfunction. *Eur J Pharmacol* **614**, 114-
- 18 121.
- 19
- 20 Verbree J, Bronzwaer AS, Ghariq E, Versluis MJ, Daemen MJ, van Buchem MA, Dahan A, van
- 21 Lieshout JJ & van Osch MJ. (2014). Assessment of middle cerebral artery diameter during
- 22 hypocapnia and hypercapnia in humans using ultra-high-field MRI. *J Appl Physiol* **117**,
- 23 1084-1089.
- 24
- 25 Wennmalm A, Carlsson I, Edlund A, Eriksson S, Kaijser L & Nowak J. (1984). Central and
- 26 peripheral haemodynamic effects of non-steroidal anti-inflammatory drugs in man. *Arch*
- 27 *Toxicol Suppl* **7**, 350-359.
- 28

- 1 Wennmalm A, Eriksson S, Hagenfeldt L, Law D, Patrono C & Pinca E. (1983). Effect of
2 prostaglandin synthesis inhibitors on basal and carbon dioxide-stimulated cerebral blood
3 flow in man. *Adv Prostaglandin Thromboxane Leukot Res* **12**, 351-355.
- 4
- 5 Wennmalm Å, Eriksson S & Wahren J. (1981). Effect of indomethacin on basal and carbon dioxide
6 stimulated cerebral blood flow in man. *Clin Physiol* **1**, 227-234.
- 7
- 8 Williams TB, Corbett J, McMorris T, Young JS, Dicks M, Ando S, Thelwell RC, Tipton MJ &
9 Costello JT. (2019). Cognitive performance is associated with cerebral oxygenation and
10 peripheral oxygen saturation, but not plasma catecholamines, during graded normobaric
11 hypoxia. *Exp Physiol* **104**, 1384-1397.
- 12
- 13 Wilson LC, Cotter JD, Fan J-L, Lucas RA, Thomas KN & Ainslie PN. (2010). Cerebrovascular
14 reactivity and dynamic autoregulation in tetraplegia. *Am J Physiol Regul Integr Comp*
15 *Physiol* **298**, R1035-R1042.
- 16
- 17 Wolters FJ, Zonneveld HI, Hofman A, van der Lugt A, Koudstaal PJ, Vernooij MW & Ikram MA.
18 (2017). Cerebral perfusion and the risk of dementia: a population-based study. *Circulation*
19 117.027448.
- 20
- 21 Xie A, Skatrud JB, Barczi SR, Reichmuth K, Morgan BJ, Mont S & Dempsey JA. (2009). Influence
22 of cerebral blood flow on breathing stability. *J Appl Physiol* **106**, 850-856.
- 23
- 24 Xie A, Skatrud JB, Morgan B, Chenuel B, Khayat R, Reichmuth K, Lin J & Dempsey JA. (2006).
25 Influence of cerebrovascular function on the hypercapnic ventilatory response in healthy
26 humans. *J Physiol* **577**, 319-329.

A **B** **C**

YOUNG

A

OLDER

B

C

D