Depositional sedimentary environment and hydrothermal controls on organic matter enrichment in the Lower Cambrian Niutitang shale reservoir, South China

Tan, Jingqiang; Wang, Zhangwu; Wang, Wenhui; Hilton, Jason; Guo, Jianhua; Wang, Xikai

DOI:

[10.1306/12222018196](https://doi.org/10.1306/12222018196)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Tan, J, Wang, Z, Wang, W, Hilton, J, Guo, J & Wang, X 2020, 'Depositional sedimentary environment and hydrothermal controls on organic matter enrichment in the Lower Cambrian Niutitang shale reservoir, South China', *AAPG Bulletin*. <https://doi.org/10.1306/12222018196>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 *Depositional environment and hydrothermal controls on organic matter enrichment in*
2 *the lower Cambrian Niutitang shale, South China*

3

4 Jingqiang Tan^{1,2}, Zhanghu Wang^{1,2}, Wenhui Wang^{1,2}, Jason Hilton³, Jianhua Guo^{1,2},
5 Xikai Wang^{1,2}

6

7 1. Key Laboratory of Metallogenic Prediction of Nonferrous Metals and Geological
8 Environment Monitoring, Ministry of Education, School of Geosciences and
9 Info-physics, Central South University, Changsha, 410083, China

10 2. Hunan Key Laboratory of Nonferrous Resources and Geological Hazards
11 Exploration, Changsha 410083, China

12 3. School of Geography, Earth and Environmental Science, University of
13 Birmingham, Birmingham, B15 2TT, UK

14

15 **ACKNOWLEDGMENTS**

16 This work was supported by the National Natural Science Foundation of China
17 (Grants 41872151, 41772001, 41402010), the Fundamental Research Funds for the
18 Central Universities of Central South University (Grant 502221901), and the
19 Innovation-driven Project of Central South University (Grant 502501005). We thank
20 Barry Katz, Robert Merrill, Thomas Algeo and two anonymous reviewers for helpful
21 and constructive comments on earlier versions of the manuscript.

22

23 **ABSTRACT**

24 The purpose of this research was to examine paleoenvironments, hydrothermal
25 activity, and seawater restriction of the lower Cambrian Niutitang Formation shale gas
26 reservoir in the eastern Xuefeng Uplift and to determine factors affecting organic
27 matter (OM) enrichment. In the studied borehole (X1) in western Hunan Province, the
28 Niutitang Formation can be subdivided into the Niu1, Niu2, and Niu3 members based
29 on geological and geochemical features, including trace element enrichment, lithology
30 and fossil content. Total organic carbon (TOC) values of the Niutitang Formation are
31 variable, averaging 1.5 wt.% in the Niu1 Member, 12.7 wt.% in the Niu2 Member,
32 and 5.1 wt.% in the Niu3 Member. Paleoclimatic changes were responsible for
33 changes in biota, which impacted patterns of OM enrichment. Climate proxies
34 (Chemical Index of Alteration (CIA)), and productivity proxies (B_{bio} , Cu/Al , and
35 Ni/Al) consistently indicate higher paleoproductivity in the Niu2 Member. The Niu1
36 and Niu2 members were possibly affected by hydrothermal events, whereas
37 hydrothermal activity was absent during deposition of the Niu3 Member.
38 Hydrothermal activity may not only provide nutrients and silica but also enhance the
39 reducing condition of the water column. In addition, hydrothermal events may have
40 possibly influenced biological survival in the different environments, which in turn
41 increased their reproduction within the early Cambrian ocean and affected OM
42 production. Redox proxies (Mo_{EF} and U_{EF}) imply that the Niu1, Niu2, and Niu3
43 members were deposited in suboxic, euxinic, and ferruginous environments,
44 respectively. Redox conditions, strongly restricted water environments, and

45 hydrothermal events were conducive to OM enrichment during the early Cambrian.

INTRODUCTION

During the early Cambrian, a marine transgression event resulted in widespread deposition of organic-rich shale across South China. Although the lower Cambrian shales were subsequently eroded in some parts of the Upper Yangtze Region, their present distribution extends across an area of more than $9.0 \times 10^5 \text{ km}^2$ ($\sim 3.5 \times 10^5 \text{ mi}^2$) (Zhang et al., 2017). These organic-rich shale intervals are 50–200 m (164–656 ft) thick in many places, with total organic carbon (TOC) contents commonly above 1.0 wt.% and can reach 30 wt.% (Tan et al., 2013; Yu et al., 2017). Over the past decade, organic-rich shale reservoirs in the Upper Yangtze Region have been widely studied for their promising petroleum potential (Hao et al., 2013; Tan et al., 2014a; Bu et al., 2015; Lyu et al., 2018a; Xiao et al., 2018; L. Li et al., 2019). Of these studies, considerable achievements have been made in our understanding of the Niutitang Formation (and its equivalent horizons, including the Qiongzhusi and Jiumenchong formations) shale gas resources. Wells targeting lower Cambrian shale intervals in Sichuan, Guizhou, and Hubei provinces have already obtained commercial scale shale gas flow. In 2017, the China Geological Survey (CGS) drilled the EYiYe-1 well in Yichang, west Hubei Province, and achieved a production rate of 2.1 MMcf per day (MMcf/d) and an open gas flow rate of 4.4 MMcf/d.

Previous studies on the Niutitang Formation have primarily focused on shale reservoir characteristics and gas potential (Tan et al., 2014b; Tian et al., 2015; T. Lin et al., 2016; J. Yan et al., 2016; Lyu et al., 2018b; Sun et al., 2018). However, the effects of hydrothermal events and depositional environment on organic matter (OM)

68 enrichment of the Niutitang shale have, by comparison, been overlooked (Liu et al.,
69 2015; Yeasmin et al., 2017; Gao et al., 2018). OM enrichment mechanisms in marine
70 shales are complex and influenced by various factors, including terrigenous detrital
71 supply, paleoclimate, primary productivity, hydrothermal activity, redox conditions,
72 and seawater restriction (Piper et al., 2004; Liu et al., 2015; Wu et al., 2016).
73 Variations in paleoclimate have a marked impact on the abundance of organisms in
74 marine settings and thus influence the supply of OM, while the preservation of OM is
75 controlled by redox conditions, terrigenous dilution, and seawater restriction. The OM
76 supply can be affected by hydrothermal activity, in addition to primary productivity
77 and influences from preservation conditions (Knoll and Carroll, 1999; Reysenbach
78 and Cady, 2001; Little and Vrijenhoek, 2003; Simoneit, 2004; R. Yang et al., 2008;
79 Lane et al., 2010; Fuchida et al., 2014; Y. Li et al., 2015; Han et al., 2017).

80 In South China, hydrothermal activity mainly occurred during the deposition of
81 the lower part of the Niutitang shale (R. Yang et al., 2008; D. Chen et al., 2009; Y. Li
82 et al., 2015; Liu et al., 2015; Han et al., 2017), whereas the upper part of the Niutitang
83 shale was not obviously affected. Submarine hydrothermal activity with abundant
84 metals and volatile gases (CH_4 and H_2S) may change the redox and chemical
85 conditions of seawater and affect the reproduction of biological communities (Jones
86 and Gislason, 2008; Chambers et al., 2013). The relationship between hydrothermal
87 activity and biological productivity is, nevertheless, not fully understood.
88 Investigations are also relatively rare on the links between seawater restriction and
89 organic-rich shales. The depositional environments during the different time intervals

90 of the early Cambrian could result in markedly different OM enrichment mechanisms.
91 Previously studied profiles were principally deposited in shallow-deep shelf
92 environments, i.e., Guizhou Province (Y. Li et al., 2015). To examine the correlation
93 between the depositional environments and OM enrichment, freshly cored shale
94 samples that were deposited in deep marine settings were investigated in detail in the
95 current research.

96 The primary objective of this study was to reconstruct sedimentary environment,
97 hydrothermal activity, and their influences on OM enrichment during shale deposition
98 from the eastern Xuefeng Uplift in the lower Cambrian Niutitang shale. To illustrate
99 the primary factors influencing OM enrichment, a series of analyses including
100 lithofacies, mineralogy, trace and major elements, and TOC were undertaken on shale
101 samples from the recently drilled X1 well in Anhua County, Hunan Province, South
102 China (Figure 1).

103

104 **GEOLOGICAL SETTING**

105 During the late Ediacaran to early Cambrian, the South China Craton consisted of the
106 Yangtze and Cathaysia blocks. From the northwest to the southeast, sedimentary
107 facies comprised marine platform, shallow shelf, deep shelf, slope, and marine basin
108 settings (Goldberg et al., 2007; S. Wang et al., 2015) (Figure 1A). The lower
109 Cambrian sedimentary rocks in the shallow-water areas were primarily composed of
110 dolomite and limestone (e.g., Zhujiaping and Madiping formations), whereas those
111 sedimentary rocks in deep-water areas principally consisted of siliceous- and

112 organic-rich shale (e.g., Liuchapo Formation and the lower part of the Niutitang
113 Formation) (Figures 1 and 2). The lower Cambrian interval includes the Cambrian
114 evolutionary radiation of small shelly fossil assemblages (including brachiopods,
115 gastropods and trilobites) and marine plankton (including algae and other microfossils)
116 (R. Yang et al., 2008), providing significant OM inputs into marine depositional
117 settings.

118 During the late Cambrian Stage 2 (Figure 2), as a result of the breakup of the
119 Rodinia Supercontinent, the Yangtze and Cathaysia blocks commenced rifting. In
120 South China, a major transgression occurred, as well as extensive hydrothermal and
121 volcanic activity (Steiner et al., 2001; D. Chen et al., 2009; Han et al., 2017). Volcanic
122 tuffs and metal ore layers (Ni-Mo, V, and Ba) were widely distributed in the
123 Yu'an-shan and Niutitang formations. Black carbonaceous shales were widely
124 developed in the research area as a result of anoxic events in the sedimentary basin, in
125 addition to phosphatic nodules, pyrite, and siliceous shales (Steiner et al., 2007; Wu et
126 al., 2017). Due to paleogeographic differences, the thicknesses and TOC contents of
127 the Niutitang shale vary considerably among its different locations (Figure 2).

128 The Xuefeng Uplift represents the transitional zone between the Yangtze and
129 Cathaysia tectonic blocks. Sedimentary strata from the Cambrian to the Silurian were
130 extensively developed in the eastern Xuefeng area. The Niutitang shales in the eastern
131 Xuefeng Uplift are characterized by their great thicknesses, high TOC contents, and
132 advanced thermal maturities (T. Lin et al., 2016; J. Chen et al., 2017). Several lower
133 Cambrian sections have been previously studied in detail for volcanic and

134 biostratigraphic analyses, including the Xiaotan section in Yunnan Province (A. Yang
135 et al., 2003; Steiner et al., 2007; Compston et al., 2008; Och et al., 2013; Jin et al.,
136 2016; C. Yang et al., 2018), the Songtao section in Guizhou Province (A. Yang et al.,
137 2003; Goldberg et al., 2007; D. Chen et al., 2015), the Longbizui section in Hunan
138 Province (J. Wang et al., 2012; C. Yang et al., 2017), and the Ganziping section in
139 Hunan Province (D. Chen et al., 2009), distributed across shallow shelf facies in the
140 west to slope facies in the east (Figure 2).

141 In 2017, a new test well was drilled in Anhua County, Hunan Province. From the
142 bottom up, the lower Cambrian consisted of the Liuchapo, Niutitang, and Wunitang
143 formations. The present investigation focused on the Niutitang Formation, which has
144 previously been subdivided into three intervals based on the distribution of metal ore
145 layers (Ni-Mo, V, and Ba), carbonaceous shale, tuff, and fossils (Jin et al., 2016). The
146 boundary between intervals I and II is primarily characterized by Ni-Mo layers,
147 carbonaceous shale, and small shelly fossil assemblages (SSF1-4 of Steiner et al.
148 (2007)). The geochronology of the Ni-Mo layer deposited in the transition period
149 between Stages 2 and 3 of the Cambrian is dated to 521 ± 5 Ma, based on the Re-Os
150 isochron age (Xu et al., 2011). Shales with high TOC contents are frequently present
151 at the bottom of the Niutitang Formation and close to the metal ore layers. The
152 Niutitang Formation can be subdivided into three members, here named the Niu1,
153 Niu2 and Niu3 members. The lowermost Niu1 Member chiefly consists of dark gray
154 calcareous shales with calcareous patches (Figures 3A and B), with dolomite present
155 at the top of the member (Table 1). The Niu2 Member consists of black siliceous shale,

156 whereas the Niu3 Member principally consists of gray-black shales (Figures 3C and
157 D). However, small shelly fossils are not obvious in the study area. The boundary of
158 the Niu1 and Niu2 members can be characterized by a Ni-Mo layer and lithology. The
159 boundary between Niu2 and Niu3 can be identified according to the relative
160 concentration of trace metals (Mo, V, and U), lithology, and biota (Jin et al., 2016).
161 Trace metal elements of the Niu2 shale are much higher than those of the Niu3 shale
162 by more than one order of magnitude. The trace element values were also consistent
163 with four other sections and wells, including the Xy1 well (J. Li et al., 2018), the
164 Jinsha (Jin et al., 2016), and Dingtai sections (Xu et al., 2012) in the deep shelf facies,
165 and the Yk1 well in slope facies (Y. Li et al., 2015). The three members of the
166 Niutitang Formation can be identified from the GR/API results (Figure 1), with the
167 Niu2 Member having higher values than the Niu1 and Niu3 members. GR/API results
168 have also been applied to the Xy1 and Yk1 wells (Y. Li et al., 2015) and may
169 represent an important method for identifying the members in other well cores.

170 Through geological history, the study area underwent multiple periods of
171 tectonic movement, resulting in thrusting, folding, uplift, and denudation. The
172 (prototype) Nanhua basin was shaped during the Ediacaran, but marine deposition
173 dominated from the early Cambrian to the Triassic. After the mid-Triassic, the basin
174 was uplifted and significantly altered, forming a series of thrust nappes and
175 para-foreland basins (Wan et al., 2018). The Indosinian (257–205 Ma), Yanshanian
176 (199.6–133.9 Ma), and Himalayan (23 Ma-present) orogenies strongly influenced the
177 study area and formed many narrow anticlines and wide synclines. At present, the

178 outcrops of sedimentary rocks in western Hunan are mainly Paleozoic and Mesozoic
179 in age, and extend in a NE-SW direction. These strata form a compound syncline,
180 with the Cambrian-Silurian rocks in the northwest and southeast, while the
181 Devonian-Middle Triassic rocks are principally located in the center.

182

183 **MATERIALS AND METHODS**

184 Samples were collected from the X1 well in west Hunan Province (Figures 1B and C).
185 The Liuchapo Formation occurs at the bottom of the well core and consists primarily
186 of a series of gray-black cherts. Pyrite is abundant within the Niutitang shale and
187 occurs as nodules, veins and star spots (Figures 3E and F). The overlying Wunitang
188 Formation consists of dark gray, muddy limestones intercalated with shales.

189 Samples were taken from the chert and organic-rich shale interval of the core at a
190 spacing of 5–6 m (16–20 ft). A total of 4 Liuchapo chert and 28 Niutitang shale
191 samples were analyzed. Individual samples were ground to less than 100 mesh for
192 analyses of mineralogy and TOC and to below 200 mesh for analyses of trace and
193 major elements.

194 X-ray diffraction (XRD) measurements were conducted at the Chongqing
195 Mineral Resources Supervision and Test Center using a Bruker D8 Advance
196 diffractometer equipped with a Cu-target tube and a curved graphite monochromator.
197 Measurements were performed at 40 kV and 40 mA. Samples were step-scanned from
198 3–45° with 0.02° (2 θ)/steps.

199 To determine TOC values, 200 mg of sample was weighed and placed into a

200 combustion crucible. Excess hydrochloric acid (volume ratio 1:7) was added to
201 remove inorganic carbon. The crucible was then dried in an oven for 1h at 105 °C
202 under vacuum conditions. The TOC content was analyzed using a Multi N/C 3100
203 Analyzer (Germany). The analytical errors were better than ± 0.2 %.

204 For elemental analysis, the sample powder was dried in an oven for 2 h at 105 °C
205 under vacuum conditions. To measure the major elements, 500 mg of dried sample
206 was oxidized with 7 g of lithium borate (mixture of 67 % $\text{Li}_2\text{B}_4\text{O}_7$ and 33 %
207 anhydrous LiBO_2) for 2 h at 200 °C, and then the mixture was melted to make a fused
208 glass disk. Measurements were conducted using an Axios MAX pw4400/40 X-ray
209 fluorescence spectrometer. Major elements were represented by oxides (SiO_2 , Al_2O_3 ,
210 FeO , CaO , Fe_2O_3 , K_2O , MgO , MnO , Na_2O , P_2O_5 and TiO_2). To determine trace
211 elements, including Th, U, V, Ni, Co, Ba, Sr, Mo, Cu and Cr, 50 mg of dried sample
212 was weighed and treated using boric acid. Trace elements were then determined with
213 quadrupole inductively coupled plasma-mass spectrometry (ICP-MS). The analytical
214 errors were better than ± 5 %.

215 Enrichment factors (EF) were calculated based on the ratio between trace
216 element and aluminum (Al) concentrations of the sample. The same ratio in the
217 Post-Archean Australian Shale (PAAS, Taylor and McLennan, 1985). The following
218 formula was used for this calculation:

219

$$220 \quad X_{\text{EF}} = (X_{\text{sample}}/\text{Al}_{\text{sample}}) / (X_{\text{PAAS}}/\text{Al}_{\text{PAAS}}) \quad (1)$$

221

222 where X_{sample} and Al_{sample} are concentrations of trace element X and Al samples,
223 respectively; X_{PAAS} and Al_{PAAS} are concentrations of trace element X and Al in the
224 PAAS, respectively. $X_{\text{EF}} > 1.0$ and $X_{\text{EF}} < 1.0$ indicate enrichment and depletion in the X
225 element, respectively.

226

227 **RESULTS**

228 **Mineral Composition**

229 Quartz, feldspar, calcite, dolomite, pyrite and clay are the principal minerals (Table 1).
230 Except for certain Niu1 samples, all samples show >60 % of brittle minerals (e.g.,
231 quartz, feldspar, and pyrite), while typically <30 % are ductile minerals (i.e., clay
232 minerals). Quartz dominates the mineral content ($\bar{x}=61.0$ %), whereas the average
233 contents of feldspar and carbonates are much lower at 4.3 %, and 7.5 %, respectively
234 (Figure 4). Two samples from the boundary between the Niu1 and Niu2 members
235 include barite (2.7 %) and glauberite (3.8 % and 29.0 wt.%). The quartz content of
236 Liuchapo chert generally exceeds 94 %.

237

238 **TOC Content**

239 The TOC content varies from 0.1–20.1 wt.% (Table 2). The TOC generally increases
240 from the Niu1 Member (0.1–4.0 wt.%, $\text{mean}(\bar{x})=1.5$ wt.%) to the Niu2 Member (8.1–
241 20.1 wt.%, $\bar{x}=12.7$ wt.%) and then decreases to the Niu3 Member (2.6–8.1 wt.%,
242 $\bar{x}=5.1$ wt.%) (Figure 5). The TOC contents of the Liuchapo cherts are lower than 0.4
243 wt.%.

244

245 **Major Element Composition**

246 The major element compositions show considerable variations among the three
247 members (Table 2). Except for a dolomite sample, the elements of the analyzed shale
248 samples are dominated by SiO_2 (30–74 %). The SiO_2 concentrations are commonly
249 higher in the Niu2 and Niu3 samples than in the Niu1 samples. The Al_2O_3
250 concentrations are variable for the Niu1 samples, ranging from 2–17 %. Most of the
251 Niu2 samples show Al_2O_3 contents below 6 %, while all Niu3 samples show Al_2O_3
252 contents higher than 6%. The Fe_2O_3 concentrations are below 5 % for all analyzed
253 samples, with the Niu3 samples showing higher values than those of the Niu1 and
254 Niu2 samples. Compared to the Niu2 and Niu3 shales, the Niu1 shale samples contain
255 relatively high MgO and CaO concentrations that can reach >10 %. In addition, the
256 K_2O concentration of in the Niu2 samples is relatively lower than that in the Niu1 and
257 Niu3 shales. Commonly, the FeO, TiO_2 , Na_2O , P_2O_5 , and MnO contents are below
258 1 %.

259

260 **Trace Element Composition**

261 Trace element values are given in Tables 3 and 4. The Cu, Ni, and Ba contents are
262 typically high in the Niu2 samples, moderate in the Niu3 samples, and low in the
263 Niu1 samples. Compared to the Liuchapo chert samples, the Niutitang samples are
264 rich in Mo, U, and V. The enrichment factors of these three redox elements vary
265 significantly (Table 4). The Niu1 samples are weakly enriched with Mo (Mo_{EF} : 3–

244), V (V_{EF} : 1–17), and U (U_{EF} : 6–31). The enrichment of these three elements is stronger in the Niu2 samples (Mo_{EF} : 168–1265, V_{EF} : 6–129, and U_{EF} : 31–657) and moderate in the Niu3 samples (Mo_{EF} : 42–161, V_{EF} : 1–4, and U_{EF} : 5–31).

DISCUSSION

Paleoclimate

Paleoclimate can greatly influence marine productivity, redox conditions, and the supply of terrestrial materials. From an analysis of paleomagnetism, South China was inferred to be located in a subtropical-tropical zone near the equator during the early Cambrian (Mckerrrow et al., 1992; Yeasmin et al., 2017). Quantitative information on the paleoclimatic conditions can be interpreted from geochemical proxies because trace elements in sedimentary rocks are sensitive paleoclimate indicators in several cases (Galloway et al., 2013; Vosoughi Moradi et al., 2016; Yeasmin et al., 2017).

The Chemical Index of Alteration (CIA) is used to identify climate conditions in deep time (Nesbitt and Young, 1982). CIA values from 50–65 indicate cold climates with weak chemical weathering, from 65–85 indicate warm and humid climates with moderate weathering, and from 85–100 represent hot and humid climates with strong weathering (Nesbitt and Young, 1982; Cole et al., 2009). Furthermore, CIA is also related to the parent rock type; when carbonate parent rocks dominate, the ratio is relatively low (Zhai et al., 2018). CIA is calculated using the molar oxides according to the following formula:

$$\text{CIA} = [\text{Al}_2\text{O}_3 / (\text{Al}_2\text{O}_3 + \text{CaO}^* + \text{Na}_2\text{O} + \text{K}_2\text{O})] \times 100 \quad (2)$$

289

290 where CaO^* was corrected by using P_2O_5 data ($\text{CaO}^* = \text{CaO} - 10/3 \times \text{P}_2\text{O}_5$). If the
 291 content of CaO was less than that of Na_2O , the CaO value was used as CaO^* . In
 292 contrast, CaO^* was assumed to be equivalent to Na_2O (McLennan, 1993; Fedo et al.,
 293 1995; Zhai et al., 2018).

294 In this research, the lithology of the Niu1 Member is principally calcareous shale
 295 (Table 2). The calculated CIA values of the Niu1 samples are relatively low
 296 ($\text{mean}(\bar{x})=71$) but should have been affected by the high carbonate contents (Figure 5).
 297 A detailed explanation of the entire Niu1 interval is thus complex based on CIA
 298 values alone. Nevertheless, glauberite occurs in the upper part of the Niu1 Member
 299 and the lower part of Niu2 Member. This suggests arid and hot climates (L. Wang et
 300 al., 2013). The average CIA value of all Niu2 shales is 75, and that of the Niu3 shales
 301 is 75. Because shale samples from these two intervals contain few carbonate minerals,
 302 CIA values indicate that the climate was warm and humid. Overall, the CIA results
 303 indicate that climate fluctuations occurred during the early Cambrian (Figure 5).
 304 Warm climates in the Niu2 and Niu3 members are more conducive to greater biomass
 305 than conditions occurring during the Niu1 Member.

306

307 **Redox Conditions**

308 Anoxic conditions are more conducive to OM preservation because of weak
 309 degradation (Tribovillard et al., 2012; Ocubalidet et al., 2018). The Niu2 Member

310 shale samples are rich in pyrite (Figures 3E and F), reflecting that the water column is
311 dominated by reducing conditions. The trace elements Mo, U and V are sensitive to
312 redox conditions in sedimentary environments and hardly migrate during diagenesis;
313 they are excellent proxies for reconstructing redox conditions (Algeo and Maynard,
314 2008; Algeo and Tribovillard, 2009; Algeo and Rowe, 2012; Wu et al., 2016). The
315 precipitation of U occurred earlier than that of Mo in the suboxic environment. In this
316 study, we rely on Mo geochemistry (Scott and Lyons, 2012) and trace metal
317 enrichment (Mo_{EF} and U_{EF}) (Tribovillard et al., 2006; Algeo and Tribovillard, 2009) as
318 paleoredox proxies.

319 Previous studies on modern oceans have shown that Mo contents less than 25
320 ppm, between 25–100 ppm, and more than 100 ppm indicate non-euxinic, intermittent,
321 and strongly euxinic environments, respectively (Scott and Lyons, 2012). In this
322 research, all Niu1 samples show Mo contents below 25 ppm ($\bar{x} = 8.3$ ppm), indicating
323 the predominance of non-euxinic environments (Figure 5). The Mo contents of the
324 Niu2 samples are variable, ranging from 18.0–209.0 ppm ($\bar{x} = 83.4$ ppm). Although
325 there are two samples below 25 ppm, most Niu2 samples have Mo contents exceeding
326 50 ppm or even 100 ppm, suggesting intermittent euxinic/euxinic environments. The
327 Mo contents of most Niu3 samples are above 25 ppm but below 100 ppm (ranging
328 from 24.3–76.2 ppm, $\bar{x} = 40.3$ ppm), indicating the predominance of intermittent
329 euxinic environments. In addition, shale samples fall into different zones on the
330 covariation plot of Mo_{EF} - U_{EF} , indicating different sedimentary environments (Figure
331 7). Niu1 shales were deposited under primarily suboxic conditions, as shown by the

332 weak enrichment of Mo and U (i.e., low Mo_{EF} and U_{EF}). In contrast, the $Mo_{EF} - U_{EF}$
333 of all Niu2 samples are high, and those of the Niu3 samples are moderate, indicating
334 that the shales were deposited under primarily oxygen-depleted conditions, i.e.,
335 euxinic and ferruginous (e.g., anoxic) environments (Algeo and Li., 2020). These
336 conclusions are consistent with the Mo geochemistry observed.

337 The redox conditions in the early Cambrian were also related to sea level
338 changes. During Niu1 shale deposition, there was a short-term marine regression with
339 seawater shallowing and becoming oxygen-rich (Figure 5). Subsequently, as the sea
340 level increased, the oxygen content decreased in the seawater; thus, reducing
341 conditions during Niu2 and Niu3 deposition increased. The ferruginous-euxinic
342 threshold of this study is 10 wt.% (e.g., TOC). Pi et al. (2013) reported a threshold
343 value of 5 wt.% because of differences in terrigenous dilution and marine
344 paleoproductivity. TOC content vs Mo/Al ($r = +0.44$, $p(a) < 0.05$, $n = 28$) (Figure 8A)
345 and TOC content vs V/Al ($r = +0.73$, $p(a) < 0.01$, $n = 28$) (Figure 8B) demonstrate
346 that redox conditions are conducive to OM enrichment in the Niutitang shale.
347 Consequently, all selected parameters suggest that the reduction strength in the early
348 Cambrian was in the order of $Niu2 > Niu3 > Niu1$.

349

350 **Terrigenous Debris**

351 All analyzed shale samples contain a high proportion of quartz, with an average
352 content of 65 % (Table 1). Siliceous minerals can be derived from terrestrial,

hydrothermal, or biogenic origins (Pi et al., 2013; X. Yang et al., 2018). Al and Ti are mainly related to terrigenous detritus (Wedepohl, 1971). In the Niutitang shale samples, the correlation between Al_2O_3 and SiO_2 is very weak (Figure 9A), indicating that the silica may not be largely derived from continental sources. In addition to terrigenous origins, a large amount of silica in the shales must therefore originate from other sources. The Al_2O_3 contents in the Niutitang shales are highly variable (Table 1), with the Niu1 Member ($\bar{x}=6.8\%$) > Niu2 Member ($\bar{x}=3.8\%$) < the Niu3 Member ($\bar{x}=9.9\%$). This suggests that the distance of Niu2 shales from the clastic source was greater than that distance of either the Niu1 or Niu3 shales. In addition, the TOC contents also show negative correlations with the TiO_2 contents, indicating the terrigenous dilution of OM (Figure 9B).

Hydrothermal Sedimentation Processes

The $\text{Al}/(\text{Al}+\text{Mn}+\text{Fe})$ ratio is used to determine siliceous origins (Adachi et al., 1986; Yamamoto, 1987), with a pure hydrothermal activity ratio of 0.01 and a pure biogenic origin having a ratio of more than 0.60 (Yamamoto, 1987; Harris et al., 2011). In addition, biogenic shales are characterized by high SiO_2 and P_2O_5 values and low Al_2O_3 , TiO_2 , and MgO values, whereas the enrichment of Fe and Mn is mainly related to hydrothermal activity (X. Wang et al., 2016; Liao et al., 2018). Similarly, the $(\text{Fe}+\text{Mn})/\text{Ti}$ ratio is an index for determining conditions of hydrothermal deposition. Typical hydrothermal deposits are characterized by $\text{Al}/(\text{Al}+\text{Mn}+\text{Fe}) < 0.4$ and $(\text{Fe}+\text{Mn})/\text{Ti} > 15$ (Adachi et al., 1986). Niutitang shale samples fall in the different

375 zones on an Al-Fe-Mn ternary diagram (Figure 10), indicating that the silica comes
376 from different origins. Cherts from the Liuchapo Formation exhibit low
377 Al/(Al+Mn+Fe) ratio values (0.1–0.3), and high (Fe+Mn)/Ti ratio values (18.1–37.7),
378 suggesting intense hydrothermal activity (Figure 6). Rare earth element geochemistry
379 of the Liuchapo cherts also suggests that intense hydrothermal activity occurred in the
380 deep water area (Guo et al., 2016). Siliceous genesis of the Niu1 and Niu2 members
381 mainly relates to intense hydrothermal activity ($\text{Al}/(\text{Al}+\text{Mn}+\text{Fe}) = 0.4\text{--}0.8$,
382 $(\text{Fe}+\text{Mn})/\text{Ti} = 5.6\text{--}17.9$), whereas that of the Niu3 Member relates more to
383 non-hydrothermal origins ($\text{Al}/(\text{Al}+\text{Mn}+\text{Fe}) > 0.60$, $(\text{Fe}+\text{Mn})/\text{Ti} < 15$). Z. Wang et al.,
384 (2020) also found that hyalophane was developed at different depths in the Niu1 and
385 Niu2 shales. The hyalophane with another minerals (e.g., spehalerite and monazite)
386 can be an important proxy for hydrothermal activity. These results are in accordance
387 with the discovery of volcanic material (Figure 2), hydrothermal vent communities
388 and hydrothermal minerals at the bottom of the Niutitang Formation (R. Yang et al.,
389 2008; Han et al., 2017). The comparison of rare earth elements (Eu anomalies) in
390 several sections from South China also confirms that hydrothermal activity occurred
391 during the early Cambrian (Guo et al., 2016; Gao et al., 2018).

392 In general, the non-detrital fraction of a given element can be estimated by
393 subtracting the detrital fraction from the total amount of the element. Regarding Si,
394 we assume that all excess Si in the non-detrital fraction and is entirely derived from
395 either hydrothermal or biological origins. The calculations for $\text{Si}_{\text{excess}}$ and $\text{SiO}_{2\text{excess}}$
396 are:

397

$$398 \quad Si_{\text{excess}} = Si_{\text{sample}} - (Si/Al)_{\text{background}} \times Al_{\text{sample}} \quad (3)$$

$$399 \quad SiO_{2\text{excess}} = SiO_{2 \text{ sample}} - [(Si/Al)_{\text{background}} \times Al_{\text{sample}}] \times 60.1/28.1 \quad (4)$$

400

401 where Si_{excess} represents the non-detrital silicon content, Si_{sample} represents the silicon
402 concentration in the sample, Al_{sample} represents the Al concentration in the sample and
403 the (Si/Al) background value is 3.11 (Wedepohl, 1971).

404 According to Figure 5, excess silica amounts increase from the top of the Niu1 to
405 Niu2 members and then decrease in the Niu3 Member. The excess silica contents of
406 the Niu1 Member vary from 0–49.0 % ($\bar{x}$ =23.8 %). The contents of the Niu2 samples
407 are commonly high, between 41.4 % and 77.6 % ($\bar{x}$ =57.5 %), while those of the Niu3
408 samples are moderate, varying from 27.6–50.3 % ($\bar{x}$ =35.3 %). This implies that a
409 large proportion of silica should be of either a hydrothermal or biogenic origin, with a
410 certain amount of silica originating from terrigenous detritus. In addition, the origin of
411 chert and the source of silica during the Ediacaran-Cambrian transition are still widely
412 disputed. It has been argued that excess silica in shales may be related to
413 hydrothermal activity and also continental weathering (D. Chen et al., 2009; D. Lin et
414 al., 2015; Liu et al., 2015). In this research, although the CIA values suggest moderate
415 weathering during deposition of the Niu2 and Niu3 members, the CIA values do not
416 correlate with the excess silica contents observed, even for samples deposited under
417 the shallowest water conditions (e.g. in the Niu1 Member) (Figure 11A). This
418 indicates insignificant influences of continental weathering on silica enrichment in the

419 analyzed samples. In contrast, the excess silica content positively correlates with the
420 ratios of $Al/(Fe+Al+Mn)$ (Figure 11B), implying that the silica contents were affected
421 by hydrothermal events. The correlations between excess silica and TOC contents are
422 significant for the Niu3 samples (Figure 11C), but are unclear for the Niu1 and Niu2
423 samples (Figure 11D). Organisms most likely thrived during the deposition of Niu3
424 and provided biogenic silica. However, the silica sources of the Niu1 and Niu2
425 members should be dominated by hydrothermal sources. Liu et al. (2015) analyzed
426 the interbedded layers of chert and shale from the Yangtiao section in Guizhou
427 Province and they concluded that chert and shale intervals were deposited during
428 intense and weak hydrothermal activity, respectively. Strong hydrothermal events
429 could release more silica into the ancient ocean than weaker events, thus forming
430 silica-rich rocks (e.g., chert). Nevertheless, in the early Cambrian hydrothermal events
431 may not only carry silica but also abundant redox-sensitive metal elements into the
432 water column. In the current study, certain redox-sensitive trace elements (Mo, U, and
433 V) from the X1 well and a typical hydrothermal plume from the Mid-Atlantic Ridge
434 were selected to illustrate whether hydrothermal activity affected the redox conditions
435 (German et al., 1991a). Through observations of the Niu3 deposition processes, no
436 obvious hydrothermal activity occurred. Thus, correlations among these three
437 elements of the Niu3 shales are still evident (Figures 12A and B). Correlations among
438 these three elements from the Niu1 and Niu2 shales are all much weaker than those
439 correlations among the three elements of the Niu3 shales (Figures 12C and D), which
440 is consistent with correlations observed in typical hydrothermal samples (Figure 12E).

441 However, the correlations of the Mo and V elements from the hydrothermal plume are
442 better than those of the X1 well samples (Figure 12F). Previous studies show that V
443 and U concentrations both exhibit pronounced linear correlations with hydrothermal
444 intensity based on samples from the Mid-Atlantic Ridge (German et al., 1991b).
445 Hydrothermal fluid may be conducive to the enrichment of these two elements. In our
446 work, V_{EF} and U_{EF} are relatively high in samples deposited in shallow water
447 conditions (e.g., Niu1 Member) but low in deep water samples (e.g., Niu3 Member).
448 This phenomenon suggests that hydrothermal activity not only provided redox trace
449 elements but also enhanced marine reducing conditions.

450

451 **Marine Productivity**

452 Paleoproductivity, referring to the amount of OM produced in a certain period of
453 geological history (Taylor, 1964; Gupta and Kawahata, 2006; Tribovillard et al., 2006;
454 Zhao et al., 2017), is typically evaluated using TOC as a proxy. However, TOC is
455 commonly affected by diagenesis, redox reactions, biological action, and the dilution
456 of macerals, which may lead to either over- or underestimates of primary productivity
457 (Gupta and Kawahata, 2006; Tribovillard et al., 2006). Since Ba is an inert element
458 and has a long residence time in oceans, it is commonly used as an auxiliary indicator
459 of paleoproductivity (Taylor, 1964; Böttcher et al., 2003; Schoepfer et al., 2015). In
460 most modern aquatic sediments, Ba mainly exists in the form of $BaSO_4$. There is a
461 significant correlation between Ba content of sediments and biological productivity in

the overlying water column. To eliminate the influence of Ba from terrestrial debris, the content of biogenic Ba (Ba_{bio}) is obtained by subtracting the terrestrial fraction from the total Ba. Ba_{bio} is calculated by the formula:

$$Ba_{bio} = Ba_{sample} - Ba_{detrital} = Ba_{sample} - Al_{sample} \times (Ba/Al_{alu}) \quad (5)$$

where Ba_{sample} is the Ba concentration in the sample, Al_{sample} is the Al concentration in the sample, and Ba/Al_{alu} is the ratio of Ba and Al in PAAS. The Ba/Al_{alu} ratio is the Ba/Al ratio of the aluminosilicate component ranging from 0.005–0.01 in crustal rocks (Taylor, 1964). A value of 0.0075 is widely used to calculate the contents of Ba_{bio} (Dymond et al., 1992; Zhao et al., 2016). In general, the Niutitang shales exhibit moderate to high degrees of calculated Ba_{bio} contents (38 to 8264 ppm; averaging 1382 ppm), compared to those values documented from the modern equatorial Pacific (~1000–5000 ppm) (Murray and Leinen, 1993). The Ba_{bio} contents are variable in the three members, showing mean values of Niu1 ($\bar{x}=1199$ ppm) < Niu2 ($\bar{x}=1642$ ppm) > Niu3 ($\bar{x}=932$ ppm). The Ba and Ba_{bio} contents slowly vary vertically throughout the shale interval, with Niu2 shale samples being slightly more enriched than Niu1 and Niu3 shale samples. The correlation between Ba_{bio} and TOC contents is unclear (Figure 6). This may be caused by the loss in Ba as a consequence of the elevated rate of bacterial sulfate reduction (Schoepfer et al., 2015). During the deposition of Niu3, Ba_{bio} contents were less than 1000 ppm, since $BaSO_4$ was easily reduced under anoxic conditions. Thus, the Ba_{bio} contents of the Niu1 samples that accumulated under

484 suboxic conditions can be reliably used to evaluate paleoproductivity.

485 Paleoproductivity was relatively high in the Niu1 and Niu2 intervals, and it was
486 moderate in the Niu3 interval.

487 In addition to biogenic Ba, Ni and Cu can also reflect initial productivity (Calvert
488 and Pedersen, 1993; Algeo and Maynard, 2004; Tribovillard et al., 2006). In modern
489 oceanic sediments, Cu and Ni show positive linear correlations with TOC. As nutrient
490 elements, they either combine with OM or form organic complexes (Froelich et al.,
491 1979; Emerson et al., 1985). High Cu and Ni contents therefore relate to high OM
492 inputs, reflecting higher paleoproductivity (Emerson et al., 1985; Piper and Perkins,
493 2004; Zhao et al., 2016). To exclude the influence of terrigenous clastic dilution,
494 Cu/Al and Ni/Al ratios were used to characterize paleoproductivity. The Cu/Al and
495 Ni/Al ratios vary to a certain degree in the three members, showing mean values of
496 Niu1 (Cu/Al = 8.8 and Ni/Al = 8.3) < Niu2 (Cu/Al = 77.3 and Ni/Al = 73.5) > Niu3
497 (Cu/Al = 10.1 and Ni/Al = 16.6) (Figure 6). Paleoproductivity in the Niu2 Member is
498 higher than that in either the Niu1 or Niu3 members, which is consistent with
499 interpretations observed from the biogenic Ba. The TOC contents of the Niu3 shales
500 are generally lower than 10 wt.% (Table 2), with an average of 5.1 wt.%, and close to
501 that of the Longmaxi Formation in the Sichuan Basin (Yan et al., 2012; Feng et al.,
502 2018). Previous studies have shown that silica genesis in the Longmaxi Formation
503 shale is dominated by biological origins, rather than hydrothermal sources (Liu et al.,
504 2017; Luo et al., 2017; Zhao et al., 2017). After demonstrating the existence of
505 hydrothermal events, there is still a need to analyze the effect of hydrothermal activity

on organismal abundance. Hydrothermal activity may not only provide nutrients, but may also affect organismal survival (Z. Wang et al., 2020). The depositional environments of the lower part of the Niutitang Formation were dominated by an intermittent sulfur-containing water column (C. Li et al., 2010; Jin et al., 2016). Chen et al. (2009) analyzed samples from the Ganziping section in Hunan Province (Figure 2) and reported that hydrothermal venting released large amounts of silica-rich hydrothermal fluids with greenhouse gases (methane) and volcanic-derived H₂S into the ocean and/or atmosphere during the early Cambrian. Gao et al. (2018) also documented that hydrothermal venting probably promoted a sulfidic ocean chemistry, which then gradually evolved into a persistent anoxic ocean. The nutrients provided by hydrothermal activity were conducive to the reproduction of phytoplankton (e.g., algae) and thus enhanced the supply of OM (Uematsu et al., 2004; Duggen et al., 2010). Toxic elements (e.g., Hg, Pb, and Cr) and volatile gases (e.g., HCl and SO₂) carried by hydrothermal events could also have inhibited the growth of zooplankton (Jones and Gislason, 2008; Chambers et al., 2013). The decomposition of OM during the deposition process could consume a large amount of oxygen in the water column, and significantly affect living organisms. The ecological imbalance probably strengthened the reducing potential of the water column and in turn was beneficial to the preservation of OM. By analyzing the evolutionary events of early Cambrian organisms, the first stage biota comprises small shelly fossils that became extinct before 521 Ma, while the Chengjiang fauna represents the second stage biota that began to expand at 518.03 ± 0.69 Ma (D. Wang et al., 2018; C. Yang et al., 2018). The

528 large-scale pervasive oxygenation of the ocean was not the primary control on animal
529 diversity (Jin et al., 2016; Xiang et al., 2017). Hydrothermal activity may inhibit
530 zooplankton reproduction and accelerate phytoplankton reproduction (Z. Wang et al.,
531 2020), as well as increase OM contents in sedimentary rocks. Under reducing
532 conditions, greater OM content in sediment can be preserved. In this research, the
533 TOC content of the Niu2 shale is generally in excess of 10 wt.% and in certain cases
534 up to 20 wt.% (Table 2). Hydrothermal activity may be one of the main reasons for
535 OM enrichment in the Niutitang Formation.

536

537 **Seawater Restriction**

538 Mo can be used to determine the degree of water restriction in ancient sediments.
539 Mo exists in a stable state with a high valence (+6) in oxygen-enriched water and is
540 difficult to deposit in sediments. However, Mo(VI) can convert from molybdates to
541 thiomolybdates, and be transferred to anoxic sediments (Taylor and McLennan, 1985;
542 Algeo and Tribovillard, 2009). The average Mo content in the upper crust is as low as
543 3.7 ppm (Zheng et al., 2000; Rowe et al., 2008; Morford et al., 2009; Tribovillard et
544 al., 2012). Furthermore, Mo has a long residence time in water (approximately 780
545 kyr). This unique feature of Mo is often used to assess the restriction of modern and
546 ancient seawater (Tribovillard et al., 2012; Zhao et al., 2016; Liu et al., 2017). The
547 relationship between Mo and TOC can reveal the degree of anoxic water restriction
548 but not oxic/suboxic conditions. In this research, the Niu1 Member is shown to be

549 dominated by suboxic sedimentary conditions (Figure 13). An oxygen-containing
550 water column was not conducive to the precipitation of Mo, so the Mo contents are
551 commonly low. In addition, the seawater was shallow, and poorly connected to the
552 surrounding ocean such that the environment should be strongly restricted. The
553 Mo/TOC ratio of the bottom part of the Niu2 Member varies from 1.2–5.8, indicating
554 that the water column was strongly restricted close to the values from recent Black
555 Sea sediments (Mo/TOC = 4.5) (Algeo and Lyons, 2006; Tribovillard et al., 2012).
556 However, the Mo/TOC ratio of the upper part of the Niu2 shale is closer to present
557 day values of the Cariaco Basin (Mo/TOC = 25) (Algeo et al., 2007; Tribovillard et al.,
558 2012), indicating a weakly to moderately restricted water environment. With sea level
559 falling, seawater became more restricted in the Niu3 Member, with the Mo/ TOC
560 value of this member becoming close to that of the present day Framvaren Fjord value
561 (Mo/TOC = 9).

562 Although the eastern Xuefeng Uplift experienced euxinic environments during
563 deposition of the Niu2 Member, the shale was also affected by hydrothermal activity.
564 However, the TOC content changes markedly in the vertical direction within the Niu2
565 shale, ranging from 8.1–20.1 wt.% (Table 1). The restricted marine environment may
566 play an important role in the enrichment of OM in shales. The water column was
567 strongly restricted in the lower part of the Niu2 Member, with the seawater isolated to
568 a large degree from the surrounding ocean. This provided a low water exchange rate
569 and allowed the water to carry less oxygen, which was conducive to preserving OM in
570 low oxygenated conditions. As the sea level rose, the water column changed to a

571 moderately/weakly restricted environment in the upper part of the Niu2 Member. The
572 water exchange rate was faster and the water carried more oxygen, resulting in more
573 OM being degraded under oxygenated conditions. Thus, seawater restriction was also
574 an important factor affecting OM enrichment.

575

576 **OM Accumulation Mechanism in the Niutitang Formation**

577 The TOC content of the Niutitang shale is variable (Table 2) for which the
578 mechanisms of OM accumulation in the three members are significantly different
579 (Figure 14). In the Niu1 interval, the climate was hot, and marine productivity was
580 comparatively moderate. Weak hydrothermal activity was present. Nutrients and
581 volatile gas entered the water column (Figure 14A), and they slightly advanced the
582 initial productivity and the reducing potential of the seawater. The weak hydrothermal
583 activity had less relative effect on the redox and chemical conditions of the water
584 column. The redox conditions were dominated by suboxic environments. This
585 interpretation is consistent with the lithology analysis (e.g., calcareous shale). OM
586 was easily decomposed and consumed in oxygen-containing seawater and thus was
587 not conducive to preservation. The TOC contents of the Niu1 shale are below 4 wt.%.
588 Thus, redox condition was the main factor controlling OM accumulation during the
589 Niu1 interval.

590 During the Niu2 interval, the climate was humid and warm, with marine
591 productivity at its highest (Figure 14B). The redox conditions of seawater were
592 dominated by an euxinic environment. Strong hydrothermal activity occurred.

593 Abundant nutrients and volatile gas entered the water column, and they strongly
594 increased the initial productivity and greatly enhanced the reducing potential of the
595 seawater. The nutrients from hydrothermal activity were conducive to algae thriving,
596 but the euxinic seawater inhibited the survival of zooplankton (e.g., radiolaria). In
597 addition, algal reproduction led to a significant increase in the OM that was deposited
598 and buried in the marine basin. The Niu2 interval can be divided into two parts due to
599 seawater restriction. In the lower part of the Niu2 interval, which is associated with
600 the sea level slowly rising, the water column was gradually transformed into euxinic
601 conditions. Seawater was also strongly restricted, and had poor connectivity with
602 surrounding oceans. The strongly restricted environment could have resulted in a slow
603 exchange rate of oxygen and an enhanced reducing potential, which is similar to the
604 current state of the Black Sea. It also reduced the consumption of OM by limiting
605 oxygen. The TOC contents of the lower Niu2 shales are generally above 13 wt.% and
606 can be up to 20 wt.%. In the upper part of the Niu2 interval, which is associated with
607 the sea level rising, the water column transformed into a weakly restricted
608 environment. The seawater had good connectivity with the surrounding oceans. A
609 rapid exchange of oxygen could have consumed a significant amount of OM. The
610 TOC contents of the upper Niu2 shales are generally greater than 10 wt.%. Overall,
611 the restricted marine environments, redox conditions, and hydrothermal activity
612 enhanced OM enrichment during the deposition of Niu2 shales.

613 During the Niu3 interval, the climate was similar to that of the Niu2 interval
614 (Figure 14C). The study area changed into a ferruginous environment due to sea level

615 decline. The water column was also moderately restricted, similar to that of the upper
616 Niu2 interval. In addition, hydrothermal activity was not obvious, and led to the
617 restoration of ecological balance. Zooplankton reproduction was not limited. More
618 algae were consumed and thus abated the supply of OM. The reducing potential of the
619 water column had not been strengthened by the other causes introduced. The TOC
620 contents of the Niu3 shales are generally less than 10 wt.%. Redox conditions and
621 moderately restricted environments both played an important role in the preservation
622 of OM.

623

624 **CONCLUSIONS**

625 Based on the analysis of lithofacies, mineralogy, and a series of proxies from major
626 and trace elements in the lower Cambrian Niutitang shale samples from the eastern
627 Xuefeng Uplift, we show that the Niutitang Formation includes three distinct
628 sedimentary periods corresponding to the Niu1, Niu2, and Niu3 members. Redox
629 proxies (Mo geochemistry, Mo_{EF} , and U_{EF}) indicate that the Niu1, Niu2 and Niu3
630 members were deposited in suboxic, euxinic, and ferruginous environments,
631 respectively. Climatic conditions and paleoproductivity proxies (Ba_{bio} , Cu/Al, and
632 Ni/Al) suggested an overall paleoproductivity trend of $Niu1 < Niu2 > Niu3$.
633 Submarine hydrothermal activity probably appeared in the Niu1 interval, strengthened
634 in the Niu2 interval, and was insignificant in the Niu3 interval. Nutrients provided by
635 hydrothermal activity were conducive to phytoplankton (e.g., algae) thriving and thus
636 increased the OM supply. However, volatile gases (e.g., HCl and SO₂) and euxinic

637 conditions may inhibit the survival of zooplankton. An ecological imbalance
 638 strengthened the reducing potential of the water column that led to a large increase in
 639 the amount of OM deposited and buried in marine basins. In addition, due to the
 640 variation in sea level, the strongly restricted environment in the early Niu2 interval
 641 was more beneficial to the preservation of OM than that occurring in the upper Niu2
 642 and Niu3 intervals. Hydrothermal activity may play a dual role in the supply and
 643 preservation of OM, which is the key factor of OM enrichment in the lower Cambrian
 644 Niutitang shale.

645

646 REFERENCES CITED

- 647 Adachi, M., K. Yamamoto, and R. Sugisaki, 1986, Hydrothermal chert and associated
 648 siliceous rocks from the northern Pacific their geological significance as
 649 indication of ocean ridge activity: *Sedimentary Geology*, v. 47, no. 1, p. 125–148,
 650 doi:10.1016/0037-0738(86)90075-8.
- 651 Algeo, T. J., and J. B. Maynard, 2004, Trace–element behavior and redox facies in
 652 core shales of Upper Pennsylvanian Kansas-type cyclothems: *Chemical Geology*,
 653 v. 206, no. 3, p. 289–318, doi:10.1016/j.chemgeo.2003.12.009.
- 654 Algeo, T. J., and T. W. Lyons, 2006, Mo–total organic carbon covariation in modern
 655 anoxic marine environments: Implications for analysis of paleoredox and
 656 paleohydrographic conditions: *Paleoceanography*, v. 21, no. 1, p. 1–23,
 657 doi:10.1029/2004PA001112.
- 658 Algeo, T. J., T. W. Lyons, R. C. Blakey, and D. J. Over, 2007, Hydrographic
 659 conditions of the Devonian–Carboniferous North American Seaway inferred from
 660 sedimentary Mo–TOC relationships: *Palaeogeography, Palaeoclimatology,*
 661 *Palaeoecology*, v. 256, no. 3, p. 204–230, doi:10.1016/j.palaeo.2007.02.035.
- 662 Algeo, T. J., and J. B. Maynard, 2008, Trace–metal covariation as a guide to water–
 663 mass conditions in ancient anoxic marine environments: *Geosphere*, v. 4, no. 5, p.
 664 872–887, doi:10.1130/GES00174.1.
- 665 Algeo, T. J., and N. Tribouillard, 2009, Environmental analysis of paleoceanographic
 666 systems based on molybdenum–uranium covariation: *Chemical Geology*, v. 268,
 667 no. 3/4, p. 211–225, doi:10.1016/j.chemgeo.2009.09.001.
- 668 Algeo, T. J., and H. Rowe, 2012, Paleoceanographic applications of trace–metal
 669 concentration data: *Chemical Geology*, v. 324, p. 6–18,

doi:10.1016/j.chemgeo.2011.09.002.

Algeo, T. J., and C. Li, 2020, Redox classification and calibration of redox thresholds in sedimentary systems: *Geochimica et Cosmochimica Acta*, in press, doi: 10.1016/j.gca.2020.01.055.

Böttcher, M. E., J. Rinna, B. Warning, R. Wehausen, M. W. Howell, B. Schnetger, R. Stein, H. J. Brumsack, and J. Rullkötter, 2003, Geochemistry of sediments from the connection between the western and the eastern Mediterranean Sea (Strait of Sicily, ODP Site 963): *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 190, p. 165–194, doi:10.1016/S0031-0182(02)00604-1.

Bu, H., Y. Ju, J. Tan, G. Wang, and X. Li, 2015, Fractal characteristics of pores in non-marine shales from the Huainan coalfield, eastern China: *Journal of Natural Gas Science and Engineering*, v. 24, p. 166–177, doi:10.1016/j.jngse.2015.03.021.

Calvert, S. E., and T. F. Pedersen, 1993, Geochemistry of Recent oxic and anoxic marine sediments: Implications for the geological record: *Marine Geology*, v. 113, no. 1, p. 67–88, doi:10.1016/0025-3227(93)90150-T.

Chambers, R., A. Candelmo, E. Habeck, M. Poach, D. Wieczorek, K. Cooper, C. Greenfield, and B. Phelan, 2013, Ocean acidification effects in the early life-stages of summer flounder, *Paralichthys dentatus*: *Biogeosciences Discussions*, v. 10, no. 8, doi:10.5194/bgd-10-13897-2013.

Chen, D., J. Wang, H. Qing, D. Yan, and L. I. Renwei, 2009, Hydrothermal venting activities in the Early Cambrian, South China: Petrological, geochronological and stable isotopic constraints: *Chemical Geology*, v. 258, no. 3, p. 168–181, doi:10.1016/j.chemgeo.2008.10.016.

Chen, D., X. Zhou, Y. Fu, J. Wang, and D. Yan, 2015, New U–Pb zircon ages of the Ediacaran–Cambrian boundary strata in South China: *Terra Nova*, v. 27, no. 1, p. 62–68, doi:10.1111/ter.12134.

Chen, J., H. Cao, and P. Sun, 2017, Fracability evaluation of shale in the Niutitang Formation in northwestern Hunan: *Geoscience Frontiers*, v. 6, no. 24, p. 390–398, doi:10.13745/j.esf.yx.2017-2-50 (in Chinese with English abstract).

Cole, J. M., S. L. Goldstein, P. B. deMenocal, S. R. Hemming, and F. E. Grousset, 2009, Contrasting compositions of Saharan dust in the eastern Atlantic Ocean during the last deglaciation and African Humid Period: *Earth and Planetary Science Letters*, v. 278, no. 3, p. 257–266, doi:10.1016/j.epsl.2008.12.011.

Compston, W., Z. Zhang, J. A. Cooper, G. Ma, and R. J. F. Jenkins, 2008, Further SHRIMP geochronology on the early Cambrian of South China: *American Journal of Science*, v. 308, no. 4, p. 399–420, doi:10.2475/04.2008.01.

Duggen, S., N. Olgun, P. Croot, L. J. Hoffmann, H. Dietze, P. Delmelle, and C. Teschner, 2010, The role of airborne volcanic ash for the surface ocean biogeochemical iron-cycle: a review: *Biogeosciences (BG)*, v. 7, no. 3, p. 827–844, doi:10.5194/bg-7-827-2010.

Dymond, J., E. Suess, and M. Lyle, 1992, Barium in deep-sea sediment: A geochemical proxy for paleoproductivity: *Paleoceanography*, v. 7, no. 2, p. 163–181, doi:10.1029/92PA00181.

Emerson, S., K. Fischer, C. Reimers, and D. Heggie, 1985, Organic carbon dynamics and preservation in deep-sea sediments: Deep Sea Research Part A. Oceanographic Research Papers, v. 32, no. 1, p. 1–21, doi:10.1016/0198-0149(85)90014-7.

Fedo, C. M., H. W. Nesbitt, and G. M. Young, 1995, Unraveling the effects of potassium metasomatism in sedimentary rocks and paleosols, with implications for paleoweathering conditions and provenance: *Geology*, v. 23, no. 10, p. 921–924, doi:10.1130/0091-7613(1995)023<0921:UTEOPM>2.3.CO;2.

Feng, W., F. Wang, J. Guan, J. Zhou, F. Wei, W. Dong, and Y. Xu, 2018, Geologic structure controls on initial productions of lower Silurian Longmaxi shale in south China: *Marine and Petroleum Geology*, v. 91, p. 163–178, doi:10.1016/j.marpetgeo.2018.01.001.

Froelich, P. N., G. P. Klinkhammer, M. L. Bender, N. A. Luedtke, G. R. Heath, D. Cullen, P. Dauphin, D. Hammond, B. Hartman, and V. Maynard, 1979, Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis: *Geochimica et Cosmochimica Acta*, v. 43, no. 7, p. 1075–1090, doi:10.1016/0016-7037(79)90095-4.

Fuchida, S., Y. Mizuno, H. Masuda, T. Toki, and H. Makita, 2014, Concentrations and distributions of amino acids in black and white smoker fluids at temperatures over 200°C: *Organic Geochemistry*, v. 66, p. 98–106, doi:10.1016/j.orggeochem.2013.11.008.

Galloway, J. M., A. R. Sweet, G. T. Swindles, K. Dewing, T. Hadlari, A. F. Embry, and H. Sanei, 2013, Middle Jurassic to Lower Cretaceous paleoclimate of Sverdrup Basin, Canadian Arctic Archipelago inferred from the palynostratigraphy: *Marine and Petroleum Geology*, v. 44, p. 240–255, doi:10.1016/j.marpetgeo.2013.01.001.

Gao, P., Z. He, S. Li, G. G. Lash, B. Li, B. Huang, and D. Yan, 2018, Volcanic and hydrothermal activities recorded in phosphate nodules from the Lower Cambrian Niutitang Formation black shales in South China: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 505, p. 381–397, doi:10.1016/j.palaeo.2018.06.019.

German, C., A. Campbell, and J. Edmond, 1991a, Hydrothermal scavenging at the Mid-Atlantic Ridge: modification of trace element dissolved fluxes: *Earth and Planetary Science Letters*, v. 107, no. 1, p. 101–114, doi:10.1016/0012-821X(91)90047-L.

German, C., A. Fleer, M. Bacon, and J. Edmond, 1991b, Hydrothermal scavenging at the Mid-Atlantic Ridge: radionuclide distributions: *Earth and Planetary Science Letters*, v. 105, no. 1–3, p. 170–181, doi:10.1016/0012-821X(91)90128-5.

Goldberg, T., H. Strauss, Q. Guo, and C. Liu, 2007, Reconstructing marine redox conditions for the Early Cambrian Yangtze Platform: Evidence from biogenic sulphur and organic carbon isotopes: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 254, no. 1, p. 175–193, doi:10.1016/j.palaeo.2007.03.015.

Guo, Q., Y. Deng, D. Hippler, G. Franz, and J. Zhang, 2016, REE and trace element

756 patterns from organic-rich rocks of the Ediacaran–Cambrian transitional interval:
757 Gondwana Research, v. 36, p. 94–106, doi:10.1016/j.gr.2016.03.012.

758 Gupta, L. P., and H. Kawahata, 2006, Downcore diagenetic changes in organic matter
759 and implications for paleoproductivity estimates: Global and Planetary Change, v.
760 53, no. 1, p. 122–136, doi:10.1016/j.gloplacha.2006.01.008.

761 Han, T., H. Fan, X. Zhu, H. Wen, C. Zhao, and F. Xiao, 2017, Submarine
762 hydrothermal contribution for the extreme element accumulation during the early
763 Cambrian, South China: Ore Geology Reviews, v. 86, p. 297–308,
764 doi:10.1016/j.oregeorev.2017.02.030.

765 Hao, F., H. Zou, and Y. Lu, 2013, Mechanisms of shale gas storage: Implications for
766 shale gas exploration in China: AAPG Bulletin, v. 97, no. 8, p. 1325–1346,
767 doi:10.1306/02141312091.

768 Harris, N. B., J. L. Miskimins, and C. A. Mnich, 2011, Mechanical anisotropy in the
769 Woodford Shale, Permian Basin: Origin, magnitude, and scale: Leading Edge, v.
770 30, no. 3, p. 284–291, doi:10.1190/1.3567259.

771 Jin, C., C. Li, T. J. Algeo, N. J. Planavsky, H. Cui, X. Yang, Y. Zhao, X. Zhang, and S.
772 Xie, 2016, A highly redox-heterogeneous ocean in South China during the early
773 Cambrian (~529–514 Ma): Implications for biota–environment co–evolution:
774 Earth and Planetary Science Letters, v. 441, p. 38–51,
775 doi:10.1016/j.epsl.2016.02.019.

776 Jones, M. T., and S. R. Gislason, 2008, Rapid releases of metal salts and nutrients
777 following the deposition of volcanic ash into aqueous environments: Geochimica
778 et Cosmochimica Acta, v. 72, no. 15, p. 3661–3680,
779 doi:10.1016/j.gca.2008.05.030.

780 Knoll, A. H., and S. B. Carroll, 1999, Early animal evolution: emerging views from
781 comparative biology and geology: Science, v. 284, no. 5423, p. 2129–2137,
782 doi:10.1126/science.284.5423.2129.

783 Lane, N., J. F. Allen, and W. Martin, 2010, How did LUCA make a living?
784 Chemiosmosis in the origin of life: Bioessays, v. 32, no. 4, p. 271–280,
785 doi:10.1002/bies.200900131.

786 Li, C., G. D. Love, T. W. Lyons, D. A. Fike, A. L. Sessions, and C. Xuelei, 2010, A
787 stratified redox model for the Ediacaran ocean: Science, v. 328, no. 5974, p. 80–
788 83, doi:10.1126/science.1182369.

789 Li, J., S. Tang, S. Zhang, Z. Xi, N. Yang, G. Yang, L. Li, and Y. Li, 2018,
790 Paleo-environmental conditions of the Early Cambrian Niutitang Formation in the
791 Fenggang area, the southwestern margin of the Yangtze Platform, southern China:
792 Evidence from major elements, trace elements and other proxies: Journal of Asian
793 Earth Sciences, v. 159, p. 81–97, doi:10.1016/j.jseaes.2018.03.013.

794 Li, L., J. Tan, D. A. Wood, Z. Zhao, D. Becker, Q. Lyu, B. Shu, and H. Chen, 2019, A
795 review of the current status of induced seismicity monitoring for hydraulic
796 fracturing in unconventional tight oil and gas reservoirs: Fuel, v. 242, p. 195–210,
797 doi:10.1016/j.fuel.2019.01.026.

798 Li, Y., T. Fan, J. Zhang, J. Zhang, X. Wei, X. Hu, W. Zeng, and W. Fu, 2015,

799 Geochemical changes in the Early Cambrian interval of the Yangtze Platform,
800 South China: Implications for hydrothermal influences and paleocean redox
801 conditions: *Journal of Asian Earth Sciences*, v. 109, p. 100–123,
802 doi:10.1016/j.jseaes.2015.05.003.

803 Liao, S., C. Tao, H. Li, G. Zhang, J. Liang, W. Yang, and Y. Wang, 2018, Surface
804 sediment geochemistry and hydrothermal activity indicators in the Dragon Horn
805 area on the Southwest Indian Ridge: *Marine Geology*, v. 398, p. 22–34,
806 doi:10.1016/j.margeo.2017.12.005.

807 Lin, D., S. Bing, C. T. A. Lee, X. J. Shu, P. Yang, Y. Sun, Z. Tang, R. Hong, X. Lang,
808 and H. Ma, 2015, Germanium/silicon of the Ediacaran-Cambrian Laobao cherts:
809 Implications for the bedded chert formation and paleoenvironment interpretations:
810 *Geochemistry Geophysics Geosystems*, v. 16, no. 3, p. 751–763,
811 doi:10.1002/2014GC005595.

812 Lin, T., S. Bao, J. Zhang, Z. Zhou, K. Yuan, B. Li, and S. Yang, 2016, Shale gas
813 accumulation conditions and gas-bearing characteristics of the Lower Cambrian
814 Niutitang Formation in Well Changye-1 in northwestern Hunan Province:
815 *Petroleum Research*, v. 1, no. 2, p. 205–212,
816 doi:10.1016/S2096-2495(17)30043-1.

817 Little, C., and R. Vrijenhoek, 2003, Are hydrothermal vent animals living fossils?:
818 *Trends in Ecology & Evolution*, v. 18, no. 11, p. 582–588,
819 doi:10.1016/j.tree.2003.08.009.

820 Liu, Z., T. J. Algeo, X. Guo, J. Fan, X. Du, and Y. Lu, 2017, Paleo-environmental
821 cyclicity in the Early Silurian Yangtze Sea (South China): Tectonic or
822 glacio-eustatic control?: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v.
823 466, p. 59–76, doi:10.1016/j.palaeo.2016.11.007.

824 Liu, Z. H., X. G. Zhuang, G. E. Teng, X. M. Xie, L. M. Yin, L. Z. Bian, Q. L. Feng,
825 and T. J. Algeo, 2015, The lower Cambrian Niutitang Formation at Yantiao
826 (Guizhou, SW China): organic matter enrichment, source rock potential, and
827 hydrothermal influences: *Journal of Petroleum Geology*, v. 38, no. 4, p. 411–432,
828 doi:10.1111/jpg.12619.

829 Luo, Q., J. Hao, C. B. Skovsted, P. Luo, I. Khan, J. Wu, and N. Zhong, 2017, The
830 organic petrology of graptolites and maturity assessment of the Wufeng–
831 Longmaxi Formations from Chongqing, China: Insights from reflectance
832 cross-plot analysis: *International Journal of Coal Geology*, v. 183, p. 161–173,
833 doi:10.1016/j.coal.2017.09.006.

834 Lyu, Q., X. Long, P. G. Ranjith, J. Tan, and Y. Kang, 2018a, Experimental
835 investigation on the mechanical behaviours of a low-clay shale under water-based
836 fluids: *Engineering Geology*, v. 233, p. 124–138,
837 doi:10.1016/j.enggeo.2017.12.002.

838 Lyu, Q., X. Long, P. G. Ranjith, J. Tan, Y. Kang, and Z. Wang, 2018b, Experimental
839 investigation on the mechanical properties of a low-clay shale with different
840 adsorption times in sub-/super-critical CO₂: *Energy*, v. 147, p. 1288–1298,
841 doi:10.1016/j.energy.2018.01.084.

842 Mckerrow, W. S., C. R. Scotese, and M. D. Brasier, 1992, Early Cambrian continental
843 reconstructions: *Journal of the Geological Society*, v. 149, no. 4, p. 599–606,
844 doi:10.1144/gsjgs.149.4.0599.

845 McLennan, S. M., 1993, Weathering and Global Denudation: *Journal of Geology*, v.
846 101, no. 2, p. 295–303, doi:10.1086/648222.

847 Morford, J. L., W. R. Martin, R. François, and C. M. Carney, 2009, A model for
848 uranium, rhenium, and molybdenum diagenesis in marine sediments based on
849 results from coastal locations: *Geochimica et Cosmochimica Acta*, v. 73, no. 10, p.
850 2938–2960, doi:10.1016/j.gca.2009.02.029.

851 Murray, R. W., and M. Leinen, 1993, Chemical transport to the seafloor of the
852 equatorial Pacific Ocean across a latitudinal transect at 135 W: tracking
853 sedimentary major, trace, and rare earth element fluxes at the Equator and the
854 Intertropical Convergence Zone: *Geochimica et Cosmochimica Acta*, v. 57, no. 17,
855 p. 4141–4163, doi:10.1016/0016-7037(93)90312-K.

856 Taylor, S.R., and McLennan, S.M., 1985, The continental crust: its composition and
857 evolution.

858 Nesbitt, H. W., and G. M. Young, 1982, Early Proterozoic climates and plate motions
859 inferred from major element chemistry of lutites: *Nature*, v. 299, no. 5885, p. 715–
860 717, doi:10.1038/299715a0.

861 Och, L. M., G. A. Shields-Zhou, S. W. Poulton, C. Manning, M. F. Thirlwall, D. Li, X.
862 Chen, H. Ling, T. Osborn, and L. Cremonese, 2013, Redox changes in Early
863 Cambrian black shales at Xiaotan section, Yunnan Province, South China:
864 *Precambrian Research*, v. 225, p. 166–189, doi:10.1016/j.precamres.2011.10.005.

865 Ocubalidet, S. G., S. M. Rimmer, and J. A. Conder, 2018, Redox conditions
866 associated with organic carbon accumulation in the Late Devonian New Albany
867 Shale, west-central Kentucky, Illinois Basin: *International Journal of Coal*
868 *Geology*, v. 190, p. 42–55, doi:10.1016/j.coal.2017.11.017.

869 Pi, D., C. Liu, G. Zhou, and S. Jiang, 2013, Trace and rare earth element geochemistry
870 of black shale and kerogen in the early Cambrian Niutitang Formation in Guizhou
871 province, South China: Constraints for redox environments and origin of metal
872 enrichments: *Precambrian Research*, v. 225, no. 2013, p. 218–229,
873 doi:10.1016/j.precamres.2011.07.004.

874 Piper, D. Z., and R. B. Perkins, 2004, A modern vs. Permian black shale—the
875 hydrography, primary productivity, and water-column chemistry of deposition:
876 *Chemical Geology*, v. 206, no. 3, p. 177–197,
877 doi:10.1016/j.chemgeo.2003.12.006.

878 Reysenbach, A. L., and S. L. Cady, 2001, Microbiology of ancient and modern
879 hydrothermal systems: *Trends in Microbiology*, v. 9, no. 2, p. 79–86,
880 doi:10.1016/S0966-842X(00)01921-1.

881 Rowe, H. D., R. G. Loucks, S. C. Ruppel, and S. M. Rimmer, 2008, Mississippian
882 Barnett Formation, Fort Worth Basin, Texas: Bulk geochemical inferences and
883 Mo–TOC constraints on the severity of hydrographic restriction: *Chemical*
884 *Geology*, v. 257, no. 1, p. 16–25, doi:10.1016/j.chemgeo.2008.08.006.

885 Schoepfer, S. D., J. Shen, H. Wei, R. V. Tyson, E. Ingall, and T. J. Algeo, 2015, Total
886 organic carbon, organic phosphorus, and biogenic barium fluxes as proxies for
887 paleomarine productivity: *Earth-Science Reviews*, v. 149, p. 23–52,
888 doi:10.1016/j.earscirev.2014.08.017.

889 Scott, C., and T. W. Lyons, 2012, Contrasting molybdenum cycling and isotopic
890 properties in euxinic versus non-euxinic sediments and sedimentary rocks:
891 Refining the paleoproxies: *Chemical Geology*, v. 324–325, p. 19–27,
892 doi:10.1016/j.chemgeo.2012.05.012.

893 Simoneit, B. R. T., 2004, Prebiotic organic synthesis under hydrothermal conditions:
894 an overview: *Advances in Space Research*, v. 33, no. 1, p. 88–94,
895 doi:10.1016/j.asr.2003.05.006.

896 Steiner, M., E. Wallis, B. D. Erdtmann, Y. Zhao, and R. Yang, 2001,
897 Submarine-hydrothermal exhalative ore layers in black shales from South China
898 and associated fossils—insights into a Lower Cambrian facies and bio-evolution:
899 *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 169, no. 3, p. 165–191,
900 doi:10.1016/S0031-0182(01)00208-5.

901 Steiner, M., G. Li, Y. Qian, M. Zhu, and B. D. Erdtmann, 2007, Neoproterozoic to
902 Early Cambrian small shelly fossil assemblages and a revised biostratigraphic
903 correlation of the Yangtze Platform (China): *Palaeogeography, Palaeoclimatology,*
904 *Palaeoecology*, v. 254, no. 1, p. 67–99, doi:10.1016/j.palaeo.2007.03.046.

905 Sun, M., B. Yu, Q. Hu, R. Yang, Y. Zhang, B. Li, Y. B. Melnichenko, and G. Cheng,
906 2018, Pore structure characterization of organic-rich Niutitang shale from China:
907 Small angle neutron scattering (SANS) study: *International Journal of Coal*
908 *Geology*, v. 186, p. 115–125, doi:10.1016/j.coal.2017.12.006.

909 Tan, J., B. Horsfield, N. Mahlstedt, J. Zhang, R. di Primio, T. A. T. Vu, C. J. Boreham,
910 G. van Graas, and B. A. Tocher, 2013, Physical properties of petroleum formed
911 during maturation of Lower Cambrian shale in the upper Yangtze Platform, South
912 China, as inferred from PhaseKinetics modelling: *Marine and Petroleum Geology*,
913 v. 48, p. 47–56, doi:10.1016/j.marpetgeo.2013.07.013.

914 Tan, J., B. Horsfield, R. Fink, B. Krooss, H.-M. Schulz, E. Rybacki, J. Zhang, C. J.
915 Boreham, G. van Graas, and B. A. Tocher, 2014a, Shale gas potential of the major
916 marine shale formations in the Upper Yangtze Platform, South China, Part III:
917 Mineralogical, lithofacial, petrophysical, and rock mechanical properties: *Energy*
918 *& Fuels*, v. 28, no. 4, p. 2322–2342, doi:10.1021/ef4022703.

919 Tan, J., P. Weniger, B. Krooss, A. Merkel, B. Horsfield, J. Zhang, C. J. Boreham, G. v.
920 Graas, and B. A. Tocher, 2014b, Shale gas potential of the major marine shale
921 formations in the Upper Yangtze Platform, South China, Part II: Methane sorption
922 capacity: *Fuel*, v. 129, p. 204–218, doi:10.1016/j.fuel.2014.03.064.

923 Taylor, S. R., 1964, Abundance of chemical elements in the continental crust: a new
924 table: *Geochimica et Cosmochimica Acta*, v. 28, no. 8, p. 1273–1285,
925 doi:10.1016/0016-7037(64)90129-2.

926 Taylor, S. R., and S. M. McLennan, 1985, The continental crust: Its composition and
927 evolution, an examination of the geochemical record preserved in sedimentary

rocks: *Journal of Geology*, v. 94, no. 4, p. 632–633,
doi:10.1016/0031-9201(86)90093-2.

Tian, H., L. Pan, T. Zhang, X. Xiao, Z. Meng, and B. Huang, 2015, Pore
characterization of organic-rich Lower Cambrian shales in Qiannan Depression of
Guizhou Province, Southwestern China: *Marine and Petroleum Geology*, v. 62, p.
28–43, doi:10.1016/j.marpetgeo.2015.01.004.

Tribouillard, N., T. J. Algeo, T. Lyons, and A. Riboulleau, 2006, Trace metals as
paleoredox and paleoproductivity proxies: An update: *Chemical Geology*, v. 232,
no. 1, p. 12–32, doi:10.1016/j.chemgeo.2006.02.012.

Tribouillard, N., T. J. Algeo, F. Baudin, and A. Riboulleau, 2012, Analysis of marine
environmental conditions based on molybdenum–uranium covariation—
Applications to Mesozoic paleoceanography: *Chemical Geology*, v. 324/325, p.
46–58, doi:10.1016/j.chemgeo.2011.09.009.

Uematsu, M., M. Toratani, M. Kajino, Y. Narita, Y. Senga, and T. Kimoto, 2004,
Enhancement of primary productivity in the western North Pacific caused by the
eruption of the Miyake-jima Volcano: *Geophysical research letters*, v. 31, no. 6,
doi:10.1029/2003GL018790.

Vosoughi Moradi, A., A. Sarı, and P. Akkaya, 2016, Geochemistry of the Miocene oil
shale (Hançili Formation) in the Çankırı-Çorum Basin, Central Turkey:
Implications for paleoclimate conditions, source–area weathering, provenance and
tectonic setting: *Sedimentary Geology*, v. 341, p. 289–303,
doi:10.1016/j.sedgeo.2016.05.002.

Wan, Y., S. Zhang, S. Tang, Z. Pan, and W. Wu, 2018, A comparative study of
characterization of lower Palaeozoic Niutitang shale in northwestern Hunan,
China: *Journal of Natural Gas Science and Engineering*, v. 53, p. 284–300,
doi:10.1016/j.jngse.2018.03.015.

Wang, D., H. Ling, S. Ulrich, X. Zhu, M. Zhu, and T. He, 2018, Coupling of ocean
redox and animal evolution during the Ediacaran-Cambrian transition: *Nature
Communications*, v. 9, no. 1, doi:10.1038/s41467-018-04980-5.

Wang, J., D. Chen, D. Yan, H. Wei, and L. Xiang, 2012, Evolution from an anoxic to
oxic deep ocean during the Ediacaran–Cambrian transition and implications for
bioradiation: *Chemical Geology*, v. 306–307, p. 129–138,
doi:10.1016/j.chemgeo.2012.03.005.

Wang, L., C. Liu, and H. Zhang, 2013, Tectonic and sedimentary settings of
evaporites in the Dengying Formation, South China Block: Implications for the
potential of potash formation: *Acta Geoscientica Sinica*, v. 34, no. 5, p. 585–593
doi:10.3975/cagsb.2013.05.09 (in Chinese with English abstract).

Wang, S., C. Zou, D. Dong, Y. Wang, X. Li, J. H. Huang, and Q. Guan, 2015, Multiple
controls on the paleoenvironment of the Early Cambrian marine black shales in
the Sichuan Basin, SW China: Geochemical and organic carbon isotopic evidence:
Marine and Petroleum Geology, v. 66, p. 660–672,
doi:10.1016/j.marpetgeo.2015.07.009.

Wang, X., N. J. Planavsky, C. T. Reinhard, H. Zou, J. J. Ague, Y. Wu, B. C. Gill, E. M.

971 Schwarzenbach, and B. Peucker-Ehrenbrink, 2016, Chromium isotope
 972 fractionation during subduction-related metamorphism, black shale weathering,
 973 and hydrothermal alteration: *Chemical Geology*, v. 423, p. 19–33,
 974 doi:10.1016/j.chemgeo.2016.01.003.

975 Wang, Z., J. Tan, R. Boyle, J. Hilton, Z. Ma, W. Wang, Q. Lyu, X. Kang, and W. Luo,
 976 2020, Episodic hydrothermal activity in South China during the early Cambrian:
 977 Implications for biotic evolution: *Marine and Petroleum Geology*, v. 117, p.
 978 104355, doi: 10.1016/j.marpetgeo.2020.104355.

979 Wedepohl, K. H., 1971, Environmental influences on the chemical composition of
 980 shales and clays: *Physics & Chemistry of the Earth*, v. 8, no. 71, p. 307–333,
 981 doi:10.1016/0079-1946(71)90020-6.

982 Wu, C., J. Tuo, M. Zhang, L. Sun, Y. Qian, and Y. Liu, 2016, Sedimentary and
 983 residual gas geochemical characteristics of the Lower Cambrian organic-rich
 984 shales in Southeastern Chongqing, China: *Marine and Petroleum Geology*, v. 75, p.
 985 140–150, doi:10.1016/j.marpetgeo.2016.04.013.

986 Wu, C., J. Tuo, L. Zhang, M. Zhang, J. Li, Y. Liu, and Y. Qian, 2017, Pore
 987 characteristics differences between clay-rich and clay-poor shales of the Lower
 988 Cambrian Niutitang Formation in the Northern Guizhou area, and insights into
 989 shale gas storage mechanisms: *International Journal of Coal Geology*, v. 178, p.
 990 13–25, doi:10.1016/j.coal.2017.04.009.

991 Xiang, L., S. D. Schoepfer, S. Shen, C. Cao, and H. Zhang, 2017, Evolution of
 992 oceanic molybdenum and uranium reservoir size around the Ediacaran–Cambrian
 993 transition: Evidence from western Zhejiang, South China: *Earth and Planetary
 994 Science Letters*, v. 464, p. 84–94, doi:10.1016/j.epsl.2017.02.012.

995 Xiao, Z., J. Tan, Y. Ju, J. Hilton, R. Yang, P. Zhou, Y. Huang, B. Ning, and J. Liu,
 996 2018, Natural gas potential of Carboniferous and Permian transitional shales in
 997 central Hunan, South China: *Journal of Natural Gas Science and Engineering*, v.
 998 55, p. 520–533, doi:10.1016/j.jngse.2018.05.024.

999 Xu, L., B. Lehmann, J. Mao, W. Qu, and A. Du, 2011, Re-Os age of polymetallic
 1000 Ni-Mo-PGE-Au mineralization in Early Cambrian black shales of South China—a
 1001 reassessment: *Economic Geology*, v. 106, no. 3, p. 511–522,
 1002 doi:10.2113/econgeo.106.3.511.

1003 Xu, L., B. Lehmann, J. Mao, T. F. Nägler, N. Neubert, M. E. Böttcher, and P. Escher,
 1004 2012, Mo isotope and trace element patterns of Lower Cambrian black shales in
 1005 South China: Multi-proxy constraints on the paleoenvironment: *Chemical
 1006 Geology*, v. 318–319, p. 45–59, doi:10.1016/j.chemgeo.2012.05.016.

1007 Yamamoto, K., 1987, Geochemical characteristics and depositional environments of
 1008 cherts and associated rocks in the Franciscan and Shimanto Terranes: *Sedimentary
 1009 Geology*, v. 52, no. 1, p. 65–108, doi:10.1016/0037-0738(87)90017-0.

1010 Yan, D., D. Chen, Q. Wang, and J. Wang, 2012, Predominance of stratified anoxic
 1011 Yangtze Sea interrupted by short-term oxygenation during the Ordo-Silurian
 1012 transition: *Chemical Geology*, v. 291, p. 69–78,
 1013 doi:10.1016/j.chemgeo.2011.09.015.

1014 Yan, J., Y. Men, Y. Sun, Q. Yu, W. Liu, H. Zhang, J. Liu, J. Kang, S. Zhang, H. Bai,
1015 and X. Zheng, 2016, Geochemical and geological characteristics of the Lower
1016 Cambrian shales in the middle–upper Yangtze area of South China and their
1017 implication for the shale gas exploration: *Marine and Petroleum Geology*, v. 70, p.
1018 1–13, doi:10.1016/j.marpetgeo.2015.11.010.

1019 Yang, A., M. Zhu, J. Zhang, and G. Li, 2003, Early Cambrian eodiscoid trilobites of
1020 the Yangtze Platform and their stratigraphic implications: *Progress in Natural
1021 Science*, v. 13, no. 11, p. 861–866, doi:10.1080/10020070312331344560

1022 Yang, C., M. Zhu, D. J. Condon, and X. Li, 2017, Geochronological constraints on
1023 stratigraphic correlation and oceanic oxygenation in Ediacaran–Cambrian
1024 transition in South China: *Journal of Asian Earth Sciences*, v. 140, p. 75–81,
1025 doi:10.1016/j.jseaes.2017.03.017.

1026 Yang, C., X. Li, M. Zhu, D. J. Condon, and J. Chen, 2018, Geochronological
1027 constraint on the Cambrian Chengjiang biota, South China: *Journal of the
1028 Geological Society*, v. 175, no. 4, p. 659–666, doi:10.1144/jgs2017-103

1029 Yang, R., H. Wei, M. Bao, W. Wang, Q. Wang, X. Zhang, and L. Liu, 2008, Discovery
1030 of hydrothermal venting community at the base of Cambrian barite in Guizhou
1031 Province, Western China: Implication for the Cambrian biological explosion:
1032 *Progress in Natural Science*, v. 18, no. 1, p. 65–70,
1033 doi:10.1016/j.pnsc.2007.07.006.

1034 Yang, X., D. Yan, X. Wei, Z. Liwei, B. Zhang, H. Xu, Y. Gong, and J. He, 2018,
1035 Different formation mechanism of quartz in siliceous and argillaceous shales: A
1036 case study of Longmaxi Formation in South China: *Marine and Petroleum
1037 Geology*, v. 94, p. 80–94, doi:10.1016/j.marpetgeo.2018.03.036.

1038 Yeasmin, R., D. Chen, Y. Fu, J. Wang, Z. Guo, and C. Guo, 2017, Climatic-oceanic
1039 forcing on the organic accumulation across the shelf during the Early Cambrian
1040 (Age 2 through 3) in the mid-upper Yangtze Block, NE Guizhou, South China:
1041 *Journal of Asian Earth Sciences*, v. 134, p. 365–386,
1042 doi:10.1016/j.jseaes.2016.08.019.

1043 Yu, B., J. Li, Q. Zeng, M. Sun, and M. Shi, 2017, *Depositional Environment and
1044 Diagenesis of Organic Shale*: East China University of Science and Technology
1045 Press: Shanghai, 234 p.

1046 Zhai, L., C. Wu, Y. Ye, S. Zhang, and Y. Wang, 2018, Fluctuations in chemical
1047 weathering on the Yangtze Block during the Ediacaran–Cambrian transition:
1048 Implications for paleoclimatic conditions and the marine carbon cycle:
1049 *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 490, p. 280–292,
1050 doi:10.1016/j.palaeo.2017.11.006.

1051 Zhang, J., Z. Huo, X. Tan, Y. Liu, and S. Han, 2017, *Shale Gas Geology in China*:
1052 East China University of Science and Technology Press: Shanghai, 263 p.

1053 Zhao, J., Z. Jin, Z. Jin, Y. Geng, X. Wen, and C. Yan, 2016, Applying sedimentary
1054 geochemical proxies for paleoenvironment interpretation of organic-rich shale
1055 deposition in the Sichuan Basin, China: *International Journal of Coal Geology*, v.
1056 163, no. 2016, p. 52–71, doi:10.1016/j.coal.2016.06.015.

1057 Zhao, J., Z. Jin, Z. Jin, X. Wen, Y. Geng, C. Yan, and H. Nie, 2017, Depositional
1058 environment of shale in Wufeng and Longmaxi Formations, Sichuan Basin:
1059 Petroleum Research, v. 2, no. 3, p. 209–221, doi:10.1016/j.ptlrs.2017.04.003.
1060 Zheng, Y., R. F. Anderson, A. V. Geen, and J. Kuwabara, 2000, Authigenic
1061 molybdenum formation in marine sediments: a link to pore water sulfide in the
1062 Santa Barbara Basin: *Geochimica et Cosmochimica Acta*, v. 64, no. 24, p. 4165–
1063 4178, doi:10.1016/S0016-7037(00)00495-6.

1064

1065 **AUTHOR VITA**

1066 JINGQIANG TAN ~ Key Laboratory of Metallogenic Prediction of Nonferrous

1067 Metals and Geological Environment Monitoring, Ministry of Education, School of

1068 Geosciences and Info-physics, Central South University, Changsha, 410083, China;

1069 tanjingqiang@csu.edu.cn

1070 Jingqiang Tan is Professor of Petroleum Geology and Head of Department of

1071 Geological Resources at the Central South University, China. He received his Ph.D.

1072 degree from the Technical University of Berlin in Germany in 2014. His primary

1073 research interests are geological and geochemical analysis of shale gas and oil

1074 systems, and characterization and stimulation of unconventional reservoirs.

1075

1076 ZHANGHU WANG ~ Key Laboratory of Metallogenic Prediction of Nonferrous

1077 Metals and Geological Environment Monitoring, Ministry of Education, School of

1078 Geosciences and Info-physics, Central South University, Changsha, 410083, China;

1079 zhanghu.wang@csu.edu.cn

1080 Zhanghu Wang is currently a Ph.D. student at Central South University. He gained his

1081 M.S. from Central South University in 2017. His research interests include sequence

1082 stratigraphy, depositional systems, and reservoir characterization. His present research
1083 includes the influence of hydrothermal activity on paleoenvironments, and
1084 unconventional shale reservoirs.

1085

1086 WENHUI WANG ~ Key Laboratory of Metallogenic Prediction of Nonferrous Metals
1087 and Geological Environment Monitoring, Ministry of Education, School of
1088 Geosciences and Info-physics, Central South University, Changsha, 410083, China;
1089 whwang@csu.edu.cn

1090 Wenghui Wang is Associate Professor of Paleontology and Stratigraphy at Central
1091 South University, China. She received her Ph.D. degree from Nanjing University in
1092 2013 where she subsequently worked as an assistant researcher until the end of 2016.
1093 Her research focusses on Palaeozoic planktonic palaeontology, palynology and
1094 stratigraphy. She is currently investigating graptolitic biostratigraphy from south
1095 China near the Cambrian/Ordovician and Ordovician/Silurian boundaries.

1096

1097 JASON HILTON ~ School of Geography, Earth and Environmental Science,
1098 University of Birmingham, Birmingham, B152TT, UK; J.M.Hilton@bham.ac.uk
1099 Jason Hilton is a Reader in Palaeobiology and Palaeoenvironments at the University
1100 of Birmingham. He received his BSc from Sheffield University in 1992, and his Ph.D.
1101 from Cardiff University in 1996. His wide-ranging, multidisciplinary research
1102 focusses on palaeobotany, sedimentology, stratigraphy and palaeoenvironments. Since
1103 2003, he has worked extensively on resource-bearing sedimentary successions in

1104 China.

1105

1106 JIANHUA GUO ~ Key Laboratory of Metallogenic Prediction of Nonferrous Metals

1107 and Geological Environment Monitoring, Ministry of Education, School of

1108 Geosciences and Info-physics, Central South University, Changsha, 410083, China;

1109 gjh796@csu.edu.cn

1110 Jianhua Guo is a Professor of Petroleum Geology at Central South University, China.

1111 He has more than 30 years experience working in the petroleum industry. His primary

1112 research interests are sedimentology and basin analysis.

1113

1114 XIKAI WANG ~ Key Laboratory of Metallogenic Prediction of Nonferrous Metals

1115 and Geological Environment Monitoring, Ministry of Education, School of

1116 Geosciences and Info-physics, Central South University, Changsha, 410083, China;

1117 852303596@qq.com

1118 Xikai Wang obtained his Masters' degree in Petroleum Geology at the Central South

1119 University in China in 2019. He is currently a Ph.D. candidate at the University of

1120 North Carolina at Chapel Hill in the United States. His research focuses on shale gas

1121 geology and basin analysis.

1123

1124 **Figure 1.** (A) Paleogeographic map of the Yangtze Platform in the early Cambrian,
1125 modified after Wang et al. (2012). Selected sections include: 1 - Xiaotan, 2 - Songtao,
1126 3 - Longbizui, 4 – Ganziping; (B) Early Cambrian paleogeographic map showing
1127 lithofacies distribution in the Xuefeng Uplift area and sampling sites; (C)
1128 Stratigraphic column of the Niutitang Formation of the X1 well and sampling depths.

1129

Figure 2. Stratigraphic correlation with biostratigraphic and tuff/ore dating of the early Cambrian (ca. 541-514 Ma) sections across South China. Intervals I, II, and III are mainly modified after Jin et al. (2016). Data source: 1 – Xiaotan (Yang et al., 2003; Och et al., 2013); 2 – Songtao (Yang et al., 2003; Goldberg et al., 2007); 3 – Longbizui (Wang et al., 2012; Yang et al., 2017); 4 –Ganziping (Chen et al., 2009); 5 – X1 (this study). U–Pb ages of the Xiaotan and Songtao sections are from adjacent Meishucun (Compston et al., 2008), Maotianshan (Yang et al., 2018a), and Bahuang (Chen et al., 2015), respectively. Abbreviations: CLP = Canglangpu Formation, DY = Dengying Formation, LCP = Liuchapo Formation, CJB = Chengjiang Biota.

1141
 1142 **Figure 3.** Lithologies of samples from X1 well. (A) Calcite patches in dark gray
 1143 calcareous shale, 869.4m (2852.4 ft); (B) Calcite patches in dark gray calcareous
 1144 shale, 864.3m (2835.6 ft); (C) Black siliceous shale, 850.1m (2789.1 ft); (D) Black
 1145 shale with intense graphite reflectance, 848.2m (2782.8 ft); (E) Pyrite bands in black
 1146 siliceous shale, 840.8m (2758.5 ft); (F) Black siliceous shale interbedded with
 1147 gray-black mudstone, 813.4m (2668.6 ft).
 1148

Figure 4. Ternary diagram of minerals from the Niutitang shale Niu1, Niu2 and Niu3 members and the underlying Liuchapo chert (LCP).

Figure 5. Stratigraphic distribution of TOC contents (wt.%), CIA, redox proxies (Mo,

1156 Mo_{EF}/U_{EF}), seawater restriction, and inferred sea-level changes. LCP = Liuchapo

1157 Formation.

1158

1159

1160 **Figure 6.** Stratigraphic distribution of TOC contents (wt.%), excess SiO_2 contents

1161 (wt.%), hydrothermal proxies, and paleoproductivity proxies (Ba_{bio} , Cu/Al, Ni/Al).

1162

1163

1164 **Figure 7.** Mo_{EF} vs. U_{EF} Plot. The solid line represents Mo/U ratios of sea water. The
 1165 dash lines represent multiples ($0.3\times$, $1\times$, and $3\times$) of Mo/U ratios of modern seawater.
 1166 The gray area represents the unrestricted marine trend and the green area represents
 1167 the particulate shuttle trend. Modified after Algeo and Tribovillard (2009) and
 1168 Tribovillard et al. (2012).

1169

1170 **Figure 8.** TOC contents vs. Mo/Al (A) and V/Al (B).
 1171

1172

1173 **Figure 9.** Al_2O_3 contents vs. SiO_2 contents (A) and TOC contents vs. TiO_2 contents

1174 (B).

1175

1176

1177 **Figure 10.** Al-Fe-Mn ternary diagram showing Liuchapo cherts (LCP) are from

1178 intense hydrothermal source, excess silica of the Niu1 and Niu2 shales are from

1179 hydrothermal and biogenic sources but that of the Niu3 shale is primarily biogenic

1180 origin. Position of hydrothermal and biogenic chert origins from Adachi et al. (1986)

1181 and Yamamoto (1987).

1182

1183
 1184 **Figure 11.** Excess SiO₂ contents vs paleoclimate proxies (A), excess SiO₂ contents vs.
 1185 hydrothermal proxies (B), TOC contents vs. excess SiO₂ of Niu3 samples (C), and
 1186 TOC contents vs. excess SiO₂ of Niu1 and Niu2 samples (D).
 1187

1188

1189 **Figure 12.** Correlations among redox metal elements. U/Al vs. Mo/Al ratios (A) and

1190 V/Al vs. Mo/Al ratios (B) in Niu3 samples; U/Al vs. Mo/Al ratios (C) and V/Al vs.

1191 Mo/Al ratios (D) in Niu1 and Niu2 samples; U concentration vs. Mo concentration (E)

1192 and V concentration vs. Mo concentration (F) in particles from the Mid-Atlantic

1193 Ridge hydrothermal plume.

1194

1195

1196 **Figure 13.** Mo concentrations vs. TOC contents. Dashed lines represent four modern
1197 anoxic basin systems in Saanich Inlet, Cariaco Basin, Framvaren Fjord, and Black sea
1198 (Algeo and Lyons, 2006; Algeo et al., 2007), showing different seawater restriction
1199 environments.

1200

Figure 14. Depositional models and inferred paleoclimates of the Lower Cambrian sedimentary rocks in the eastern Xuefeng Uplift through the Niutitang Formation.

1208

Table 1. Mineral compositions of the analyzed samples.

Sample	Formation	Member	Lithology	Mineral Composition (vol.%)								
				Quartz	Feldspar	Calcite	Dolomite	Pyrite	Barite	Anhydrite	Glauberite	Clay
A28	Niutitang	Niu3	Shale	59.6	2.0	0.0	0.0	7.5	0.0	0.9	0.0	30.0
A27	Niutitang	Niu3	Shale	58.3	1.3	0.9	0.0	5.9	0.0	2.5	0.0	31.1
A25	Niutitang	Niu3	Shale	62.1	3.0	2.8	0.0	5.1	0.0	0.9	0.0	26.1
A23	Niutitang	Niu3	Shale	60.9	3.0	0.0	0.0	7.5	0.0	1.2	0.0	27.4
A20	Niutitang	Niu3	Shale	71.7	0.0	5.9	0.0	6.0	0.0	0.0	0.0	16.4
A18	Niutitang	Niu2	Siliceous shale	81.8	0.0	0.5	0.0	5.2	0.0	0.0	0.0	12.5
A16	Niutitang	Niu2	Shale	75.7	0.0	10.3	0.0	7.2	0.0	0.0	0.0	6.8
A14	Niutitang	Niu2	Shale	68.1	5.1	1.4	4.9	6.8	0.0	0.0	0.0	13.7
A12	Niutitang	Niu2	Siliceous shale	79.8	4.6	1.9	0.7	4.6	0.0	0.0	0.0	8.4
A10	Niutitang	Niu2	Siliceous shale	78.5	3.3	2.1	0.0	7.7	0.0	0.0	0.0	8.4
A08	Niutitang	Niu2	Shale	49.3	1.1	3.8	0.0	9.5	2.7	0.0	29.0	4.6
A06	Niutitang	Niu1	Dolomite	28.2	3.3	1.3	56.3	5.5	0.0	0.0	0.0	5.4
A05	Niutitang	Niu1	Calcareous shale	28.2	16.3	10.1	12.4	12.2	0.0	0.0	3.8	17.0
A03	Niutitang	Niu1	Siliceous shale	68.2	2.4	7.2	0.0	1.9	0.0	0.0	0.0	20.3
A01	Niutitang	Niu1	Calcareous shale	49.1	0.0	0.0	30.5	5.6	0.0	0.0	0.0	14.8
B03	Liuchapo		Chert	94.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.6
B01	Liuchapo		Chert	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

1209

1210

1211

1212 **Table 2.** TOC contents and major elements of the analyzed samples.

Sample	Formation	Member	Lithology	Depth/m	TOC (wt.%)	Major element (wt.%)										
						SiO ₂	Al ₂ O ₃	FeO	Fe ₂ O ₃	TiO ₂	Na ₂ O	MgO	K ₂ O	CaO	P ₂ O ₅	MnO
A28	Niutitang	Niu3	Shale	730.0	2.61	72.92	10.81	0.51	3.40	0.41	0.11	1.43	3.22	0.35	0.05	0.01
A27	Niutitang	Niu3	Shale	737.3	4.59	67.25	11.27	0.41	4.28	0.50	0.31	1.29	2.70	0.79	0.07	0.01
A26	Niutitang	Niu3	Shale	740.6	5.83	74.22	9.49	0.26	2.66	0.41	0.20	1.10	2.70	0.41	0.06	0.01
A25	Niutitang	Niu3	Shale	746.9	4.99	69.68	9.75	0.21	2.88	0.43	0.36	1.07	2.54	2.00	0.07	0.02
A24	Niutitang	Niu3	Shale	756.5	3.70	69.50	11.80	0.36	2.88	0.52	0.15	1.40	3.38	0.89	0.08	0.01
A23	Niutitang	Niu3	Shale	761.5	4.02	71.70	10.87	0.31	2.71	0.43	0.08	1.34	2.74	0.44	0.06	0.01
A22	Niutitang	Niu3	Shale	772.0	4.20	65.62	10.15	0.31	4.13	0.47	0.18	2.10	2.83	2.73	0.12	0.04
A21	Niutitang	Niu3	Shale	775.7	7.99	69.03	8.94	0.26	3.18	0.39	0.06	1.20	2.02	0.60	0.09	0.01
A20	Niutitang	Niu3	Shale	783.4	8.05	72.24	6.22	0.26	2.54	0.29	0.06	1.13	1.48	2.50	0.08	0.02
A19	Niutitang	Niu2	Shale	790.6	11.10	67.74	6.25	0.46	2.84	0.29	0.03	1.11	1.43	1.42	0.08	0.01
A18	Niutitang	Niu2	Siliceous shale	796.8	11.40	75.60	4.66	0.26	2.90	0.23	0.06	0.85	1.25	0.46	0.06	0.01
A17	Niutitang	Niu2	Shale	798.6	11.98	73.60	4.14	0.21	2.67	0.20	0.01	0.76	1.11	2.35	0.82	0.03
A16	Niutitang	Niu2	Shale	804.7	8.05	58.23	3.12	0.31	2.85	0.21	0.09	4.17	0.28	3.00	0.34	0.02
A15	Niutitang	Niu2	Siliceous shale	810.3	13.88	80.52	0.82	0.36	0.42	0.06	0.07	0.41	0.20	1.50	0.03	0.01
A14	Niutitang	Niu2	Shale	815.3	11.78	70.88	6.18	0.36	3.03	0.34	0.51	0.90	1.66	1.28	0.21	0.01
A13	Niutitang	Niu2	Shale	820.1	18.63	73.34	1.79	0.26	0.40	0.09	0.03	1.02	0.28	1.06	0.02	0.02
A12	Niutitang	Niu2	Siliceous shale	824.3	11.30	74.63	4.40	0.46	1.75	0.28	0.15	0.78	1.35	0.74	0.09	0.01
A11	Niutitang	Niu2	Siliceous shale	828.7	12.30	75.59	2.91	0.36	1.17	0.24	0.14	1.00	0.75	1.81	0.24	0.02
A10	Niutitang	Niu2	Siliceous shale	833.6	12.63	76.81	2.89	0.26	2.93	0.22	0.39	1.29	0.95	0.56	0.16	0.01
A09	Niutitang	Niu2	Shale	840.2	10.60	60.29	4.90	0.46	1.69	0.27	0.08	1.58	0.56	0.56	0.08	0.01
A08	Niutitang	Niu2	Shale	848.2	20.10	57.07	4.46	0.21	2.02	0.22	0.14	0.85	0.67	4.62	2.87	0.01
A07	Niutitang	Niu2	Siliceous shale	851.1	11.15	78.07	2.97	0.26	1.03	0.12	0.08	0.60	0.17	0.34	0.04	0.01

A06	Niutitang	Niu1	Dolomite	859.0	0.18	8.29	1.52	0.21	0.85	0.09	0.02	17.70	0.55	25.89	0.05	0.02
A05	Niutitang	Niu1	Calcareous shale	861.2	4.03	49.72	16.86	0.31	2.75	0.65	0.18	1.86	6.01	1.38	0.06	0.01
A04	Niutitang	Niu1	Calcareous shale	864.0	1.80	62.23	3.76	0.31	1.83	0.21	0.06	5.65	1.63	7.35	0.03	0.02
A03	Niutitang	Niu1	Siliceous shale	866.2	1.96	74.21	7.76	0.41	1.48	0.41	0.07	1.52	2.21	3.17	0.04	0.03
A02	Niutitang	Niu1	Calcareous shale	869.3	0.87	30.07	4.17	0.41	3.38	0.30	0.01	10.10	1.35	17.35	2.03	0.24
A01	Niutitang	Niu1	Calcareous shale	872.0	0.11	52.99	6.97	0.36	3.68	0.42	0.01	7.60	2.12	7.95	0.1	0.12
B04	Liuchapo		Siliceous shale	873.2	0.16	70.32	9.34	0.31	3.49	0.34	0.03	3.54	2.99	2.00	0.01	0.00
B03	Liuchapo		Chert	876.1	0.28	95.92	0.41	0.21	0.51	0.03	0.10	0.21	0.13	0.15	0.01	0.01
B02	Liuchapo		Chert	881.9	0.30	98.59	0.19	0.26	0.32	0.04	0.18	0.27	0.15	0.11	0.01	0.01
B01	Liuchapo		Chert	885.3	0.35	95.43	0.11	0.31	0.29	0.02	0.01	0.26	0.07	0.22	0.13	0.01

1213

1214

1215

1216

1217 **Table 3.** Trace element composition of the analyzed samples.

Sample	Formation	Member	Lithology	Depth/m	Trace element (ppm)									
					V	Cr	Co	Ni	Cu	Sr	Mo	Th	U	Ba
A28	Niutitang	Niu3	Shale	730.0	190	57	13	70	53	25	24	9	11	1304
A27	Niutitang	Niu3	Shale	737.3	127	59	13	76	63	54	31	10	14	532
A26	Niutitang	Niu3	Shale	740.6	103	47	11	61	56	25	28	7	8	1450
A25	Niutitang	Niu3	Shale	746.9	118	45	12	53	49	76	37	9	12	1456
A24	Niutitang	Niu3	Shale	756.5	348	61	13	91	44	40	45	7	13	1902
A23	Niutitang	Niu3	Shale	761.5	308	56	12	91	46	29	41	6	10	1732
A22	Niutitang	Niu3	Shale	772.0	257	65	11	95	57	107	31	8	14	911
A21	Niutitang	Niu3	Shale	775.7	292	60	12	119	55	31	76	6	46	1058
A20	Niutitang	Niu3	Shale	783.4	145	38	9	85	44	82	50	4	20	1588
A19	Niutitang	Niu2	Shale	790.6	308	46	10	132	45	42	78	5	32	494
A18	Niutitang	Niu2	Siliceous shale	796.8	2510	80	8	221	272	23	145	3	40	720
A17	Niutitang	Niu2	Shale	798.6	1449	60	8	206	301	82	118	3	92	824
A16	Niutitang	Niu2	Shale	804.7	151	35	10	100	60	166	209	3	207	257
A15	Niutitang	Niu2	Siliceous shale	810.3	227	11	1	43	38	90	18	1	89	2160
A14	Niutitang	Niu2	Shale	815.3	3364	133	9	186	169	69	128	2	78	482
A13	Niutitang	Niu2	Shale	820.1	908	32	2	63	107	64	22	1	39	3352
A12	Niutitang	Niu2	Siliceous shale	824.3	1033	52	8	125	39	74	56	2	66	1426
A11	Niutitang	Niu2	Siliceous shale	828.7	2978	102	5	124	131	160	71	1	82	3387
A10	Niutitang	Niu2	Siliceous shale	833.6	746	41	10	247	201	56	115	1	73	401
A09	Niutitang	Niu2	Shale	840.2	1273	62	10	197	49	39	62	2	74	673
A08	Niutitang	Niu2	Shale	848.2	4075	744	4	223	193	434	40	1	56	752
A07	Niutitang	Niu2	Siliceous shale	851.1	2426	181	2	96	191	44	29	1	30	8382

A06	Niutitang	Niu1	Dolomite	859.0	210	185	3	14	12	544	20	2	8	6939
A05	Niutitang	Niu1	Calcareous shale	861.2	936	258	7	43	32	45	24	5	38	2760
A04	Niutitang	Niu1	Calcareous shale	864.0	113	67	2	8	27	117	2	2	3	2196
A03	Niutitang	Niu1	Siliceous shale	866.2	102	84	4	11	19	51	2	5	4	1735
A02	Niutitang	Niu1	Calcareous shale	869.3	49	36	7	26	17	400	1	6	5	556
A01	Niutitang	Niu1	Calcareous shale	872.0	70	51	8	35	29	162	1	5	5	315
B04	Liuchapo		Siliceous shale	873.2	230	129	4	22	37	46	2	6	7	1035
B03	Liuchapo		Chert	876.1	16	15	1	4	9	5	1	1	1	324
B02	Liuchapo		Chert	881.9	17	14	1	5	4	5	1	1	1	382
B01	Liuchapo		Chert	885.3	41	21	1	15	225	8	2	1	2	287

1218

1219

1220

Table 4. Element ratios of the analyzed samples.

Sample	Formation	Member	Lithology	Depth/m	CIA	Al/(Al+Fe+Mn)	U _{EF}	Mo _{EF}	V _{EF}	Ni/Al	Cu/Al	Ba _{bio}	Excess-SiO ₂
A28	Niutitang	Niu3	Shale	730.00	74	0.67	6.03	42.46	2.21	12.16	9.19	875	34.85
A27	Niutitang	Niu3	Shale	737.30	74	0.64	7.30	52.46	1.42	12.80	10.58	85	27.56
A26	Niutitang	Niu3	Shale	740.60	73	0.71	5.44	55.73	1.37	12.04	11.05	1073	40.80
A25	Niutitang	Niu3	Shale	746.90	71	0.70	7.56	72.46	1.52	10.35	9.51	1069	35.35
A24	Niutitang	Niu3	Shale	756.50	74	0.73	6.56	71.55	3.71	14.52	7.06	1433	27.95
A23	Niutitang	Niu3	Shale	761.50	77	0.73	5.72	70.38	3.57	15.83	8.03	1300	33.42
A22	Niutitang	Niu3	Shale	772.00	73	0.63	8.34	56.95	3.19	17.75	10.51	508	29.88
A21	Niutitang	Niu3	Shale	775.70	79	0.66	31.28	161.00	4.11	25.14	11.58	703	37.55
A20	Niutitang	Niu3	Shale	783.40	78	0.62	19.59	150.63	2.94	25.72	13.48	1341	50.34
A19	Niutitang	Niu2	Shale	790.60	79	0.58	31.00	235.13	6.21	39.89	13.63	246	45.73
A18	Niutitang	Niu2	Siliceous shale	796.80	75	0.52	52.69	587.74	67.83	89.58	110.25	535	59.19
A17	Niutitang	Niu2	Shale	798.60	77	0.52	135.40	538.38	44.07	93.99	137.33	660	59.02
A16	Niutitang	Niu2	Shale	804.70	84	0.42	404.26	1265.31	6.09	60.36	36.32	133	47.24
A15	Niutitang	Niu2	Siliceous shale	810.30	65	0.43	657.62	414.63	34.86	98.82	87.07	2127	77.63
A14	Niutitang	Niu2	Shale	815.30	64	0.58	76.61	391.23	68.55	56.85	51.65	237	49.12
A13	Niutitang	Niu2	Shale	820.10	82	0.66	133.10	230.04	63.88	66.80	112.91	3281	67.04
A12	Niutitang	Niu2	Siliceous shale	824.30	69	0.59	91.40	217.22	29.56	53.66	16.61	1251	59.14
A11	Niutitang	Niu2	Siliceous shale	828.70	70	0.58	171.49	458.92	128.87	80.49	85.03	3271	65.34
A10	Niutitang	Niu2	Siliceous shale	833.60	63	0.40	153.28	751.63	32.51	161.44	131.37	286	66.63
A09	Niutitang	Niu2	Shale	840.20	85	0.63	91.65	237.46	32.72	75.94	19.00	478	43.03
A08	Niutitang	Niu2	Shale	848.20	82	0.60	76.78	168.56	115.06	61.31	81.74	575	41.36
A07	Niutitang	Niu2	Siliceous shale	851.10	87	0.63	60.52	185.07	102.86	16.90	121.47	8264	67.61
A06	Niutitang	Niu1	Dolomite	859.00	70	0.51	30.67	244.81	17.40	16.90	15.29	6879	2.94

A05	Niutitang	Niu1	Calcareous shale	861.20	70	0.80	13.66	27.34	6.99	4.85	3.54	2091	0.00
A04	Niutitang	Niu1	Calcareous shale	864.00	66	0.56	4.34	11.00	3.78	4.15	13.61	2047	48.99
A03	Niutitang	Niu1	Siliceous shale	866.20	75	0.75	2.77	3.70	1.66	2.58	4.67	1427	46.88
A02	Niutitang	Niu1	Calcareous shale	869.30	74	0.43	7.42	4.12	1.46	11.82	7.84	390	15.39
A01	Niutitang	Niu1	Calcareous shale	872.00	75	0.56	4.63	3.47	1.26	9.49	7.83	38	28.45
B04	Liuchapo		Siliceous shale	873.2	74	0.65	4.75	4.23	3.10	4.49	7.54	664	37.43
B03	Liuchapo		Chert	876.1	47	0.29	7.88	45.61	4.98	19.40	39.99	308	94.48
B02	Liuchapo		Chert	881.9	20	0.19	21.17	76.55	11.40	45.93	35.39	375	97.92
B01	Liuchapo		Chert	885.3	50	0.12	108.57	348.59	47.05	260.13	178.52	282	95.04

1222