

Accrual of atherosclerotic vascular events in a multicentre inception SLE cohort

Urowitz, Murray B.; Gladman, Dafna D.; Farewell, Vernon; Su, Jiandong; Romero-diaz, Juanita; Bae, Sang-cheol; Fortin, Paul R.; Sanchezguerrero, Jorge; Clarke, Ann Elaine; Bernatsky, Sasha; Gordon, Caroline; Hanly, John G.; Wallace, Daniel J.; Isenberg, David A.; Rahman, Anisur; Merrill, Joan T.; Ginzler, Ellen; Alarcón, Graciela S.; Chatham, W. Winn; Petri, Michelle A.

DOI:
[10.1002/art.41392](https://doi.org/10.1002/art.41392)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):

Urowitz, MB, Gladman, DD, Farewell, V, Su, J, Romero-diaz, J, Bae, S, Fortin, PR, Sanchezguerrero, J, Clarke, AE, Bernatsky, S, Gordon, C, Hanly, JG, Wallace, DJ, Isenberg, DA, Rahman, A, Merrill, JT, Ginzler, E, Alarcón, GS, Chatham, WW, Petri, MA, Bruce, IN, Khamashta, MA, Aranow, C, Dooley, MA, Manzi, S, Ramseygoldman, R, Nived, O, Jönsen, A, Steinsson, K, Zoma, AA, Ruiz-irastorza, G, Lim, SS, Kalunian, KC, nanç, M, Vollenhoven, R, Ramoscasals, M, Kamen, DL, Jacobsen, S, Peschken, CA, Askanase, A & Stoll, T 2020, 'Accrual of atherosclerotic vascular events in a multicentre inception SLE cohort', *Arthritis and Rheumatology*, vol. 72, no. 10, pp. 1734-1740. <https://doi.org/10.1002/art.41392>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Urowitz, M.B., Gladman, D.D., Farewell, V., Su, J., Romero-Diaz, J., Bae, S., Fortin, P.R., SanchezGuerrero, J., Clarke, A.E., Bernatsky, S., Gordon, C., Hanly, J.G., Wallace, D.J., Isenberg, D.A., Rahman, A., Merrill, J.T., Ginzler, E., Alarcón, G.S., Chatham, W.W., Petri, M.A., Bruce, I.N., Khamashta, M.A., Aranow, C., Dooley, M.A., Manzi, S., Ramsey Goldman, R., Nived, O., Jönsen, A., Steinsson, K., Zoma, A.A., Ruiz-Irastorza, G., Lim, S.S., Kalunian, K.C., nanç, M., van Vollenhoven, R., RamosCasals, M., Kamen, D.L., Jacobsen, S., Peschken, C.A., Askanase, A. and Stoll, T. (2020), Accrual of Atherosclerotic Vascular Events in a Multicenter Inception Systemic Lupus Erythematosus Cohort. *Arthritis Rheumatol*, 72: 1734-1740, which has been published in final form at <https://doi.org/10.1002/art.41392>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

ACCRUAL OF ATHEROSCLEROTIC VASCULAR EVENTS IN A MULTICENTRE INCEPTION SLE COHORT

Short Title: AVE in SLE

AUTHORS:

M B Urowitz, MD, FRCPC,¹ D D Gladman, MD, FRCPC,¹ V. Farewell, PhD,² J Su, MB, BSc¹ J Romero-Diaz, MD, MSc,³ S C Bae, MD, PhD,⁴ P R Fortin, MD, MPH,⁵ J Sanchez-Guerrero, MD, MSc,¹ C Gordon, MD,⁶ A E Clarke, MD, MSc,⁷ S Bernatsky, MD, PhD,⁸ J G Hanly, MD,⁹ D J Wallace, MD,¹⁰ D Isenberg, MD,¹¹ A Rahman, MD, PhD,¹¹ J Merrill, MD,¹² E Ginzler, MD, MPH,¹³ I N Bruce, MD,^{14,15} M Petri, MD, MPH,¹⁶ G S Alarcón, MD, MPH,¹⁷ W W Chatham, MD,¹⁷ C Aranow, MD,¹⁸ M Dooley, MD, MPH,¹⁹ R Ramsey-Goldman, MD, DrPH,²⁰ M Khamashta, MD,²¹ O. Nived, MD, PhD,²² A. Jönsen, MD, PhD,²² K Steinsson, MD,²³ S Manzi, MD, MPH,²⁴ A Zoma, MD,²⁵ G Ruiz-Irastorza, MD,²⁶ S Lim, MD,²⁷ M İnanç, MD,²⁸ R van Vollenhoven, MD,²⁹ M Ramos-Casals, MD,³⁰ D L Kamen, MD,³¹ K C Kalunian, MD,³² S Jacobsen, MD, DMSc,³³ C Peschken, MD, MSc,³⁴ A Askanase, MD, MPH,³⁵ T Stoll, MD³⁶

AUTHORS' AFFILIATION:

¹Centre for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital and University of Toronto, Toronto Ontario, Canada

²MRC Biostatistics Unit, Cambridge Institute of Public Health, University of Cambridge, Cambridge, UK

³Instituto Nacional de Ciencias Medicas y Nutrición, Mexico City, Mexico

⁴Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul, Korea

⁵Division of Rheumatology, Centre Hospitalier Universitaire de Québec et Université Laval, Quebec City, Canada

⁶ Rheumatology Research Group, Institute of Inflammation and Ageing, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

⁷Division of Rheumatology, Cumming School of Medicine University of Calgary, Calgary, Alberta, Canada

⁸Divisions of Clinical Immunology/Allergy and Clinical Epidemiology, Montreal General Hospital, McGill University Health Centre, Montreal, Quebec, Canada

⁹Division of Rheumatology, Department of Medicine and Department of Pathology, Queen Elizabeth II Health Sciences Centre and Dalhousie University, Halifax, Nova Scotia, Canada

¹⁰Cedars-Sinai/David Geffen School of Medicine at UCLA, Los Angeles, California, USA

¹¹Centre for Rheumatology Research, University College, London, UK

¹²Department of Clinical Pharmacology, Oklahoma Medical Research Foundation, Oklahoma City, Oklahoma, USA

¹³Department of Medicine, SUNY Downstate Medical Center, Brooklyn, New York, USA

¹⁴Arthritis Research UK Centre for Epidemiology, Centre for Musculoskeletal Research, Institute of Inflammation and Repair, Manchester Academic Health Sciences Centre, The University of Manchester, Manchester, UK

¹⁵NIHR Manchester Musculoskeletal Biomedical Research Unit, Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, London, UK

¹⁶Department of Rheumatology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

¹⁷Department of Medicine, Division of Clinical Immunology and Rheumatology, The University of Alabama at Birmingham, Birmingham, Alabama, USA

¹⁸Feinstein Institute for Medical Research, Manhasset, New York, USA

¹⁹Thurston Arthritis Research Center, Division of Rheumatology and Immunology, Department of Medicine, University of North Carolina, Chapel Hill, North Carolina, USA

²⁰Northwestern University and Feinberg School of Medicine, Chicago, Illinois, USA

²¹Lupus Research Unit, The Rayne Institute, St Thomas' Hospital, King's College London School of Medicine, London, UK

²² Department of Clinical Sciences Lund, Rheumatology, Lund University, Skåne University Hospital, Lund, Sweden

²³Department of Rheumatology, Center for Rheumatology Research, Fossvogur Landspítali University Hospital, Reykjavik, Iceland

²⁴Lupus Center of Excellence, Autoimmunity Institute Allegheny Health Network, Pittsburgh, Pennsylvania, USA

²⁵Lanarkshire Centre for Rheumatology, Hairmyres Hospital, East Kilbride, Scotland, UK

²⁶Autoimmune Disease Unit, Department of Internal Medicine, BioCruces Health Research Unit, Hospital Universitario Cruces, University of the Basque Country, Barakaldo, Spain

²⁷Division of Rheumatology, Emory University School of Medicine, Atlanta, Georgia, USA

²⁸Division of Rheumatology, Department of Internal Medicine, Istanbul Medical Faculty, Istanbul University, Istanbul, Turkey

²⁹Department of Rheumatology and Clinical Immunology, Amsterdam University Medical Centers, Amsterdam, Holland

³⁰Josep Font Autoimmune Diseases Laboratory, IDIBAPS, Department of Autoimmune Diseases, Hospital Clínic, Barcelona, Spain

³¹Division of Rheumatology, Medical University of South Carolina, Charleston, South Carolina, USA

³²UCSD School of Medicine, La Jolla, California, USA

³³Copenhagen Lupus and Vasculitis Clinic, 4242, Department of Rheumatology Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark

³⁴Department of Medicine and Community Health Sciences, University of Manitoba, Winnipeg, Manitoba, Canada

³⁵Hospital for Joint Diseases, NYU, Seligman Center for Advanced Therapeutics, New York, NY

³⁶Department of Rheumatology, Kantonsspital, Schaffhausen, Switzerland

ACKNOWLEDGMENTS OF FUNDING

Dr. John G. Hanly was supported by the Canadian Institutes of Health Research grant MOP-88526.

Dr. Sang-Cheol Bae's work was supported in part by NRF-2017M3A9B4050335, Republic of Korea.

Dr Caroline Gordon is supported by Lupus UK, Sandwell and West Birmingham Hospitals NHS Trust and the National Institute for Health Research (NIHR)/Wellcome Trust Birmingham Clinical Research Facility. The views expressed are those of the authors(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

The Hopkins Lupus Cohort is supported by the NIH (grant AR43727 and 69572).

The Montreal General Hospital Lupus Clinic is partially supported by the Singer Family Fund for Lupus Research.

Dr. Clarke holds The Arthritis Society Chair in Rheumatic Diseases at the University of Calgary.

Dr. Paul R. Fortin holds a tier 1 Canada Research Chair on Systemic Autoimmune Rheumatic Diseases at Université Laval.

Dr. Bruce is a National Institute for Health Research (NIHR) Senior Investigator and is supported by Arthritis Research UK, the NIHR Manchester Biomedical Centre and the NIHR/ Wellcome Trust Manchester Clinical Research Facility. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

Dr. Soren Jacobsen is supported by the Danish Rheumatism Association (A3865) and the Novo Nordisk Foundation (A05990).

Dr. Ramsey-Goldman's work was supported by the NIH (grants 5UL1TR001422-02, formerly 8UL1TR000150 and UL-1RR-025741, K24-AR-02318, and P60AR064464 formerly P60-AR-48098).

Dr. Mary Anne Dooley's work was supported by the NIH grant RR00046.

Dr. Ruiz-Irastorza is supported by the Department of Education, Universities and Research of the Basque Government.

Dr Isenberg and Dr Rahman are supported by and supported by the National Institute for Health Research University College London Hospitals Biomedical Research Center.

CORRESPONDING AUTHOR:

Murray B. Urowitz, MD, FRCPC

University of Toronto Lupus Clinic

Centre for Prognosis Studies in the Rheumatic Diseases

Toronto Western Hospital, 399 Bathurst St. 1E-410B, Toronto, Ontario, M5T 2S8, Canada

Tel: 416-603-5828; Fax 416-602-9387. Email: m.urowitz@utoronto.ca

Keywords: Lupus, SLE, atherosclerotic vascular events, incidence, risk factors

Word Count: 4,146

ABSTRACT

Background/Purpose: In previously published work, atherosclerotic vascular events (AVE) occurred in approximately 10% of patients with SLE. We aimed to investigate the annual occurrence and potential risk factors for AVE in a multinational, multiethnic inception cohort of patients with SLE.

Methods: A large 33-centre cohort of SLE patients was followed yearly between 1999-2017. AVEs were attributed to atherosclerosis on the basis of SLE being inactive at the time of the AVE, and typical atherosclerotic changes on imaging or pathology, and/or evidence of atherosclerosis elsewhere. Analysis included descriptive statistics, rate of AVE's per 1000 patient-years, and univariable and multivariable relative risk regression models.

Results: Of the 1848 patients enrolled in the cohort, 1710 had at least one follow up visit after enrolment, for a total of 13,666 patient-years. Of 1710, 3.6% had one or more AVEs attributed to atherosclerosis, for an event rate of 4.6 per 1000 patient-years. In multivariable analyses, lower AVE rates were associated with antimalarials (HR: 0.54[95% CI 0.32, 0.91]) while higher AVE rates were associated with any prior vascular event (VE) (HR: 4.00[1.55,10.30]) and body mass index (BMI) >40 (HR: 2.74[1.04,7.18]) A prior AVE increased the risk for subsequent AVE (HR 5.42[3.17,9.27], p<0.001).

Conclusion: The prevalence of AVE and rate of AVE accrual in this study is much lower than that seen in previously published data. This may be related to better control of both the disease activity and classic risk factors.

INTRODUCTION

Atherosclerotic disease has been recognized as a major cause of morbidity and mortality among patients with systemic lupus erythematosus (SLE)¹. In previously published work, the fraction of patients with atherosclerotic vascular events (AVE) was approximately 10% in patients with SLE, after an average follow-up of 9 years². Women with SLE have been observed to have a 5 to 50-fold increase in their risk of coronary artery disease (CAD)^{3,4} compared to the general population. This has triggered a concentrated effort to identify and treat traditional cardiovascular risk factors^{5,6}, facilitated by the development of cholesterol-lowering medications and newer antihypertensive medications. Furthermore, the cardioprotective effect of antimalarial medications has been recognized in conjunction with their effect on prevention of disease flares, thrombosis, and damage accrual and improved survival^{7,8,9,10}, leading to increased use in patients with SLE¹¹. This prompted us to reexamine the prevalence of AVE in the current era.

The Systemic Lupus International Collaborating Clinics (SLICC) group has developed an inception cohort of SLE patients to study atherosclerosis with the following specific objectives: 1. To determine the incidence, prevalence and nature of AVE in SLE; 2. To identify associated risk factors for the development of AVEs; 3. To develop interventional approaches to modify the identified risk factors^{12,13,14}.

We report on the prevalence, incidence rate, and risk factors for AVE in this multinational, multiethnic, inception cohort of patients with SLE, the SLICC inception cohort.

METHODS

Patients: An inception cohort of SLE patients has been assembled and followed according to a standardized protocol between October 1999-October 2017. Patients were enrolled in the cohort

within 15 months of SLE diagnosis (≥ 4 American College of Rheumatology (ACR) criteria)¹⁵. Clinical and laboratory features of SLE and comorbidities were recorded at yearly intervals¹². Disease activity was measured by the Systemic Lupus Erythematosus Disease Activity Index 2000 (SLEDAI-2K¹⁶). Accumulated damage was recorded according to the SLICC-ACR Damage Index (SDI)¹⁷. Data were entered and stored on an Oracle database. Patients who had at least one follow-up visit were included in analyses reported here.

The SLICC Registry Ethics approval with the University Health Network (UHN) Research Ethics Board (REB) is #00-0279-A under the title *The Systemic Lupus Collaborating Clinics Registry for Atherosclerosis in SLE*. All participating sites have their own local ethics approvals.

Outcome: The diagnosis of AVE was confirmed using standard clinical criteria, relevant laboratory data and imaging and includes (1) Myocardial infarction, defined as one of: definite electrocardiographic (ECG) abnormalities; typical symptoms with probable ECG abnormalities and abnormal enzymes ($> 2X$ upper limit of normal). (2) Angina, defined as severe pain or discomfort over the upper or lower sternum or anterior left chest and left arm, of short duration, relieved by rest or vasodilators. (3) Transient ischemic attack (TIA), defined as a brief episode of neurological dysfunction without residua. (4) Stroke, defined as an abrupt onset of neurological dysfunction resulting in neurological damage. (5) Congestive heart failure due to ischemic heart disease requiring treatment. (6) Bradyarrhythmia due to ischemic heart disease requiring pacemaker insertion. (7) Peripheral vascular disease (based on the presence of intermittent claudication). Attribution of vascular events to atherosclerosis was based on SLE being inactive (clinical SLEDAI-2K of 0) at the time of the event, and the presence of typical atherosclerotic changes on imaging or pathology and/or evidence of atherosclerosis elsewhere. Only vascular events that occurred after enrolment into the cohort were included. All required information for an

AVE diagnosis was recorded by the local SLE clinical investigator on a standardized data retrieval form. Information was obtained from the attending physician and recorded on the data retrieval form. Each event had an event date associated with it which was not necessarily the same as the annual follow up date and the assessment of lack of disease activity occurred at the time of the event.

The following variables were recorded: baseline demographics, classic cardiovascular risk factors (smoking, hypertension [$>140/90$], high cholesterol, random glucose levels, body mass index (BMI), alcohol use, family history of cardiovascular disease), disease-related (SLEDAI-2K, SDI, antiphospholipid antibodies [anticardiolipin antibody (aCL) and/or lupus anticoagulant (LA)] ever in those who were tested), previous non-atherosclerotic vascular events (VEs), and SLE treatment (glucocorticosteroids, antimalarials, immunosuppressive agents).

Statistical Analysis: We calculated the incidence rate per 1000 patient-years for AVEs. Demographic information was summarized with mean and standard deviations (SD) for normally distributed data and count (percentage) for categorical data. For missing values for patients' weight, height, and family history of cardiovascular disease, carried-forward imputations were used.

Kaplan-Meier curves were calculated for time-to-event outcomes. Demographic information at enrolment was used to define time-fixed covariates in Cox relative risk regression models while clinical, treatment, and classic AVE risk factor information were updated at the time of each visit¹⁸.

The primary outcome of interest was the time to the first AVE. Other patients were deemed to be event-free until the date of their last visit. The time scale used for analysis was age, to provide the most general adjustment for age, with the time of entry into follow-up defined as the age at the enrolment visit. We carried out univariable and multivariable time-dependent Cox regressions for time to first AVE, with all analyses adjusted for the prior occurrence of non-AVEs. Because aCL

and/or LA tests were not performed in 23% of the patients, we initially performed these analyses on all patients but excluded antiphospholipid antibodies as a risk factor. We then repeated the analyses in the 77% of the patients who had these antibodies tested. Multivariable models were developed based on univariable results and the fitting of models with the most relevant clinical variables. Additional analyses incorporated recurrent AVEs and were based on relative risk modelling with stratification by previous AVE¹⁹.

The analyses were performed using SAS, version 9.3 (SAS Institute, Inc., Cary, North Carolina), and the program coxph in R²⁰.

RESULTS

Of the 1848 patients in the cohort, 1710 patients had at least one follow-up visit for a total of 13,366 patient years, and constitute the study population. The 138 patients excluded from the study were not different in terms of initial visit information from the 1710 patients who formed this study cohort. This is an early, multicenter, multiethnic inception cohort followed for a mean of 8.3 and standard deviation of 4.3 years. **Table 1** illustrates the demographics, clinical, and therapeutic features of this cohort. Of the 1710 patients enrolled, 85 have died, giving a crude mortality of 5.0% (85/1710) or 6.4 per thousand person years (85/13366). Three hundred and fifty-seven patients were lost to follow-up giving a crude rate of 20.9%).

One hundred and seventy vascular events were identified in 113 patients after their enrolment visit. Of these, 86 were attributed to AVE, and 84 were VEs not attributed to atherosclerosis. Of these 84, 53 were attributed to SLE, and 31 were attributed to other causes, including vasospasm, trauma, fluid-overload associated with chronic renal disease, vascular malformations, and sepsis. Since the purpose of our study was to identify and determine risk factors for AVEs, we focused our analyses on AVEs. Of the 86 AVEs, 61 were first events and 25 were recurrent events (**Table 2**).

Sixty-one (3.57%) of the 1710 patients had at least one AVE. The mean (SD) duration from diagnosis to the first event was 6 (3.9) years. The total number of patient-years of follow-up among these patients from entry into the cohort to AVE, last visit or death was 13,366 years providing an incidence rate of 4.56 per 1000 patient-years (95% confidence interval, CI 3.6 -5.9). **Figure 1** presents Kaplan-Meier estimates of the probability of being event free from first AVEs.

We analyzed individually the predictors for first AVE. **Table 3** summarizes these analyses, all adjusted for prior other VEs which is highly significant as a predictor itself (HR[95%CI]: 4.13[1.61,10.6]). Missing values in the predictors led to fewer events in some analyses but with the exception of aCL/LA information, this was minimal. There was some evidence that female sex, antimalarial treatment and never having smoked were protective and a high BMI and ever having a positive aCL/LA conferred increased risk. The coding for BMI was influenced by non-parametric estimation of the effect of BMI which indicated that only very high values of BMI (>40) were associated with a higher risk of AVEs (not statistically significant). Similarly, there was a trend for average steroid dose to be associated with increased risk of AVE.

Table 4 presents results from two multivariable models including the predictors with significant effects in the single factor analyses, one, Model (a), without the aCL/LA variable and one, Model (b), with this variable. The inclusion of aCL/LA led to the exclusion of 405 patients, of whom 7 had AVEs, for whom this variable was missing. In addition to the predictive effect associated with prior other VEs, high BMI was also predictive of first AVE while only antimalarial therapy demonstrated a highly significant protective effect, [HR (95%CI): 0.54 (0.32, 0.91)], after adjustment for the other factors in the model. While 67.8% of all patients took antimalarials at enrolment, 86.7% were using antimalarials during the course of follow-up. The other effects were broadly consistent with the single factor analyses although the estimated effects were generally

smaller in multivariable models with the exception of the increased risk associated with very high BMI (>40) which is somewhat more significant in these models.

We performed a univariable analysis to determine whether previous AVE was a risk for recurrent AVE. Previous AVE was found to increase the risk of subsequent AVEs [HR(95% CI): 5.42(3.17,9.27), $p<0.001$].

Table 5 presents an additional multivariable analysis when recurrent AVEs are included in the analysis, with time dependent stratification on prior occurrence of an AVE. Initial investigations (results not shown) indicated that there was some evidence of differential effects for first AVEs and recurrent AVEs for female sex and anti-malarial therapy. Both variables appear to have protective effect only for first AVEs (although not significant for female sex). This is reflected in Table 5 by presenting separate effects for these variables in the two strata. Broadly speaking, the estimated effects associated with other variables are consistent with those found in analyses of first AVEs, although the effect of never smoking is somewhat more marked. A stratified analysis of all AVEs including the aCL/LA variable is not presented as there was insufficient data for some variables to allow reliable estimation. The results for other variables were consistent with the results shown in **Table 5**.

DISCUSSION

Atherosclerotic cardiovascular events have been recognized as a major comorbidity in SLE. Initial estimates of clinical prevalence of AVE in SLE were 6.7-10%^{2,3,21}.

The SLICC inception cohort was established in 1999 initially to determine the incidence, prevalence and nature of AVEs in SLE. Patient accrual to this cohort continued until 2011 when 1848 patients had been recruited. Patients continue to be followed to the present time. Thus, identification and prevalence of AVEs in the SLICC inception cohort represents the modern era.

The prevalence of AVE in this study is much lower than in previously published data. In the Toronto Cohort, reported in 2007 (followed from 1970 to 2004), 10% of the patients in both the prevalent and inception cohorts had the AVEs². In the current study, only 3.6% of patients had an AVE. Furthermore, the rate of AVEs in the Hopkins Cohort reported in 2012, which included the period 1987 to 2010, was 14.1/1000 patient-years²². It should be noted that in the Baltimore study, VEs in patients with and without active SLE were both counted whereas our analysis excluded VEs in patients with SLEDAI > 0. Thus one must be cautious in comparing event rates between the two studies. However, this is similar to AVE rate in the Toronto Lupus Cohort followed from 1975 to 1999, which was 19/1000 patient-years²³. In the SLICC cohort including patients from 1999 to 2017 the incidence rate was 4.56 per 1000 patient-years, similar to the incidence of 4.4/1000 seen in the Toronto cohort followed during the same period both using similar definitions for AVEs²³. Taking out the Toronto patients from the SLICC cohort provided the same frequency. This decline in rate of AVEs mirrors the declining incidence of AVEs noted in the general population^{24,25}. The decline in AVEs in the SLICC cohort may also be due to better control of lupus disease activity, more judicious use of glucocorticosteroids²⁶ as well as improvements in the treatment of cardiac risk factors in the modern era²³.

With regards to predictors of first AVEs, in the total cohort we demonstrated that prior other VE and BMI >40 were predictive of AVEs while antimalarial treatment was protective. These results were consistent whether the whole cohort or the sub-cohort that included only patients tested for aCL/LA were analyzed. Although aCL/LA, never having smoked and female sex were predictive in univariable analyses they were less marked in the multivariable analysis. In a study of 182 patients from Sweden, published in 2009, any aCL was predictive of first cardiovascular event²⁷, and another study performed in Northern Sweden including 277 patients also identified aCL as a

prognostic factor for cardiovascular events²⁸. Also, in a study from Baltimore including 1874 patients with SLE, a history of LA was predictive of cardiovascular events²². These reports did not provide information on the attribution of AVEs to aCL.

When we analyzed multiple AVEs we found that a prior AVE was predictive of a recurrent AVE and a multivariable analysis also demonstrated that a prior other vascular events was predictive for recurrent events, never having smoked was protective for all events and antimalarial treatment was protective for first AVEs.

Hanly et al.²⁹ investigated cerebrovascular events in the SLICC cohort and found that 94% were due to SLE. Moreover, 40% of events were identified at the enrollment visit, and the remainder over the mean follow-up of 6.6 years, almost 2 years shorter than the current study. These early events due to SLE are in contrast to the events described here that are later in onset and attributed to atherosclerosis.

The strength of the study is that it is based on a multinational multiethnic large inception cohort followed over 18 years with an average follow-up of 8 years. Definitions for AVE were specified a priori and known risk factors for AVE were collected prospectively.

There are a number of limitations for the study. The adjudication of AVEs vs non-atherosclerotic VEs was subject to potential inaccuracy as it was determined by each local SLE expert/site investigator although attempts were made for standardization of this procedure by using clear definitions of each AVE and SLE disease activity was taken into account. Silent atherosclerotic events might have been missed but those ascertainment criteria are similar to those used in the general population. As well, more AVEs may accrue with further follow-up. Another possible limitation is the fact that the study was designed as yearly evaluation and the exact timing of possible discontinuation of therapies could not be analyzed. However, the time dependent analyses

do inform on the effect of taking a medication prior to the event. Since SLE activity was tied into the outcome definition, it may have impacted the observed association with potential predictors that may also be associated with SLE disease activity, although antimalarials are generally continued regardless of disease activity.

With regards to the SLICC atherosclerosis registry aims, we have now demonstrated that the incidence of AVEs in the current era is lower than expected. While it was anticipated that there would be both classic risk factors (e.g. smoking) and disease related risk factors, we were unable to establish clear relationships between AVEs and many of the variables studied. We hypothesize that more effective control of lupus disease activity and treatment of classic atherosclerotic risk factors may have both contributed to controlling AVEs in this inception cohort. However, in all our analyses we found that antimalarial treatment was protective for AVE emphasizing the importance of this therapy in the management of SLE.

REFERENCES

¹ Sheane BJ, Urowitz MB, Gladman DD. Atherosclerosis in Systemic Lupus Erythematosus – Epidemiology, Risk Factors, Subclinical Assessment and Future Study. *Rheumatology Current Research* 2013; S5

² Urowitz MB, Ibañez D, Gladman DD. Atherosclerotic vascular events (AVE) in a single large lupus cohort: prevalence and risk factors. *J Rheumatol* 2007;34:70-75.

³ Manzi S, Meilahn EN, Rairie JE, et al. Age specific incidence rates of myocardial infarction and angina in women with systemic lupus erythematosus: Comparison with the Framingham Study. *Am J Epidemiol* 1997;145:408-5.

- ⁴ Bruce IN, Gladman DD, Urowitz MB. Premature Atherosclerosis in SLE. *Rheumatic Disease Clinics of North America* 2000;26:257-78.
- ⁵ Tselios K, Sheane BJ, Gladman DD, Urowitz MB. Optimal Monitoring For Coronary Heart Disease Risk in Patients with Systemic Lupus Erythematosus: A Systematic Review. *J Rheumatol* 2016;43(1):54-65.
- ⁶ Schoenfeld SR, Kasturi S, Costenbader KH. The epidemiology of atherosclerotic cardiovascular disease among patients with SLE: a systematic review. *Semin Arthritis Rheum* 2013;43(1):77-95.
- ⁷ Ruiz-Irastorza G, Ramos-Casals M, Brito-Zeron P, Khamashta MA. Clinical efficacy and side effects of antimalarials in systemic lupus erythematosus: a systematic review. *Ann Rheum Dis* 2010;69(1):20-28.
- ⁸ Bykerk V, Sampalias JS, Esdaile JM, Shoquette D, Senecal JL, Danoff D, Smith CD, Cividino A, Osterland CK, Yeadaon C. A randomized study of the effect of withdrawing hydroxychloroquine sulfate in systemic lupus erythematosus. *New Engl J Med* 1991;224:150-4.
- ⁹ Bruce IN, O'Keefe AG, Farewell V, Hanly JG, Manzi S, Su L, Gladman DD, et al. Factors associated with damage accrual in patients with systemic lupus erythematosus: results from the Systemic Lupus International Collaborating Clinics (SLICC) Inception Cohort. *Ann Rheum Dis* 2015;74:1706-13.
- ¹⁰ Jung H, Bobba R, Su J, Shariati-Sarabi Z, Gladman DD, Urowitz M, Lou W, Fortin PR. Antimalarials and Thromboembolic Disease in SLE. *Arthritis Rheum* 2010;62:863-868..
- ¹¹ Pakchotanon R, Gladman DD, Su J, Urowitz MB. More Consistent Antimalarial Intake in First 5 Years of Disease Is Associated with Better Prognosis in Patients with Systemic Lupus Erythematosus. *J Rheumatol* 2018;45:90-94.

- ¹² Urowitz MB, Gladman DD. The SLICC Inception Cohort for Atherosclerosis. *Current Rheumatology Reports* 2008; 10:281-285.
- ¹³ Urowitz MB, Gladman D, Ibañez D, et al. Accumulation of coronary artery disease risk factors over three years: data from an international inception cohort. *Arthritis Rheum* 2008;59:176-180.
- ¹⁴ Urowitz MB, Gladman DD, Anderson NM, Su J, Romero-Diaz J, Bae SC, et al. Cardiovascular events prior to or early after diagnosis of systemic lupus erythematosus in the systemic lupus international collaborating clinics cohort. *Lupus Sci.Med.* 2016;3: e000143.
- ¹⁵ Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1725.
- ¹⁶ Gladman DD, Ibañez D, Urowitz MB. SLE Disease Activity Index 2000. *J Rheumatol* 2002;29:288-91.
- ¹⁷ Gladman D, Ginzler E, Goldsmith C, Fortin P, Liang M, Urowitz M, et al. The development and initial validation of the SLICC/ACR damage index for SLE. *Arthritis Rheum* 1996;39:363-9.
- ¹⁸ Hosmer, DWJ.; Lemeshow, S. New York, NY: John Wiley & Sons; 1999. Applied survival analysis. Regression modeling of time to event data.
- ¹⁹ Therneau T.M., Grambsch P.M. (2000) The Cox Model. In: *Modeling Survival Data: Extending the Cox Model.* Statistics for Biology and Health. Springer, New York, NY OR
- ²⁰ R Core Team. *R: A Language and Environment for Statistical Computing.* Vienna, Austria: R Foundation for Statistical Computing 2015.
- ²¹ Petri M, Spence D, Bone LR, Hochberg MC. Coronary artery disease risk factors in the Johns Hopkins Lupus Cohort: prevalence, recognition by patients, and preventive practices. *Medicine* 1992;71:291-302.
- ²² Magder LS, Petri M. Incidence of and Risk Factors for Adverse Cardiovascular Events Among

Patients With Systemic Lupus Erythematosus. *Am J Epidemiol.* 2012;176:708–719

²³ Urowitz MB, Su J, Gladman DD. Atherosclerotic vascular events in SLE – an evolving story. *J Rheumatol Online* 2019/02/01

²⁴ Hardoon SL, Whincup PH, Lennon LT, Wannamethee SG, Capewell S, Morris RW. How much of the recent decline in the incidence of myocardial infarction in British men can be explained by changes in cardiovascular risk factors? Evidence from a prospective population-based study. *Circulation* 2008;117:598-604.

²⁵ Mannsverk J, Wilsgaard T, Mathiesen EB et al. Trends in Modifiable Risk Factors Are Associated With Declining Incidence of Hospitalized and Nonhospitalized Acute Coronary Heart Disease in a Population. *Circulation* 2016;133:74-81.

²⁶ Ruiz-Arruza I, Lozano J, Cabezas-Rodriguez I, Medina JA, Ugarte A, Erdozain JG, Ruiz-Irastorza G. Restrictive Use of Oral Glucocorticoids in Systemic Lupus Erythematosus and Prevention of Damage Without Worsening Long-Term Disease Control: An Observational Study. *Arthritis Care Res* 2018;70:582-91.

²⁷ Gustafsson J, Gunnarsson I, Borjesson O et al. Predictors of the first cardiovascular event in patients with systemic lupus erythematosus - a prospective cohort study. *Arthritis Res Ther* 2009;11(6):R186.

²⁸ Bengtsson C, Ohman ML, Nived O, Rantapaa DS. Cardiovascular event in systemic lupus erythematosus in northern Sweden: incidence and predictors in a 7-year follow-up study. *Lupus* 2012;21:452-9.

²⁹ Hanly JG, Qiuju LI, Su LI, Urowitz MB, Gordon C, Bae SC, et al. Cerebrovascular Events in Systemic Lupus Erythematosus: Results From an International Inception Cohort Study. *Arthritis Care Res* 2018;70:1478-87.

FIGURE LEGEND

Figure 1. Kaplan Meier survival curve for first AVE in 1710 SLE patients including 95% confidence intervals.

Table 1: Characteristics of SLE population N=1710	
Mean Age at SLE Diagnosis (years)*	34.7 ± 13.4
Mean disease duration at enrolment (months)*	5.7 ± 4.2
Females	1515 (88.6%)
Race (%) Caucasian / African descendant / Hispanic / Asian / Other	49.4 / 16.4 / 15.5/ 15.0 / 3.7
Mean duration of follow-up (yrs)*	8.3 ± 4.3
Mean SLEDAI-2K at first visit*	5.4 ± 5.4
SLICC Damage Index at year 1 *	0.3 ± 0.8
SLICC Damage Index at year 1 for those SDI >0* (N = 314)	1.5 ± 0.8
Glucocorticosteroid treatment at enrolment	1186 (69.4%)
Average daily glucocorticosteroid dose at enrolment (of those treated with glucocorticosteroids)*mg/day	23.9 ± 16.5
Antimalarial treatment at enrolment	1159 (67.8%)
Immunosuppressive treatment at enrolment	692(40.5%)
SBP>140 or DBP>90 mmHg at enrolment	204 (11.9%)
BP treatment	505 (29.5%)
Abnormal total cholesterol regardless of treatment	517 (30.2%)
Statins treatment at enrolment	155 (9.1%)
BMI at enrolment	25.3 ± 5.8
BMI > 40	38 (2.2%)

Smoking (currently or prior) at enrolment	598 (35.0%)
Alcohol consumption Yes (%)	470 (27.5%)
Units per week in those consumed	3.7 ± 4.9
Family history of sudden death, MI, angina or stroke at enrolment	512 (29.9%)
Ever ACL/LA positive ever (N = 1305)	469 (35.9%)
Anticoagulants treatment at enrolment	92 (5.4%)
Aspirin treatment at enrolment	287 (16.8%)
*Mean and standard deviation (SD)	

Table 2: Nature and attribution of vascular event

Event name	Attributed to atherosclerosis	Non Atherosclerotic
Myocardial infarction	21	4
Angina	23	4
TIA	5	13
Stroke	10	30
CHF	12	19
Pacemaker insertion	5	4
Peripheral vascular disease	10	10
Total number of events	86	84

Table 3. Cox relative risk regression for the outcome first AVE, adjusted for prior other VEs			
Predictor	Univariate		
	# events*	HR (95% CI)	p value
Age at SLE diagnosis (years)	61	1.03(0.95,1.11)	0.47
Female vs. Male	61	0.53(0.29,0.97)	0.04
Caucasian vs. Non-Caucasian	61	1.36(0.76,2.44)	0.30
SLEDAI score	61	1.03(0.97,1.10)	0.30
Adjusted mean SLEDAI	61	1.05(0.99,1.13)	0.13
Smoking: Never vs. ever smoked	61	0.46(0.22,0.95)	0.04
Ex-smoker vs. current-smoker		0.69(0.31,1.50)	0.35
Elevated random glucose compared to normal	61	0.97(0.45,2.07)	0.94
SBP >140 or diastolic > DBP 90 regardless of treatment	61	0.91(0.49,1.72)	0.77
Elevated cholesterol regardless of treatment vs. normal	61	1.17(0.66,2.08)	0.59
Body Mass Index: 30-40	60	0.82(0.42,1.61)	0.57
>40		2.51(0.97,6.47)	0.06
Alcohol consumption (yes/no)	56	0.76 (0.42-1.37)	0.36
Family history of sudden death, MI, angina or stroke	59	1.26(0.72,2.21)	0.43
Average oral steroid dose (mg/day)	61	1.02(1.00,1.04)	0.08
Antimalarials treated vs. not treated	61	0.51(0.30,0.87)	0.01
Immunosuppressives treated vs. not treated	61	0.99(0.59,1.68)	0.97
Ever ACL/LA positive	54	2.09(1.18,3.71)	0.01

SBP – systolic blood pressure, DBP – diastolic blood pressure. Except age, sex, and Caucasian variables, which were time-fixed, the remaining variables were time-dependent. *. Missing values in the predictors led to fewer events in some analyses.

Table 4: Multivariate Models. Outcome: first AVE				
	Model (a) [60 events]*		Model (b) [53 events]*	
Predictor	HR(95%CI)	p-value	HR(95%CI)	p-value
Prior other VEs	4.00(1.55,10.3)	0.004	4.76(1.80,12.60)	0.002
Female	0.60(0.32,1.24)	0.12	0.57(0.28,1.13)	0.11
Never vs ever smoked	0.51(0.24,1.06)	0.07	0.48(0.23,1.03)	0.06
Ex-smoker vs current smoker	0.67(0.30,1.48)	0.33	0.53 (0.23,1.21)	0.13
Anti-malarials	0.54(0.32,0.91)	0.02	0.52(0.29,0.92)	0.02
BMI 30-40	0.95(0.48,1.88)	0.89	0.88(0.42,1.87)	0.74
≥ 40	2.74(1.04,7.18)	0.04	3.10(1.17,8.23)	0.02
aCL/LA ever			1.73(0.95,3.12)	0.07
<p>*Two multivariable models including the predictors with significant effects in the single factor analyses: Model (a), without the aCL/LA variable and, Model (b), with this variable. Missing values in the predictors led to fewer events in some analyses. BMI – body mass index; aCL/LA – anticardiolipin/lupus anticoagulant; VEs – vascular events</p>				

Table 5: Multivariable model including first and recurrent AVEs in 82 events with stratification on prior AVE

Predictor	HR(95%CI)	p-value
Prior other VEs	2.85(1.20,6.78)	0.02
Female (no prior AVE)	0.63(0.34,1.21)	0.16
Females (with prior AVE)	1.25(0.39,4.00)	0.70
Never vs ever smoked	0.40(0.21,0.76)	0.005
Ex-smokers vs current smokers	0.63(0.32,1.25)	0.19
Antimalarials (no prior AVE)	0.53(0.31,0.91)	0.02
Antimalarials (with prior AVE)	1.90(0.67,5.35)	0.23
BMI 30-40	1.00(0.55,1.80)	0.99
BMI ≥ 40	1.96(0.75,5.14)	0.17

* Missing values in the predictors led to fewer events in some analyses. BMI – body mass index; aCL/LA – anticardiolipin/lupus anticoagulant; VEs – vascular events