RNA-sequencing and somatic mutation status of adrenocortical tumors

Dalmazi, Guido Di; Altieri, Barbara; Scholz, Claus; Sbiera, Silviu; Luconi, Michaela; Waldmann, Jens; Kastelan, Darko; Ceccato, Filippo; Chiodini, Iacopo; Arnaldi, Giorgio; Riester, Anna; Osswald, Andrea; Beuschlein, Felix; Sauer, Sascha; Fassnacht, Martin; Appenzeller, Silke; Ronchi, Cristina

DOI:

[10.1210/clinem/dgaa616](https://doi.org/10.1210/clinem/dgaa616)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Dalmazi, GD, Altieri, B, Scholz, C, Sbiera, S, Luconi, M, Waldmann, J, Kastelan, D, Ceccato, F, Chiodini, I, Arnaldi, G, Riester, A, Osswald, A, Beuschlein, F, Sauer, S, Fassnacht, M, Appenzeller, S & Ronchi, C 2020, 'RNA-sequencing and somatic mutation status of adrenocortical tumors: novel pathogenetic insights', *Journal of Clinical Endocrinology and Metabolism*, vol. 105, no. 12, pp. e4459–e4473. <https://doi.org/10.1210/clinem/dgaa616>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Guido Di Dalmazi, Barbara Altieri, Claus Scholz, Silviu Sbiera, Michaela Luconi, Jens Waldman, Darko Kastelan, Filippo Ceccato, Iacopo Chiodini, Giorgio Arnaldi, Anna Riester, Andrea Osswald, Felix Beuschlein, Sascha Sauer, Martin Fassnacht, Silke Appenzeller, Cristina L Ronchi, RNA-Sequencing and Somatic Mutation Status of Adrenocortical Tumors: Novel Pathogenetic Insights, *The Journal of Clinical Endocrinology & Metabolism*, , dgaa616, <https://doi.org/10.1210/clinem/dgaa616>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

RNA-Sequencing and Somatic Mutation Status of Adrenocortical Tumors: Novel Pathogenetic Insights.

Guido Di Dalmazi^{1*}, Barbara Altieri^{2*}, Claus Scholz³, Silviu Sbiera², Michaela Luconi⁴, Jens Waldman⁵, Darko Kastelan⁶, Filippo Ceccato⁷, Iacopo Chiodini^{8,9}, Giorgio Arnaldi¹⁰, Anna Riester¹¹, Andrea Osswald¹¹, Felix Beuschlein^{11,12}, Sascha Sauer¹³, Martin Fassnacht², Silke Appenzeller^{14*}, Cristina L. Ronchi^{2,15*}.

Affiliations

¹Endocrinology Unit, Department of Medical and Surgical Sciences, University of Bologna (Italy). ²Division of Endocrinology and Diabetes, Department of Internal Medicine I, University Hospital, University of Würzburg, Würzburg (Germany). ³Life and Medical Sciences Institute, University of Bonn (Germany). ⁴Department of Experimental and Clinical Biomedical Sciences, University of Florence (Italy). ⁵Mivendo Klinik Hamburg (Germany). ⁶Department of Endocrinology, University Hospital Center Zagreb (Croatia). ⁷Endocrinology Unit, Department of Medicine DIMED, University-Hospital of Padua (Italy). ⁸Istituto Auxologico Italiano, IRCCS, Unit for Bone Metabolism Diseases and Diabetes & Lab of Endocrine and Metabolic Research, Milan (Italy). ⁹University of Milan, Milan (Italy). ¹⁰Division of Endocrinology, Department of Clinical and Molecular Sciences, Polytechnic University of Marche, Ancona (Italy). ¹¹Medizinische Klinik und Poliklinik IV, Klinikum der Universität München, Munich (Germany). ¹²Klinik für Endokrinologie Diabetologie und Klinische Ernährung, Universitäts Spital Zürich, Zürich (Switzerland). ¹³Max Delbrück Center for Molecular Medicine/Berlin Institute of Health, Berlin (Germany). ¹⁴Core Unit Bioinformatics, Comprehensive Cancer Center Mainfranken, University of Würzburg (Germany). ¹⁵Institute of Metabolism and Systems Research, University of Birmingham (United Kingdom).

***Authors contributed equally to the work**

Keywords: adrenocortical adenoma, Cushing syndrome, mild autonomous cortisol excess, transcriptome, gene fusions, long non-coding RNA.

Short title: Pathogenesis of adrenocortical tumors.

Word count: 4663

Corresponding author/reprint requests:

Cristina L Ronchi, MD, PhD, Division of Endocrinology and Diabetes, Department of Internal Medicine I, University Hospital, University of Würzburg, Oberduerrbacher-Str 6, 97080 Würzburg (Germany). Tel 0049-931-201-39704. E-Mail: Ronchi_C@ukw.de

Funding:

This work has been supported by the Deutsche Forschungsgemeinschaft (DFG) within the CRC/Transregio (project number: 314061271 – TRR 205 to A.R., A.O., F.B., M.F., and C.L.R.), and project FA-466/8-1 and RO-5435/3-1 (M.F and C.L.R., respectively), the Deutsche Krebshilfe (70113526 S.A., M.F. and C.L.R., and 70112617 to F.B. and M.F.), the Else Kröner-Fresenius-Stiftung (2015_A228 to A.R.), the Italian Ministry of Health (RF-GR-2018-12365398 to G.D.D. and RF-2013-02356606 to I.C.) and the Italian Ministry of University and Research (PRIN 2017 to G.D.D.).

Disclosure summary: The authors have nothing to disclose.

ABSTRACT

Context. Pathogenesis of autonomous steroid secretion and adrenocortical tumorigenesis remains partially obscure.

Objective. To investigate the relationship between transcriptome profile and genetic background in a large series of adrenocortical tumors and identify new potential pathogenetic mechanisms.

Design. Cross-sectional study.

Setting. University Hospitals of the European Network for the Study of Adrenal Tumors (ENSAT).

Patients. We collected snap-frozen tissue from patients with adrenocortical tumors (n=59) with known genetic background: 26 adenomas with Cushing syndrome (CS-CPA), 17 adenomas with mild autonomous cortisol secretion (MACS-CPA), 9 endocrine-inactive adenomas (EIA), and 7 adrenocortical carcinomas (ACC).

Intervention. RNA-sequencing.

Main Outcome Measures. Gene expression, long non-coding RNA (lncRNA) expression, and gene fusions. Correlation with genetic background defined by targeted Sanger sequencing or whole-exome sequencing.

Results. Transcriptome analysis identified two major clusters for adenomas: Cluster 1 (n=25) consisting of EIAs and MACS-CPAs with *CTNNB1* or no driver mutations, and Cluster 2 (n=25) consisting of CS-CPAs and MACS-CPAs with *GNAS* mutations and CS-CPAs with *PRKACA* or no drivers mutation. CS-CPA with *CTNNB1* mutations (n=2) clustered close to ACC. In Cluster 2, *FATE1* was among most common overexpressed genes. lncRNA analysis confirmed that patients clustered in four groups. Novel gene fusions were found, including *AKAP13-PDE8A* in one CS-CPA sample with no identified driver mutation.

Conclusions. MACS-CPAs and EIAs showed a similar transcriptome profile, independently of the genetic background, except for MACS-CPAs with *GNAS* mutations that clustered with most CS-CPAs. Still unrevealed molecular alterations in the cAMP/PKA or Wnt/beta catenin pathways might be involved in the pathogenesis of adrenocortical tumors.

Introduction

Over the last few years, the molecular and genetic events underlying the pathogenesis of adrenocortical tumors have been investigated by applying classical genetic approaches and next-generation sequencing techniques including whole-exome sequencing (WES). Alterations of several components of the cAMP/PKA pathway have been identified as a causative factor for tumorigenesis and cortisol hypersecretion in adrenocortical adenoma (ACA) [reviewed in (1,2)]. Independent studies have reported somatic mutations in the *PRKACA* gene as the pivotal pathogenetic event in approximately half of the ACAs associated with Cushing syndrome (CS), due to a constitutive activation of the catalytic subunit α of PKA (3-7). In a smaller proportion of patients with CS, additional mutations in members of the cAMP-PKA signaling pathway have newly been described, such as somatic mutations of the gene encoding the regulatory subunit II beta of PKA (*PRKACB*, very rare) (8), or confirmed, such as those occurring in the gene encoding the regulatory subunit 1 α of PKA (*PRKARIA*) or the protein Gs α (*GNAS*) (4,5,7,9). Activating mutations in the gene encoding β -catenin (*CTNNB1*) have also been identified as an important contributor of adrenocortical growth. Interestingly, *CTNNB1* mutations had been reported in ACA and adrenocortical carcinoma (ACC) with a similar prevalence (10-30%) (9,10) with the highest frequency in endocrine inactive adenomas (EIA) (9,11). Despite these substantial advances, the molecular mechanisms underlying autonomous steroid secretion and adrenocortical tumorigenesis remain obscure for around 2/3 of cases that cannot be linked to known driver mutations.

Another important issue, is that, depending on the definition used, up to 40% of patients with apparently **non-functioning** ACA might present a mild autonomous cortisol secretion (MACS), a condition previously termed “subclinical CS”, since the classical clinical manifestations of overt hypercortisolism are absent (12,13). However, several evidences suggest that MACS could be associated with cortisol-dependent comorbidities, including higher risk of cardiovascular events, hypertension, dyslipidemia, diabetes mellitus type 2, obesity, osteoporosis and higher mortality rate (12,14-18). A previous study on transcriptome analysis on a small series of ACAs identified two different clusters according to hormone secretion: cluster 1, including only cortisol-producing adenomas (CPAs) associated to CS (CS-CPAs), and cluster 2 including non-functioning ACAs, as

well as MACS- and CS-CPAs, revealing an association between cortisol secretion and expression of a subset of genes implicated in steroid secretion (19). However, as to this point, the association between transcriptome and mutational status of benign adrenocortical tumors has not been investigated.

Our first aim was to analyze the association between the genetic background and the transcriptome profile assessed by RNA-sequencing (RNA-seq) in a large series of ACAs. Our second aim was to identify novel molecular events that may be involved in cortisol hypersecretion and adrenocortical tumorigenesis through the analysis of alternative gene spliced transcripts, long non-coding RNAs (lncRNAs), and gene fusions.

Materials and methods

Study protocol

This is a European multicentric retrospective study among centers belonging to the European Network for the Study of Adrenal Tumors (ENSAT), designed and conducted in accordance with the Declaration of Helsinki. The local ethic committees of each Institution approved the study protocol. Written informed consent was obtained from all subjects prior to study enrollment.

Selection of patient cohort

We selected snap-frozen samples from the previously published series of 99 non-aldosterone producing ACAs without exon 7 hot-spot *PRKACA* mutations, already characterized by WES (9). Among these, we took 54 samples with still available tumor material for RNA isolation. After excluding seven cases due to poor quality RNA (see paragraph “sample preparation”), we obtained a final series of 47 ACAs for RNA-seq analysis (**Figure 1**). According to the results of WES, we identified 22/47 (46.8%) ACAs with somatic mutations in known driver genes, such as *GNAS*, *CTNNB1*, and *PRKACA* (outside the hot-spot region). These included 7 CS-CPAs, 11 MACS-CPAs, and 4 endocrine inactive adenomas (EIAs). The remaining 25/47 (53.2%) tumors (n=14 CS-CPAs, 6 MACS-CPAs, and 5 EIAs) did not have somatic mutations in known driver genes at WES. In addition, we included 5 further CS-CPAs samples with known hot-spot L206R *PRKACA* mutations from the previously published cohort (6) and 7 early-stage/low aggressive ACCs (ENSAT tumor stage

1-3, Ki67 proliferation index <20%) that served as controls. The final cohort included 59 samples (Figure 1). Four normal adrenal glands (NAGs) obtained from surgery for renal carcinoma (n=1) or from tissue surrounding EIA (n=3) were used as controls for gene expression profiling. The dissection of the adrenal tissues from the snap-frozen specimen was made by expert pathologists at each Center participating in the study.

At variant calling performed on our RNA-seq data (see methods below), we could detect two additional somatic mutations in two CS-CPA cases classified as without driver mutations based on previous WES (see details shown in Figure 1). These mutations were in genes *PRKACA* (hot spot L206R found in CS-CPA2) and *GNAS* (hot spot R201S in CS-CPA6). In addition, two previously identified alterations in *CTNNB1* genes could not be verified, i.e. a 1602bp splice indel in CS-CPA22 and a missense mutation T41A in CS-CPA9. According to these findings, we classified our cohort as follows: 27 ACA with known driver mutations (7 CS-CPA *PRKACA*, 4 CS-CPA *GNAS*, 2 CS-CPA *CTNNB1*, 9 MACS *CTNNB1*, 1 MACS-CPA *GNAS*, 4 EIA *CTNNB1*) and 25 ACA without driver mutations (13 CS-CPA, 7 MACS-CPA and 5 EIA).

Clinical data collection

Clinical and hormonal data such as sex, age at diagnosis, initial tumor size, steroid secreting pattern, symptoms of overt CS were already available from previous studies (6,9).

Overt CS and MACS were defined according to the Endocrine Society guidelines (13), in order to maintain the same classification of previous studies (6,9). These guidelines recommend the use of the 1 mg overnight dexamethasone test (DST) with a cortisol cutoff of <1.8 µg/dL (50 nmol/L) as the primary test to exclude autonomous cortisol secretion. However, the newest European Society of Endocrinology (ESE)/ ENSAT guidelines from 2016 (12) recommend interpreting the 1 mg overnight DST as a continuous variable, considering post-dexamethasone serum cortisol levels between 1.9–5.0 µg/dL (51 and 138 nmol/L) as ‘possible autonomous cortisol secretion’ and levels >5.0 µg/dL (>138 nmol/L) as confirmed ‘autonomous cortisol secretion’. Both guidelines recommend the use of additional biochemical tests to confirm cortisol secretory autonomy and assess the degree of cortisol

secretion, and MACS were defined in absence of clinical and catabolic signs of overt CS (12,13). To avoid bias in the interpretation of our results, we performed a second analysis classifying the patients according to the ESE/ENSAT guidelines. According to that, 2 MACS-CPAs (MACS-CPA2 and MACS-CPA11) without driver mutation were reclassified as EIAs, and one EIA with *CTNNB1* mutation (EIA6) was reclassified as MACS-CPA. Differences between the results deriving from the two classifications were evaluated. In all our patients with MACS with values between 1.8 and 5 mcg/dL, ACTH was below 2.2 pmol/L (10 pg/mL) or midnight cortisol was above the reference range.

Clinical and histopathological data of ACC cases, including ENSAT tumor stage, Weiss score, and Ki67 index, were available from a previous publication (20) or were collected through the ENSAT registry (<https://registry.ensat.org/>).

Sample preparation and RNA-sequencing

RNA was isolated from snap-frozen tumor tissue by RNEasy Lipid Tissue Mini Kit (Qiagen, Hilden, Germany) (n=23) or by Maxwell® 16 Total RNA Purification Kit used with the Maxwell® 16 Instrument (n=36), according to the manufacturers' instruction. Initial RNA quality control was performed by spectrophotometry at 260 nm and by running a denaturing agarose gel. Additional analysis was performed using an Agilent Technologies 2100 Bioanalyzer and an RNA Integrity Number (RIN) value ≥ 8 was required to ensure efficient mRNA sequencing. TruSeq RNA Library Prep Kits was applied prior Illumina sequencing (NextSeq500) of pooled normalized libraries. Specifically, a paired-end 75nt mode (high-output flow-cells) was used for a minimum of 40-100 million reads per sample.

Data analysis

An initial quality assessment was performed using FastQC, v0.11.5 (Available online at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>). Adapter- and quality trimming was done with Cutadapt, v1.1.5 (<https://cutadapt.readthedocs.io/en/stable/>). STAR v2.5.3a (21) was used to map the trimmed reads to the GENCODE human reference genome GRCh37 release 29. We used

Samtools v1.3 (22) utilizing htlib, v1.3 for sam-to-bam-conversions as well as sorting and indexing of the alignment files. For gene annotation, the GENCODE human reference genome GRCh37 release 29 was used. FPKM-values and differential gene expression as well as differential isoform expression were calculated with Cufflinks package, v2.2.1 (23). For principal component analysis (PCA) analysis, one was added to all FPKM values. The samples were normalized to the median of the four sequenced NAGs and log2-transformed. FPKM-values of all genes with a coefficient of variation ≥ 0.5 . PCA was performed with the prcomp function from the R stats package, v3.4.4. Subsequently, 500 protein coding genes and 250 lncRNA genes with the highest PC loadings in one of the first three components were selected for unsupervised complete linkage clustering, respectively. The clustering was performed on the rows and columns using the Euclidian distance metric with the heatmap.2 function in the gplots R-package, v3.0.1. Arriba v1.1.0 was used for fusion detection with default settings (21-23). In the heatmap, we used log2(FC). The value 1 was added to each FPKM in order to make dividing by the median possible.

For pathway analysis, we used the Gene Set Enrichment Analysis (GSEA) software by Broad Institute v.MSigDB 7.0 (24,25) using gene sets from both Reactome and KEGGS pathways.

Variant calling using RNAseq data

Variant calling was performed following the GATK Best Practices workflow for SNP and Indel calling on RNAseq data (<https://software.broadinstitute.org/gatk/documentation/article.php?id=3891>). Trimmed reads TrimGalore, v06.6.1 (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) utilizing Cutadapt, v2.3 (26) were mapped to the reference using STAR, v2.5.4b (21). The index and the dictionary of the reference sequence were created using Samtools v1.3 (22) and Picard, v2.18.11 (<https://broadinstitute.github.io/picard/>), respectively. Picard was also used to add read groups, to sort alignment files, to mark duplicates, and to create index files. The Genome Analysis Toolkit v3.8 (27) was used to reassign mapping qualities, indel realignment, base recalibration, variant calling, as well as variant filtering. Variants were annotated using ANNOVAR (v2019-10-24) (28). Somatic protein-altering variants and splice site variants were reported if they alter the protein sequence or affect a

splice site, are rare in the population (below a frequency of 2 % in 1000g2015aug_all, ExAC_nontcga_ALL, gnomAD_exome_ALL and gnomAD_genome_ALL) and the position is covered by at least 20 reads and the alternative allele is covered by at least 8 reads and comprised at least 10 %.

Validation of gene fusions

We selected a total of 7 high-confidence gene fusions to be validated by Polymerase Chain Reaction (PCR) and Sanger-Sequencing. Reverse transcription of RNA was performed using the QuantiTect Reverse Transcription Kit (Qiagen, Hilden, Germany) as previously described (29). PCR amplification of 10 ng of cDNA was performed in a final volume of 25 µl using specific primer sets designed to span the fusion breakpoints, whereas GAPDH was used as internal control gene (Supplementary Table 2 (30)). Cycling conditions were: 96°C for 2 min followed by 30 cycles of denaturing at 96°C (30 sec), annealing at 58°C (30 sec) and elongation at 72°C (1 min), and a final step of 72°C for 5 min. The PCR products were run on a 2.4% low-melt agarose gel with ethidium-bromide. PCR products were purified (ExoSAP-IT, Amersham Biosciences, Munich, Germany) and sequenced using the QuickStart Cycle Sequencing Kit (AB Sciex, Beckman Coulter, #608120, Krefeld, Germany) on a Genetic Analysis System (CEQ) 8000 (AB Sciex, #A16637-AA, Krefeld, Germany) capillary sequencer according to the manufacturer instructions. Data were analyzed using the CEQ8000 DNA Analyzer (AB Sciex).

Statistical analysis

To correlate the results of the unsupervised clustering analysis with the clinical features, the following parameters were evaluated: sex, age (cutoff 50 years old), tumor size (cutoff 35 mm), diagnosis (EIA, MACS-CPA, CS-CPA, and ACC), and mutational status (no driver mutations, *CTNNB1* mutation, mutations of the cAMP/PKA pathway, other driver mutations). A Fisher exact or Chi-square (χ^2) tests were used to investigate dichotomic variables, while a two-sided *t*-test or Mann-Whitney U-test was used to compare continuous variables, as appropriate. Kruskal-Wallis test, followed by Bonferroni

post-hoc test or Dunn's multiple comparison test, was used for comparison among several groups for non-normally distributed variables. Correlations and 95% confidence intervals (95% CI) between different parameters were evaluated by linear regression analysis. Statistical analyses were made using GraphPad Prism (v.6.0, La Jolla, CA, USA) and SPSS Software (v.23, SPSS Inc., Chicago, IL, USA). Data are presented as median with interquartile range in parentheses or mean $\pm$ standard deviation (SD). A *p* value <0.05 was considered statistically significant.

Results

Patient characteristics

Clinical parameters and hormonal levels at the time of diagnosis are reported in **Table 1**. Patients with CS-CPAs were mostly females and significantly younger than those of the other groups. As expected, they had significantly higher post-dexamethasone cortisol levels and 24-h urinary free cortisol, as well as lower ACTH levels in comparison to other groups. No other significant differences were observed in clinical parameters.

Among ACCs, 2 patients had non-secreting tumors and 5 had secreting ACC, including 2 overt-Cushing, 1 associated with mild autonomous cortisol secretion, and 2 associated with androgens excess.

Gene expression profile (overview)

The entire list of differentially expressed genes is reported in the **Supplementary Table 1**. Protein coding genes recognized at RNA-seq represented 86% of all detected genes, whereas the remaining gene types included processed and unprocessed pseudogenes, mitochondrial rRNA, and lncRNA (**Supplementary Figure 1** (30)). The number of significantly over- and under-expressed protein-coding genes in each group and the most frequently altered pathways are reported in **Table 2**. CS-CPAs showed 67 overexpressed genes, being mostly related to *GPCR ligand binding* and *G alpha (q) signaling*, with *NTS*, *TJP3*, *MMP7*, *ITIH1*, *GPRC6A*, *TNNT1*, *FATE1*, and *GIPR* among the top 40 genes. Conversely, MACS-CPAs and EIAs showed only few overexpressed genes (not enough for pathway analysis). ACCs samples presented 472 significantly overexpressed genes, mostly involved

in *Cell Cycle* and *Mitosis*. *DUSP9*, *PAX9*, *SULT1C2*, *PMAIP1*, *DHRS2*, *CPNE4*, *PBK*, and *DSG2* were among the top 40 overexpressed genes. Under-expressed genes in ACCs, like *GNG4*, *GNB3*, *CAMK2B*, *ADCY2*, and *SLC17A7*, were observed mostly in *Transport of small molecules*, *Neuronal System*, and *GPCR ligand binding*.

Relationship between gene expression profile and mutational status

At group 3D PCA (Figure 2A) and individual unsupervised clustering analysis (represented as heatmap in Figure 2B) for protein-coding genes we could classify tumor samples in four clusters: 1) “Cluster 1” (n=32), including MACS-CPAs and EIAs, either with *CTNNB1* mutations (MACS-CPA_CTNNB1, n=9, and EIA_CTNNB1, n=4) or without mutation in a known driver gene (MACS-CPA_none, n=6, and EIA_none, n=4), CS-CPA without known driver mutations (n=7) or with *GNAS* somatic mutations (n=2); 2) “Cluster 2”, comprising mostly CS-CPAs with or without driver mutation (CS-CPA_PRKACA, n=7; CS-CPA_GNAS, n=2; and CS-CPA_none, n=6); 3) CS-CPAs with *CTNNB1* mutations (CS-CPA_CTNNB1, n=2), which show a gene expression pattern closer to ACC than most of the remaining benign tumors, and 4) “ACC”, clustering separately from all other groups.

The single MACS-CPA_GNAS case (MACS-CPA8) belonging to Cluster 2 was a 29-years old female patient, with a small ACA of 2 cm, with not suppressed post-dexamethasone serum cortisol levels (397.3 nmol/L), and 24-hour urine free cortisol (UFC) and ACTH levels within the normal range. The only MACS_no driver in Cluster 2 was a 34-years old male with 3.4 cm adrenal nodule, failed dexamethasone suppression test (100.2 nmol/L), and normal 24-hour urine free cortisol (UFC) and ACTH levels. Both had no clinical signs of CS.

Particularly, cases in Cluster 1 were mostly old male patients having large tumor, whereas those in Cluster 2 were predominantly young patients female with small tumor. The individual 3D PCA plots for protein-coding genes in the entire cohort (52 ACA, 7ACC, 4 NAG) is shown in the Supplementary Figure 2A.

Results from PCA and unsupervised clustering analysis did not change by classifying the adenomas according to the ESE/ENSAT guidelines (12), when considering all the differentially expressed genes. Therefore, further analyses were performed according the Endocrine Society guidelines.

Comment [GDD1]: Indeed, in the individual heatmap 1 of the 2 *ctnnb1*-cpas is included in cluster 2. Conversely, there is a *gnas*-cpa which cluster together with the other *cpa*-*ctnnb1*, which are both closer to ACC. Nonetheless, I would leave it as it is, since 3DPCA and individual heatmap are generated from different data (the whole expression data fro PCA and the top 500 genes for individual heatmap). Is it correct, Silke?

Comment [GDD2]: If we describe this case in detail, should we also describe the MACS-CPA13_none, which belongs to cluster 2? Otherwise, we can remove this detailed description. What do you think about it?

Comment [CR3]: I would add also the MACS-CPA13

Looking at specifically over-expressed genes in the different clusters (boxes in **Figure 2B**), we identified *IGFN1*, *CXCL2*, *DPEP1*, *PITX1*, *SHISA3* and *CXCL14* in Cluster 2. Among Cluster 1, no genes were homogeneously over-expressed. Even if not highlighted in the heatmap, among the most overexpressed genes in Cluster 2, we observed *FATE1* (Fetal and Adult Testis Expressed 1). Targeted data analysis of *FATE1* confirmed that CPAs with *GNAS* or *PRKACA* mutations had significantly higher *FATE1* mRNA expression than MACS-CPAs or EIAs with or without *CTNNB1* mutations ($p \leq 0.001$ and $p \leq 0.001$, respectively) (**Figure 3A**).

Moreover, genes like *PTTG1*, *TOP2A*, *CDK1*, *CCNB1*, *UBE2C*, and *PMAIP1* were highly expressed selectively in ACC (boxes in **Figure 2B**), most of them being involved in cell cycle regulation. Of note, the cluster with CS-CPAs_ *CTNNB1* showed some overexpressed genes in common with ACC (*SPINK13*, *GPX8*, *TMEM26*, *NPTX2*, and *HSPB8*). Targeted data analysis of *TOP2A* (Topoisomerase 2A) and *CCNB1* (cyclin B1) confirmed that ACC had significantly higher mRNA expression than other groups ($p \leq 0.001$ compared to MACS-CPAs or EIAs with *CTNNB1* mutation or without driver mutation and compared to MACS-CPAs and CS-CPAs with *GNAS* or *PRKACA* mutation, and $p \leq 0.05$ CS-CPAs_none, **Figure 3B**). Targeted analysis of *HSPB8* showed higher mRNA levels in both ACC and CS-CPA_ *CTNNB1* compared to other groups (**Figure 3B**). Moreover, as expected, ACC showed lower mRNA expression of *ZNFR3* than all other groups: -1.1 ± 0.89 FKPM - $P < 0.05$ only vs EIA/MACS_ *CTNNB1* (0.1 ± 0.4) and EIA/MACS_no driver (-0.01 ± 0.55) (**Figure 3A**). One CPA with *CTNNB1* mutation as well as few cases of EIA/MACS/CPA without driver mutations showed relatively low *ZNFR3* levels, even if this was observed also in some CPA with known mutations in cAMP/PKA related genes. As shown in **Figure 3A**.

Considering other genes related to the steroidogenesis, *NR5A1* (Nuclear Receptor Subfamily 5 Group A Member 1, known also as steroidogenic factor 1, *SFI*) mRNA expression was significantly lower in ACC in comparison to MACS-CPAs or EIAs with *CTNNB1* mutation or without driver mutations ($p \leq 0.001$ and $p \leq 0.05$, respectively), CS-CPA and MACS-CPA with *GNAS* or *PRKACA* mutation ($p \leq 0.001$) or CS-CPA_none ($p \leq 0.05$) (**Figure 3A**). Of note, CS-CPA with *GNAS* or *PRKACA* mutation showed significantly higher levels of *CYP11A1* and *CYP21A2* than EIA and MACS with *CTNNB1* mutations (**Figure 3A**). On the other side, ACCs showed also significantly lower levels of

Comment [CR4]: I modified this

Comment [GDD5]: I cannot really say that by looking at the heatmap. Silke, could you please confirm these data?

Comment [CR6]: I modified this

Comment [CR7]: Silke/Guido please check P values for both genes.

Comment [CR8]: Add P value for HSPB8

Comment [CR9]: Add P values

several genes encoding for steroidogenic enzymes, including *CYP11B1* and *CYP11B2*, when compared to remaining tissues (data not shown). No other significant differences could be detected among other groups.

The analysis of clinical data among clusters identified by mRNA gene expression showed significant differences in sex ($\chi^2=9.90$, $p=0.02$), age ($\chi^2=11.87$, $p=0.008$), tumor size ($\chi^2=16.44$, $p=0.001$), diagnosis ($\chi^2=113.74$, $p<0.001$), and mutational status ($\chi^2=52.28$, $p<0.001$) (Figure 2A). Particularly, patients in Cluster 1 were older (61% of the cases had ≥ 55 years) and had larger tumors (58% of the cases had tumors ≥ 35 mm) either inactive (all but one inactive tumors grouped in Cluster 1) or associated with MACS (47%). Except two CPAs, all remaining tumors with *CTNNB1* somatic mutations were also included in Cluster 1 (Figure 2B). On the contrary, in Cluster 2, all patients were female, except one, with diagnosis of overt Cushing syndrome in 15/18 cases, among which 77% of cases had <55 years. No driver mutation and mutation in the cAMP/PKA pathway were found in 44% of the cases. We could distinguish three sub-groups in Cluster 1 [1A (n=12), 1B (n=12), and 1C (n=8)] and two in Cluster 2 [2A (n=11) and 2B (n=7)] with slightly different clinical phenotype (Supplementary Figure 2B (30)). In particular, the homogeneous sub-cluster 2A includes all seven CS-CPAs with *PRKACA* mutations and three CS-CPAs without driver mutations, together with one MACS-CPA_none.

Differential exon usage

We investigated the alternative gene spliced transcripts in all our samples. Comparing the individual groups (ACCs, CS-CPAs, EIAs, MACS-CPAs each against NAGs) according both guidelines (12,13), no significant differential exon usage was detected (results not shown).

Long non-coding RNA

Several lncRNAs were differentially represented among the groups of tumors. As shown in Supplementary Figure 4 (30), the PCA and the unsupervised clustering analysis confirmed that

Comment [GDD10]: 19/31 + 1 unknown

Comment [GDD11]: 18/31 + 1 unknown

Comment [GDD12]: From Barbara's email

Comment [GDD13]: This was the original sentence, before changing. Particularly, patients in Cluster 1 were older (68% of cases) and had larger tumors (64% of cases) either inactive (36%) or associated with MACS (64%) and with *CTNNB1* mutations (52%) or unknown driver mutations (48%) (Figure 4 and Supplementary Figure 2B (30)). By looking at the data, there are several differences between the one sent by Barbara and the one reported in new figure 2b (individual heatmap). Am I missing something? Barabara, could you please double check?

Comment [GDD14]: 14/18 + 1 unknown

patients clustered in four groups: ACCs, CS-CPAs_CTNNB1, Inc_Cluster 1 and Inc_Cluster 2 tumors, in analogy to the results of the coding gene expression profile. *IGFL2-ASI* was overrepresented in ACCs, whereas *LINC02539* and *LINC02104* were clearly overrepresented in CS-CPA_CTNNB1 tumors. Additionally, *LINC00400* and *DIO3OS* were overrepresented in Inc_Cluster 2. Both PCA and the heatmap for Inc showed similar results when we classified the tumors according to the ESE/ENSAT guidelines (12).

Gene fusions

All ACC samples provided evidence for gene fusions. The number of high- and medium-confidence fusions per sample was significantly higher in ACCs than ACAs (13.0 ± 6.14 vs 1.67 ± 1.95 , $p < 0.0001$). In contrast, no difference in the number of fusions was observed among CS-CPAs, MACS-CPAs and EIAs ($p = 0.65$; **Supplementary Figure 3A** (30)). By considering only high-confidence gene fusions, ACCs had a significantly higher number of fusions per sample (8.14 ± 5.43) compared to those found in Cluster 1 (0.76 ± 1.13 , $p = 0.0005$) and Cluster 2 (0.72 ± 0.89 , $p = 0.001$), but not to those observed in CS-CPA_CTNNB1 (2.00 ± 2.82 , $p = 0.77$; **Figure 4A**). After restricting the analysis to CS-CPAs only, there was no significant difference in number of high-confidence fusions *per* samples considering the mutational status (**Figure 4B**). However, CS-CPA_CTNNB1 presented a trend towards higher number of high-confidence fusions compared to other subgroups (**Figure 4B**). We did not observe any significant difference in number of gene fusions when we classified the tumor according to the ESE/ENSAT guidelines.

The entire list of high confidence gene fusions detected in the present cohort is reported in the **Supplementary Table 3** (30). No recurrent fusions were detected in genes involved in cAMP/PKA nor Wnt/beta catenin pathways. However, we identified a “private” fusion between the gene coding for the A-kinase anchor protein 13 (*AKAP13*) and the gene coding the phosphodiesterase 8A (*PDE8A*) in one CS-CPA sample without known driver mutations (CS-CPA26; **Figure 4C-D**). This was validated by PCR. Regarding the other investigated fusions (**Supplementary Table 3** (30)), we were able to find 4 out of 6 gene fusions by PCR and Sanger-Sequencing, including *MAPKAPK5-ACAD10*,

NUP155-WDR70, *GTF2B-GBP5* and *SMC6-MSGN*, in the corresponding samples that were positive at RNA-seq. *GADPH*, used as control gene, was found in all the evaluated samples.

Discussion

This is the first study employing RNA-seq to investigate the relationship between transcriptome and the somatic genetic tumor background in a large, well-characterized cohort of benign adrenocortical tumors. We found that transcriptome profiling was able to classify clearly distinct groups that also differed in their hormonal profile and genetic background. Moreover, by applying deep RNA-seq, we identified additional molecular alterations, including gene fusions and differentially expressed lncRNAs associated with the genetic background of the tumors.

The transcriptome profile of benign adrenocortical tumors has been described in a previous publication (19). By applying microarray analysis on a small number of ACAs (n=22), Wilmout Roussel and colleagues identified a specific molecular signature of CS-CPAs, different from that of EIAs or MACS-CPAs, which clustered together. The results of the unsupervised clustering analysis performed in our study partly confirms those findings, showing that most benign adrenocortical tumors could be divided in two groups, named Cluster 1, including EIAs and MACS-CPAs, and Cluster 2, comprising CS-CPAs. The clinical features of patients in the two clusters revealed that patients in Cluster 2 were mostly young females, with small hyperfunctioning tumors, whereas those in Cluster 1 had mostly inactive or “low-functional” large tumors. Therefore, MACS-CPAs and EIAs samples showed a similar transcriptome profile, which was clearly different from that of CS-CPAs and ACCs. Those findings are consistent with previous clinical observations, showing that MACS-CPA tumors seem to be a separate clinical entity from overt CS. Although patients with MACS could present comorbidities associated to mild cortisol hypersecretion, such as risk of cardiovascular events, hypertension, dyslipidemia, diabetes mellitus type 2, obesity, osteoporosis (12,14-18), the long-term data did not provide evidence for an evolution from MACS to overt CS (31). Additionally, recent analysis by liquid-chromatography tandem mass spectrometry in patients with adrenal incidentalomas

showed that the steroid profiling of subjects with EIA showed some similarities with those having MACS-CPA(32,33).

The combined analysis of transcriptome profiling with a tumor whole genome background performed in our cohort allowed us to pinpoint additional important findings. The similar transcriptome profile of MACS-CPA and EIA samples was independent of the presence of *CTNNB1* mutations, which were the most common alterations detected in tumors of this group. Additionally, the similar transcriptome profile among CS-CPAs is consistent with the activation of the cAMP/PKA pathway, independently of the underlying putative mutation. Conversely, the expression profile of the MACS-CPA associated with *GNAS* mutation was similar to that of CS-CPAs.

Among differentially expressed genes, it is worth mentioning *FATE1* as the most over-expressed gene in tumors belonging to Cluster 2, and specifically, in CS-CPAs samples carrying *PRKACA* and *GNAS* somatic mutations. *FATE1* encodes a known transcription factor involved in regulation of apoptosis and cell proliferation, regulated by steroidogenic factor 1 (SF1) at the promoter level (34). Considering the pivotal role of SF1 in steroid producing cells and the interplay between FATE1 and SF1 (35), our data provide additional evidence for the molecular events caused by cAMP/PKA pathway disruption due to *PRKACA* and *GNAS* somatic mutations at a transcriptional level.

Considering that the most evident difference between cluster 1 and cluster 2 is the steroid secreting pattern, we analyzed the expression of several genes involved in steroidogenesis. In particular, we could confirm that the expression of genes encoding for key enzymes such as *CYP11A1* (cytochrome P450 cholesterol monooxygenase) and *CYP21A2* (21-hydroxylase) was significantly higher in *CYP11B2* was lower between CPA carrying mutations in the cAMP-PKA genes compared to both ACC and EIA/MACS-CPA with *CTNNB1* mutations.

Overall, the mRNA expression profile of the subgroup of CS-CPAs associated with *CTNNB1* mutations was more similar to that of ACCs than the remaining CS-CPAs and tumors included in Cluster 1, which were mostly carrying *CTNNB1* mutations. However, when analyzed at an individual level, CS-CPAs with *CTNNB1* mutations showed heterogeneity of the transcriptome profiling at unsupervised clustering. Notably, *TOP2A*, the gene encoding Topoisomerase II Alpha, was among the overexpressed genes in both CS-CPA samples with *CTNNB1* mutations and ACCs, different from the

Comment [CR15]: Guido please add one sentence about the relevance of this.

Comment [CR16]: I deleted this entire part and rewrote it. Those data are consistent with a suppression of the renin-angiotensin system observed in severe hypercortisolism (as it is in tumors carrying cAMP-PKA pathway alterations), uncovered by lower levels of aldosterone, as shown in previous studies (Eisenhofer Clin Chem 2017). On the contrary, we could not demonstrate a difference in *CYP11B1* mRNA levels between CPA and EIAs. The expression level of *CYP11B1* in CPA has been investigated by different methods in several previous studies, with discordant results (Bassett, JCEM, 2005; Enberg, J Endocrinol Invest, 2009; Cao, Int J Urol, 2011; Kubota-Nakayama, Hum Pathol, 2016). In our study, the lack of difference of *CYP11B1* expression in CPA vs EIAs or MACS-CPA may be due by the lack of selective *CYP11B1* immunohistochemistry-driven RNA tissue extraction.

remaining benign tumors. TOP2A is a cell cycle-related protein, involved in cell proliferation by controlling the topologic states of DNA during transcription in several types of cancer and is being studied as a potential druggable target for several anticancer agents (36). Several studies in ACC tissues have shown an overexpression of *TOP2A*, which was tightly correlated with Ki67, a marker of cell proliferation, and poor survival (37,38). Taken together, those observations reinforce the hypothesis proposed in several previous studies of a potential adenoma-carcinoma sequence in a subgroup of adrenocortical tumors. Indeed, studies on animal models and human tissues have shown that alterations of the Wnt-beta catenin pathway, IGF2 signaling and Notch signaling may boost the transformation of an ACA into ACC (39-41). Even though such an event is exceedingly rare (42), the alterations identified in benign adrenocortical tumors described in our study and in previous investigations may represent a preliminary phase in the multi-step potential evolution towards ACC.

The analysis of fusion genes, performed in benign adrenocortical tumors for the first time, showed that CS-CPAs samples carrying *CTNNB1* mutations had a number of high-confidence gene fusions that was set in between ACCs and remaining ACAs belonging to Cluster 1 and 2. Previous studies have identified fusion transcripts, a product of chromosomal rearrangements, as useful markers for several hematological malignancies and epithelial cancers, with important implications in treatment strategies (43). A recent pan-genomic characterization of ACCs has described a few private fusion transcripts, namely *EXOSC10-MTOR* and *MLL-ATP5L*, involving genes with well-known roles in cancer development (44), even though the impact of those events in the pathogenesis of ACC should be confirmed by further targeted studies. The results of the analysis of fusion transcripts in our cohort reinforce the hypothesis that CS-CPAs with *CTNNB1* mutations may have a different biological behavior than the remaining functioning and non-functioning benign adrenocortical tumors. As an additional interesting finding, we could identify the fusion transcript *AKAP13-PDE8A* in one case of CS-CPA without known driver somatic mutations. In a recent study, a similar fusion transcript has been frequently detected in colorectal cancer (45). The predicted PDE8B protein retained the conserved catalytic domain located at the C-terminal region, leading to the hypothesis that the promoter region of AKAP13, which is an A-kinase anchoring protein, might alter cAMP signaling through PDE8B dysregulation (45). Considering the involvement of phosphodiesterases in the

pathogenesis of Cushing syndrome (46), the involvement of *AKAP13-PDE8A* fusion transcript in the pathogenesis of a subset of CS-CPAs without driver mutations at WES is an intriguing mechanism that should be investigated in future functional studies.

Finally, the analysis of lncRNAs provided interesting insights into the molecular signature of adrenocortical tumors. LncRNAs are non-coding RNA transcripts >200 nucleotides in length, involved in epigenetic silencing, regulation of splicing and transcription (47). LncRNAs dysregulation has been described as a potential marker of ACC, with clinical implications. Indeed, recent studies have shown that the lncRNA molecular signature may discriminate ACCs from ACAs and NAGs and identify patients affected by ACC with poor prognosis (48,49). Our lncRNA signature analysis reinforced the results of the transcriptome analysis, showing that tumors could be divided in four clusters, including ACC, CS-CPA with *CTNNB1* mutations, lnc_Cluster 1 (“low-functional” tumors) and lnc_Cluster 2 (“functional tumors”). Indeed, ACC clustered separately from ACA and NAG tissue, confirming previous results (48,49). Notably, CS-CPAs with *CTNNB1* mutations clustered as a separate entity than CS-CPAs with known driver mutations affecting cAMP/PKA pathway and CS-CPAs with no driver mutations, giving further support to the different biological behavior of those tumors.

The main limitation of this study is represented by the rather small number of patients in selected groups of interest, which may limit the interpretation of the results and the possibility to perform correlations between transcriptome and clinical data. Nonetheless, this is the result of a strict selection process that relied on the genetic mutations identified in previous studies (6,9) and availability of high-quality RNA for RNA-seq. To avoid bias in the evaluation of our results, we performed the analysis following the definition of MACS-CPA and CS-CPA according to both the Endocrine Society and the ESE/ENSAT guidelines (12,13). The classification of three samples changes between the two guidelines: 2 MACS-CPAs without driver mutations, which are classified as EIAs according to the newest ESE/ENSAT guidelines, and 1 EIA with *CTNNB1* mutation MACS-CPS that was reclassified as EIA. By comparing the results, we found few differences only in the protein-coding heatmap, where CS-CPAs without known driver mutations do not cluster together with the CS-CPA with *GNAS* or *PRKACA* mutation as previously. In our opinion, this difference depends on the

relatively low number of cases in some diagnostic groups, leading to small changes in gene expression patterns and different clustering. Moreover, gene fusions, PCA, and heatmaps for lnc genes do not change between the tumor classification according the two guidelines. Therefore, we decide to shoe into details the results obtained by classifying the tumor according to the Endocrine Society guidelines, in order to keep the same tumor classification of the previous studies (6,9), allowing a better characterization of the tumor samples.

Moreover, another possible explanation for the mutation-negative cases might be the sensitivity of sequencing methods in detecting mutations with low allele frequencies, as shown in recent studies highlighting the possibility of increasing the detection of mutations by applying targeted immunohistochemistry (IHC)-driven deep-sequencing in primary aldosteronism (Nanba, JCEM, 2018; Nanba, Hypertension, 2019). Nonetheless, our retrospective cohort of fresh-frozen tumors was heterogeneous and an IHC-driven approach would not have been possible. Finally, another limitation of our study is that tumors were dissected macroscopically. Therefore, even though the proceudre was made by expert pathologists, our analysis may carry the chance of sampling error, in analogy to other studies dealing with snap-frozen tissues. More accurate data may derive by an *in situ* methodology, such as laser-capture microdissection. Of note, the strength of our study is that the characterization of transcriptome profile and additional RNA alterations (gene fusions and differential expression of lncRNA) is evaluated in a large cohort of adrenocortical tumors (n=59) and analyzed according to both the functional status and the presence of driver mutations.

Conclusion

In conclusion, our findings suggest that still unrevealed molecular modifications (in cAMP/PKA pathway) might be involved in the pathogenesis of autonomous steroid secretion, some of them depending on the underlying genetic background. At variance, MACS-CPAs seem to be molecularly related to EIAs, independently from the genetic background. Finally, the similarity of molecular alterations between CS-CPAs carrying *CTNNB1* mutations and ACCs may pave the way for further targeted investigation aimed at defining the potential adenoma-carcinoma sequence in adrenocortical tumors.

Comment [CR17]: Barbara please Add reference

Comment [CR18]: Silke please check thi part, do you agree?

Acknowledgements

The authors are grateful to Mrs. Sonja Steinhauer and Ms. Martina Zink for excellent technical support and Ms. Michaela Haaf for coordinating the ENSAT Registry. This work has been carried out with the help of the Interdisciplinary Bank of Biomaterials and Data of the University Hospital of Wuerzburg and the Julius Maximilian University of Wuerzburg (IBDW). The implementation of the IBDW has been supported by the Federal Ministry for Education and Research (Grant number FKZ: 01EY1102).

Data Availability

The raw data have been uploaded into the platform EGA (<https://www.ebi.ac.uk/ega/home>, accession number EGAS00001004533). Further datasets analyzed during the current study are available from the corresponding author on reasonable request.

References

- Di Dalmazi G. Cushing Syndrome-Unilateral Adrenal Adenoma. Encyclopedia of Endocrine Diseases. 2nd ed: Academic Press; 2018:4252.
- Ronchi CL. cAMP/protein kinase A signalling pathway and adrenocortical adenomas. *Curr Opin Endocr Metab Res* 2019;8:15-21.
- Beuschlein F, Fassnacht M, Assie G, Calebiro D, Stratakis CA, Osswald A, Ronchi CL, Wieland T, Sbiera S, Faucz FR, Schaak K, Schmittfull A, Schwarzmayer T, Barreau O, Vezzosi D, Rizk-Rabin M, Zabel U, Szarek E, Salpea P, Forlino A, Vetro A, Zuffardi O, Kisker C, Diener S, Meitinger T, Lohse MJ, Reincke M, Bertherat J, Strom TM, Allolio B. Constitutive activation of PKA catalytic subunit in adrenal Cushing's syndrome. *N Engl J Med*. 2014;370(11):1019-1028.
- Cao Y, He M, Gao Z, Peng Y, Li Y, Li L, Zhou W, Li X, Zhong X, Lei Y, Su T, Wang H, Jiang Y, Yang L, Wei W, Yang X, Jiang X, Liu L, He J, Ye J, Wei Q, Li Y, Wang W, Wang J, Ning G. Activating hotspot L205R mutation in PRKACA and adrenal Cushing's syndrome. *Science*. 2014;344(6186):913-917.
- Goh G, Scholl UI, Healy JM, Choi M, Prasad ML, Nelson-Williams C, Kunstman JW, Korah R, Suttrop AC, Dietrich D, Haase M, Willenberg HS, Stalberg P, Hellman P, Akerstrom G, Bjorklund P, Carling T, Lifton RP. Recurrent activating mutation in PRKACA in cortisol-producing adrenal tumors. *Nat Genet*. 2014;46(6):613-617.
- Di Dalmazi G, Kisker C, Calebiro D, Mannelli M, Canu L, Arnaldi G, Quinkler M, Rayes N, Tabarin A, Laure Jullie M, Mantero F, Rubin B, Waldmann J, Bartsch DK, Pasquali R, Lohse M, Allolio B, Fassnacht M, Beuschlein F, Reincke M. Novel somatic mutations in the catalytic subunit of the protein kinase A as a cause of adrenal

- 554 Cushing's syndrome: a European multicentric study. *J Clin Endocrinol Metab.*
555 2014;99(10):E2093-2100.
- 556 7. Sato Y, Maekawa S, Ishii R, Sanada M, Morikawa T, Shiraishi Y, Yoshida K, Nagata
557 Y, Sato-Otsubo A, Yoshizato T, Suzuki H, Shiozawa Y, Kataoka K, Kon A, Aoki K,
558 Chiba K, Tanaka H, Kume H, Miyano S, Fukayama M, Nureki O, Homma Y, Ogawa
559 S. Recurrent somatic mutations underlie corticotropin-independent Cushing's
560 syndrome. *Science.* 2014;344(6186):917-920.
- 561 8. Espiard S, Knape MJ, Bathon K, Assie G, Rizk-Rabin M, Faillot S, Luscap-Rondof
562 W, Abid D, Guignat L, Calebiro D, Herberg FW, Stratakis CA, Bertherat J.
563 Activating PRKACB somatic mutation in cortisol-producing adenomas. *JCI Insight.*
564 2018;3(8).
- 565 9. Ronchi CL, Di Dalmazi G, Faillot S, Sbiera S, Assie G, Weigand I, Calebiro D,
566 Schwarzmayer T, Appenzeller S, Rubin B, Waldmann J, Scaroni C, Bartsch DK,
567 Mantero F, Mannelli M, Kastelan D, Chiodini I, Bertherat J, Reincke M, Strom TM,
568 Fassnacht M, Beuschlein F, European Network for the Study of Adrenocortical T.
569 Genetic Landscape of Sporadic Unilateral Adrenocortical Adenomas Without
570 PRKACA p.Leu206Arg Mutation. *J Clin Endocrinol Metab.* 2016;101(9):3526-3538.
- 571 10. Tissier F, Cavard C, Groussin L, Perlemoine K, Fumey G, Hagnere AM, Rene-Corail
572 F, Jullian E, Gicquel C, Bertagna X, Vacher-Lavenu MC, Perret C, Bertherat J.
573 Mutations of beta-catenin in adrenocortical tumors: activation of the Wnt signaling
574 pathway is a frequent event in both benign and malignant adrenocortical tumors.
575 *Cancer Res.* 2005;65(17):7622-7627.
- 576 11. Bonnet S, Gaujoux S, Launay P, Baudry C, Chokri I, Ragazzon B, Libe R, Rene-
577 Corail F, Audebourg A, Vacher-Lavenu MC, Groussin L, Bertagna X, Dousset B,
578 Bertherat J, Tissier F. Wnt/beta-catenin pathway activation in adrenocortical
579 adenomas is frequently due to somatic CTNNB1-activating mutations, which are
580 associated with larger and nonsecreting tumors: a study in cortisol-secreting and -
581 nonsecreting tumors. *J Clin Endocrinol Metab.* 2011;96(2):E419-426.
- 582 12. Fassnacht M, Arlt W, Bancos I, Dralle H, Newell-Price J, Sahdev A, Tabarin A,
583 Terzolo M, Tsagarakis S, Dekkers OM. Management of adrenal incidentalomas:
584 European Society of Endocrinology Clinical Practice Guideline in collaboration with
585 the European Network for the Study of Adrenal Tumors. *Eur J Endocrinol.*
586 2016;175(2):G1-G34.
- 587 13. Nieman LK, Biller BM, Findling JW, Newell-Price J, Savage MO, Stewart PM,
588 Montori VM. The diagnosis of Cushing's syndrome: an Endocrine Society Clinical
589 Practice Guideline. *J Clin Endocrinol Metab.* 2008;93(5):1526-1540.
- 590 14. Di Dalmazi G, Vicennati V, Garelli S, Casadio E, Rinaldi E, Giampalma E, Mosconi
591 C, Golfieri R, Paccapelo A, Pagotto U, Pasquali R. Cardiovascular events and
592 mortality in patients with adrenal incidentalomas that are either non-secreting or
593 associated with intermediate phenotype or subclinical Cushing's syndrome: a 15-year
594 retrospective study. *Lancet Diabetes Endocrinol.* 2014;2(5):396-405.
- 595 15. Di Dalmazi G, Vicennati V, Rinaldi E, Morselli-Labate AM, Giampalma E, Mosconi
596 C, Pagotto U, Pasquali R. Progressively increased patterns of subclinical cortisol
597 hypersecretion in adrenal incidentalomas differently predict major metabolic and
598 cardiovascular outcomes: a large cross-sectional study. *Eur J Endocrinol.*
599 2012;166(4):669-677.
- 600 16. Altieri B, Muscogiuri G, Paschou SA, Vryonidou A, Della Casa S, Pontecorvi A,
601 Fassnacht M, Ronchi CL, Newell-Price J. Adrenocortical incidentalomas and bone:
602 from molecular insights to clinical perspectives. *Endocrine.* 2018;62(3):506-516.

- 603 17. Reimondo G, Castellano E, Grosso M, Priotto R, Puglisi S, Pia A, Pellegrino M,
604 Borretta G, Terzolo M. Adrenal Incidentalomas are Tied to Increased Risk of
605 Diabetes: Findings from a Prospective Study. *J Clin Endocrinol Metab.* 2020;105(4).
- 606 18. Athimulam S, Delivanis D, Thomas M, Young WF, Khosla S, Drake MT, Bancos I.
607 The Impact of Mild Autonomous Cortisol Secretion on Bone Turnover Markers. *J*
608 *Clin Endocrinol Metab.* 2020;105(5).
- 609 19. Wilmot Roussel H, Vezzosi D, Rizk-Rabin M, Barreau O, Ragazzon B, Rene-Corail
610 F, de Reynies A, Bertherat J, Assie G. Identification of gene expression profiles
611 associated with cortisol secretion in adrenocortical adenomas. *J Clin Endocrinol*
612 *Metab.* 2013;98(6):E1109-1121.
- 613 20. Lippert J, Appenzeller S, Liang R, Sbiera S, Kircher S, Altieri B, Nanda I, Weigand I,
614 Gehrig A, Steinhauer S, Riemens RJM, Rosenwald A, Muller CR, Kroiss M, Rost S,
615 Fassnacht M, Ronchi CL. Targeted Molecular Analysis in Adrenocortical
616 Carcinomas: A Strategy Toward Improved Personalized Prognostication. *J Clin*
617 *Endocrinol Metab.* 2018;103(12):4511-4523.
- 618 21. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson
619 M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics.*
620 2013;29(1):15-21.
- 621 22. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
622 Durbin R, Genome Project Data Processing S. The Sequence Alignment/Map format
623 and SAMtools. *Bioinformatics.* 2009;25(16):2078-2079.
- 624 23. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg
625 SL, Wold BJ, Pachter L. Transcript assembly and quantification by RNA-Seq reveals
626 unannotated transcripts and isoform switching during cell differentiation. *Nat*
627 *Biotechnol.* 2010;28(5):511-515.
- 628 24. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,
629 Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment
630 analysis: a knowledge-based approach for interpreting genome-wide expression
631 profiles. *Proc Natl Acad Sci U S A.* 2005;102(43):15545-15550.
- 632 25. Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J,
633 Puigserver P, Carlsson E, Ridderstrale M, Laurila E, Houstis N, Daly MJ, Patterson
634 N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES, Hirschhorn JN,
635 Altshuler D, Groop LC. PGC-1alpha-responsive genes involved in oxidative
636 phosphorylation are coordinately downregulated in human diabetes. *Nat Genet.*
637 2003;34(3):267-273.
- 638 26. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
639 reads. *EMBnetjournal.* 2011;17(1):10-12.
- 640 27. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella
641 K, Altshuler D, Gabriel S, Daly M, DePristo MA. The Genome Analysis Toolkit: a
642 MapReduce framework for analyzing next-generation DNA sequencing data. *Genome*
643 *Res.* 2010;20(9):1297-1303.
- 644 28. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants
645 from high-throughput sequencing data. *Nucleic Acids Res.* 2010;38(16):e164.
- 646 29. Altieri B, Sbiera S, Della Casa S, Weigand I, Wild V, Steinhauer S, Fadda G, Kocot
647 A, Bekteshi M, Mambretti EM, Rosenwald A, Pontecorvi A, Fassnacht M, Ronchi
648 CL. Livin/BIRC7 expression as malignancy marker in adrenocortical tumors.
649 *Oncotarget.* 2017;8(6):9323-9338.
- 650 30. Di Dalmazi G, Altieri B, Scholz C, Sbiera S, Luconi M, Walsman J, Kastelan D,
651 Ceccato F, Chiodini I, Arnaldi G, Riester A, Osswald A, Beuschlein F, Sauer S,
652 Fassnacht M, Appenzeller S, Ronchi CL. Data from: RNA-Sequencing and Somatic

- Mutation Status of Adrenocortical Tumors: Novel Pathogenetic insights. OSF - Open Science Framework 2020. Deposited 06/05/2020. https://osf.io/fhqay/?view_only=854a3a749b6d4f17835f54c858726ca1.
31. Elhassan YS, Alahdab F, Prete A, Delivanis DA, Khanna A, Prokop L, Murad MH, O'Reilly MW, Arlt W, Bancos I. Natural History of Adrenal Incidentalomas With and Without Mild Autonomous Cortisol Excess: A Systematic Review and Meta-analysis. *Ann Intern Med*. 2019;171(2):107-116.
 32. Di Dalmazi G, Fanelli F, Mezzullo M, Casadio E, Rinaldi E, Garelli S, Giampalma E, Mosconi C, Golfieri R, Vicennati V, Pagotto U, Pasquali R. Steroid Profiling by LC-MS/MS in Nonsecreting and Subclinical Cortisol-Secreting Adrenocortical Adenomas. *J Clin Endocrinol Metab*. 2015;100(9):3529-3538.
 33. Di Dalmazi G, Fanelli F, Zavatta G, Ricci Bitti S, Mezzullo M, Repaci A, Pelusi C, Gambineri A, Altieri P, Mosconi C, Balacchi C, Golfieri R, Cosentino ER, Borghi C, Vicennati V, Pasquali R, Pagotto U. The Steroid Profile of Adrenal Incidentalomas: Subtyping Subjects With High Cardiovascular Risk. *J Clin Endocrinol Metab*. 2019;104(11):5519-5528.
 34. Doghman M, Karpova T, Rodrigues GA, Arhatte M, De Moura J, Cavalli LR, Virolle V, Barbry P, Zambetti GP, Figueiredo BC, Heckert LL, Lalli E. Increased steroidogenic factor-1 dosage triggers adrenocortical cell proliferation and cancer. *Mol Endocrinol*. 2007;21(12):2968-2987.
 35. Doghman-Bouguerra M, Granatiero V, Sbiera S, Sbiera I, Lacas-Gervais S, Brau F, Fassnacht M, Rizzuto R, Lalli E. FATE1 antagonizes calcium- and drug-induced apoptosis by uncoupling ER and mitochondria. *EMBO Rep*. 2016;17(9):1264-1280.
 36. Lee JH, Berger JM. Cell Cycle-Dependent Control and Roles of DNA Topoisomerase II. *Genes (Basel)*. 2019;10(11).
 37. Jain M, Zhang L, He M, Zhang YQ, Shen M, Kebebew E. TOP2A is overexpressed and is a therapeutic target for adrenocortical carcinoma. *Endocr Relat Cancer*. 2013;20(3):361-370.
 38. Gupta D, Shidham V, Holden J, Layfield L. Value of topoisomerase II alpha, MIB-1, p53, E-cadherin, retinoblastoma gene protein product, and HER-2/neu immunohistochemical expression for the prediction of biologic behavior in adrenocortical neoplasms. *Appl Immunohistochem Mol Morphol*. 2001;9(3):215-221.
 39. Heaton JH, Wood MA, Kim AC, Lima LO, Barlasakar FM, Almeida MQ, Fragoso MC, Kuick R, Lerario AM, Simon DP, Soares IC, Starnes E, Thomas DG, Latronico AC, Giordano TJ, Hammer GD. Progression to adrenocortical tumorigenesis in mice and humans through insulin-like growth factor 2 and beta-catenin. *Am J Pathol*. 2012;181(3):1017-1033.
 40. Drelon C, Berthon A, Ragazzon B, Tissier F, Bandiera R, Sahut-Barnola I, de Joussineau C, Batisse-Lignier M, Lefrancois-Martinez AM, Bertherat J, Martinez A, Val P. Analysis of the role of Igf2 in adrenal tumour development in transgenic mouse models. *PLoS One*. 2012;7(8):e44171.
 41. Ronchi CL, Sbiera S, Leich E, Henzel K, Rosenwald A, Allolio B, Fassnacht M. Single nucleotide polymorphism array profiling of adrenocortical tumors--evidence for an adenoma carcinoma sequence? *PLoS One*. 2013;8(9):e73959.
 42. Belmihoub I, Silvera S, Sibony M, Dousset B, Legmann P, Bertagna X, Bertherat J, Assie G. From benign adrenal incidentaloma to adrenocortical carcinoma: an exceptional random event. *Eur J Endocrinol*. 2017;176(6):K15-K19.
 43. Neckles C, Sundara Rajan S, Caplen NJ. Fusion transcripts: Unexploited vulnerabilities in cancer? *Wiley Interdiscip Rev RNA*. 2020;11(1):e1562.

44. Zheng S, Cherniack AD, Dewal N, Moffitt RA, Danilova L, Murray BA, Lerario AM, Else T, Knijnenburg TA, Ciriello G, Kim S, Assie G, Morozova O, Akbani R, Shih J, Hoadley KA, Choueiri TK, Waldmann J, Mete O, Robertson AG, Wu HT, Raphael BJ, Shao L, Meyerson M, Demeure MJ, Beuschlein F, Gill AJ, Sidhu SB, Almeida MQ, Fragoso M, Cope LM, Kebebew E, Habra MA, Whitsett TG, Bussey KJ, Rainey WE, Asa SL, Bertherat J, Fassnacht M, Wheeler DA, Cancer Genome Atlas Research N, Hammer GD, Giordano TJ, Verhaak RGW. Comprehensive Pan-Genomic Characterization of Adrenocortical Carcinoma. *Cancer Cell*. 2016;30(2):363.
45. Nome T, Thomassen GO, Bruun J, Ahlquist T, Bakken AC, Hoff AM, Rognum T, Nesbakken A, Lorenz S, Sun J, Barros-Silva JD, Lind GE, Myklebost O, Teixeira MR, Meza-Zepeda LA, Lothe RA, Skotheim RI. Common fusion transcripts identified in colorectal cancer cell lines by high-throughput RNA sequencing. *Transl Oncol*. 2013;6(5):546-553.
46. Leal LF, Szarek E, Faucz F, Stratakis CA. Phosphodiesterase 8B and cyclic AMP signaling in the adrenal cortex. *Endocrine*. 2015;50(1):27-31.
47. Morris KV, Mattick JS. The rise of regulatory RNA. *Nat Rev Genet*. 2014;15(6):423-437.
48. Glover AR, Zhao JT, Ip JC, Lee JC, Robinson BG, Gill AJ, Soon PS, Sidhu SB. Long noncoding RNA profiles of adrenocortical cancer can be used to predict recurrence. *Endocr Relat Cancer*. 2015;22(1):99-109.
49. Buishand FO, Liu-Chittenden Y, Fan Y, Tirosh A, Gara SK, Patel D, Meerzaman D, Kebebew E. Adrenocortical tumors have a distinct, long, non-coding RNA expression profile and LINC00271 is downregulated in malignancy. *Surgery*. 2020;167(1):224-232.

Legend for Figures

Figure 1. Flow-chart of the selection process of adrenocortical tumor samples, including diagnosis and criteria for group classification. WES, whole-exome sequencing, ACA, adrenocortical adenomas. NAG, normal adrenal glands. CS-CPA, cortisol-producing adenomas with overt Cushing Syndrome. MACS-CPA, cortisol-producing adenomas with mild autonomous cortisol secretion. EIA, endocrine-inactive adenoma. ACC, adrenocortical carcinoma. Old and new classification of samples according to previous whole-exome sequencing data (Ronchi et al JCEM 2016) and to new variant calling from RNA-seq data, respectively, is reported.

Comment [CR19]: Add reference

Figure 2. Transcriptome profile analysis of adrenocortical tumor samples.

Comment [CR20]: Silke check if everything correct

A) Three-dimensional principal component analysis for protein-coding genes in adrenocortical carcinomas (ACCs) (n=7) and adenomas (ACAs, n=52) divided by functional status according the Endocrine Society guidelines (CS-CPA, cortisol-producing adenoma; MACS-CPA, mild autonomous cortisol secretion; EIA, endocrine-inactive adenoma) as well as mutational status (*PRKACA*, *GNAS* or *CTNNB1* mutations or no known driver mutations).

B) Heatmap of the unsupervised clustering analysis of protein-coding genes in adrenocortical carcinomas (ACCs) (n=7) and adenomas (ACAs, n=52) divided by functional status according the Endocrine Society guidelines (CS-CPA, cortisol-producing adenoma; MACS-CPA, mild autonomous cortisol secretion; EIA, endocrine-inactive adenoma) as well mutational status (*PRKACA*, *GNAS* or *CTNNB1* mutations or no known driver mutations). Individual IDs can be seen at the bottom of the Figure. Different clusters are recognized and highlighted as Cluster 1 (blue), divided in 1A (n=12), 1B (n=12) and 1C (n=8), and Cluster 2 (pink), divided in 2A (n=11) and 2B (n=7). 2 CS-CPAs (one with *CTNNB1* and one without driver mutations) were not included in the previous clusters. Statistical analysis to evaluate the differences on sex, age (cutoff 50-years old), tumor size (cutoff 3.5 cm), diagnosis (EIA, MACS-CPA, CS-CPA, ACC) and mutation status (no driver vs mutation of cAMP/PKA pathway vs *CTNNB1* mutation vs other driver mutation) among the clusters was performed by Chi-square test. A *p* value<0.05 was considered statistically significant.

In box a) protein-coding genes which are upregulated in ACC are enlarged. In box b) and c) protein coding genes upregulated in Cluster2. The analysis was performed including the top 500 differentially expressed protein-coding genes using log₂ fold changes in principal component 1, 2, and 3. Color key bar on the top right corner shows log₂(FC).

Figure 3. Targeted data analysis evaluating the mRNA expression (log₂FC) of A) steroidogenic genes: *FATE1*, *NR5A1*, *CYP11A1*, *CYP21A2*, and B) cell cycle-related genes: *TOP2A*, *ZNRF3*, *HSPB8*, *CCNB1*. The samples were grouped in adrenocortical carcinomas (ACC, n=7) and adrenocortical adenomas (ACAs, n=52) divided by functional and mutational status: cortisol-producing adenomas (CPAs) associated with Cushing syndrome (CS-CPAs) with *CTNNB1* mutation (n=2); mild autonomous cortisol secretion (MACS)-CPAs or endocrine-inactive adenomas (EIAs) with *CTNNB1* or without driver mutations (n=13 and n=12, respectively); CS-CPAs or MACS-CPAs with *PRKACA* or *GNAS* mutations (n=9); CS-CPA without driver mutations (n=13). Functional status was evaluated according the Endocrine Society guidelines. A non-parametric Kruskal-Wallis test was used significant to detect differences in the distributions of values among groups. The Dunn post-hoc analysis was performed using the dunnTest function of R's FSA package, v0.8.23.

Figure 4. Analysis of gene fusions by RNA-seq in adrenocortical tumours.

A) High confidence gene fusions recognized at transcriptome profile analysis in ACC compared to those found in cortisol-producing adenoma (CPA) associated to Cushing syndrome (CS-CPA) with *CTNNB1* mutation (CS-CPA_*CTNNB1*), and those in Cluster 2 and Cluster 1. ACCs had a significantly higher number of high-confidence gene fusions compared to the other groups. B) High confidence gene fusions among CS-CPAs according to the mutational status, showing that CS-CPA_*CTNNB1* had a trend toward a higher number of high-confidence fusions (mean 2.00±2.82) than the other subgroups (mean 1.07±1.04 in CS-CPA_no driver, 0.43±0.53 in CS-CPA_*PRKACA*). To note, no one of the 4 CS-CPA_*GNAS* had highe-confidence gene fusions. In both A) and B) the

783 histograms represent the number of fusions $\pm$ SEM. Statistical analysis was performed by non-
784 parametric test with Dunn's multiple comparison test.
785 C) Specific breakpoint of the detected A-kinase anchor protein 13 (*AKAP13*)-phosphodiesterase 8A
786 (*PDE8A*) fusion verified by Sanger sequencing. D) Predicting fusion by Arriba software including the
787 coding sequence of the *AKAP13* gene and an intronic sequence of the *PDE8A* gene.
788

789 **Table 1. Clinical parameters, hormonal and genetic data of patients with adrenocortical tumors.**

	EIA (n=9)	MACS-CPA (n=17)	CS-CPA (n=26)	ACC (n=7)
Clinical and hormonal parameters				
Age, years	59.0 (55.0-71.5)	57 (47-62)	41.5 (35.0-47.3)**	51.0 (47.0-60.0)
Females, n (%)	4 (44%)	13 (76%)	25 (96%)**	4 (57%)
Tumor size, cm	3.3 (2.1-5.2)	4.0 (3.4-4.5)	3.3 (2.8-3.6)	11.0 (8.5-15.0)
Post-dexamethasone cortisol, nmol/L [n.a.]	35.9 (24.8-63.4)	84.1 (75.2-168.3)**	449.7 (373.8-551.8)**	59.3 (36.5-539.4) [3]
24-h UFC, nmol/day [n.a.]	n.a.	568 (174-1060) [4]	1327 (682-2118)* [12]	105.3 (74.2-248.8) [2]
ACTH, pmol/L [n.a.]	5.1 (1.9-7.3) [4]	0.8 (0.4-1.8)* [1]	0.7 (0.2-1.1)** [11]	14.1 (3.0-33.3) [1]
Ki67 index, % - median (range)	n.appl.	n.appl.	n.appl.	10 (2-15)
Genetic data				
Number of somatic mutations [§] - median (range)	4 (1-15)	2 (0-17)	4 (0-40)	0 (0-1) [2]
Samples with known driver mutations ^{§†} , n (%)	4 (44%) <i>CTNNB1</i> n=4	11 (65%) <i>CTNNB1</i> n=10 <i>GNAS</i> n=1	12 (46%) <i>PRKACA</i> n=6 <i>GNAS</i> n=3 <i>CTNNB1</i> n=3	2 (40%) <i>TP53</i> n=1 <i>ZNRF3</i> n=1

Data are expressed as median and interquartile ranges in parentheses, if not otherwise specified. CPA-CS: cortisol-producing adenoma associated with overt Cushing syndrome, CPA-MACS: cortisol-producing adenoma associated with mild autonomous cortisol secretions, EIA: endocrine inactive adenoma, ACC: adrenocortical carcinoma, n.a.: number of patients with not available, n.appl., not applicable, UFC: urinary free cortisol.

Overt CS and MACS were defined according to the Endocrine Society guidelines (13).

[§]Number of somatic mutations were counted according to previously published data deriving from whole-exome sequencing for ACAs (9) and from targeted next generation sequencing of 160 genes for 5/7 ACCs (20), whereas for 2 ACCs this information was not available. Therefore, the number of mutations were not comparable between ACAs and ACCs. The use of a panel of 160 genes and the inclusion of only early-stage/low aggressive ACCs in this study, explains the very low number of found mutations.

[†]According to targeted sequencing of *PRKACA* exon 7 hot-spot mutations (6) or targeted next-generation sequencing for ACC (20).

* $P < 0.05$ vs EIA (Kruskal-Wallis test followed by Bonferroni); for 24-h UFC MACS-CPA vs CS-CPA (Mann-Whitney test). ** $P < 0.005$ vs EIA (Kruskal-Wallis test followed by Bonferroni).

792 **Table 2. Biological pathways in protein-coding genes differentially expressed in**
 793 **adrenocortical tumors.**

Under-expressed genes			Over-expressed genes	
	Total n genes	Pathways	Total n genes	Pathways
CS-CPA (n=26)	125	1) Regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)	67	1) GPCR ligand binding 2) G alpha (q) signalling 3) Peptide ligand-binding receptors 4) Class A/1 (Rhodopsin-like receptors) 5) Signalling by GPCR
MACS-CPA (n=17)	338	1) Neuronal system 2) Muscle contraction 3) Transmission across Chemical Synapses 4) Cardiac conduction 5) Potassium channel	7	No pathways significantly represented among the overexpressed genes
EIA (n=9)	246	1) Neuronal system 2) Transmission across chemical synapses 3) Neurotransmitter receptor binding	1	No pathways significantly represented among the overexpressed genes
ACC (n=7)	591	1) Transport of small molecules 2) Neuronal System 3) GPCR ligand binding 4) Signalling by GPCR 5) Class B/2 (Secretin family receptors)	471	1) Cell cycle 2) Mitosis 3) Rho GTPase 4) Glucuronidation 5) Kinesins

CPA-CS: cortisol-producing adenoma associated with overt Cushing syndrome, CPA-MACS: cortisol-producing adenoma associated with mild autonomous cortisol secretions, EIA: endocrine inactive adenoma, ACC: adrenocortical carcinoma. Under-expressed genes by $\log_2$ fold change values >2 and over-expressed genes by $\log_2$ fold change values <-2 . Pathway analysis was performed by GSEA (Gene sets from Reactome and KEGGS).