Temporal trends in radiometrically dated sediment cores from English lakes show polybrominated diphenyl ethers correlate with brominated but not mixed bromo/chloro dioxins and furans

Peters, Leon I.; Rose, Neil L.; Yang, Handong; Klánová, Jana; Moehring, Thomas; Harrad, Stuart

DOI:

[10.1016/j.scitotenv.2020.143118](https://doi.org/10.1016/j.scitotenv.2020.143118)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Peters, LI, Rose, NL, Yang, H, Klánová, J, Moehring, T & Harrad, S 2021, 'Temporal trends in radiometrically dated sediment cores from English lakes show polybrominated diphenyl ethers correlate with brominated but not mixed bromo/chloro dioxins and furans', *Science of the Total Environment*, vol. 762, 143118.
<https://doi.org/10.1016/j.scitotenv.2020.143118>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Download date: 26. Apr. 2023

**Temporal Trends in Radiometrically Dated Sediment Cores
from English Lakes Show Polybrominated Diphenyl Ethers
Correlate with Brominated but not Mixed Bromo/Chloro
Dioxins and Furans**

Leon I. Peters^{a#}, Neil L. Rose^b, Handong Yang^b, Jana Klánová^c, Thomas Mochring^d, Stuart
Harrad^{a*}

^aSchool of Geography, Earth, and Environmental Sciences, University of Birmingham,
Birmingham, B15 2TT, U.K.

^bEnvironmental Change Research Centre, Department of Geography, University College London,
London, WC1E 6BT, U.K.

^cRECETOX, Masaryk University, Brno, Czech Republic

^dThermo Fisher Scientific (GmbH) Bremen, Hanna-Kunath-Str. 11, 28199 Bremen, Germany

[#]Deceased

^{*}Email address: S.J.Harrad@bham.ac.uk

ABSTRACT

This paper reports concentrations between ~1950 and present, of polybrominated diphenyl ethers (PBDEs) and polybrominated dibenzo-*p*-dioxins and furans (PBDD/Fs), in radiometrically-dated sediment cores from three English lakes. Mixed bromo/chloro dibenzo-*p*-dioxins and furans (PXDD/Fs) were measured in two of the same lakes. Concentrations of PXDD/Fs decreased over time to the present. To our knowledge, this is the first report of temporal trends of PXDD/Fs in the environment. In contrast, concentrations of PBDEs increased towards the present and were significantly correlated ($R = 0.88-0.98$; $p < 0.05$) with concentrations of PBDFs in all three lakes. These observations suggest that the sources of PXDD/Fs are not related to PBDEs and differ from those of PBDFs. We also report for the first time the presence of octabromodibenzofuran (OBDF) in the two most recent core slices at one lake. The source of OBDF in these samples is unclear. While OBDF has been reported previously as a significant contaminant of some commercial formulations of Deca-BDE, it is also present in Octa-BDE products and in emissions from a variety of combustion activities. Overall, while the positive correlation between PBDEs and PBDFs suggests increased use of PBDEs has contributed substantially to environmental contamination with PBDFs; examination of PBDF homologue patterns implies emissions from combustion activities are likely also important.

36 **Keywords**

37 PBDD/Fs; PXDD/Fs; PBDEs; Time Trends; Sources; Lakes

Introduction

Polybrominated diphenyl ethers (PBDEs) are industrial chemicals that have found extensive global use as flame retardants added to a wide range of consumer items such as electrical and electronic equipment and furniture fabrics and foams. As a consequence of their environmental persistence, potential for long-range atmospheric transport and bioaccumulation, and adverse effects on humans and/or wildlife, PBDEs are listed under the United Nations Environment Program's Stockholm Convention on Persistent Organic Pollutants (POPs). Moreover, manufacture and new use of PBDEs is either banned or restricted by many jurisdictions. Against this backdrop, there is a clear need to evaluate the efficacy of such actions; for example, by monitoring temporal trends in environmental contamination. Several studies worldwide have previously delineated the increase in concentrations of PBDEs in the environment from the onset of their widespread use in the 1980s to mid-2000s when the first restrictions on them were introduced (Kohler et al, 2008; Zegers et al, 2003; Vane et al, 2010; Webster et al, 2008). Subsequent trends up to the present, point to concentrations levelling off and even declining as actions to eliminate PBDEs take effect (Yang et al, 2016). One concern with respect to PBDEs is that commercial PBDE products have been shown to contain polybrominated dibenzo-*p*-dioxins and furans (PBDD/Fs) as contaminants at levels that amount to a substantial mass (0.43 - 2.2 t globally) of PBDD/Fs (Hanari et al, 2006; Ren et al, 2011). This is consistent with a recent report that temporal trends of PBDFs correlate with those of PBDEs in sediment cores from Tokyo Bay (Goto et al, 2017). This is concerning owing to evidence of the toxicity of PBDFs that is considered by the WHO to be on a par with that of their chlorinated analogs (Van den Berg et al, 2013). Moreover, there exists emerging concern about environmental contamination with mixed bromo-/chloro-dibenzo-*p*-dioxins and furans (PXDD/Fs) (Wall et al, 2015). Although little is known about the toxicity of PXDD/Fs, their structural similarity to

PBDD/Fs means that establishing the level at which they are present in the environment and elucidating their sources is desirable. Studies exist that demonstrate that combustion activities such as iron ore sintering and waste incineration are sources of PXDD/Fs (Chatkittikunwong and Creaser, 1994; Weber et al, 2003), with their presence in air in Japan (Hayakawa et al, 2004) as well as soil impacted by a recycling plant fire (Myers et al, 2012) and by informal e-waste handling reported (Leung et al, 2007; Ma et al, 2008; Tue et al, 2013; Yu et al, 2008; Zennegg et al, 2009). Currently however, very few data exist on concentrations of PXDD/Fs in sediments. Concentrations of Σ PXDD/Fs were between 0.03-0.1 ng/g dry weight in freshwater sediment taken from a former chlor-alkali plant in the USA, with monobromoheptachloro dibenzo-*p*-dioxin the only congener detected (Kannan et al, 1998). In Osaka Bay, Japan, a wider range of PXDD/Fs were detected in marine sediment, most prominently 2-Br-3,7,8-ClCDD detected between 0.84-6.5 pg/g dry weight (Ohta et al, 2002), while concentrations of monobromo-polychlorinated dibenzo-*p*-dioxins/dibenzofurans which ranged between below detection limit and 1.8 ng/g dry weight were reported for marine surficial sediments from Hong Kong and Korea (Terauchi et al, 2009).

In this study we exploit the fact that sediment deposited in lakes over time can provide a reliable record of contaminant inputs into lacustrine systems. We thus collected sediment cores from three English lakes for which data on concentrations of PBDEs, hexabromocyclododecane (HBCDD), and polychlorinated biphenyls (PCBs) have been measured previously (Yang et al, 2016) and used radiometric techniques to assign dates to core slices representing different depths. Concentrations of PBDEs, PBDD/Fs, and PXDD/Fs were measured in individual core slices and the data used to test the hypothesis that these contaminant classes will display similar temporal trends.

2. Experimental

2.1 Sampling locations

A map of our sampling locations is given as supplementary data (Figure SD-1), with additional information about individual sites supplied in Table SD-1. Locations studied were 3 of the same seepage lakes for which we have previously reported temporal trends in concentrations of PBDEs, HBCDD, and PCBs in sediment cores (Yang et al, 2016). These were: Edgbaston Pool, Holt Hall Lake, and Wake Valley Pond. These lakes were selected from locations with a range of population densities as a surrogate indicator of anthropogenic inputs. One sediment core from each lake was collected between 23rd – 26th June 2015, covering sedimentation from at least ~1950 to the date of sampling. Consistent with our previous study of these lakes (Yang et al, 2016), each core was collected from a flat area of the lake basin near the deepest point.

2.2 Sampling and sectioning of lake sediment cores

Cores were sampled from a flat area close to the maximum depth at each site and were collected from a purpose built pontoon to a depth of between 0.75-0.95 m below the benthic surface using a large diameter sediment core apparatus ('Big-Ben') (Patmore et al, 2014). The sediment corer, piston and core covers were all thoroughly decontaminated with hexane before and after use. The 'Big-Ben' corer having an internal diameter of 140 mm, is considerably larger than conventional piston corers (typically 50 - 80 mm i.d.) and provides sediment cores with a cross-sectional area of 154 cm² resulting in far greater sample for analysis, and facilitating the analysis of PBDD/Fs and PXDD/Fs at the anticipated ultra-trace levels. Sediment cores were extruded vertically in the field at 10 mm intervals, with each sample stored at -20° C in individually sealed Whirl-PackTM sampling bags until extraction analysis. Sample contamination derived from use of Whirl-PackTM sampling bags manufactured from low density polyethylene (LDPE) was controlled for with the use of sampling blanks, which consisted of 30 g pre- cleaned Na₂SO₄ spiked with 10 µL

¹³C₁₂-BDE-138. Sampling control blanks were opened to the atmosphere for approximately 30 min to allow the sampling spike solvent to evaporate, before being homogenised and sealed until analysis. Three sampling blanks were employed per site and treated analogously to sediment samples, including extraction and analysis. Recoveries of ¹³C₁₂-BDE-138 ranged between 50 - 110 % with a mean of 85.4 ± 35.2 % (Mean ± 1SD) and all cases (n= 9) yielded BFR concentrations below limits of quantification confirming that the sample collection procedure did not contribute to sediment contamination with PBDEs.

Each sediment sample was divided into two during the core-sectioning procedure: two thirds kept for analysis of our target contaminants; with the remaining one third used for radiometric dating and determination of sediment water and total organic carbon (TOC) content. The latter two metrics were determined gravimetrically by mass loss from a 2 g (wet weight) sample, oven dried at 105 °C for 3 h to determine water content and a further 2 h at 550 °C to determine TOC by loss-on-ignition. After freeze-drying, each 10 mm core slice was homogenized. Thereafter, based on information from our radiometric dating results, we combined 10 mm core slices from the same core to generate a series of pooled core slice samples that each represented around five years of sedimented material. In total, 31 such pooled samples were prepared for analysis. A further core slice was analyzed from the bottom of the core taken from each of our lakes, to provide an indication of concentrations of our target contaminants prior to ~1950.

Analytical methods

Details of methods used to radiometrically date (using ²¹⁰Pb) sediment core slices are given as SD, with sediment chronologies and sedimentation rates provided as Tables SD-2-SD-4 and Figures SD-2-SD-7. For determination of concentrations of target contaminants in sediment samples, 5 g each of both hydromatrix and sodium sulfate (both pre-cleaned) were loaded into pre-cleaned

stainless steel extraction cells, along with an aliquot of freeze-dried and homogenized sediment core slice (typically 5 g, accurately weighed). This was treated with ^{13}C -labeled internal standards (details below) and subjected to pressurized liquid extraction using an ASE-350 (Dionex, CA). Extraction was first with hexane:dichloromethane (3:2, v/v; 2 cycles), followed by toluene (2 cycles) at 90°C, 1500 psi and hold time= 5 min. Following concentration to ~ 1 mL using a Turbovap solvent evaporator, crude sediment extracts were purified prior to instrumental analysis via elution through a sequence of acid silica and activated carbon columns (Cape Technologies, Maine, USA). This yielded two concentrated extracts (F1b and F2b) for instrumental analysis of PBDEs (F1b) and PBDD/Fs + PXDD/Fs (F2b) for each sediment core slice. Detailed descriptions of the complex procedures used to purify and fractionate sediment core extracts prior to instrumental analysis are provided as SD (Figures SD-8-10 and accompanying text).

Commercially available standards were used for the analysis of BFRs. These comprised native PBDEs and $^{13}\text{C}_{12}$ -BDEs 28, 47, 99, 100, 153, 154, 183, 207, and 209 (Wellington Laboratories) and native and $^{13}\text{C}_{12}$ - analogues of the following PBDD/Fs (Cambridge Isotope Laboratories): 2,3,7,8-TBDD, 1,2,3,7,8-PeBDD, 1,2,3,4,7,8-HxBDD, 1,2,3,6,7,8-HxBDD, 1,2,3,7,8,9-HxBDD, 1,2,3,4,6,7,8-HpBDD, OBDD, 2,3,7,8-TBDF, 2,4,6,8-TBDF, 1,2,3,7,8-PeBDF, 2,3,4,7,8-PBDF, 1,2,3,4,7,8-HxBDF, 1,2,3,4,6,7,8-HpBDF, and OBDF. The following native standards of PXDD/Fs were kindly provided by Dr. Alwyn Fernandes, FERA, UK: 2-Br-7,8-CDD, 2-Br-3,7,8-CDD, 2,3-Br-7,8-CDD, 1-Br-2,3,7,8-CDD+2-Br-1,3,7,8-CDD, 2-Br-3,6,7,8,9-CDD, 2-Br-7,8-CDF, 2-Br-6,7,8-CDF+3-Br,2,7,8-CDF, and 1-Br-2,3,7,8-CDF. In the absence of $^{13}\text{C}_{12}$ -labeled PXDD/Fs, $^{13}\text{C}_{12}$ -2,4,6,8-TBDF was used as internal (surrogate) standard for quantification of PBDD/Fs.

Instrumental Analysis

Quantification of target compounds and congener groups was conducted on a Thermo Scientific GC Q Exactive GC/orbitrap MS. Concentrations of PBDEs were determined via injection of 2 μ L of F1 onto a Thermo Fisher Scientific Trace Gold Dioxin capillary column (12 m x 0.25 mm x 0.1 μ m), with the GC-MS operated in selected ion monitoring (SIM) mode. Concentrations of PBDD/Fs and PXDD/Fs were determined by injecting 2 μ L of F2 onto the same GC column, with the GC-MS operated in full scan mode. For both F1 and F2, extracts were injected via a PTV injector operating in splitless mode. A programmed temperature of injection of 120 $^{\circ}$ C followed by a ramp profile from 150 $^{\circ}$ C to 320 $^{\circ}$ C at 14.5 $^{\circ}$ C/sec was deployed. A constant He flow of 1.3 mL/min was deployed with a temperature program of: 120 $^{\circ}$ C held for 3 min before ramping to 250 $^{\circ}$ C at 6.5 $^{\circ}$ C/min, followed by a second ramp from 250 $^{\circ}$ C to 305 $^{\circ}$ C at 8 $^{\circ}$ C/min held for 7 min. The orbitrap MS was operated at 60,000 mass resolution, which facilitated confirmation of analyte identity via accurate mass. In all cases, where a standard for a given compound was included in our calibration standard, identification of that compound was via comparison of retention time to the standard, while quantification of that compound was made using its response factor relative to the appropriate internal (surrogate) standard. For PBDD/F and PXDD/F compounds for which we did not have standards, quantification was achieved using an averaged relative response factor for compounds of the same homologue group. For example, hexabromo-PBDFs were quantified using the relative response factor for 1,2,3,4,7,8-HxBDF. Full details of mass spectrometric parameters are provided as SD.

QA/QC

Recoveries for all target analyte internal standards ranged between 50 and 110 %, with the exception of $^{13}\text{C}_{12}$ -OBDD and $^{13}\text{C}_{12}$ -OBDF for which recoveries in some samples were as low as 40%. Method accuracy was assessed for PBDEs by replicate (n=7) analysis of NIST SRM1944

(New York/ New Jersey Waterway Sediment). This revealed good agreement between concentrations measured in this study with the certified values reported by NIST. Full details are provided in SD (Table SD-5). While certified or indicative values have not previously been reported for PBDD/Fs and PXDD/Fs, they are also provided as SD (Tables SD-6 and SD-7) for comparison with future studies.

Statistical methods

All statistics in this study were conducted using IBM SPSS Statistics 21, and Microsoft Excel 2007.

Results and discussion

Concentrations of PBDEs

Concentrations of all target PBDEs and PBDD/Fs in each individual core slice from each sampling location are provided as supporting data (Tables SD-8 – SD-13). Table 1 provides a summary of concentrations of Σ PBDEs detected in this study together with those reported in selected relevant previous studies for comparison. Most pertinently, concentrations of Σ PBDEs reported for surficial sediments for our three lakes match closely with those reported previously by our group for surficial sediments (albeit dating from 6-7 years earlier) for the same lakes (Yang et al, 2016). While the lake situated in the local authority with the highest population density (Edgbaston Pool) displayed the greatest concentrations of Σ PBDEs, concentrations at Wake Valley Pond and Holt Hall Lake are similar despite Wake Valley Pond being located in a more densely populated local authority area (Table SD-1 – i.e. Edgbaston Pool 2,500 people/km² > Wake Valley Pond 250-499 people/km² > Holt Hall Lake 100-249 people/km²). This is partly consistent with previous findings showing higher concentrations of PBDEs in air and soil from more densely populated areas due to

emissions from the built environment (Harrad and Hunter, 2006). Concentrations of Σ PBDEs in this study at Edgbaston Pool ca 3 km from the center of the city of Birmingham are close to the average reported for surficial sediments collected in 2011 from 45 locations along the River Thames in the UK (Ganci et al, 2019). By comparison, those at Holt Hall Lake and Wake Valley Pond are around an order of magnitude lower than at Edgbaston Pool. When compared to Σ PBDE concentrations in surficial sediments collected in 2002-2003 as part of two studies conducted in industrial areas of the River Clyde in Scotland (Vane et al, 2010; Webster et al, 2008), concentrations at all three lakes in our study are substantially lower.

Relative Abundance of PBDE Congeners

The PBDE congener pattern in all our samples is dominated by BDE-209. This is consistent with industry figures that in Europe, Deca-BDE production far exceeded that of both Penta- and Octa-BDE (BSEF, 2003). It is also in line with previous reports for the same lakes (Yang et al, 2016), with sediment cores from the Clyde Estuary in Scotland, UK (Vane et al, 2010; Webster et al, 2008), and with surficial sediments from the River Thames running through London, UK (Ganci et al, 2019). Consistent with the Clyde Estuary, the relative contribution of the nona-BDEs-206, -207, and -208 to Σ PBDEs in this study exceeds that observed in commercial formulations of Deca-BDE. Specifically, in the top three core slices from all three locations, Σ nona-BDEs comprises 8-27% of Σ BDEs. This compares to 2.5 and 9.3% respectively in the commercial Deca-BDE formulations Saytex 102E and Bromkal 82-0DE (La Guardia et al, 2006). While the contribution of Σ nona-BDEs to Σ BDEs is higher in the commercial Octa-BDE products DE-79 and Bromkal 79-8DE at 12.1 and 18.9% respectively (La Guardia et al, 2006); the contribution of the main indicator congener for Octa-BDE (BDE-183) to Σ PBDEs in our sediments is much lower (0.45-2.1% in the top three slices at all lakes) than in DE-79 and Bromkal 79-8DE (42% and 12.6%

respectively). On this evidence, the presence of the three nona-BDEs in this study seem most likely attributable to debromination of BDE-209. This is supported by the fact that BDE-209 and Σ nona-BDEs are positively correlated ($p < 0.05$ at Edgbaston Pool; $p < 0.1$ at Wake Valley Pond and Holt Hall Lake) at each lake studied here. Similar conclusions were drawn based on the observation that nona-BDEs were the second most abundant homologue group after BDE-209 in river sediments from China (Mai et al, 2005; Zhang et al, 2009). However, we note a report of elevated abundances of BDE-208 relative to BDE-209 in TV/PC display casings, and PC components sampled in South China. In this study the authors attributed to the decomposition of higher brominated PBDEs during the process of manufacturing use and/or recycling of PBDE-containing products (Chen et al., 2010). Specifically, the ratios of average concentrations of BDE-208: average concentrations of BDE-209 in TV and PC housing were 5.4% and 2.8% respectively. By comparison, BDE-208:BDE-209 ratios in the uppermost 4 core slices in our study ranged from 0.2% to 0.6% at Edgbaston Pool; 1.3% to 2.0% at Wake Valley Pond; and 1.3% to 9.5% at Holt Hall Lake. It is thus possible that the nona-BDEs detected in our study have arisen at least in part to emissions from treated products as well as any post-emission debromination (Law et al., 2008).

Temporal Trends in PBDEs in English Lake Sediments

Figure 1 plots the temporal trends in concentrations of Σ PBDEs at all three locations. Given the predominant contribution of BDE-209 to Σ PBDEs, the trends in Σ PBDEs reflect those of BDE-209. Importantly, BDE-209 concentrations in the cores from Edgbaston Pool and Wake Valley Pond do not peak in the uppermost core slice (2015); instead the highest concentrations are found in the slice immediately below (2012 and 2009 for Edgbaston Pool and Wake Valley Pond respectively). The situation differs for Holt Hall Lake, where the highest BDE-209 concentration is

in the surficial sediment (2015). Overall, this suggests that while the full impact of recent restrictions on the manufacture and use of Deca-BDE has yet to manifest itself at the lakes studied here; there are tentatively encouraging signs that contamination is levelling off.

Concentrations, Relative Abundance, and Possible Sources of PBDD/Fs in English Lake Sediments

PBDD/Fs were detected in all samples in this study at concentrations two orders of magnitude below those of PBDEs. Table 1 compares Σ PBDD/F concentrations in this study compared to those reported in a variety of previous studies from other locations. Concentrations in sediment core slices from the less urbanised Wake Valley Pond and Holt Hall Lake are in line with those reported for Swedish lakes and rivers (Hagberg et al, 2005; Lundstedt, 2016). However, those at Edgbaston Pool exceed those reported elsewhere. Similar to the situation for PBDEs, while the lake situated in the local authority with the highest population density (Edgbaston Pool) has noticeably the highest Σ PBDD/F concentrations; levels of these contaminants are similar at Wake Valley Pond and Holt Hall Lake despite the former being located in a more densely populated local authority area. PBDFs were overwhelmingly dominant, with OBDD the only PBDD detected and that only in the uppermost two layers in the cores taken at Wake Valley Pond and Holt Hall Lake. This is consistent with previous observations for surficial sediments and atmospheric deposition (Goto et al, 2017; Hayakawa et al, 2004), as well as in emissions from waste incinerators (Wang et al, 2010), and iron ore sintering (Drage et al, 2014). It is also pertinent to note that while OBDD and to a far lesser extent penta- and tetra-BDDs have been detected in commercial PBDE formulations (Ren et al, 2011), PBDFs – especially OBDF and 1,2,3,4,6,7,8-HpBDF - were in comparison detected in such products at far higher concentrations (Hanari et al, 2006; Ren et al, 2011).

Figure 2 shows the homologue profiles of PBDFs in sediment core slices from each lake. It is interesting to note the spatial variation in these profiles, for example TBDFs are far less abundant at Edgbaston Pool than at the other two lakes. A striking observation is that OBDF was detected at Wake Valley Pond - albeit only in the two uppermost core slices (2015 and 2009). To our knowledge, this is the first report of OBDF in sediments. The low detection frequency of OBDF in sediments is perhaps surprising given previous data that reports OBDF to be an impurity in both Octa-BDE and Deca-BDE commercial formulations (Hanari et al, 2006; Ren et al, 2011). However, this may be because OBDF has been reported to be easily debrominated (Goto et al, 2017). Overall, the PBDF homologue pattern in our sediment cores (expressed as a percentage of Σ PBDFs) more closely resembles that reported for the Octa-BDE formulation which comprises 0.3 % TBDFs, 1.8 % PeBDFs, 44.9 % HxBDFs, 27.1 % HpBDFs, and 25.8 % OBDF, than the Deca-BDE formulation which comprises 0.1% for each of TBDFs, PeBDFs, and HxBDFs, 3.6% for HpBDFs, and 96.1% OBDF. While this might suggest the Octa-BDE commercial formulation is a more significant source of the PBDFs observed in our sediments than the Deca-BDE product; as noted above, the major PBDE formulation used in the UK was Deca-BDE. Moreover, we note that concentrations of Σ PBDFs are more strongly correlated with those of BDE-209 ($R=0.96$, $p<0.05$ and $R=0.85$, $p<0.1$) than those of BDE-183 at both Edgbaston Pool and Wake Valley Pond ($R=0.48$, $p>0.1$ and $R=0.64$, $p>0.1$) - no meaningful inference could be drawn in this regard at Holt Hall Lake as there were too few samples in which PBDFs and both BDEs-183 and -209 were detected.

In addition to the potential contribution of PBDE use, both OBDF and 1,2,3,4,6,7,8-HpBDF (the predominant HpBDF detected in this study) have been reported to be emitted from combustion processes (Wang et al, 2015). We also note that we did not target

2,7-/2,8-dibromodibenzo-p-dioxins or 1,3,7-/1,3,8-tribromodibenzo-p-dioxins, which have been identified as being of biogenic origin (Goto et al, 2017) and can therefore not rule out possible contributions from biogenic sources to the burden of PBDD/Fs in our lakes. In summary, drawing firm conclusions about PBDD/F source attribution based on homologue and congener patterns in environmental matrices like lake sediments, is complicated by post-emission modification of such profiles by weathering processes that favour some PBDD/Fs more than others. Notwithstanding this, the evidence of congener/homologue profiles presented here, suggests that the PBDD/Fs detected in this study are predominantly a complex integral of combustion source emissions and their presence as contaminants of the Deca-BDE formulation.

Temporal Trends in concentrations of Σ PBDFs and PBDF homologue patterns

Temporal trends in concentrations of Σ PBDFs are shown in Figure 1. Overall, they reveal a steady rise in concentrations from the 1980s onwards, peaking at the surface (2015) at Edgbaston Pool and Wake Valley Pond, and in the second most recent core slice (dated to 2009) at Holt Hall Lake. Prior to the onset of rising concentrations in the 1980s, concentrations were much lower. Inspection of Figure 1 suggests temporal trends in Σ PBDEs and Σ PBDFs are positively correlated and indeed the correlation coefficient for the three sites is significant, ranging between 0.88-0.98 ($p < 0.05$ at each site). With respect to changes in the homologue pattern over time, Figure 2 reveals no clear trend at Holt Hall Lake. In contrast, there is a marked shift from HpBDFs to HxBDFs in more recent sediment layers at Edgbaston Pool. Meanwhile at Wake Valley Pond, there is a noticeable decline in the abundance of TBDFs, along with increased abundance of HpBDFs in more recent core slices, as well as the aforementioned presence of OBDF in the top two layers (dated to 2009 and 2015) only. These variable homologue patterns and temporal trends suggest that changes in the relative contributions of different sources of PBDFs over time differ between our three lakes. To

our knowledge there is only one study with which our data on PBDF homologue patterns in sediment cores can be compared. In this study of a sediment core from Osaka Bay in Japan that covered the period 1904-2000, OBDF was not detected and in declining order of abundance the other homologues followed the order HpBDF>HxBDF>PeBDF>TBDF (Takigami et al, 2005). This homologue pattern most closely resembled that observed in this study at Holt Hall Lake.

Concentrations, Relative Abundance, and Temporal Trends of PXDD/Fs in English Lake Sediments

Table 2 reports concentrations of individual PXDD/Fs and homologue groups in each sediment core slice analyzed in this study. Also included in Table 2 are concentrations of 2-Br-3,7,8-CDD, 2-Br-3,6,7,8,9-CDD, 2-Br-2,7,8-CDF, and 1-Br-2,3,7,8-CDF reported previously in surficial sediments collected from Osaka Bay, Japan (Ohta et al, 2002). Concentrations of these PXDD/Fs in our study are of a similar range to those reported for Osaka Bay. As observed for both PBDEs and PBDD/Fs, concentrations of Σ PXDD/Fs at the two lakes studied for these contaminants do not correspond with the population density of the local authority within which the lake was submitted (Table SD-1). Specifically, t-test comparison shows PXDD/F concentrations at Wake Valley Pond to be statistically indistinguishable ($p>0.05$) from those at Holt Hall Lake. For PXDD/Fs, the concentrations detected were typically two orders of magnitude below those of PBDD/Fs. PXDFs were more abundant than PXDDs in every core slice analysed from both locations where PXDD/Fs were determined, with PXDFs typically 2-3 times more abundant. There was no clear temporal trend in the ratio of PXDFs:PXDDs at either lake. Moreover, the homologue pattern is broadly similar at both lakes and does not appear subject to temporal variation.

Temporal trends in concentrations of PXDD/Fs differ between the two lakes in which they were measured. Concentrations at Wake Valley Pond peak in the oldest slice (1954) analyzed at this location and then steadily decrease through to the present (2015). In slight contrast, concentrations at Holt Hall Lake rise from 1935 to 1976, before declining through to the uppermost layer (2015). In contrast to the significant positive correlation between Σ PBDEs and Σ PBDFs at all three lakes, concentrations of Σ PBDEs are not significantly correlated ($p>0.05$) with those of Σ PXDD/Fs. Likewise, Σ PBDD/Fs are not correlated with Σ PXDD/Fs ($p>0.05$). Overall, this suggests that the sources of PXDD/Fs are different to those of PBDD/Fs, and that the manufacture and use of PBDEs does not appear to be a source of PXDD/Fs.

Conclusions

Concentrations of PXDD/Fs in radiometrically-dated sediment cores from 3 English lakes decreased over time to the present. In contrast, concentrations of PBDEs increased towards the present and were significantly correlated with concentrations of PBDFs in all three lakes. These findings suggest that PBDEs do not appear to be a source of PXDD/Fs and that the sources of PXDD/Fs are different to those of PBDFs. Moreover, while the correlation between PBDEs and PBDFs implies that increased use of PBDEs has contributed substantially to environmental contamination with PBDFs; examination of PBDF homologue patterns implies emissions from combustion activities such as waste incineration and iron ore sintering are likely also important. Concentrations of PBDEs in all samples are dominated by BDE-209 which is the main constituent of the Deca-BDE formulation most widely used in the UK. At two of the three lakes studied, the concentrations of BDE-209 were highest in the sediment layers dated to 2012 and 2009, rather than in the surface layer dated to 2015. This may indicate that recent bans and restrictions on the

manufacture and use of Deca-BDE may be starting to take effect and reduce concentrations in the environment.

Acknowledgements

The research leading to these results has received funding from the Marie Curie Actions of the European Union's FP7 Programme under REA grant agreement # 606857 (ELUTE project). The analytical work was supported by the RECETOX research infrastructure (LM2018121). We thank Ian Patmore for his help in the field, Aristide Ganci for field and lab-work assistance, and Alwyn Fernandes for the kind gift of PXDD/F standards. This paper is dedicated to the memory of Dr Leon Peters.

Supplementary Data

Supplementary data to this article can be found online at

References

1. Bromine Science and Environmental Forum (BSEF) 2003. Major Brominated Flame Retardants Volume Estimates: Total Market Demand by Region in 2001, www.bsef.com (accessed 21st January 2003).
2. Chattrakittikunwong, W., Creaser, C. S. 1994. Bromo-, Bromochloro- and Chloro-Dibenzo-p-Dioxins and Dibenzofurans in Incinerator Flyash. *Chemosphere* 29, 559-566.

- 378 3. Chen, S. J., Ma, Y. J., Wang, J., Tian, M., Luo, X-J., Chen, D., Mai, B-X. 2010 Measurement
379 and human exposure assessment of brominated flame retardants in household products from
380 South China. J. Hazard Mater. 176, 979–984.
- 381 4. Choi, J. W., Fujimaki, S., Kitamura, K., Hashimoto, S., Ito, H., Sakurai, T., Suzuki, N.,
382 Nagasaka, H., Tanabe, K., Sakai, S., Morita, M. 2003. Historical trends of PBDD/Fs,
383 PBDEs, PCDD/Fs, and dioxin - like PCBs in sediment cores from Tokyo Bay.
384 Organohalogen Compd. 61, 119-122.
- 385 5. Drage, D. S., Aries, E., Harrad, S. 2014. Studies into the formation of PBDEs and PBDD/Fs
386 in the iron ore sintering process. Sci. Tot. Environ. 485-486, 497-507.
- 387 6. Ganci, A. P., Vane, C. H., Abdallah, M. A.-E., Moehring, T., Harrad, S. 2019. Legacy PBDEs
388 and NBFRs in sediments of the tidal River Thames using liquid chromatography coupled to a
389 high resolution accurate mass Orbitrap mass spectrometer. Sci. Tot. Environ. 658, 1355–
390 1366.
- 391 7. Goto, A., Tue, N. M., Someya, M., Isobe, T., Takahashi, S., Tanabe, S., Kinisue, T. 2017.
392 Spatio-temporal trends of polybrominated dibenzo-*p*-dioxins and dibenzofurans in archived
393 sediments from Tokyo Bay, Japan. Sci.Tot. Environ. 599-600, 340-347.
- 394 8. Hagberg, J., Grahn, E., van Bavel, B., Lindström, G. 2005. Occurrence and levels of
395 PCDD/Fs and PBDD/Fs in two Swedish lake sediments. Organohalogen Compd. 67, 2030-
396 2032.
- 397 9. Hanari, N., Kannan, K., Okazawa, T., Kodavanti, P. R. S., Aldous, K. M., Yamashita, N.
398 2006. Occurrence of Polybrominated Biphenyls, Polybrominated Dibenzo-*p*-dioxins, and

- 399 Polybrominated Dibenzofurans as Impurities in Commercial Polybrominated Diphenyl Ether
400 Mixtures. *Environ. Sci. Technol.* 40, 4400–4405.
- 401 10. Harrad, S., Hunter, S. 2006. Concentrations of Polybrominated Diphenyl Ethers in Air and
402 Soil on a Rural-Urban Transect Across a Major UK Conurbation. *Environ. Sci. Technol.* 40,
403 4548-4553.
- 404 11. Hayakawa, K., Takatsuki, H., Watanabe, I., Sakai, S. 2004. Polybrominated diphenyl ethers
405 (PBDEs), polybrominated dibenzo-p-dioxins/dibenzofurans (PBDD/Fs) and
406 monobromo-polychlorinated dibenzo-p-dioxins/dibenzofurans (MoBPXDD/Fs) in the
407 atmosphere and bulk deposition in Kyoto, Japan. *Chemosphere* 57, 343–56.
- 408 12. Kannan, K., Watanabe, I., Giesy, J. P. 1998. Congener profile of polychlorinated/brominated
409 dibenzo-p-dioxins and dibenzofurans in soil and sediments collected at a former chlor-alkali
410 plant. *Toxicol. Environ. Chem.* 67, 135-146.
- 411 13. Law, R. J.; Herzke, D.; Harrad, S.; Morris, S.; Bersuder, P.; Allchin, C. R. 2008. Levels and
412 trends of brominated flame retardants in the European environment. *Chemosphere* 73, 223–
413 241.
- 414 14. Kohler, M., Zennegg, M., Bogdal, C., Gerecke, A. C., Schmid, P. V., Heeb, N., Sturm, M.,
415 Vonmont, H., Kohler, H-P. E., Giger, W. 2008. Temporal Trends, Congener Patterns, and
416 Sources of Octa-, Nona-, and Decabromodiphenyl Ethers (PBDE) and
417 Hexabromocyclododecanes (HBCD) in Swiss Lake Sediments. *Environ. Sci. Technol.* 42,
418 6378–6384.

- 419 15. La Guardia, M. J., Hale, R. C., Harvey, E. 2006. Detailed Polybrominated Diphenyl Ether
420 (PBDE) Congener Composition of the Widely Used Penta-, Octa-, and Deca-PBDE Technical
421 Flame-retardant Mixtures. Environ. Sci. Technol. 40, 6247–6254.
- 422 16. Leung, A. O. W., Luksemburg, W. J., Wong, A. S., Wong, M. H. 2007. Spatial Distribution
423 of Polybrominated Diphenyl Ethers and Polychlorinated Dibenzo-*p*-dioxins and
424 Dibenzofurans in Soil and Combusted Residue at Guiyu, an Electronic Waste Recycling Site
425 in Southeast China. Environ. Sci. Technol. 41, 2730–2737.
- 426 17. Lundstedt, S. 2016. Sources and levels of PBDD/Fs in the Swedish environment
427 <https://www.diva-portal.org/smash/get/diva2:945535/FULLTEXT01.pdf>. (accessed March 3rd
428 2020).
- 429 18. Ma, J., Kannan, K., Cheng, J., Hori, Y., Wu, Q., Wang, W. 2008. Concentrations, profiles,
430 and estimated human exposures for polychlorinated dibenzo - *p* - dioxins and dibenzofurans
431 from electronic waste recycling facilities and a chemical industrial complex in Eastern China.
432 Environ. Sci. Technol. 42, 8252 - 8259.
- 433 19. Ma, J., Addink, R., Yun, S. H., Cheng, J. P., Wang, W. H., Kannan K. 2009. Polybrominated
434 dibenzo-*p* - dioxins/dibenzofurans and polybrominated diphenyl ethers in soil, vegetation,
435 workshop - floor dust, and electronic shredder residue from an electronic waste recycling
436 facility and in soils from a chemical industrial complex in Eastern China. Environ. Sci.
437 Technol. 43, 7350 - 7356.

- 438 20. Mai, B., Chen, S., Luo, X., Chen, L., Yang, Q., Sheng, G., Peng, P., Fu, J., Zeng, E. Y. 2005.
439 Distribution of Polybrominated Diphenyl Ethers in Sediments of the Pearl River Delta and
440 Adjacent South China Sea. *Environ. Sci. Technol.* 39, 3521-3527.
- 441 21. Myers, A. L., Mabury, S. A., Reiner, E. J. 2012. Analysis of mixed halogenated
442 dibenzo-*p*-dioxins and dibenzofurans (PXDD/PXDFs) in soil by gas chromatography tandem
443 mass spectrometry (GC-MS/MS). *Chemosphere* 87, 1063-1069.
- 444 22. Naturvårdsverket. 2011. Recycling and disposal of electronic waste - health hazards and
445 environmental impacts. 6417.
446 <http://naturvardsverket.se/Documents/publikationer6400/978-91-620-6417-4.pdf> (accessed
447 March 3rd 2020)
- 448 23. Ohta, S., Nakao, T., Nishimura, H., Okumura, T., Aozasa, O., Miyata, H. 2002.
449 Contamination levels of PBDEs, TBBPA, PCDD/Fs, PBDD/Fs and PXDD/Fs in the
450 environment of Japan. *Organohalogen Compd.* 57, 57-60.
- 451 24. Patmore, I. R., Sayer, C. D., Goldsmith, B., Davidson, T. A., Rawcliffe, R., Salgado, J. 2014.
452 Big Ben: A new wide-bore piston corer for multi-proxy palaeolimnology. *J. Paleolimnol.* 51,
453 79-86.
- 454 25. Ramu, K., Isobe, T., Takahashi, S., Subramanian, A., Parthasarathy, P., Tanabe S. 2008.
455 Brominated flame retardants and dioxins in soil from electronic waste recycling sites in India.
456 *Organohalogen Compd.* 70, 2058 - 2061.
- 457 26. Ren, M., Peng, P., Cai, Y., Chen, D., Zhou, L., Chen, P., Hu, J. 2011. PBDD/F impurities in
458 some commercial deca-BDE. *Environ. Pollut.* 159, 1375-1380.

- 459 27. Takigami, H., Sakai, S., Brouwer, A. 2005. Bio/chemical Analysis of Dioxin-like Compounds
460 in Sediment Samples from Osaka Bay, Japan. *Environ. Technol.* 26, 459–70.
- 461 28. Terauchi, H., Takahashi, S., Lam, P. K. S., Min, B-Y., Tanabe, S. 2009. Polybrominated,
462 polychlorinated and monobromo-polychlorinated dibenzo-*p*-dioxins/dibenzofurans and
463 dioxin-like polychlorinated biphenyls in marine surface sediments from Hong Kong and
464 Korea. *Environ. Pollut.* 157, 724-730.
- 465 29. Tue, N. M., Takahashi, S., Subramanian, A., Sakai, S., Tanabe, S. 2013. Environmental
466 contamination and human exposure to dioxin-related compounds in e-waste recycling sites of
467 developing countries. *Environ. Sci. Proc. Imp.* 15, 1326–1331.
- 468 30. Van den Berg, M., Denison, M. S., Birnbaum, L. S., DeVito, M. J., Fiedler, H., Falandysz, J.,
469 Rose, M., Schrenk, D., Safe, S., Tohyama, C., Tritscher, A., Tysklind, M., Peterson, R. E.
470 2013. Polybrominated Dibenzo-*p*-Dioxins, Dibenzofurans, and Biphenyls: Inclusion in the
471 Toxicity Equivalency Factor Concept for Dioxin-Like Compounds. *Toxicol. Sci.* 133, 197-
472 208.
- 473 31. Vane, C. H., Ma, Y-J., Chen, S-J., Mai, B-X. Increasing polybrominated diphenyl ether
474 (PBDE) contamination in sediment cores from the inner Clyde Estuary, UK. 2010. *Environ.*
475 *Geochem. Hlth.* 32, 13–21.
- 476 32. Wall, R. J., Fernandes, A., Rose, M., Bell, D. R., Mellor, I. R. 2015. Characterisation of
477 chlorinated, brominated and mixed halogenated dioxins, furans and biphenyls as potent and
478 as partial agonists of the Aryl hydrocarbon receptor. *Environ. Int.* 76, 49–56.
- 479 33. Wang, L-C., Hsi, H-C., Wang, Y-F., Lin, S-L, Chang-Chien, G-P. 2010. Distribution of
480 polybrominated diphenyl ethers (PBDEs) and polybrominated dibenzo-*p*-dioxins and

481 dibenzofurans (PBDD/Fs) in municipal solid waste incinerators. *Environ. Pollut.* 158, 1595–
 482 602.

483 34. Wang, M., Liu, G., Jiang, X., Liu, W., Li, L., Li, S., Zheng, M., Zhan, J. 2015. Brominated
 484 dioxin and furan stack gas emissions during different stages of the secondary copper smelting
 485 process. *Atmos. Pollut. Res.* 6, 464-468.

486 35. Weber, R., Kuch, B. 2003. Relevance of BFRs and thermal conditions on the formation
 487 pathways of brominated and brominated–chlorinated dibenzodioxins and dibenzofurans.
 488 *Environ. Int.* 29, 699– 710.

489 36. Webster, L., Russell, M., Adefehinti, F., Dalgarno, E. J., Moffat, C. F. 2008. Preliminary
 490 assessment of polybrominated diphenyl ethers (PBDEs) in the Scottish aquatic environment,
 491 including the Firth of Clyde. *J. Environ. Monit.* 10, 463–473.

492 37. Yang, C., Rose, N.L., Turner, S.D., Yang, H., Goldsmith, B., Losada, S., Barber, J. L., Harrad,
 493 S. 2016. Hexabromocyclododecanes, polybrominated diphenyl ethers, and polychlorinated
 494 biphenyls in radiometrically dated sediment cores from English lakes, ~ 1950–present. *Sci.*
 495 *Tot. Environ.* 541, 721–728.

496 38. Yu, X., Zennegg, M., Engwall, M., Rotander, A., Larsson, M., Wong, M. H., Weber, R. 2008.
 497 E-Waste Recycling Heavily Contaminates a Chinese City with Chlorinated, Brominated and
 498 Mixed Halogenated Dioxins. *Organohalogen Compd.* 70, 813-816.

499 39. Zegers, B. N., Lewis, W. E., Booij, K., Smittenberg, R. H., Boer, W., de Boer, J., Boon, J. P.
 500 2003. Levels of polybrominated diphenyl ether flame retardants in sediment cores from
 501 Western Europe. *Environ. Sci. Technol.* 37, 3803–3807.

- 502 40. Zennegg, M., Yu, X., Wong, M. H., Weber, R. 2009. Fingerprints of Chlorinated, Brominated
503 and Mixed Halogenated Dioxins at Two E-Waste Recycling Sites in Guiyu/China.
504 *Organohalogen Compd.* 71, 2248-2252.
- 505 41. Zhang, X-L., Luo, X-J., Chen, S-J., Wu, J-P., Mai, B-X. 2009. Spatial distribution and
506 vertical profile of polybrominated diphenyl ethers, tetrabromobisphenol A, and
507 decabromodiphenylethane in river sediment from an industrialized region of South China.
508 *Environ. Pollut.* 157, 1917–1923
- 509 42. Zhou, L., Li, H., Yu, Z., Ren, M., Zeng, X., Peng, P., Sheng, G., Fu, J. 2012. Chlorinated and
510 brominated dibenzo-p-dioxins and dibenzofurans in surface sediment from Taihu Lake, China.
511 *J. Environ. Monit.* 14, 1935-1942.

512 **Table 1: Concentrations of Σ PBDEs and Σ PBDD/Fs (ng/g dry weight) in surficial sediments**
513 **from this study with those in soil and sediments from selected other studies**

Matrix, Location	Σ PBDD/F	Σ PBDE	Reference
Fresh water sediments			
Edgbaston Pool, UK	7.1	210	This Study
Holt Hall Lake, UK	0.49	18	This Study
Wake Valley Pond, UK	0.49	21	This Study
Edgbaston Pool, UK ^a	-	120	Yang et al, 2016
Holt Hall Lake, UK ^a	-	10	Yang et al, 2016
Wake Valley Pond, UK ^a	-	9.8	Yang et al, 2016
Surficial Sediments, River Thames, UK	-	180 (mean)	Ganci et al, 2019
Clyde Estuary, UK (core slices)	-	1-2,600	Vane et al, 2010
Scotland, UK (core slices)	-	2.3-98,000	Webster et, 2008
Rural/urban lakes, Sweden	0.44-0.54	-	Hagberg et al, 2005
Urban river, Sweden	0.41-1.7	29-62	Lundstedt, 2016
Rural lake Sweden	0.082-0.085	4.4-16	Lundstedt, 2016
Urban lake, China	0.00048-0.0057	-	Zhou et al, 2012
Stream at dump site, Peru	0.012-0.074	3.7-6.1	Naturvårdsverket, 2011
Lake, industrial area, Thailand	0.037-1.5	3.4-58	Naturvårdsverket, 2011
Marine sediments			
Hong Kong/Korea	nd.-0.46	-	Terauchi et al, 2009
Tokyo Bay, Japan	0.0052-0.070	10-78,050	Choi et al, 2003
Osaka, Japan	0.0041-0.077	8.0-352	Ohta et al, 2002
Osaka, Japan	0.0024-0.59	53-910	Takigami et al, 2005
Coastal and offshore, Sweden	0.050-10	-	Lundstedt, 2016
Rural Soil			
Lanna, Sweden	0.028-0.054	0.065-1.3	Lundstedt, 2016
Urban soil			
Umeå and Norrköping, Sweden	0.0011-0.22	0.18-66	Lundstedt, 2016
Bangalore and Chennai, India	0.0060-0.31	-	Ramu et al, 2008
Kyoto, Japan	0.28	-	Hayakawa et al, 2004
Industrial area, China	nd.-0.43	2.03-269	Ma et al, 2008; Ma et al, 2009
Industrial area, Thailand	0.019-0.16	1.8-13	Naturvårdsverket, 2011
Dump site, Peru	0.0086-0.32	3.6-92	Naturvårdsverket, 2011

514 ^aData for core slices dated to 2008 for Edgbaston Pool and for 2009 for both Holt Hall Lake and
515 Wake Valley Pond.

516 **Table 2: Concentrations (pg/g OC) of PXDD/Fs in Sediment Core Slices from Wake Valley Pond and Holt Hall Lake with Comparative**
517 **Data from Osaka Bay, Japan**

Location	Congener/ Homologue/ Year	2015	2009	2005	2004	1999	1993	1985	1977	1969	1965	1954	1935
Wake Valley Pond	2-Br-7,8-CDD	0.023	0.16	0.24		0.32	0.4	0.035			0.079	<0.032	
Holt Hall Lake	2-Br-7,8-CDD	1.1	2.8		2.6	3.4		1.1	<0.025	<0.024		<0.021	<0.021
Wake Valley Pond	<i>Total Br-2CDD</i>	0.73	1.9	1.9		2.7	2.3	3.6			0.8	9.3	
Holt Hall Lake	<i>Total Br-2CDD</i>	1.8	4.1		4.0	5.3		2.2	11	7.1		2.3	<0.021
Wake Valley Pond	2-Br-3,7,8-CDD	0.081	<0.028	<0.027		0.12	0.22	0.22			0.29	<0.047	
Holt Hall Lake	2-Br-3,7,8-CDD	0.02	0.049		0.054	0.16		0.08	1.4	<0.035		<0.031	<0.03
Osaka Bay, Japan ^a	2-Br-3,7,8-CDD	0.84-6.5											
Wake Valley Pond	<i>Total Br-3CDD</i>	1.5	2.2	2.8		3.8	3	3.5			6.2	9.3	
Holt Hall Lake	<i>Total Br-3CDD</i>	0.77	2.1		2.4	2.7		2.4	18	<0.027		<0.019	<0.018
Wake Valley Pond	2,3-Br-7,8-CDD	<0.015	<0.017	<0.016		<0.016	<0.017	<0.016			<0.016	<0.029	
Holt Hall Lake	2,3-Br-7,8-CDD	<0.013	0.095		0.062	<0.018		0.034	<0.022	<0.021		<0.019	<0.018
Osaka Bay, Japan ^a	2,3-Br-7,8-CDD	Not detected											
Wake Valley Pond	<i>Total 2Br-2CDD</i>	0.48	0.64	0.87		1.2	1.5	1.8			1.2	4.4	
Holt Hall Lake	<i>Total 2Br-2CDD</i>	0.33	0.79		0.82	1.1		0.97	5.0	2.2		0.62	<0.017
Wake Valley Pond	1-Br-2,3,7,8-CDD + 2-Br-1,3,7,8-CDD [*]	<0.018	<0.021	0.37		<0.02	<0.021	<0.02			<0.021	<0.032	
Holt Hall Lake	1-Br-2,3,7,8-CDD + 2-Br-1,3,7,8-CDD [*]	<0.016	<0.019		<0.02	<0.023		<0.027	<0.028	<0.027		<0.024	<0.023
Osaka Bay, Japan ^a	1-Br-2,3,7,8-CDD	Not detected											
Wake Valley Pond	<i>Total Br-4CDD</i>	1.7	1.8	2.8		3.8	5.9	4.6			7	5.7	
Holt Hall Lake	<i>Total Br-4CDD</i>	0.47	1.9		1.0	1.9		<0.021	<0.022	<0.021		0.71	<0.018
Wake Valley Pond	2-Br-3,6,7,8,9-CDD	<0.015	<0.017	<0.016		<0.016	<0.017	<0.016			<0.016	<0.029	
Holt Hall Lake	2-Br-3,6,7,8,9-CDD	<0.013	<0.015		<0.016	<0.018		0.6	<0.022	<0.021		<0.019	<0.018
Osaka Bay, Japan ^a	2-Br-3,6,7,8,9-CDD	Not detected - 0.32											
Wake Valley Pond	<i>Total Br-5CDD</i>	1	0.82	3.5		2.2	5.1	7			7.5	15	
Holt Hall Lake	<i>Total Br-5CDD</i>	<0.018	0.11		<0.01	<0.012		0.6	<0.022	<0.021		<0.019	<0.018

Location	Congener/ Homologue/ Year	2015	2009	2005	2004	1999	1993	1985	1977	1969	1965	1954	1935
Wake Valley Pond	2-Br-7,8-CDF	0.77	1	0.88		<0.01	2	2.7			<0.01	<0.018	
Holt Hall Lake	2-Br-7,8-CDF	<0.008	<0.01		<0.01	<0.012		0.53	<0.014	<0.013		<0.012	<0.012
Wake Valley Pond	<i>Total Br-2CDF</i>	6.9	16	16		26	17	25			22	62	
Holt Hall Lake	<i>Total Br-2CDF</i>	4.7	7.9		7.7	12		9.6	88	89		15	9.5
Wake Valley Pond	2-Br-6,7,8-CDF + 3-Br-2,7,8-CDF*	<0.012	0.13	0.17		0.3	0.65	0.46			0.076	<0.023	
Holt Hall Lake	2-Br-6,7,8-CDF + 3-Br-2,7,8-CDF*	0.07	0.28		0.18	0.40		1.3	5.1	<0.017		0.63	<0.015
Osaka Bay, Japan ^a	3-Br-2,7,8-CDF	Not detected-2.0											
Wake Valley Pond	<i>Total Br-3CDF</i>	4.3	6	6.4		8.4	8.7	10			11	57	
Holt Hall Lake	<i>Total Br-3CDF</i>	2.5	4.8		5.3	8.8		5.0	44	49		6.6	3.1
Wake Valley Pond	1-Br-2,3,7,8-CDF	0.022	<0.014	0.047		0.042	0.66	<0.012			<0.012	<0.021	
Holt Hall Lake	1-Br-2,3,7,8-CDF	<0.009	<0.011		<0.012	<0.014		<0.016	<0.017	<0.016		<0.014	<0.014
Osaka Bay, Japan ^a	1-Br-2,3,7,8-CDF	Not detected											
Wake Valley Pond	<i>Total Br-4CDF</i>	1.5	1.6	2.6		3.6	6.4	5.8			2	20	
Holt Hall Lake	<i>Total Br-4CDF</i>	1.8	5.0		2.4	5.2		0.81	7.3	<0.016		3.5	4.4
Wake Valley Pond	$\sum$ PXDD	5.4	7.5	12		14	18	20			23	43	
Holt Hall Lake	$\sum$ PXDD	3.3	9.1		8.2	11		6.0	34	9.3		7.1	<0.03
Wake Valley Pond	$\sum$ PXDF	13	24	25		38	32	41			34	140	
Holt Hall Lake	$\sum$ PXDF	9.0	18		15	26		15	140	140		25	17
Wake Valley Pond	$\sum$ PXDD/F	18	31	37		51	50	61			57	180	
Holt Hall Lake	$\sum$ PXDD/F	12	27		24	37		21	170	150		32	17

518 ^aRange (pg/g dry weight) surficial sediments from Osaka, Japan (n=6) (Ohta et al, 2002) – detection limits not reported

519

520 **Figure 1: Temporal Trends in Concentrations (ng/g OC) of Σ PBDE, Σ PBDF, and**
521 **Σ PXDD/F in 3 English Lake Sediment Cores**
522

523

524

525

Figure 2: Temporal Trends in Relative Contributions of PBDF Homologues to Σ PBDF Concentrations in 3 English Lake Sediment Cores

531

532