

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements

Pant, Pallavi; Harrison, Roy

DOI:

[10.1016/j.atmosenv.2013.04.028](https://doi.org/10.1016/j.atmosenv.2013.04.028)

Document Version

Early version, also known as pre-print

Citation for published version (Harvard):

Pant, P & Harrison, RM 2013, 'Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements: A review' *Atmospheric Environment*, vol. 77, pp. 78-97. DOI: [10.1016/j.atmosenv.2013.04.028](https://doi.org/10.1016/j.atmosenv.2013.04.028)

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

[10.1016/j.atmosenv.2013.04.028](https://doi.org/10.1016/j.atmosenv.2013.04.028) NOTICE: this is the author's version of a work that was accepted for publication in *Atmospheric Chemistry*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Atmospheric Chemistry*, [vol. 77, 2013] DOI: <http://dx.doi.org/10.1016/j.atmosenv.2013.04.028>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1
2
3
4
5
6
7
8
9
10
11
12
13

**Estimation of the Contribution of Road Traffic
Emissions to Particulate Matter Concentrations
from Field Measurements: A Review**

Pallavi Pant and Roy M. Harrison^{*1}

**Division of Environmental Health & Risk Management,
School of Geography, Earth & Environmental Sciences
University of Birmingham, Edgbaston**

**Birmingham B15 2TT
United Kingdom**

^{*} Corresponding Author: Tel: 00 44 121 414 3494; Fax: 00 44 121 414 3709;
Email: r.m.harrison@bham.ac.uk

¹ Also at: Centre of Excellence in Environmental Studies, King Abdulaziz University, PO Box 80203,
Jeddah (21589) Saudi Arabia

14 **ABSTRACT**

15 Road traffic is one of the main sources of particulate matter in the atmosphere. Despite its
16 importance, there are significant challenges in quantitative evaluation of its contribution to airborne
17 concentrations. This article first reviews the nature of the particle emissions from road vehicles
18 including both exhaust and non-exhaust (abrasion and resuspension sources). It then briefly
19 reviews the various methods available for quantification of the road traffic contribution. This
20 includes tunnel/roadway measurements, twin site studies, use of vehicle-specific tracers and other
21 methods. Finally, the application of receptor modelling methods is briefly described. Based on the
22 review, it can be concluded that while traffic emissions continue to contribute substantially to
23 primary PM emissions in urban areas, quantitative knowledge of the contribution, especially of non-
24 exhaust emissions to PM concentrations remain inadequate.

25

26 **Keywords:** Particulate matter; road traffic; exhaust emissions; non-exhaust emissions

27

28 1. INTRODUCTION

29 Emissions due to road traffic are known to make a large contribution to total particulate matter (PM)
30 concentrations in urban areas and exposure to PM from vehicular emissions has been demonstrated
31 to have detrimental impacts on human health (Mauderly, 1994; Buckeridge et al., 2002; Fan et al.,
32 2006; HEI, 2010; Masiol et al., 2012; Rissler et al., 2012). An improved understanding of the
33 characteristics of emissions from different traffic-related sources is therefore vital for conducting
34 source apportionment and health effects studies (Schauer et al., 2006). PM emissions from road
35 vehicles include emissions from the tailpipe (exhaust emissions) and emissions due to wear and tear
36 of vehicle parts such as brake, tyre and clutch and re-suspension of dust (non-exhaust emissions).
37 Non-exhaust emissions contribute mainly to the coarse mode of PM (PM_{2.5-10}) while exhaust
38 emissions contribute predominantly to fine PM (aerodynamic diameter < 2.5 µm) (Abu-Allaban et
39 al., 2003; Tervahattu et al., 2006; Thorpe et al., 2007; Kam et al., 2012) but this is a far from clear
40 distinction. A number of different properties of traffic emissions are studied including physical
41 shape and structure, particle size distributions, chemical composition, and temporal and spatial
42 variation (Watson and Chow, 2007). Most of the research as well as policy action in the last few
43 decades has largely focused on exhaust emissions, and stringent regulations and technological
44 upgrades have resulted in a decline of the percentage contribution of vehicle tailpipe emissions to
45 total ambient PM concentrations (Allen et al., 2006; Thorpe and Harrison, 2008; Mathissen et al.,
46 2011). As a result, the contribution of non-exhaust PM is becoming more important although
47 detailed information on non-exhaust PM emissions is relatively scarce (Kuhlbusch et al., 2009;
48 Amato et al., 2012; Harrison et al., 2011; 2012b; Denier van der Gon et al., 2013). It has been
49 shown that even with zero tail-pipe emissions, traffic will continue to contribute to fine and
50 ultrafine particles through non-exhaust emissions (Dahl et al., 2006; Kumar et al., 2013) and it is
51 estimated that nearly 90% of the total emissions from road traffic will come from non-exhaust
52 sources by the end of the decade (Rexeis and Hausberger, 2009).

53 In order to understand the contribution of road traffic emissions to the environment, it is important
54 to understand various aspects including sources, chemical characteristics and quantitative
55 contributions. In the last three decades, a significant amount of research has been conducted on
56 characterization and analysis of exhaust emissions and as a result, there is a very strong body of
57 knowledge in the field. Non-exhaust emissions, however, are becoming more important now, and
58 further research is anticipated in this field in the coming years. This review aims to reconcile the
59 various studies that have been reported around this issue, and tease out some common
60 characteristics, and highlight areas with future research potential.

61 This paper reviews the current information on primary road traffic emissions including exhaust and
62 non-exhaust emissions. Key characteristics of the major sources are discussed. Various
63 methodologies used for measurement and analysis of such emissions are also discussed with a brief
64 discussion on the merits and shortcomings of each method. In the final section, various source
65 apportionment techniques useful for the estimation of traffic source contributions are discussed
66 together with a summary of some of the reported studies. A number of in-depth review articles have
67 been published in the last few years on various issues related to road traffic emissions on topics
68 such as properties and characteristics of non-exhaust emissions (Thorpe and Harrison, 2008),
69 toxicological impacts of tyre wear emissions (Wik and Dave, 2009), chemical composition of brake
70 pads (Chan and Stachowiak, 2004), development of emission factors (Franco et al., 2013),
71 nanoparticle emissions from non-exhaust sources (Kumar et. al., 2013). However, each of these
72 reviews focuses on one or a few specific aspects of road traffic emissions. This review builds on the
73 information provided in those review articles, and repetition is avoided. In summary, the key
74 questions that we hope to address through this review article are as follows:

- 75 1. What are the key characteristics/sources/contributions of road traffic emissions? [What and
76 how much do we know about each of the road traffic emission sources?]
- 77 2. How can such emissions be characterized? [Various methods, pros and cons of the methods]

78 3. What have we learnt so far?

79 4. What are the key issues that are not resolved yet?

80

81 The literature search for this review was primarily performed using Web of Knowledge, Science
82 Direct and Google Scholar using various combinations of keywords such as traffic emissions, non-
83 exhaust/exhaust emissions, resuspension, road dust, traffic PM etc. However, the authors
84 acknowledge that not all articles published on the topic have been included in the review.

85 2. NON-EXHAUST EMISSIONS

86 Non-exhaust PM comprises the various emissions that do not derive from the tailpipe of a vehicle
87 including particles generated due to brake and tyre wear, road surface abrasion, wear and
88 tear/corrosion of other vehicle components such as the clutch, and re-suspension of road surface
89 dusts (Table 1). Key reasons for needing to understand non-exhaust emissions include their inherent
90 toxicity including their tendency to act as carriers of heavy metals and carcinogenic components
91 (Adachi and Tainosho, 2004; Hjortenkrans et al., 2007; Johansson et al., 2009; Amato et al., 2011a)
92 and their contribution to exceedences of air quality guidelines and standards (Amato et al., 2011a;
93 Denier van der Gon, 2013).

94 Harrison et al. (2012b) showed that the size distributions of trace metals were indicative of particle
95 sources. Keuken et al. (2010) concluded that re-suspension of accumulated PM and road wear
96 related particles are the primary contributors to non-exhaust emissions, and tyre wear and brake
97 wear contribute to the fine and coarse fraction respectively. However, Narvaez et al. (2008)
98 reported that although a majority of abrasion particles are in the coarse fraction, abrasion can
99 contribute significantly to the fine fraction of PM. In Taiwan, EFs for elements such as Ba and Se
100 were found to be higher in the PM_{2.5-10} fraction indicating mechanical friction and brake wear as
101 potential sources (Chiang and Huang, 2009). Road traffic was found to be a key contributor to
102 fine/ultrafine and nano mode particles of Ba, Zn and Pb (Lin et al., 2005) and in Barcelona (Spain),

103 62-96% of elements including Cu, Sb, Ba, Mn and Zn were found to be present in the PM₁₋₁₀
104 fraction (Perez et al., 2010). A detailed review of the nanoparticle emissions from non-exhaust
105 sources including traffic and non-traffic sources is presented in Kumar et al. (2013).

106 Non-exhaust emissions are typically characterized by trace metals (e.g. Cu, Zn, Ba, Sb, Mn)
107 although organic markers (e.g. polycyclic aromatic hydrocarbons (PAHs), n-alkanes) have been
108 used in some cases (Wahlin et al., 2006; Amato et al., 2009; El Haddad et al., 2009; Gietl et al.,
109 2010; Oliviera et al., 2011; Kwon and Castaldi, 2012). A list of key tracers used for source
110 characterization of non-exhaust emissions is presented in Table 2. However, emissions of trace
111 metal markers are reported to vary with the fleet composition, with higher emissions reported for
112 some of the elements for heavy duty vehicles (HDVs) (Grieshop et al., 2006; Mancilla and
113 Mendoza, 2012). In addition, the profile of trace metal concentrations in non-exhaust particulate
114 matter is unique for every region and varies based on parameters such as traffic volume and pattern,
115 vehicle fleet characteristics, driving and traffic patterns and climate and geology of the region
116 (Omstedt et al., 2005; Amato et al., 2011a,b; Duong and Lee, 2011). Another important aspect is the
117 variability of tyre and brake composition depending on the manufacturer which makes it very
118 difficult to ascertain fleet-wide composition other than from environmental measurements
119 (Canepari et al., 2008; Gietl et al., 2010; Denier van der Gon et al., 2013).

120
121 Several factors are reported to affect non-exhaust emissions including increase in vehicle speed
122 (Chen et al., 2006; Gustafsson et al., 2008; Hussein et al., 2008; Mathissen et al., 2011) although
123 Keuken et al. (2010) reported a non-linear relationship between traffic volume and non-exhaust
124 emissions. Metal emissions due to road dust show a low correlation to metal emissions due to
125 abrasion/brake wear and combustion (Johansson et al., 2009). One of the major problems in
126 analysis of non-exhaust PM using field data has been the difficulty in distinguishing between wear
127 and tear emissions and road dust since the chemical composition is often very similar (Bukowiecki
128 et al., 2010). This may, in any case, not be a clear distinction as wear emissions may deposit to the

129 road surface, only to be re-suspended subsequently. Gehrig et al. (2010) used mobile load
130 simulators to compare the contributions of road abrasion and resuspension emissions, and
131 concluded that particle emissions due to abrasion are a function of the state of the pavement
132 whereas resuspension is normally higher, and increases with higher dust loads on the road. Thorpe
133 and Harrison (2008) present a more elaborate account of sources and properties of non-exhaust
134 emissions.

135 Mixed results have been obtained for the impact of street washing activities on PM and while
136 Amato et al. (2009) observed a reduction, albeit statistically non-significant, in the concentration of
137 trace metals known to be contributed by non-exhaust traffic sources in Spain, Keuken et al. (2010)
138 reported a lack of significant reduction in the non-exhaust emissions after street sweeping or
139 washing in the Netherlands.

140 **2.1 Tyre Wear**

141 Tyre wear particles are generated either by shear forces between the tyre tread and the road surface
142 and are predominantly coarse (PM_{2.5-10}) or by volatilization (Rogge et al., 1993; Kupiainen et al.,
143 2005; Allen et al., 2006; Thorpe and Harrison, 2008; Aatmeeyata and Sharma, 2010). Tyre wear
144 particles are reported to be generally elongated with rough surfaces based on microscopic analysis
145 (Gunawardana et al., 2011). Tyre tread, a source of airborne particles, contains natural rubber
146 copolymers such as styrene-butadiene rubber and polyisoprene rubber, and zinc (Zn) is added to
147 tyre tread as zinc oxide and organozinc compounds to facilitate the vulcanization process. Passenger
148 car tyres in EU are known to contain nearly 1% zinc oxide, and rubber, metals and carbon black
149 make up typically 47%, 16.5% and 21.5% respectively (Milani et al., 2004). The key tracer
150 components of tyre wear include n-alkanes, n-alkanoic acids, PAHs, benzothiazoles and trace
151 metals (Rogge et al., 1993; Reddy et al., 1997; Camatani et al., 2001; Adachi and Tainosho, 2004;
152 Boulter et al., 2005; Aatmeeyata and Sharma, 2010). Zn is reported to be nearly 1% by weight in
153 rubber tyres (Councell et al., 2004) and tyre wear has been reported to be a significant source of Zn

154 (Adachi and Tainosho, 2004; Hjortenkrans et al., 2007; Ondracek et al., 2011). The concentration of
155 Zn was found to be approximately 15 times higher in tyres compared to brakes while concentrations
156 of other heavy metals such as copper (Cu) and barium (Ba) were higher for brake materials in
157 Massachusetts (USA) (Apaegyeyi et al., 2011). Differences have also been observed between tyre
158 tread composition (primarily Zn) and tyre wear (Al, Si, Ca, Fe, Zn, Ti) (Adachi and Tainosho,
159 2004). However, the spatial and temporal trend for Zn was found to be different from other roadside
160 tracers in New York (USA), and a very small portion of airborne Zn concentration was attributed to
161 tyre wear (Peltier et al., 2011). However, Zn is emitted from brake wear, motor oil and other
162 sources and cannot be used as the only tracer for tyre wear (Lough et al., 2005). Benzothiazoles are
163 also used as markers for tyre wear, particularly benzothiazole (BT), 2-hydroxy benzothiazole
164 (HOBT), 2-(4-morpholinyl)benzothiazole (24MoBT) and N-cyclohexyl-2-benzothiazolamine
165 (NCBA) (Kumata et al., 2002; Allen et al., 2006; Wik and Dave, 2009). Among the PAHs, pyrene,
166 benzo(ghi)perylene, fluoranthene and phenanthrene are known to be emitted from tyres
167 (Boonyatumanond et al., 2007; Kwon and Castaldi, 2012). Aatmeeyata and Sharma (2010) reported
168 that small cars tested in India emitted 378 ng of total PAH/tyre/km while in the USA, 200 µg/g of
169 PAHs were found in tyre wear with pyrene, fluoranthene and phenanthrene being present in the
170 highest concentrations (Rogge et al., 1993). Tyre tread wear has also been found to contain an
171 average of 0.53 µg/g of dibenzopyrenes, indicating that tyre dust may be significant source of
172 dibenzopyrenes in the environment (Sadiktis et al., 2012).

173
174 Average tread wear for tyres is reported to be between 0.006-0.009 g/km based on the road, tyre and
175 vehicle conditions (Rogge et al., 1993) and tyres can lose up to 10% of their mass during their
176 lifetime (Milani et al., 2004). Speed has been reported as an influential parameter for particle mass
177 and number concentrations for tyre wear emissions (Gustafsson et al., 2008; Mathissen et al., 2011).
178 Other important parameters for tyre tread wear include road surface type, and tyre and driving
179 conditions (Allen et al., 2006; Gustafsson et al., 2008). For example, asphalt surfaces have been

180 reported to cause less tyre wear than concrete pavements and in Arizona, USA, the emission rate for
181 tyre wear was found to be 1.4-2 times lower for asphalt pavement compared to concrete pavement
182 (Allen et al., 2006). The type of tyre also impacts the magnitude of tyre wear emissions and studded
183 tyres are known to cause more emissions compared with summer and friction tyres (Kupiainen et al.,
184 2005; Gustafsson et al., 2008; Hussein et al., 2008; Schaap et al., 2009). Unimodal (70-90 nm) and
185 bimodal (< 10 and 30-60 nm) peaks in the nano size range have been reported for tyre particles by
186 number under low and high speed conditions respectively (Mathissen et al., 2011).

187

188 **2.2 Brake Wear**

189 Brake wear, including abrasion of brake lining material and brake discs, caused by grinding of
190 brake pad constituents (coarse range particles) or volatilization and condensation of brake pad
191 materials (fine range particles), is known to release PM directly into the atmosphere and to
192 contribute to the trace metal concentration in airborne PM, particularly less than 10 μm (Garg et al.,
193 2000; Blau and Meyer, 2003; Ingo et al., 2004; Wahlin et al., 2006; Gietl et al., 2010; Varrica et al.,
194 2012). Key components of brake pads include fillers, frictional additives, reinforcing fibres and
195 binder (Chan and Stachowiak, 2004) and the key chemical species used include sulphides of metal,
196 abrasives (e.g. silica), barium silicate/sulphate (particularly in brake linings) and other metallic
197 particles (as filler material), carbon fibres and lubricant (e.g. graphite) (Ingo et al., 2004; Dongarra
198 et al., 2009). Chan and Stachowiak (2004) present a detailed review of the chemical constituents of
199 brake pads. Garg et al. (2000) observed an average mass median diameter (median particle
200 diameter based on mass) of 1.49 μm for brake wear particles using tests on a brake dynamometer
201 whereas Sanders et al. (2003) reported a mass median diameter of 6 μm for brake debris generated
202 during urban driving conditions. Higher brake wear related emissions have been reported during
203 rush hour . A higher number of particles are released in the braking phase as compared to the
204 acceleration phase (Hussein et al., 2008; Mathissen et al., 2011) and various studies have
205 corroborated this observations including Greishop et al. (2006) who reported a higher particle

206 number concentration during rush hour where the traffic often operates in the stop-and-go mode and
207 Abu-Allaban et al. (2003) who observed higher contribution of brake wear at freeway exit sites
208 compared to other types of roadside sites (Abu-Allaban et al., 2003).

209 Passenger cars have been estimated to emit nearly 44 g/car/year brake dust (Iijima et al., 2007).
210 Sanders et al. (2003) reported high concentrations of Fe, Ba and Cu in brake lining wear using
211 dynamometer and track tests whereas Adachi and Tainosho (2004) reported Fe to be the most
212 abundant metal in brake dust along with other metals such as Ba, Cu, Sb, Zr and Zn, and
213 Hjortenkrans et al. (2007) reported Cu, Zn and Sb to be present in brake wear emissions. Varrica et
214 al. (2012) undertook a detailed study on Sb in brake dust and reported the most commonly released
215 forms of Sb due to brake abrasion to be Sb (III) and Sb (V). However, brake pads are often found to
216 have different composition based on the brand, particularly with respect to Cu and Sb (Hjortenkrans
217 et al., 2007; Canepari et al., 2008). Polyalkylene glycol ethers are also reported to be present in
218 brake wear particles with small concentrations of n-alkanes and n-alkanoic acids (Rogge et al.,
219 1993).

220
221 Iijima et al. (2007) reported a unimodal number-based distribution for brake abrasion dust with the
222 mode at 1-2 μm while the mass-based distribution had a peak at 3-6 μm . EFs for brake wear have
223 been reported by several studies including Garg et al. (2000) who reported a brake wear EF of 3-9
224 mg/km for light duty gasoline vehicles (LDGV) and Abu-Allaban et al. (2003) who reported an EF
225 of 0-80 mg/km. Wahlin et al. (2006), on the other hand, reported a Cu emission factor of 0.7 ± 0.2
226 mg/km/vehicle in Copenhagen.

227 Sternbeck et al. (2002) proposed the ratio of Cu:Sb (4.6 ± 2.3) as characteristic of brake wear
228 particles and Table 3 lists the Cu: Sb ratios as reported in the literature. Differences in the ratios can
229 be attributed to the difference in the brake pad composition, and contributions of metals from other
230 sources. In some cases, concentrations in ambient air can also vary due to the site characteristics.

231 2.3 Road Dust and Road Surface Wear

232 Road dust, of which crustal dust is a key component, consists of primarily coarse-sized particles
233 derived from different sources such as traffic, industrial emissions, mineralogical dust etc.
234 (Kupianinen et al., 2005; Tanner et al., 2008). Composition of road dust shows spatial as well as
235 temporal variation and it is often difficult to classify dust into crustal/resuspended/direct emission
236 etc. In Monterrey (Mexico), re-suspended dust was found to be contributing nearly 20-25% to the
237 PM_{2.5} EF (Mancilla and Mendoza, 2012). The amount of re-suspended road dust particles depends
238 on a number of factors including vehicle movement (particularly traffic speed), street maintenance,
239 season and associated meteorological parameters and speed of traffic (Etyemezian et al., 2003;
240 Gertler et al., 2006; Thorpe et al., 2007; Bhaskar and Sharma, 2008; Kaunhaniemi et al., 2011;
241 Laidlaw et al., 2012; Majumdar et al., 2012). Nicholson (1988) presents a detailed review of the
242 mechanisms of resuspension of road dust. Thorpe et al. (2007) reported a strong association
243 between heavy duty traffic and re-suspension in the UK with wind speed not found to be a strong
244 influence. In addition, precipitation was found to have no influence on the amount of re-suspension.
245 In somewhat related research, street-washing has been reported to be ineffective for PM control
246 based on experiments in Spain (Karanasiou et al., 2012). However, in Sweden, road wetness was
247 found to be an important factor in the amount of re-suspension (Omstedt et al., 2005), but this
248 appears related to road sanding and the use of studded tyres in Sweden. It is also important to note
249 that the residence time of PM₁₀ on paved roads (travel lanes) has been estimated as a few hours
250 (Etyemezian et al., 2003). In the case of studded tyres, interaction between tyres and the road
251 pavement can generate particles less than 0.1 µm (Gustafsson et al., 2008) and such tyres have been
252 shown to increase PM₁₀ emissions by a factor of 1.5 (Tervahattu et al., 2006). The type of pavement
253 is another important factor for road dust emissions. Granite pavements have been reported to emit
254 more PM compared to quartzite pavements (Kupiainen et al., 2005; Gustafsson et al., 2009).
255 Tervahattu et al. (2006) also explained the higher PM₁₀ concentration attributed to use of anti-skid

256 aggregate using the *sandpaper effect*, where the aggregates used to prevent skidding generate PM₁₀
257 particles and lead to further particle emissions from the pavement.

258
259 Pristane, phytane, hopanes, steranes in addition to unresolved complex mixture (UCM) were
260 reported in road dust particles which were found to be enriched in biogenic component by Omar et
261 al. (2007). High concentrations of phenanthrene, fluoranthene and pyrene along with high
262 molecular weight (4-7 ring) PAHs were reported in road dust samples in Taiwan (Fang et al., 2004),
263 China (Liu et al., 2007; Han et al., 2009) and Egypt (Hassanien and Abdel-Latif, 2008) respectively.
264 The spatial distribution of PAHs was reported to be affected by wind direction Liu et al. (2007) and
265 total organic carbon content by Liu et al. (2007) and Han et al. (2009). Mono- (α - and β -glucose)
266 and disaccharides (sucrose and mycose) were used as markers for road dust re-suspension for
267 source apportionment by Omar et al. (2007).

268 Han et al. (2007) analysed suspended dust in Beijing and reported concentrations of elements such
269 as Ca, S, Cu, Zn, Ni, Pb, and Cd to be much higher than the crustal abundances. Cu, Zn, Ni, and Pb
270 were attributed to traffic emissions together with coal burning, and ions such as Ca²⁺, SO₄²⁻, Cl⁻,
271 NO₃⁻, K⁺, and Na⁺ were also found to be present in re-suspended dust. Luo et al. (2011) observed
272 that the concentrations of trace metals contributed by vehicle exhaust and tyre abrasion, i.e. Pb, Zn
273 and Cu have been reported to be much higher in roadside samples than the concentrations of these
274 elements in background soils in China. Wei and Yang (2010) analysed urban soil, urban road dust
275 and agricultural soil in Chinese cities and found relatively higher concentrations of Cd, Cu, Pb and
276 Zn in urban soil samples and Cu, Cr, Pb, Zn, Ni and Cd in urban road dusts. The high
277 concentrations of metals were attributed to anthropogenic sources - road traffic and industrial
278 emissions. Higher concentration levels for several trace elements were reported for roadside dust
279 compared to dust collected at a distance of 100 m in Bosnia, and particularly high concentrations
280 were reported for Hg and Co and were attributed to vehicle exhaust and brake pads respectively.
281 Elements such as platinum, palladium and rhodium have also been reported to be present in road

282 dust and are attributed to the catalytic converters (Prichard and Fisher, 2012). Duong and Lee (2011)
283 analyzed heavy metal contamination in road dust in high traffic areas in Korea and concluded that
284 the concentrations of heavy metals are much higher in high traffic areas in relation to a background
285 site. Their analysis also confirmed a high degree of correlation between the concentration of heavy
286 metals at a specific location and the traffic throughput in the area. Martuzevicius et al. (2011)
287 analysed street dust and reported that it can be a major source for particle-bound PAHs together
288 with other PM. Han et al. (2009) reported a high correlation between PAHs and OC in road dust,
289 and traffic emissions were identified as a major contributor to PAHs in road dust using diagnostic
290 ratio analysis. Mathematical models have been proposed for estimation of PM emissions due to
291 road dust although they do not include emissions due to wear of brake/tyre materials (Omstedt et al.,
292 2005; Ketzler et al., 2007; Berger and Denby, 2011; Kaunhaniemi et al., 2011).

293 Aatmeeyata et al. (2009) reported a bimodal number and mass distribution (0.3 μm and 4-5 μm) for
294 PM_{10} generated due to surface (concrete pavement) and tyre wear . Duong and Lee (2011) reported
295 a multi-modal distribution for road dust particles (based on weight) collected from the roadside with
296 a majority of the particles between 180-850 μm in Ulsan, Republic of Korea. Chen et al. (2006)
297 reported a bimodal (5-10 μm and $> 30 \mu\text{m}$) mass size distribution for road dust particles in Beijing
298 (China).

299 **3. EXHAUST EMISSIONS**

300 Motor vehicles are an important source for carbonaceous aerosols particularly for the particles in
301 the fine size range (aerodynamic diameter $< 2.5 \mu\text{m}$) (El Haddad et al., 2009; Kam et al., 2012;
302 Keuken et al., 2012). PM emissions from vehicles depend on the engine type and age and
303 maintenance and contain carbon in the form of OC and EC with smaller amounts of trace metals
304 and ions (Brook et al., 2007; Robert et al., 2007a; Fulper et al., 2010; Peltier et al., 2011). For
305 example, Robert et al. (2007a,b) reported that that emission rates for ultrafine and fine particles for
306 heavy-duty diesel vehicles (HDDVs) are one order of magnitude higher than LDGVs. Similarly,

307 Chen et al. (2013) reported 4-times higher diesel EFs compared to gasoline EFs in a tunnel
308 environment. Higher emissions from the high-emitting (smoker) vehicles have previously been
309 attributed to lubricating oil (Fujita et al., 2006). Emissions from diesel and gasoline vehicles are
310 different in terms of composition as diesel engines emit both a greater mass of PM and a larger
311 number of ultrafine particles (UFPs) compared to gasoline vehicles (Rose et al., 2006). Gasoline
312 engines are known to release a higher fraction of OC while diesel engines emit more EC (Watson et
313 al., 1994; Weingartner et al., 1997; Ntziachristos et al., 2007) although Sodeman et al. (2005)
314 demonstrated a dominance of EC in the ultrafine range in gasoline vehicles. Zhu et al. (2010)
315 reported PM_{2.5} emissions from diesel and gasoline vehicles to be rich in different fraction of EC and
316 OC; while EC₂ and OC₂ fraction were higher in case of diesel vehicles, EC₃ and OC₃ were higher
317 in case of gasoline vehicle emissions. Vehicles are also a major source of n-alkanes, and diesel
318 engines are known to emit more n-alkanes compared to gasoline engines (Rogge et al., 1993).
319 Results from tunnel studies and chassis dynamometer analyses have shown that lower molecular
320 weight PAHs such as phenanthrene and fluoranthene are characteristic of diesel-derived aerosols
321 whereas gasoline engines contributed more to higher molecular weight PAHs such as
322 benzo(ghi)perylene (Smith and Harrison, 1996). PAH emissions from gasoline engines are reported
323 to be one order of magnitude higher than the diesel engine emissions integrated across the U.S.
324 vehicle fleet (Lough et al., 2005). A detailed description of diesel particulate matter is presented in
325 Maricq (2007). The situation in relation to exhaust emissions from traffic is constantly changing as
326 abatement technologies develop. In particular the adoption of diesel particle filters is greatly
327 reducing mass emissions.

328 Both trace elements and organic compounds are used as source markers for vehicle emissions.
329 However, since the removal of lead from gasoline, trace metals have proved far less useful as a
330 tracer of engine exhaust (Harrison et al., 2003). Organic compounds released from vehicles are
331 particularly useful markers for conducting receptor modelling analyses since such markers help in
332 distinguishing the vehicular emissions from other sources. The most commonly used molecular

333 markers for vehicular emissions are hopanes and steranes and PAHs such as benzo(ghi)perylene
334 which are attributed to lubricating oil (Rogge et al., 1993; Schauer et al., 1996; Cass, 1998; Lough
335 et al., 2007) and due to their source, the distribution of hopane emissions from vehicles is suggested
336 to be independent of fleet composition (He et al., 2008). It has also been suggested that hopanes and
337 steranes (derived from lubricating oil) and PAHs (reactions at high temperature and fuel
338 combustion) are indicative of different processes during the organic carbon formation in vehicle
339 engines (Lee et al., 1995; Fujita et al., 2006; Riddle et al., 2007; Hanedar et al., 2008).

340
341 A number of PAHs have been reported to be present in vehicle exhaust including benzo(e)pyrene,
342 benzo(a)pyrene, indeno(1,2,3-cd) pyrene, coronene and benzo(ghi)perylene (Rogge et al., 1993;
343 Lough et al., 2007; Riddle et al., 2007). However, Gao et al. (2011) attributed indeno(1,2,3-
344 cd)pyrene and benzo(ghi)perylene to biomass combustion in Guangzhou (China). PAHs are found
345 predominantly in the smallest size fraction ($<0.4 \mu\text{m}$), and most of the high molecular weight PAHs
346 are found in the fine particle fraction (Hien et al., 2007). Diagnostic ratio analysis including the
347 ratio between concentrations of indeno(1,2,3-cd) pyrene to the sum of indeno(1,2,3-cd) pyrene and
348 benzo(ghi)perylene is also used for distinguishing between gasoline and diesel emissions (Chellam
349 et al., 2005; He et al., 2006; Shen et al., 2010; Ancelet et al., 2011; Kim et al., 2012). Ratios
350 between methylphenanthrenes/phenanthrene has also been used for estimation of the contribution of
351 traffic to PAH concentrations in ambient air (Lim et al., 1999). However, it is important to
352 remember that PAHs can be altered by atmospheric chemical reactions (Wu et al., 2007). Also,
353 emission profiles for PAHs are difficult to generalize since they vary across the different vehicle
354 classes (Miguel et al., 1998). Congener profiles of PAH from road traffic have been derived from
355 tunnel studies (e.g. Smith and Harrison, 1996) and from roadside concentrations from which the
356 local background is subtracted (Mari et al., 2010). Nielsen (1996) also prepared a traffic PAH
357 profile for Copenhagen (Denmark) using a roadside study.

358

359 Elemental markers which have been used for vehicular emissions include Cu, Mn, Fe, Zn, Ba, Sn,
360 Ni, Mo and Sb (Lough et al., 2005; Almeida et al., 2006; Birmili et al., 2006; Crawford et al., 2007;
361 Dongarra et al., 2009; Fabretti et al., 2009; Gietl et al., 2010; Amato et al., 2011a,b). Metals can be
362 emitted from various exhaust-related sources including fuel and lubricant combustion, catalytic
363 converters, particulate filters and engine corrosion (Lough et al., 2005; Pulles et al., 2012; Sysalova
364 et al., 2012; Varrica et al., 2012) but many of these appear most likely to arise from non-exhaust
365 sources. Ni and V have also been reported to be present in emissions due to oil combustion (Pey et
366 al., 2010). Cheng et al. (2010) reported higher emission rates for elements such as Br, Ba, Sb and V
367 for gasoline and liquefied petroleum gas (LPG) engines compared to diesel engines in Hong Kong
368 and indicated potential use of these species as markers for gasoline/LPG engines. Ba, Cd, Zn, Sb
369 and V in nanoparticles have been reported to be strongly associated with diesel fuel whereas Cu,
370 Mn and Sr in the particles $<0.1 \mu\text{m}$ have been found to be associated with gasoline (Lin et al., 2005).
371 Metal concentrations from diesel and gasoline vehicles are reported to vary over two orders of
372 magnitude (Pulles et al., 2012). In a traffic emissions study in New York (USA), Ba, Br and black
373 carbon (BC) were used as tracers for gasoline/diesel combustion (Peltier et al., 2011). However, the
374 levels of trace elements emitted in the exhaust are very low, and great care is needed to distinguish
375 them from non-exhaust traffic emissions and other sources. In addition, different authors have
376 attributed metals to different sources which renders precise source attribution with trace metals
377 alone very difficult unless detailed local information on source particle composition is available (e.g.
378 Pant and Harrison, 2012).

379 **4. PARTICLE SIZE DISTRIBUTIONS**

380 The size distribution of a particular element or compound not only influences the potential health
381 impact (in terms of respiratory deposition – Harrison et al., 2010), but also influences the extent of
382 atmospheric dispersion (Allen et al., 2001). Size distributions can be measured in terms of mass, number
383 or surface area (Harrison et al., 2000). A number of factors can influence the size distribution

384 including, but not limited to vehicle fleet mix, road type and grade, meteorological conditions (e.g.
385 season type) and distance from road (Zhu et al., 2006; Beddows et al., 2009; Hays et al, 2011; Song
386 and Gao, 2011). For example, differences were observed in the modal structure of particles close to
387 the freeway and at a crossroad in Prague and the difference was attributed to the higher
388 concentration of UFPs in the freeway due to the high volume of traffic (Ondracek et al., 2011).
389 However, in a study in the USA, Padro-Martinez et al. (2012) concluded that distance from the
390 highway does not impact the particle size distribution. Many other studies (e.g. Shi et al., 1999)
391 have reported the opposite. The behaviour of nanoparticles after emission is discussed later in this
392 section. Size distributions can be useful in ascertaining the sources of exhaust and non-exhaust
393 particle emissions. For example, sampling on a busy London (UK) highway, Harrison et al. (2011)
394 used Positive Matrix Factorization (PMF) to disaggregate the particle size distribution which
395 demonstrated a nucleation mode centered on around 20 nm and a solid particle mode at 50-60 nm in
396 the number distribution. However, in using such data, it should be borne in mind that the size
397 distributions measured by cascade impactors may be modified from those in ambient air as an
398 artefact caused by the semi-volatile nature of the PAHs, hopanes and steranes.

399

400 **4.1 Mass-Size Distributions**

401 Based on size distribution analysis for trace metals found in atmospheric aerosols at urban and
402 background sites in the UK, it was concluded that there are three key categories; metals with most
403 of the mass in the accumulation (fine) mode (e.g. Cs, Sn, Pb) with an additional minor mode, metals
404 with mass distributed between fine, coarse and intermediate mode (e.g. Ni, Zn, Cu, Mn) and metals
405 with most of the mass in the coarse range (Fe, Ba, Sr) (Allen et al., 2001; Birmili et al., 2006). Song
406 and Gao (2011) reported similar results for size-segregated PM at a highway site in the USA with a
407 bimodal mass-based distribution with the two peaks at 0.32-0.56 μm and 3.2-5.6 μm Table 4
408 summarizes some of the published data for traffic-related elements.

409

410 Similar observations regarding mass-based size distributions for trace metals were reported by
411 Samara and Voutsas (2005) for a traffic site in Greece. Metals such as Ba, Pb and Zn were found to
412 exhibit bimodal distributions whereas some others such as Cr and V showed trimodal distributions
413 in Southern Taiwan (Lin et al., 2005). The presence of an Aitken mode in the ultrafine range may
414 be indicative of a contribution from vehicular emissions, both diesel and gasoline. Hays et al. (2011)
415 reported erratic mass size distributions for various elements including Ni, Cu, Cd, and Zn and
416 attributed this to a range of emission sources near highways in North Carolina (USA). A peak was
417 observed at 80 nm for soot particles in a street canyon corresponding to diesel engine emissions in
418 Leipzig, Germany and the concentration of soot particles was reported to be influenced by traffic
419 volume (Rose et al., 2006)

420

421 **4.2 Number-Size Distributions**

422 Ultrafine particles are characterized as particles with diameter less than 100 nm diameter and are
423 typically measured as particle number concentration (PNC) (Charron and Harrison, 2009). Such
424 particles may have a short lifetime and as a result, the spatial variability of the particles differs
425 based on proximity to the emission source (Puustinen et al., 2007; Cyrys et al., 2008; Charron and
426 Harrison, 2009). Traffic has been reported as one of the major sources of UFPs (Abu-Allaban et al.,
427 2002; Nanzetta and Holmen, 2004; Westerdahl et al., 2009) and particle number is linked to fresh
428 vehicle exhaust emissions (Rodriguez et al., 2007). Number size distributions are heavily influenced
429 by proximity to traffic (Rose et al., 2006; Kerminen et al., 2007; Beddows et al., 2009; Keuken et
430 al., 2012). As a consequence of this, concentrations of organic compounds have also been reported
431 to be highly variable for near-highway sites (Sun et al., 2012). Highest PNC has been reported
432 during morning rush hours in Germany (Wehner et al., 2002) and Spain (Perez et al., 2010). A
433 significant difference has also been reported between weekdays and weekends, highlighting the
434 influence of traffic (Morawska et al., 2002; Wehner et al., 2002; Van Poppel et al., 2012). A
435 significant difference was observed in the particle number size distribution in the Kingsway tunnel

436 in the UK (peak at 30-40 nm) and Plabutsch tunnel in Austria (80-100 nm) and this was attributed
437 to the differences in traffic intensity and the fleet composition, i.e. proportion of HDVs and LDVs
438 (Imhof et al., 2005a). It is important to note that although PNC is heavily influenced by road traffic,
439 it is also influenced by other sources including atmospheric nucleation processes (Imhof et al.,
440 2005a; Reche et al., 2011).

441
442 The typical particle number distribution of a vehicle has 3 modes: nucleation mode (particles < 50
443 nm), consisting of particles formed due to condensation of exhaust particles; accumulation mode
444 (50-300 nm with a peak at ~ 80-100 nm) consisting of soot particles, and a coarse mode (1µm-
445 10µm) consisting of particles generated due to abrasion and wear and tear of brake pads and tyres
446 and re-suspension of road dust (Kittleson et al., 2004; Imhof et al., 2005a). The number size
447 distribution for diesel engine emissions has been reported as lognormal with mean diameter
448 between 60-120 nm whereas gasoline engine emissions have asymmetric distributions with a mean
449 diameter between 40-80 nm (Harris and Maricq, 2001). PNC shows higher variability compared to
450 particle mass (Weijers et al., 2004). While nuclei mode particles only constitute about 10% of the
451 total particle mass, they contribute more than 90% to PNC (Kittleson et al., 2004). Fushimi et al.
452 (2008) reported the presence of a nuclei mode at 20 nm in the number distribution for PM at
453 roadside while no such mode was observed at the background site in Kawasaki, Japan. PNC was
454 reported to be significantly higher in a street canyon compared to an urban background location in a
455 study in Leipzig, Germany (Wehner et al., 2002).

456
457 The particle number distribution for vehicular exhaust emissions measured in ambient air is
458 typically bimodal with one peak at 10-40 nm and another one at 50-90 nm (Charron and Harrison,
459 2009). Gasoline exhaust particles fall in the size range of 20-60 nm whereas diesel exhaust particles
460 are in the size range of 20-130 nm (accumulation mode between 50-90 nm and a smaller nucleation
461 mode) (Zhu et al., 2002b; Rissler et al., 2012). Such bimodal distributions have been reported for
462 vehicular emissions in many parts of the world including Korea (30-50 nm and 100-400 nm for

463 diesel engines) (Lim et al., 2008), Czech Republic (25-30 and 60-70 nm) (Ondracek et al., 2011)
464 and United Kingdom (13.3 nm and 86.6 nm) (Kumar et al., 2008). On the other hand, multi-modal
465 distributions have been reported in several cities and the larger number of modes has been attributed
466 to mixing of traffic and background aerosol (Ondracek et al., 2011). Lingard et al. (2006) reported
467 a four-mode particle number size distribution for an urban roadside site in Leeds (UK) with
468 particles in modes II and III to be highest during the rush hours indicating a traffic source. The four
469 modes were classified as sub-11 nm nucleation mode, super-11 nm nucleation mode, Aitken mode
470 and accumulation mode and were attributed to photolysis, diesel engine emissions and secondary
471 aerosols (Lingard et al., 2006). Rosenbohm et al. (2006) observed a bimodal (20 nm and 80 nm)
472 size distribution at a site downwind of the motorway whereas a unimodal distribution (80 nm) was
473 observed for an upwind site. El Haddad et al. (2009) reported a single mode (50-90 nm) distribution
474 for PM in a road tunnel in Marseille, France and Abu-Allaban et al. (2002) observed the highest
475 PNC in the nucleation mode (11-17 nm) in a roadway tunnel in USA.. A bimodal (11 nm and 80 nm)
476 distribution was reported for PM_{0.5} emissions during a vehicle chase experiment (Wang et al., 2011).
477 Although the magnitude of PNC emissions from a gasoline spark ignition engine exhaust is smaller
478 than from a diesel engine, emissions from gasoline engines are comparable to diesel engines in the
479 cases of acceleration and freeway cruising (Kittleson et al., 2004). Klems et al., (2010) also reported
480 higher PNC at the time of acceleration at traffic lights.

481 Significant differences have been reported between dynamometer and real-world studies on PNC. A
482 mean diameter of 15 nm for PM emissions was measured in Leipzig, Germany in a street canyon
483 though some lab-studies on gasoline and diesel engines have shown mean diameters between 40-70
484 nm (Wehner et al., 2002). It is important for dynamometer studies to reflect adequately the dilution
485 properties of the atmosphere or they may suppress the nucleation mode which is formed during the
486 initial dilution of the exhaust plume (Shi and Harrison, 1999).

487

488 For traffic emissions, distance from the road is also reported to cause changes in PNC due to
489 processes other than dilution (Zhu et al., 2006). This feature is of considerable interest; particularly
490 from an epidemiological perspective as this would mean that exposure from near-roadway sites can
491 be much different from other sites. HEI (2010) reported the key exposure zone for traffic emissions
492 to be within 300-500 m of a roadway. In Los Angeles, both PNC and size distribution changed with
493 an increase in distance from the road, particularly for particles with a diameter less than 30 nm and
494 it has been suggested that this drastic change in PNC is caused by coagulation and atmospheric
495 dilution (Zhu et al., 2002a). Padro-Martinez et al. (2012) found PNC to be at least two times higher
496 at 0-50 m from the highway compared to further distances while Zhu et al. (2002b) observed the
497 PNC to be 25 times higher near the freeway compared to background locations. Rose et al. (2006)
498 also observed a decrease in soot particle number concentration with increasing distance from the
499 street canyon which was attributed to coagulation processes. Kittleson et al. (2004) observed a
500 decline in the nuclei mode particles as the distance from the highway increased and recently,
501 Canagaratna et al. (2010) also reported a decrease in PNC with increasing distance away from the
502 roadway. Dilution processes will inevitably cause a reduction in PNC with distance from the road,
503 as they do for other traffic-generated pollutants (e.g. NO_x, BC, etc.). Coagulation causes an
504 increase in particle size, and a decrease in number, but is unlikely to be a significant influence close
505 to the road due to an insufficient number concentration to drive the process, and short travel times.
506 An observed increase in particle diameters with distance from a road is most probably the result of
507 mixing of traffic emissions with a coarser background aerosol leading to a large modal diameter
508 (Shi et al., 1999). The nucleation mode traffic particles are semi-volatile and this can lead to shifts
509 in size distribution (typically to smaller sizes) and particle loss due to evaporation (Dall'Osto et al.,
510 2011; Harrison et al., 2012a).

511
512 Charron and Harrison (2003) reported that increasing wind speeds and lower temperatures can
513 increase the abundance of smaller nucleation mode particles (diameter of sub-30 nm) and this could

514 be due to the greater dilution and lower condensation sink. Ruellan and Cachier (2001) also
515 reported an influence of wind speed and boundary layer height on PNC. Kumar et al. (2008)
516 reported traffic volume and wind speed as the key variables influencing PNC in street canyon
517 conditions and they observed a linear dependence of PNC on traffic volume with a higher
518 correlation in the case of smaller particles (diameter < 300 nm) and a weaker correlation in the case
519 of larger particles for which other sources are large relative contributors. Highest PNC was
520 observed for calm winds and lowest for wind speeds above 1.6 m/s in Somerville (USA) (Padro-
521 Martinez et al., 2012). PNC was found to be correlated with traffic intensity in France (Ruellan and
522 Cachier, 2001; El Haddad et al., 2009; Perez et al., 2010) and in Los Angeles, PNC was reported to
523 be influenced by traffic density and wind speed and direction (Zhu et al., 2002b). In a study in
524 London, Milan and Barcelona, Rodriguez et al. (2007) concluded that vehicular exhaust emissions
525 largely influence ultrafine particles and found a unimodal distribution for particle number with a
526 mode at 37 nm. Kittleson et al. (2004) observed a direct influence of increasing vehicle speed on
527 nuclei-mode particles. Decreasing the engine load from 100% to about 60% was reported to
528 decrease the PNC as well as EC and OC emissions from diesel vehicles (Lim et al., 2008).
529 Morawska et al. (2002), on the other hand, indicated a 70% increase in PNC with a 50% increase in
530 traffic flow. While roadside PNC measurements have indicated an influence of temperature on the
531 concentration, similar studies in laboratory conditions using chassis dynamometers have not shown
532 such influence (Abu-Allaban et al., 2002; Kittleson et al., 2004). Abu-Allaban et al. (2002) reported
533 an inverse relationship between ambient air temperature and the diameter of nucleation mode
534 particles. Using modelling analysis, Jayaratne et al. (2009) observed an increase in PNC at a
535 signalized pedestrian crossing due to the stop and go nature of traffic at the signal and the increase
536 was found to be higher for HDVs compared to LDVs. This is similar to the observations from Abu-
537 Allaban et al. (2002) and Wang et al. (2010) who observed higher concentrations arising from
538 HDVs compared to LDVs in USA and Denmark respectively. Rosenbohm et al. (2006) reported a
539 correlation between PNC and NO in Heidelberg (Germany) although no correlation was observed

540 for PNC and PM₁₀ concentration. Similarly, Padro-Martinez et al. (2012) also reported a similar
541 spatial trend for PNC, NO_x, particulate PAHs and BC in the USA while a different trend was
542 reported for PM_{2.5}.

543
544 Thus, key factors that influence the particle number concentration measurements include wind
545 speed, traffic density and the distance from source at which the measurements are made. Nanzetta
546 and Holmen (2004) observed differences in the PNC across different seasons and concluded that
547 results for one season cannot be used for generalization of PNC.

548

549 **5. METHODOLOGIES FOR ASSESSMENT OF TRAFFIC EMISSIONS**

550 There are several ways of evaluating emissions from road vehicles including tests with
551 dynamometers, measurements in tunnels/highways and near roadside measurements and road
552 simulator tests (Figure 1) (Mclaren et al., 1996; Phuleria et al., 2007; Handler et al., 2008;
553 Gustafsson et al., 2008; El Haddad et al., 2009; Yan et al., 2009; Sjodin et al., 2010). Table 5
554 presents a qualitative description of measurements using dynamometers under lab conditions and
555 tunnel/roadway/twin-site studies under ambient conditions. In recent years, significant differences
556 have been observed between laboratory-tested and real-world mixed source traffic emissions
557 (Gertler et al., 2002; Yan et al., 2009; Ancelet et al., 2011). Near roadside/kerbside, traffic tunnel
558 and highway measurements are reported to be more realistic since they represent mixed-fleet
559 emissions under real-world driving conditions (Phuleria et al., 2007). Beddows and Harrison (2008)
560 compared particle number EFs measured in dynamometer studies with those estimated in roadside
561 studies, finding generally good agreement overall. However, Canagaratna et al. (2010) reported the
562 EFs calculated using on-road measurements to be nearly 50% lower than EFs measured using a
563 dynamometer.

564

565 An emission factor is defined as “the mass (or number) of pollutant released per unit time/distance
566 travelled or mass of fuel used” and typically increases with vehicle speed for LDVs as well as
567 HDVs (Jones and Harrison, 2006). Characterization of emissions from road vehicles includes
568 analysis of size distributions and chemical composition (organic, inorganic and trace metal) and can
569 lead to generation of EFs. EFs typically depend on conditions at the sampling site, traffic intensity
570 and modal shares of HDVs and LDVs (Staehelin et al., 1995; Imhof et al., 2005a). A recent review
571 article on development of emission factors delves into this issue in greater detail (Franco et al.,
572 2013).

573
574 EFs can be measured using a number of different methods including dynamometer studies (Oanh et
575 al., 2010; Chiang et al., 2012), roadway tunnel studies (Gertler et al., 2002; Handler et al., 2008;
576 Mancilla and Mendoza, 2012), on-road chase experiments (Wang et al., 2011) and remote sensing
577 (Zhang et al., 1995). Phuleria et al. (2007) reported overall agreement between EFs determined at
578 tunnel and roadway sites in USA.

579
580 Two of the most common methods for EF expression are fuel-based EF (Ban-Weiss et al., 2010;
581 Wang et al., 2011) and vehicle kilometre travelled (VKT)-based calculation (Abu-Allaban et al.,
582 2002; Kittleson et al., 2004; Jones and Harrison, 2006; Amato et al., 2012). Phuleria et al. (2006,
583 2007) and Bukowiecki et al. (2010) describe the methodology of EF derivation in greater detail.
584 Fuel-based EF can be defined as “grams of pollutant emitted for every kilogram of fuel burned” and
585 can be calculated for various pollutants using CO₂ and CO as a measure of fuel burned (Liacos et al.,
586 2012). The concentration of pollutant with respect to the concentration of CO₂ + CO can be used to
587 calculate the fuel burnt using the following formula (Miguel et al., 1998; Liacos et al., 2012):

$$EF_P = 10^6 \times \left(\frac{\Delta P}{\Delta CO_2 + \Delta CO} \right) W_c \quad (1)$$

588 where EF_p is the emission factor for the pollutant ($\mu\text{g}/\text{kg}$ fuel burned), ΔP (ng/m^3), ΔCO_2 and ΔCO
589 ($\mu\text{gC}/\text{m}^3$) are the increments over background concentration and w_c is the fraction of carbon weight
590 in the fuel.

591
592 One of the other most commonly used methods is the derivation of PM EFs using NO_x EFs, which
593 are calculated as described in the following equation (Gehrig et al., 2004; Imhof et al., 2005b). It is
594 assumed in this case that both NO_x and PM have similar dilution.

$$595 \quad D = \frac{N_{\text{LDV}} \times EF_{\text{NOx LDV}} + N_{\text{HDV}} \times EF_{\text{NOx HDV}}}{\Delta\text{NOx}} \quad (2)$$

596 where D is the dilution rate ($\text{m}^2 \text{h}^{-1}$), n is the number of vehicles (h^{-1}), and EF_{NO_x} is the emission
597 factor for NO_x in mg/km .

598
599 Using the dilution rate, the EF (per hour) can be calculated per vehicle based on:

$$EF_x = \frac{\Delta\text{Concentration}_x \times D}{n_{\text{total}}} \quad (3)$$

600 where $\Delta\text{Concentration}_x$ is the difference in concentration of species x between roadside and
601 background and n is the total number of vehicles crossing the sampling point per hour. The
602 respective contribution of HDVs and LDVs can then be estimated with multiple linear regression
603 (MLR) (Gertler and Pierson, 1996; Imhof et al., 2005b). Table 6 presents some EFs reported for
604 PM_{10} , $\text{PM}_{2.5}$ and particle number concentration in the literature.

605

606 **6.1 Tunnel/Roadway Measurements**

607 EF calculations from tunnel studies are typically based on calculation of the difference in
608 concentration between entrance and exit (Weingartner et al., 1997; Abu-Allaban et al., 2002), and
609 Gertler and Pierson (1996) describe the methodology for calculation of emission factors using
610 tunnel measurements. Chang et al. (1981) developed a model relating air quality in the tunnel to

611 vehicular emissions which can also be used for calculation of emission factors using tunnel
612 measurements. For the tunnel studies, “the velocity and concentration are assumed to be the same
613 across the tunnel’s cross-section and the difference between the influx and outflux is assumed to be
614 the sum of emission rates of vehicles in the tunnel” (Rogak et al., 1998). The following formula is
615 typically used for calculation of the EF (Gertler and Pierson, 1996; He et al., 2008):

$$EF = \frac{C_{out}V_{out} - C_{in}V_{in}}{NL} \quad (4)$$

616 where C is the concentration of the pollutant, V is the volume of air calculated using duration of
617 sampling, cross-sectional area of tunnel and wind speed; N is the number of vehicles and; L is the
618 distance between sampling locations at inlet and outlet of the tunnel.

619

620 Several studies have been conducted for measurement of PM emissions in roadway tunnels in
621 different parts of the world including the USA (Fraser et al., 1998; Rogak et al., 1998; Gillies et al.,
622 2001; Abu-Allaban et al., 2002; Gertler et al., 2002), China (He et al., 2006; He et al., 2008; Chiang
623 and Huang, 2009), Europe (Stechmann and Dannecker, 1990; Smith and Harrison, 1996;
624 Weingartner et al., 1997; Imhof et al., 2005a; Handler et al., 2008; El Haddad et al., 2009; Oliviera
625 et al. 2011), New Zealand (Ancelet et al., 2011), Japan (Funasaka et al., 1998), Taiwan (Chiang and
626 Huang, 2009), Chile (Caceres et al., 1998) and Mexico (Mancilla and Mendoza, 2012).

627

628 Although tunnel studies provide an opportunity to conduct measurements on a mixed vehicle fleet,
629 it is important to remember that factors such as variations in speed, aerodynamic conditions in the
630 tunnel and the fleet characteristics (i.e. proportion of HDVs and LDVs) can cause variability in
631 measurements (Rogak et al., 1998; He et al., 2008). In addition, vehicles in the tunnel are often
632 driving at a steady speed which does not happen under other road conditions where traffic follows a
633 stop-and-go pattern (Gertler et al., 2002). Chirico et al. (2011) have shown that the distribution of
634 organic compounds between particles and vapour is heavily affected by the high PM concentrations

635 in a road tunnel, and thus may influence the estimate of emission factors for semi-volatile
636 components.

637
638 PM₃ (particles with diameter < 3 µm) was analysed in the Gubrist Tunnel in Zurich, Switzerland
639 and it was concluded that most of the particles are emitted from the tailpipe with a small
640 contribution from tyre wear and road dust (Weingartner et al., 1997). While BC accounted for 31%
641 of the total emissions, particulate PAHs accounted for 0.86% of PM₃ (Weingartner et al., 1997). In
642 New Zealand, PAHs were reported to contribute nearly 0.10% of the total PM mass with a higher
643 proportion of low molecular weight PAHs indicating a higher contribution of diesel emissions
644 (Ancelet et al., 2011). In roadway tunnel studies in China, EC was found to contribute 63% and 45%
645 to PM mass whereas OC contributed 34% and 31% (He et al., 2006; He et al., 2008). Ancelet et al.
646 (2011), however, found OC to contribute between 32-46% to PM mass. Gillies et al. (2001) and He
647 et al. (2008) have attributed higher PM_{2.5} emissions to HDVs compared to light duty vehicles
648 (LDVs) in USA and China respectively. Caceres et al. (1998) prepared a PAH source signature for
649 traffic using tunnel measurements in Santiago de Chile whereas Smith and Harrison (1996)
650 prepared a source profile fingerprint using tunnel measurements for traffic in Birmingham, UK.
651 Funasaka et al. (1998) analysed emissions in a traffic tunnel in Osaka, Japan and EC and NO_x (NO
652 as the main component) were found to be correlated with diesel traffic volume whereas OC did not
653 show good correlation. In addition, the concentration of EC and SPM was observed to increase
654 linearly with an increase in traffic volume. In Lisbon, (Portugal), nearly 84% of the total PAH mass
655 in a road tunnel was reported to be less than 0.49 µm diameter by Oliviera et al. (2011).

656 Concentrations of trace elements were not found to be correlated with EC concentration in Zhujiang
657 Tunnel (China) and the authors concluded non-exhaust sources to be contributing to the trace
658 element mass (He et al., 2008). A similar conclusion was reported by Funasaka et al. (1998) and
659 Ancelet et al. (2011) who attributed coarse particles in a road tunnel in Osaka, Japan and inorganic
660 elements such as Fe, Zn etc. respectively to non-exhaust emissions, primarily re-suspension of road

661 dust. It should be noted, however, that the contributors to road dust in a tunnel will differ from
662 those in a normal roadway environment and that both chemical composition and emission factors
663 will be different. Partitioning of the semi-volatile components between particles and vapour can
664 also be influenced by the high concentrations present (Chirico et al., 2011).

665

666 **6.2 Twin-Site Studies**

667 A number of papers have reported the estimation of the contribution of traffic emissions to total PM
668 using twin-site studies in recent years (Yan et al., 2009; Bukowiecki et al., 2010; Gietl et al., 2010;
669 Oliviera et al., 2010; Pey et al., 2010). With the assumption that all sources other than traffic
670 (including any local or regional sources) have the same impact at both roadside and background
671 sites, the increment at the roadside site obtained using the following equation is used as a local
672 traffic increment estimate (Harrison, 2009; Yan et al., 2009; Wang et al., 2010).

$$673 \quad \textit{Concentration of } X_{\text{traffic}} = \textit{Concentration of } X_{\text{roadside}} - \textit{Concentration of } X_{\text{background}} \quad (5)$$

674 Typically, the difference between rural and urban concentration observation provides an estimate
675 for the urban increment while the difference between roadside and urban background concentration
676 observations provides an estimate for the traffic increment (Harrison, 2009). However, results from
677 such studies can be influenced by street geometry. Harrison et al. (2004) observed different
678 behaviour for particle mass based on the site geometry, where the traffic contribution increases for
679 larger daily background mass concentrations at open sites, whereas the magnitude of the traffic
680 mass increment is independent of daily background concentration for enclosed sites, thus making
681 selection of a suitable sampling location critical for coherent results. They also found the pattern of
682 air circulation to be an important determinant of ambient PM concentration at an enclosed street site.
683 Roadway studies are useful in providing a reasonable estimate of pollutant concentrations that
684 commuters are exposed to (Phuleria et al., 2007) but if the sampling is carried out only at roadside

685 sites, the data generated is representative only of that part of the total vehicle fleet that was sampled
686 (Liacos et al., 2012).

687
688 Harrison et al. (2003) concluded that the trace element enrichment observed at roadside sites is
689 contributed largely by wear and tear of tyres and brakes. Significant roadside enrichment has been
690 reported for elements, ions and organic compounds in various cities including London (UK) (Gietl
691 et al., 2010) Los Angeles (USA) (Ntziachristos et al.,~~2008~~2007), Sicily, Rome and Milan (Italy)
692 (Manno et al., 2006; Canepari et al., 2008; Perrone et al., 2012), Barcelona (Spain) (Amato et al.,
693 2011), Portugal (Oliviera et al., 2011) and Queensland (Australia) (Gunawardana et al., 2011). . In
694 New York City, Fe, Al and Si were found to contribute between 0.5- 1.0 $\mu\text{g}/\text{m}^3$ to the ambient
695 $\text{PM}_{2.5}$ loading (Peltier et al., 2011) whereas Lin et al., 2008 reported Ca, Al and Fe to constitute
696 between 65-77% of suspended particles in Taiwan. Nearly 90% of the road traffic emissions of Cu
697 were reported to be due to brake wear in Stockholm (Sweden) (Johansson et al., 2009). Amato et al.
698 (2011b) analyzed roadside PM_{10} enrichment (RE) of trace elements in Barcelona (Spain) reported
699 more than 70% roadside enrichment ratio for key traffic tracers including EC, Fe, Ba, Cu, Sb, Cr
700 and Sn. Gietl et al. (2010) estimated that based upon published emission factors brake dust
701 comprises nearly 13% of the total PM at a roadside site in London (UK) and Amato et al. (2011b)
702 reported approximately 60% enrichment of brake wear particles in a roadside location compared to
703 an urban background location in Barcelona (Spain). Pey et al. (2010) reported that the
704 concentration of EC was found to be 3-8 times higher for the urban site in comparison with the
705 background site in Barcelona (Spain) and Perrone et al. (2012) found EC to be contributing 11-15%
706 of $\text{PM}_{2.5}$ in urban areas compared to 3-5% at a rural site and 1-2% at the background site in Milan
707 (Italy). Vertical concentration gradients have also been reported for trace elements with highest
708 concentrations at kerbside as compared with rooftop (Johansson et al., 2009). Analysis of size-
709 resolved $\text{PM}_{2.5}$ at a roadside site in Beijing revealed that re-suspended dust and construction dust

710 contribute 32.7-50.4% of total PM_{2.5} (Song et al., 2012) and a reduction was observed in
711 concentrations of species associated with traffic during the 2008 Olympic Games).

712
713 Comparatively higher concentrations of hopanes, steranes, n-alkanes and PAHs have been reported
714 for highway/near-roadway sites compared to other sites (Yan et al., 2009; Aldabe et al., 2011;
715 Peltier et al., 2011; Perrone et al., 2012). Yan et al. (2009) analysed PM_{2.5} concentrations at urban
716 and rural sites and observed a higher concentration for EC and primary organic compounds released
717 by motor vehicle exhaust such as hopanes and steranes. They found organic matter to be
718 contributing between 51-72% to PM_{2.5} mass at the urban site compared with 40-54% at the rural
719 site. Aldabe et al. (2011) reported a concentration gradient for total carbon with the highest
720 concentration at the roadside site and lowest concentration at a rural site, and a decreasing trend of
721 OC concentration was observed in Prague with a decrease in traffic intensity (Ondracek et al., 2011).
722 Gao et al. (2011) also reported higher concentrations for EC and hopanes at the roadside site
723 compared to a rooftop site, highlighting the contribution of traffic emissions. Jones and Harrison
724 (2006) used ratios of concentration to NO_x for determination of EFs for road traffic or particle
725 number and mass metrics with the assumption that particle dispersion is similar to dispersion of
726 NO_x in the atmosphere. The EFs that they obtained from this study for particle mass and number
727 (Table 6) were comparable with emission factors calculated using other methods such as
728 dynamometer and tunnel/roadway studies. Liacos et al. (2012) observed higher concentrations of
729 trace metals using on-road measurements and postulated that studies using sampling from near
730 roadside conditions might not always capture the pollutant concentrations that drivers are exposed
731 to (Liacos et al., 2012).

732

733 **6.3 NO_x as Tracer**

734 With the assumption that road traffic is the primary urban source for NO_x in the urban atmosphere,
735 Harrison et al. (1997) used regression analysis between particle mass fractions and NO_x to obtain

736 contributions from the traffic source for Birmingham (UK) with a total of 32% of the fine PM
737 concentration attributed to vehicular exhaust emissions for the winter season. The method is very
738 simple, using the assumption that the regression intercept in PM at zero NO_x is due to non-traffic
739 sources, with the traffic contribution estimated by difference from the measured mean concentration.
740 The same approach was also used by Fuller et al. (2002) who estimated PM₁₀ and PM_{2.5}
741 concentrations using NO_x and PM emissions for London (UK) and derived a typical PM₁₀: NO_x
742 ratio of 0.045± 0.003. Lim et al. (1999) also used this approach to determine the contribution of
743 PAHs to traffic in Birmingham and concluded that traffic contributed 80-82% and 61-67% of PAH
744 concentrations in the city centre and an urban background site (university campus) respectively.
745 Thorpe et al. (2007) used this method to calculate re-suspension EFs. The PM₁₀ emission factor was
746 calculated using NO_x as a tracer and coarse particle source strength and abrasion emissions were
747 subtracted from total PM₁₀ emission to obtain a re-suspension emission factor. However, it is
748 important to note that this method is only useful if the predominant source for local ground-level
749 NO_x is traffic, unless used in a twin-site approach.

750

751 **6.4 Other Methods**

752 Several studies have reported lower levels of vehicular emissions on weekends (Rose et al., 2006;
753 Barmpadimos et al., 2011). Analysis of the weekly cycle of coarse PM concentrations (2.5-10 µm)
754 with the weekly PM coarse traffic emission was carried out in Zurich (Switzerland) and it was
755 estimated that nearly 70% and 53% of the ambient coarse PM mass is contributed by traffic on
756 weekdays and weekends respectively (Barmapadimos et al., 2011). With the assumption that the
757 difference in PM concentrations between weekdays and weekends at an urban background site is
758 due to traffic movement, it is possible to calculate the traffic increment to urban PM concentrations,
759 although the results show large uncertainties (Jones et al., 2008).

760 Vehicle-chase experiments have also been used where vehicles are chased for a specific period of
761 time to collect pollutant concentrations allowing data to be collected for real-world conditions of
762 dilution (Canagaratna et al. 2010; Wang et al., 2011). However, the EFs are only valid for the
763 specific sampling conditions and there is no control over driving conditions (Wang et al., 2011).

764 6. RECEPTOR MODELLING OF TRAFFIC PM

765 Watson and Chow (2007) describe receptor models as models that “interpret measurements of
766 physical and chemical properties taken at different times and places to infer the possible sources of
767 excessive concentrations and to quantify the contributions from those sources”. A number of
768 receptor models are used for source apportionment including the Chemical Mass Balance (CMB)
769 model, statistical models such as PCA and PMF, Multilinear Engine (ME), Constrained Physical
770 Receptor Model (COPREM) and UNMIX. With the assumption that the relative concentrations of
771 chemical species are preserved between sources and receptors, receptor models use the principle of
772 mass conservation for apportionment of PM mass to different air pollution sources. Thus, the
773 concentration of a species measured in a particular sample can be described as (Hopke, 2003):

$$X_{ij} = \sum_{p=1}^p g_{ip} f_{pj} + e_{ij} \quad (6)$$

774 where X_{ij} is the measured concentration of the j th species in the i th sample, f_{pj} is the concentration of
775 the j th species in material emitted by the source p , g_{ip} is the contribution of the p th source to the i th
776 sample and e_{ij} is the portion of the measurement that cannot be fitted by the model. Different
777 receptor models use different approaches to solve this equation, for e.g., the CMB model uses the
778 effective-variance least squares method whereas UNMIX uses eigenvector analysis. Concentrations
779 of trace elements (e.g. Si, Fe, Cu), ions (e.g. SO_4^{2-}), EC/OC and organic compounds (e.g. PAHs,
780 hopanes, alkanolic acids) are used as inputs for the models.

781

782 In recent years, there have been many studies for source apportionment of road traffic-generated
783 PM, some of which appear in Table 7. There have also been studies that have used both CMB and
784 factor analysis models (Larsen and Baker, 2003; Bullock et al., 2008).

785

786 **6.1 Multivariate Statistical Methods**

787 There are a number of different methods based on factor analysis including PCA, PMF, UNMIX
788 and ME. Such methods do not require a priori information about source emission characteristics and
789 are useful in cases where relevant source profiles are not available (Hopke, 2003; Viana et al.,
790 2008). In the simplest matrix form, the equation 6 can be represented as (Hopke, 2003)

$$X = GF'$$

791 A number of studies have been conducted using the different factor analysis methods and studies in
792 different locations have often reported different magnitudes of exhaust and non-exhaust emissions.
793 In Taiwan, gasoline, diesel and fuel oil emissions together with dust and brake wear were found to
794 be key contributors to traffic-generated PM using PCA (Lin et al., 2005). Using COPREM, Wahlin
795 et al. (2006) apportioned the local traffic emissions to exhaust, road dust, brake wear and winter
796 salting in Copenhagen (Denmark). Amato et al. (2009) analysed road dust in Barcelona (Spain)
797 using ME-2, and re-suspension was found to contribute 37%, 15% and 3% of total traffic emissions
798 respectively in PM₁₀, PM_{2.5} and PM₁ respectively. Other identified sources included secondary
799 aerosols and a soil source. Sjodin et al. (2010) found road wear to be the most important source of
800 PM₁₀ emissions in streets as well as background locations in Sweden, which is probably a reflection
801 of road sanding and use of studded tyres. Using PMF, three distinct traffic sources including vehicle
802 abrasion (36%), re-suspension (43%) and fuel combustion (21%) were resolved for traffic sources
803 using trace metals in Nice (France) from data collected in a road tunnel (Fabretti et al., 2009). Based
804 on PMF analysis in Switzerland, Bukowiecki et al. (2010) reported 21%, 38% and 41% of the
805 traffic emissions to be contributed by brake wear, re-suspended road dust and exhaust emissions

806 respectively for street canyons. For freeways, while the contribution of exhaust emissions remained
807 the same, the contribution of brake wear was much less at 3% whereas re-suspended road dust
808 contributed as much as 56%. Movement of traffic also contributes to re-suspension of road dust and
809 re-suspended road dust has been estimated to contribute as much as 38% of the total PM in a street
810 canyon and 56% on a highway (Bukowiecki et al., 2010; Laidlaw et al., 2012). Analysis of PM in
811 Madrid (Spain) using PMF estimated that nearly 29% of the PM₁₀ in the city is contributed by road
812 dust while another 31% is contributed by direct vehicular emission (Karanasiou et al., 2011). In
813 Oporto (Portugal), exhaust emissions and re-suspension due to traffic were found to contribute 12-
814 55% of the coarse PM and 14-66% of the fine PM based on PCA- multiple regression analysis and
815 inter-site mass balance analysis (Oliviera et al., 2010). They used inter-site mass balance for
816 separating the traffic increment in the road dust and concluded that nearly 30% of the fine fraction
817 and 50-70% of the coarse fraction PM is contributed by road dust. They also used a roadside impact
818 factor (defined as the fractional difference between the concentrations at the roadside and urban
819 background) and the highest factors were observed for Cu, Zr, Fe, Cr and Sn indicating a traffic
820 source for these metals. Thorpe et al. (2007) reported re-suspended road dust, abrasion source
821 emissions and exhaust and fine-range abrasion emissions to contribute 20-22%, 19% and 60% to
822 roadside incremental PM₁₀ concentration respectively in a major highway in London. Four major
823 sources for road surface dust particles of diameter less than 10 µm including motor exhaust, brake
824 and tyre wear and road surface wear together with background urban dust were reported for
825 European cities with an increase in the concentrations of trace metals and OC/EC with an increase
826 in traffic volume (Amato et al., 2011a). Song and Gao (2011) identified three distinct sources of
827 trace metals at a highway site in New Jersey (USA) using factor analysis: brake wear and fuel
828 combustion, primary fuel combustion and tyre wear and fuel combustion. PCA analysis was used
829 for determining PM_{2.5} sources in Incheon (Korea) and motor vehicles/sea salt and soil emissions
830 were found to contribute 39% and 7% respectively (Choi et al., 2012). Similar results were obtained

831 by Cheng et al. (2011) in Hong Kong where traffic emissions and re-suspended dust emissions were
832 found to contribute 38% and 5% using PMF.

833 Dreyfus et al. (2009) conducted PMF analysis with time-resolved organic molecular marker data
834 and identified diesel and road dust/car emissions factors which accounted for 2/3rd of the total OC
835 concentration. Receptor modelling analysis for PM_{2.5} was conducted using PMF and UNMIX, and
836 PMF distinguished between diesel and gasoline engine emissions using temperature resolved
837 carbon fractions (Sahu et al., 2011). Pey et al. (2009) performed source apportionment using PCA
838 and regression with a large dataset including information on particle number concentration; size
839 distribution and chemical composition for an urban background site in Barcelona, and road traffic
840 was identified as the most significant source (52- 86%) for aerosols in the sub-micrometer range.

841 Liu et al. (2007) attributed the PAHs in road dust in Shanghai (China) to traffic emissions and coal
842 combustion using principal component analysis (PCA) and compositional analysis. Using the same
843 technique, Han et al. (2009) attributed PAHs in road dust in Anshan (China) to emissions from
844 traffic, coal combustion, steel industry and cooking. Using MLR, Boonyatumanond et al. (2007)
845 reported tyres, diesel vehicle soot and used engine oil to be the key sources for PAHs found in street
846 dust in Bangkok (Thailand). Hassanien and Abdel-Latif (2008) attributed PAHs in road dust in
847 Cairo (Egypt) to emissions from traffic, industry and incomplete combustion of waste based on
848 correlation analysis. Teixeira et al. (2012) reported vehicular emissions and coal and wood
849 combustion to be the key sources for PAHs in Porte Alegre (Brazil) using PCA.

850 Other approaches have also been used for estimation of source contributions. Recently, Harrison et
851 al. (2012b) used field data for estimating the contribution of brake wear, tyre wear and road dust
852 using size distributions of Ba, Zn and Si as the tracers for the sources respectively. By merging size
853 distributions measured with a Scanning Mobility Particle Sizer (SMPS) and an Aerodynamic
854 Particle Sizer (APS), Harrison et al. (2011) were able to generate continuous particle size
855 distributions for a roadside site from 15 nm – 10 µm diameter. By applying PMF, they were able to

856 separately identify nucleation mode and accumulation mode exhaust particles as well as brake dust
857 and re-suspension particles, and several particle classes arising from the urban background.

858

859 **7.2 Chemical Mass Balance (CMB) Model**

860 The CMB model uses the ambient measurement data for chemical species together with the
861 associated uncertainty and source profiles for different sources as inputs and the output consists of
862 estimates of contribution of each source to the total mass. The CMB model has several assumptions
863 including (Watson et al., 2002):

- 864 • chemical species do not react with one another and the species composition does not change
865 between the source(s) and the receptor(s)
- 866 • number of species are more than the number of sources
- 867 • all of the potential sources are included the model
- 868 • source profiles are not collinear
- 869 • random and non-correlated uncertainties

870

871 It is important to note that while the multivariate models do not require any information on source
872 emission composition, the CMB model requires the species abundances (in the form of source
873 profiles) in each source as a model input. Source profiles are “the mass abundances, i.e. fraction of
874 total mass of chemical species in source emissions, and such profiles are generally representative of
875 source categories rather than individual emitters” (Watson et al., 2002). There are several ways of
876 preparing source profiles for road vehicles including tests with chassis dynamometers,
877 measurements in tunnels/ highways and near roadside measurements (Phuleria et al., 2007; Handler
878 et al., 2008; Yan et al., 2009) as discussed in section 6. Source profiles are used for quantification
879 of source contributions to particulate matter using the chemical mass balance (CMB) model as well
880 as to compare and validate results obtained from factor analysis models. The CMB model relies to a
881 large extent on the accuracy of the source profiles used as an input. The typical components of any

882 source profile are found to be more-or-less similar although the mass abundances vary depending
883 on location and emitter characteristics. Thus, different combinations of source profiles can provide
884 statistically valid yet completely different solutions and selection of a source profile comparable to
885 ambient data is one of the key prerequisites for a good CMB solution (Robinson et al., 2006).
886 Absence of locally-representative source profiles and inter-region variability can result in errors in
887 estimation of source contributions from the CMB model (Yan et al., 2009; Rutter et al., 2011).
888 Significant differences have also been reported between real-world emissions and laboratory-tested
889 emissions (Yan et al., 2009; Ancelet et al., 2011). Analysis of vehicular source profiles in Bangkok,
890 Thailand led to the conclusion that while source profiles for different vehicle categories vary
891 significantly, ranges of the light duty and heavy duty average source profiles overlap and can be
892 represented by a single composite profile (Oanh et al., 2010). Fraser et al. (2003) and Fujita et al.
893 (2007) used the CMB model to analyse the apportionment of PM mass between diesel and gasoline
894 vehicles. While high molecular weight (4-7 ring) PAHs (coronene, benzo(ghi)perylene), hopanes
895 and steranes were found to be influential in case of gasoline engine emissions, 2-4 ring methylated
896 PAHs and EC were reported to be key species in the case of diesel engine emissions. Hanedar et al.
897 (2008) used the CMB model to resolve the sources of PAHs in Istanbul (Turkey) and identified
898 vehicle emissions (gasoline and diesel), natural gas combustion, coal and wood burning as major
899 sources with vehicular emissions contributing more than 50% at both urban and rural location.
900 Perrone et al. (2012) used CMB analysis to resolve PM sources in Milan (Italy) and reported traffic
901 emissions to contribute between 17-24% to PM_{2.5} mass in an urban location compared to 7-9% at a
902 rural site and El Haddad et al. (2011) reported vehicular emissions to contribute 17% to the PM
903 mass in Marseille (France). In Detroit (USA), traffic emissions and road dust emissions were
904 reported to contribute 24-36% and 3-5% respectively of mass using CMB analysis of PM_{2.5} data.

905
906 In recent years, there have been several studies focusing on characterization of trace metals and/or
907 organic marker compounds in PM contributed by road traffic emissions and preparation of mixed

908 traffic source profiles (El Haddad et al., 2009; Fabretti et al., 2009; Yan et al., 2009; Ancelet et al.,
909 2011). El Haddad et al. (2009) prepared a source profile for the vehicle fleet in France using tunnel
910 measurements and nearly 70 organic markers, whereas Fabretti et al. (2009) used trace element
911 marker data from tunnel-based measurements to derive chemical fingerprints for different traffic-
912 related sources including combustion, re-suspended dust and abrasion-related particles. Tunnel
913 studies were also used in USA for preparation of source profiles for gasoline and diesel engines
914 using inorganic and organic markers (Phuleria et al., 2006; Landis et al., 2007). Yin et al. (2010)
915 used organic molecular source tracers to provide separate quantification of gasoline, diesel and
916 smoker vehicle emissions at two UK sites. Recently, an attempt has also been made to characterize
917 the contribution of lubricating oil and fuel to the PM emissions (Sonntag et al., 2012).

918 919 **7. CONCLUSIONS**

920 Road traffic frequently makes a very significant contribution (5-80% depending on site and location)
921 to airborne concentrations of PM, whether expressed in terms of particle mass or number
922 concentration (Thorpe and Harrison, 2008; Belis et al., 2013). Particles arise not only from the
923 engine exhaust but also from abrasion of tyres, road surface and brake components as well as from
924 resuspension from the road surface. The sizes of particles emitted from these sources differ
925 considerably and size association may be a useful guide to particle source, as is chemical
926 composition. Fleet mix, type of road, traffic characteristic and fuel type are the factors that impact
927 the emission type and the relative contributions from exhaust and non-exhaust emissions. The
928 determination of particle mass and number increments due to traffic emissions can be based on a
929 number of methods including measurements in well controlled environments such as a road tunnel,
930 or twin site studies such as a highway and urban background (allowing for subtraction of the
931 background contribution), as well as using vehicle-specific tracers and the more sophisticated
932 receptor modelling methods based on multi-variate statistics or Chemical Mass Balance which have
933 been used in many studies to estimate the contributions to particulate matter concentrations, not

934 only from road vehicles but also from other sources. However, significant gaps still exist in our
935 understanding of traffic emissions, particularly:

- 936 1. Characterization and quantification of different non-exhaust sources.
- 937 2. Impacts of non-exhaust emissions upon human health: While there are many studies reporting
938 links between exposure to air pollution and adverse health impacts, detailed information on
939 the components that contribute to toxicity is missing (Kelly and Fussell, 2012).
- 940 3. Aerosol chemistry in high-traffic environments, and the evolution of particles emitted from
941 vehicles

942 Further, most of the reported analyses have been carried out in the USA or Europe, and there is a
943 lack of reliable information on traffic emissions in areas with high population density in Africa,
944 Asia and South America. The field offers much scope for future research , including the
945 development of enhanced methods for quantification of non-exhaust contributions to airborne
946 concentrations.

947

948 **REFERENCES**

949

950 Aatmeeyata, K. D. S. and Sharma, M. (2009). Traffic generated non-exhaust particulate emissions
951 from concrete pavement: A mass and particle size study for two-wheelers and small cars.
952 *Atmospheric Environment*, 43(35), 5691-5697.

953

954 Aatmeeyata, K. D. S., and Sharma, M. (2010). Polycyclic aromatic hydrocarbons, elemental and
955 organic carbon emissions from tire-wear. *Science of the Total Environment*, 408(20), 4563-4568.

956

957 Abu-Allaban, M., Coulomb, W., Gertler, A.W., Gillies, J., Pierson, W.R., Rogers, C.F., Sagebiel,
958 J.C. and Tarnay, L. (2002). Exhaust particle size distribution measurements at the Tuscarora
959 Mountain Tunnel. *Aerosol Science and Technology*, 36(6), 771-789.

960

961 Abu-Allaban, M., Gillies, J.A., Gertler, A.W., Clayton, R. and Proffitt, D. (2003). Tailpipe,
962 resuspended road dust, and brake wear emission factors from on-road vehicles. *Atmospheric
963 Environment*, 37, 5283-5293.

964

965 Adachi, K., and Tainosho, Y. (2004). Characterization of heavy metal particles embedded in tire
966 dust. *Environment International*, 30(8), 1009-1017.

967

968 Aldabe, J., Elustondo, D., Santamaría, C., Lasheras, E., Pandolfi, M., Alastuey, A. and Querol, X.
969 (2011). Chemical characterisation and source apportionment of PM_{2.5} and PM₁₀ at rural, urban
970 and traffic sites in Navarra (North of Spain). *Atmospheric Research*, 102(1-2), 191-205.

971

972 Allen, A. G., Nemitz, E., Shi, J. P., Harrison, R. M., and Greenwood, J. C. (2001). Size distributions
973 of trace metals in atmospheric aerosols in the United Kingdom. *Atmospheric Environment*, 35(27),
974 4581-4591.

975

976 Allen, J. O., Alexandrova, O. and Kaloush, K.E. (2006). Tire wear emissions for asphalt rubber and
977 portland cement concrete pavement surfaces: Arizona State University. Report submitted to Arizona
978 Department of Transportation (Contract KR-04-0720-TRN).

979

980 Almeida, S. M., Pio, C. A., Freitas, M. C., Reis, M. A., and Trancoso, M. A. (2006). Source
981 apportionment of atmospheric urban aerosol based on weekdays/weekend variability: evaluation of
982 road re-suspended dust contribution. *Atmospheric Environment*, 40(11), 2058-2067.

983

984 Amato, F., Pandolfi, M., Viana, M., Querol, X., Alastuey, A., and Moreno, T. (2009). Spatial and
985 chemical patterns of PM₁₀ in road dust deposited in urban environment. *Atmospheric Environment*,
986 43(9), 1650-1659.

987

988 Amato, F., Pandolfi, M., Moreno, T., Furger, M., Pey, J., Alastuey, A., Bukowiecki, N., Prevot,
989 A.S.H., Baltensperger, U. and Querol, X. (2011a). Sources and variability of inhalable road dust
990 particles in three European cities. *Atmospheric Environment*, 45(37), 6777-6787.

991

992 Amato, F., Viana, M., Richard, A., Furger, M., Prévôt, A. S. H., Nava, S., Lucarelli, F., Bukowiecki,
993 N., Alastuey, A., Reche, C., Moreno, T., Pandolfi, M., Pey, J. and Querol, X. (2011b). Size and
994 time-resolved roadside enrichment of atmospheric particulate pollutants. *Atmospheric Chemistry
995 and Physics*, 11(6), 2917-2931.

996

997 Amato, F., Karanasiou, A., Moreno, T., Alastuey, A., Orza, J.A.G., Lumbreras, J., Borge, R., Boldo,
998 E., Linares. C. and Querol, X. (2012). Emission factors from road dust resuspension in a
999 Mediterranean freeway. *Atmospheric Environment*, 61, 580-587.

1000 Ancelet, T., Davy, P. K., Trompetter, W. J., Markwitz, A., and Weatherburn, D. C. (2011).
1001 Carbonaceous aerosols in an urban tunnel. *Atmospheric Environment*, 45(26), 4463-4469.
1002
1003 Apeageyi, E., Bank, M. S. and Spengler, J. D. (2011). Distribution of heavy metals in road dust
1004 along an urban-rural gradient in Massachusetts. *Atmospheric Environment*, 45(13), 2310-2323.
1005
1006 Ban-Weiss, G.A., Lunden, M.M, Kirchstetter, T.W. and Harley, R.A. (2010). Size-resolved particle
1007 number and volume emission factors for gasoline and diesel motor vehicles. *Aerosol Science*, 41, 5-
1008 12.
1009
1010 Barmpadimos, I., Nufer, M., Oderbolz, D. C., Keller, J., Aksoyoglu, S., Hueglin, C., Baltensperger,
1011 U. and Prevot, A.S.H. (2011). The weekly cycle of ambient concentrations and traffic emissions of
1012 coarse (PM₁₀–PM_{2.5}) atmospheric particles. *Atmospheric Environment*, 45(27), 4580-4590.
1013
1014 Beddows, D. C. S. and Harrison, R. M. (2008). Comparison of average particle number emission
1015 factors for heavy and light duty vehicles derived from rolling chassis dynamometer and field studies.
1016 *Atmospheric Environment*, 42, 7954-7966.
1017
1018 Beddows, D. C. S., Dall'Osto, M. and Harrison, R. M. (2009). Cluster analysis of rural, urban and
1019 curbside atmospheric particle size data. *Environmental Science and Technology*, 43, 4694-4700.
1020
1021 Belis, C.A., Karagulian, F., Larsen, B.R. and Hopke, P.K. (2013). Critical review and meta-analysis
1022 of ambient particulate matter source apportionment using receptor models in Europe. *Atmospheric*
1023 *Environment*, 69, 94-108.
1024
1025 Berger, J. and Denby, B. (2011). A generalized model for traffic induced road dust emissions.
1026 Model description and evaluation. *Atmospheric Environment*, 45, 3692-3703.
1027
1028 Bhaskar, V.S. and Sharma, M. (2008). Assessment of fugitive road dust emissions in Kanpur, India.
1029 *Transportation Research Part D*, 13 , 400-403.
1030
1031 Birmili, W., Allen, A. G., Bary, F., and Harrison, R. M. (2006). Trace Metal Concentrations and
1032 Water Solubility in Size-Fractionated Atmospheric Particles and Influence of Road Traffic.
1033 *Environmental Science and Technology*, 40(4), 1144-1153.
1034
1035 Blau, P.J . and Meyer, H. M. III (2003). Characteristics of wear particles produced during friction
1036 tests of conventional and unconventional disc brake materials. *Wear*, 255, 1261-1269.
1037
1038 Boonyatumanond, R., Murakami, M., Wattayakorn, G. Togo, A. and Takeda, H. (2007). Sources of
1039 polycyclic aromatic hydrocarbons (PAHs) in street dust in a tropical Asian mega-city, Bangkok,
1040 Thailand. *Science of the Total Environment*, 384, 420-432.
1041
1042 Boulter, P. G., Thorpe, A. J., Harrison, R. M. and Allen, A. G. (2005). Road vehicle non-exhaust
1043 particulate matter: Final report on emission modelling. Published Project Report PPR110. Available
1044 at http://uk-air.defra.gov.uk/reports/cat15/0706061624_Report2_Emission_modelling.PDF.
1045
1046 Brook, J. R., Graham, L., Charland, J. P., Cheng, Y., Fan, X., Lu, G., Li, S.M., Lillyman, C.,
1047 MacDonald, P., Caravaggio, G. and MacPhee, J.A. (2007). Investigation of the motor vehicle
1048 exhaust contribution to primary fine particle organic carbon in urban air. *Atmospheric Environment*,
1049 41(1), 119-135.
1050

- 1051 Buckeridge, D.L., Glazier, R., Harvey, B.J., Escobar, M., Amrhein, C. and Frank, J. (2002). Effect
1052 of motor vehicle emissions on respiratory health in an urban area. *Environmental Health*
1053 *Perspectives*, 110(3), 293-300.
1054
- 1055 Bukowiecki, N., Lienemann, P., Hill, M., Furger, M., Richard, A., Amato, F., Prevot, A.S.H.,
1056 Baltensperger, U., Buchmann, B. and Gehrig, R. (2010). PM10 emission factors for non-exhaust
1057 particles generated by road traffic in an urban street canyon and along a freeway in Switzerland.
1058 *Atmospheric Environment*, 44(19), 2330-2340.
1059
- 1060 Bullock, K.R., Duvall, R.M., Norris, G.A., McDow, S.R. and Hays, M.D. (2008). Evaluation of the
1061 CMB and PMF models using organic molecular markers in fine particulate matter collected during
1062 the Pittsburgh Air Quality Study. *Atmospheric Environmet*, 42, 6897-6904.
1063
- 1064 Caceres, D., Adonis, M., Gil, L., and Miguel, A. H. (1998). Exhaust PAH in Chilean vehicles:
1065 Levels, source signatures and receptor modeling. *Journal of Aerosol Science*, 29, Supplement 2(0),
1066 S741-S742.
1067
- 1068 Camatani, M., Crosta, G. F., Dolukhanyan, T., Sung, C., Giuliani, G., Corbetta, G. M., Cencetti, S.
1069 and Regazzoni, C. (2001). Microcharacterization and identification of tire debris in heterogenous
1070 laboratory and environmental specimens. *Materials Characterization*, 46, 271-283.
1071
- 1072 Canagaratna, M. R., Onasch, T. B., Wood, E. C., Herndon, S. C., Jayne, J. T., Cross, E. S., Miake-
1073 Lye, R. C., Kolb, C. E. and Worsnop, D. R. (2010). Evolution of vehicle exhaust particles in the
1074 Atmosphere. *Journal of Air and Waste Management Association*, 60, 1192-1203.
1075
- 1076 Canepari, S., Perrino, C., Olivieri, F., and Astolfi, M. L. (2008). Characterisation of the traffic
1077 sources of PM through size-segregated sampling, sequential leaching and ICP analysis.
1078 *Atmospheric Environment*, 42(35), 8161-8175.
1079
- 1080 Cass, G. R. (1998). Organic molecular tracers for particulate air pollution sources. *Trends in*
1081 *Analytical Chemistry*, 17(6), 356-366.
1082
- 1083 Chan, D. and Stachowiak, G. W. (2004). Review of automotive brake friction materials.
1084 *Proceedings of the Institution of Mechanical Engineers, Part D: Journal of Automobile Engineering*,
1085 218 (9), 953-966.
1086
- 1087 Chan, T. W. and Mozurkewich, M. (2007). Application of absolute principal component analysis to
1088 size distribution data: identification of particle origins. *Atmospheric Chemistry and Physics*, 7, 887-
1089 897.
1090
- 1091 Chang, T. Y., Modzelewski, S. W., Norbeck, J. M., and Pierson, W. R. (1981). Tunnel air quality
1092 and vehicle emissions. *Atmospheric Environment*, 15(6), 1011-1016.
1093
- 1094 Charron, A. and Harrison, R. M. (2003). Primary particle formation from vehicle emissions during
1095 exhaust dilution in the roadside atmosphere. *Atmospheric Environment*, 37(29), 4109-4119.
1096
- 1097 Charron, A. and Harrison, R. M. (2009). *Atmospheric Nanoparticles. Environmental and Human*
1098 *Health Impacts of Nanotechnology*, John Wiley and Sons, Ltd, 63-209.
1099
- 1100 Chellam, S., Kulkarni, P. and Fraser, M. P. (2005). Emissions of organic compounds and trace
1101 metals in fine particulate matter from motor vehicles: a tunnel study in Houston, Texas. *Journal of*
1102 *Air and Waste Management Association*, 55, 60-72.

1103 Chen, J., Zheng, H., Wang, W., Liu, H., Lu, L., Bao, L. and Ren, L. (2006). Resuspension method
1104 for road surface dust collection and aerodynamic size distribution characterization. *China*
1105 *Particuology*, 4(6), 300-303.
1106

1107 Chen, F., Hu, W. and Zhong, Q. (2013). Emissions of particle-phase polycyclic aromatic
1108 hydrocarbons (PAHs) in the Fu Gui-Shan Tunnel of Nanjing, China. *Atmospheric Research*, 124,
1109 53-60.
1110

1111 Cheng, Y., Lee, S. C., Ho, K. F., Chow, J. C., Watson, J. G., Louie, P. K. K., Cao, J. J. and Hai, X.
1112 (2010). Chemically-specified on-road PM_{2.5} motor vehicle emission factors in Hong Kong.
1113 *Science of the Total Environment*, 408(7), 1621-1627.
1114

1115 Cheng, Y., Zou, S. C., Lee, S. C., Chow, J. C., Ho, K. F., Watson, J. G., Han, Y. M., Zhang, R. J.,
1116 Zhang, F., Yau, P.S., Huang, Y., Bai, Y. and Wu, W. J. (2011). Characteristics and source
1117 apportionment of PM₁ emissions at a roadside station. *Journal of Hazardous Materials*, 195, 82-91.
1118

1119 Chiang, H.-L. and Huang, Y.-S. (2009). Particulate matter emissions from on-road vehicles in a
1120 freeway tunnel study. *Atmospheric Environment*, 43, 4014-4022.
1121

1122 Chiang, H.-L., Lai, Y.-M. and Chang, S.-Y. (2012). Pollutant constituents of exhaust emitted from
1123 light-duty diesel vehicles. *Atmospheric Environment*, 47, 399-406.
1124

1125 Chirico, R., Prevot A. S. H., DeCarlo, P. F., Heringa, M. F., Richter, R., Weingartner, E. and
1126 Baltensperger, U. (2011). Aerosol and trace vehicle emission factors measured in a tunnel using an
1127 Aerosol Mass Spectrometer and other on-line instrumentation. *Atmospheric Environment*, 45,
1128 2182-2192.
1129

1130 Choi, J.-K., Heo, J.-B., Ban, S.-J., Yi, S.-M. and Zoh, K.-D. (2012). Chemical characteristics of
1131 PM_{2.5} aerosol in Incheon, Korea. *Atmospheric Environment*, 46, 583-592.
1132

1133 Councill, T. B., Duckenfield, K. U., Landa, E. R. and Callender, E. (2004). Tire wear particles as a
1134 source of Zn to the environment. *Environmental Science and Technology*, 38(15), 4206-4214.
1135

1136 Crawford, J., Chambers, S., Cohen, D. D., Dyer, L., Wang, T. and Zahorowski, W. (2007).
1137 Receptor modelling using Positive Matrix Factorisation, back trajectories and Radon-222.
1138 *Atmospheric Environment*, 41(32), 6823-6837.
1139

1140 Cyrus, J., Pitz, M., Heinrich, J., Wichmann, H. E. and Peters, A. (2008). Spatial and temporal
1141 variation of particle number concentration in Augsburg, Germany. *Science of the Total*
1142 *Environment*, 401(1-3), 168-175.
1143

1144 Dahl, A., Gharibi, A., Swietlicki, E., Gudmundsson, A., Bohgard, M., Ljungman, A., Blomqvist, G.
1145 and Gustafsson, M. (2006). Traffic-generated emissions of ultrafine particles from pavement-tire
1146 interface. *Atmospheric Environment*, 40(7), 1314-1323.
1147

1148 Dall'Osto, M., Thorpe, A., Beddows, D. C. S., Harrison, R. M., Barlow, J. F., Dunbar, T., Williams,
1149 P. I. and Coe, H. (2011). Remarkable dynamics of nanoparticles in the urban atmosphere.
1150 *Atmospheric Physics and Chemistry*, 11, 6623-6637.
1151
1152
1153

1154 Denier van der Gon, H., Gerlofs-Nijland, M.E., Gehrig, R., Gustafsson, M., Janssen, N., Harrison,
1155 R.M., Hulskotte, J., Johansson, C., Jozwicka, M., Keuken, M., Krijgheld, K., Ntziachristos, L.,
1156 Riediker, M. and Cassee, F.R. (2013). The Policy Relevance of Wear Emissions from Road
1157 Transport, Now and in the Future- An International Workshop Report and Consensus Statement.
1158 Journal of the Air and Waste Management Association, 63(2), 136-149.
1159

1160 Dongarra, G., Manno, E. and Varrica, D. (2009). Possible markers of traffic-related emissions.
1161 Environmental Monitoring and Assessment, 154, 117-125.
1162

1163 Dreyfus, M. A., Adou, K., Zucker, S. M. and Johnston, M. V. (2009). Organic aerosol source
1164 apportionment from highly time-resolved molecular composition measurements. Atmospheric
1165 Environment, 43(18), 2901-2910.
1166

1167 Duong, T. T. and Lee, B. K. (2011). Determining contamination level of heavy metals in road dust
1168 from busy traffic areas with different characteristics. Journal of Environmental Management, 92(3),
1169 554-562.
1170

1171 El Haddad, I., Marchand, N., Dron, J., Temime-Roussel, B., Quivet, E., Wortham, H., Jaffrezo, J.L.,
1172 Baduel, C., Voisin, D., Besombes, J.L. and Gille, G. (2009). Comprehensive primary particulate
1173 organic characterization of vehicular exhaust emissions in France. Atmospheric Environment,
1174 43(39), 6190-6198.
1175

1176 El Haddad, I., Marchand, N., Wortham, H., Piot, C., Besombes, J.-L., Cozic, J., Chauvel, C.,
1177 Armengaud, A., Robin, D. and Jaffrezo, J.-L. (2011). Primary sources of PM_{2.5} organic aerosol in
1178 an industrial Mediterranean city, Marseille. Atmospheric Chemistry and Physics, 11, 2039- 2058.
1179

1180 Etyemezian, V., Kuhns, H., Gillies, J., Chow, J., Hendrickson, K., McGown, M. and Pitchford, M.
1181 (2003). Vehicle-based road dust emissions measurement (III): effect of speed, traffic volume,
1182 location, and season on PM₁₀ road dust emissions in the Treasure Valley, ID. Atmospheric
1183 Environment, 37, 4583-4593.
1184

1185 Fabretti, J.-F., Sauret, N., Gal, J.-F., Maria, P.-C. and Schärer, U. (2009). Elemental characterization
1186 and source identification of PM_{2.5} using Positive Matrix Factorization: The Malraux road tunnel,
1187 Nice, France. Atmospheric Research, 94(2), 320-329.
1188

1189 Fan, Y., Meng, Q., Weisel, C., Shalat, S., Laumbach, R., Ohman-Strickland, P., Black, K.,
1190 Rodriguez, M. and Bonanno, L. (2006). Acute Short-Term Exposures to PM_{2.5} Generated by
1191 Vehicular Emissions and Cardiopulmonary Effects in Older Adults. Epidemiology, 17(6), S213-
1192 S214.
1193

1194 Fang, G.-C., Chang, C.-N., Wu, Y.-S., Fu, P. P.-C., Yan, I.-L. and Chen, M.-H. (2004).
1195 Characterization, identification of ambient air and road dust polycyclic aromatic hydrocarbons in
1196 central Taiwan, Taichung. Science of the Total Environment, 327, 135-146.
1197

1198 Franco, V., Kousoulidou, M., Muntean, M., Ntziachristos, L., Hausberger, S. and Dilara, P. (2013).
1199 Road vehicle emission factors development: a review. Atmospheric Environment, 70, 84-97.
1200

1201 Fraser, M.P., Cass, G.R. and Simoneit, B.R.T. (1998). Gas-Phase and Particle-Phase Organic
1202 Compounds Emitted From Motor Vehicle Traffic in a Los Angeles Roadway Tunnel.
1203 Environmental Science and Technology, 32, 2051-2060.
1204

1205 Fraser, M.P., Buzcu, B., Yue, Z.W., Mcgaughey, G.R., Desai, N.R., Allen, D.T., Seila, R.L.,
1206 Lonneman, W.A. and Harley, R.A.(2003). Separation of Fine Particulate Matter Emitted from
1207 Gasoline and Diesel Vehicles Using Chemical Mass Balancing Techniques. *Environmental Science*
1208 *and Technology*, 37, 3904-3909.

1209

1210 Fujita, E. M., Campbell, D. E. and Zielinska, B. (2006). Chemical analysis of lubrication oil
1211 samples from a study to characterize exhaust emissions from light-duty gasoline vehicles in the
1212 Kansas City metropolitan area (Final Report).

1213

1214 Fujita, E. M., Campbell, D. E., Arnott, W. P., Chow, J. C. and Zielinska, B. (2007). Evaluations of
1215 the Chemical Mass Balance Method for Determining Contributions of Gasoline and Diesel Exhaust
1216 to Ambient Carbonaceous Aerosols. *Journal of Air and Waste Management Association*, 57, 721-
1217 740.

1218

1219 Fuller, G. W., Carslaw, D. C. and Lodge, , H. W. (2002). An empirical approach for the prediction
1220 of daily mean PM10 concentrations. *Atmospheric Environment*, 36, 1431-1441.

1221

1222 Fulper, C.R., Kishan, S., Baldauf, R.W., Sabisch, R., Warila, J., Fujita, E.M., Scarbro, C., Crews,
1223 W.S., Snow, R., Gabele, P., Sanots, R., Tierney, E. and Cantrell, B. (2010). Methods of
1224 Characterizing the Distribution of Exhaust Emissions from Light-Duty, Gasoline-Powered Motor
1225 Vehicles in the U.S. Fleet. *Journal of Air and Waste Management Association*, 60, 1376-1387.

1226

1227 Funasaka, K., Miyazaki, T., Kawaraya, T., Tsuruho, K. and Mizuno, T. (1998). Characteristics of
1228 particulates and gaseous pollutants in a highway tunnel. *Environmental Pollution*, 102(2-3), 171-
1229 176.

1230

1231 Furujo, E., Sternbeck, J. and Cousins, A.P. (2007). PM10 source characterization at urban and
1232 highway roadside locations. *Science of the Total Environment*, 387, 206-219.

1233

1234 Fushimi, A., Hasegawa, S., Takahashi, K., Fujitani, Y., Tanabe, K. and Kobayashi, S. (2008).
1235 Atmospheric fate of nuclei-mode particles estimated from the number concentrations and chemical
1236 composition of particles measured at roadside and background sites. *Atmospheric Environment*,
1237 42(5), 949-959.

1238

1239 Gao, B., Yu, J.-Z., Li, S.-X., Ding, X., He, Q.-F. and Wang, X.-M. (2011). Roadside and rooftop
1240 measurements of polycyclic aromatic hydrocarbons in PM2.5 in urban Guangzhou: Evaluation of
1241 vehicular and regional combustion source contributions. *Atmospheric Environment*, 45(39), 7184-
1242 7191.

1243

1244 Garg, B. D., Cadle, S. H., Mulawa, P. A., Groblicki, P. J., Laroo, C. and Parr, G. A. (2000). Brake
1245 Wear Particulate Matter Emissions. *Environmental Science and Technology*, 34(21), 4463-4469.

1246

1247 Gehrig, R., Hill, M., Buchmann, B., Imhof, D., Weingartner, E. and Baltensperger, U. (2004).
1248 Separate determination of PM10 emission factors of road traffic for tailpipe emissions and
1249 emissions from abrasion and resuspension processes. *International Journal of Environment and*
1250 *Pollution*, 22(3): 312-325.

1251

1252 Gehrig, M., Zeyer, K., Bukowiecki, N., Lienemann, P., Poulidakos, L.D., Furger, M. and
1253 Buchmann, B. (2010). Mobile load simulators- A tool to distinguish between the emissions due to
1254 abrasion and resuspension of PM10 from road surfaces. *Atmospheric Environment*, 44, 4937-4943.

1255

1256 Gertler, A.W. and Pierson, W.R. (1996). Recent measurements of mobile source emission factors in
1257 North American tunnels. *The Science of the Total Environment*, 189/190, 107-113.
1258
1259 Gertler, A.W., Gillies, J.A., Pierson, W.R., Rogers, C.F., Sagebiel, J.C., Abu-Allaban, M.,
1260 Coulombe, W., Tarnay, L. and Cahill, T.A. (2002). Real-World Particulate Matter and Gaseous
1261 Emissions from Motor Vehicles in a Highway Tunnel. Research Report 107, Health Effects
1262 Institute.
1263
1264 Gertler, A., Kuhns, H., Abu-Allaban, M., Damm, C., Gilies, J., Etyemezian, V., Clayton, R. and
1265 Proffitt, D. (2006). A case study of the impact of winter road sand/salt and street sweeping on road
1266 dust re-entrainment. *Atmospheric Environment*, 40, 5976-5985.
1267
1268 Gietl, J. K., Lawrence, R., Thorpe, A. J. and Harrison, R. M. (2010). Identification of brake wear
1269 particles and derivation of a quantitative tracer for brake dust at a major road. *Atmospheric
1270 Environment*, 44(2), 141-146.
1271
1272 Gillies, J.A., Gertler, A.W., Sagebiel, J.C. and Dippel, W.A. (2001). On-Road Particulate Matter
1273 (PM_{2.5} and PM₁₀) Emissions in the Sepulveda Tunnel, Los Angeles, California. *Environmental
1274 Science and Technology*, 35, 1054-1063.
1275
1276 Grieshop, A. P., Lipsky, E. M., Pekney, N. J., Takahama, S. and Robinson, A. L. (2006). Fine
1277 particle emission factors from vehicles in a highway tunnel: Effects of fleet composition and season.
1278 *Atmospheric Environment*, 40, 287-298.
1279
1280 Gunawardana, C., Goonetilleke, A., Egodawatta, P., Dawes, L. and Kokot, S. (2011). Source
1281 characterisation of road dust based on chemical and mineralogical composition. *Chemosphere*,
1282 87(2), 163-170.
1283
1284 Gustafsson, M., Blomqvist, G., Gudmundsson, A., Dahl, A., Swietlicki, E., Bohgard, M., Lindbom,
1285 J. and Ljungman, A. (2008). Properties and toxicological effects of particles from the interaction
1286 between tyres, road pavement and winter traction material. *The Science of the Total Environment*,
1287 393(2-3), 226-240.
1288
1289 Gustafsson, M., Blomqvist, A., Gudmundsson, A., Dahl, A., Jonsson, P. and Swietlicki, E. (2009).
1290 Factors influencing PM₁₀ emissions from road pavement wear. *Atmospheric Environment*, 49,
1291 226-240.
1292
1293 Han, L., Zhuang, G., Cheng, S., Wang, Y. and Li, J. (2007). Characteristics of re-suspended road
1294 dust and its impact on the atmospheric environment in Beijing. *Atmospheric Environment*, 41(35),
1295 7485-7499.
1296
1297 Han, B., Bai, Z., Guo, G., Wang, F., Li, F., Liu, Q., Ji, Y., Li, X. and Hu, Y. (2009).
1298 Characterization of PM(10) fraction of road dust for polycyclic aromatic hydrocarbons (PAHs)
1299 from Anshan, China. *Journal of hazardous materials*, 170(2-3), 934-940.
1300
1301 Handler, M., Puls, C., Zbiral, J., Marr, I., Puxbaum, H. and Limbeck, A. (2008). Size and
1302 composition of particulate emissions from motor vehicles in the Kaisermühlen-Tunnel, Vienna.
1303 *Atmospheric Environment*, 42(9), 2173-2186.
1304
1305 Hanedar, A., Alp, K., Kaynak, B., Baek, J., Avsar, E. and Odman, M.T. (2008). Concentrations and
1306 sources of PAHs at three stations in Istanbul, Turkey. *Atmospheric Research*, 99, 391-399.
1307

1308 Harris, S. J. and Maricq, M. M. (2001). Signature size distributions for diesel and gasoline engine
1309 exhaust particulate matter. *Journal of Aerosol Science*, 32(6), 749-764.
1310
1311 Harrison, R. M., Deacon A. R. and Jones, M. R. (1997). Sources and processes affecting
1312 concentrations of PM10 and PM2.5 Particulate Matter in Birmingham (U.K.). *Atmospheric*
1313 *Environment*, 31, 4103-4117.
1314
1315 Harrison, R.M., Shi, J.P., Xi, S., Khan, A., Mark, D., Kinnersley, R. and Yin, J. (2000).
1316 Measurement of number, mass and size distribution of particles in the atmosphere, *Phil. Trans. R.*
1317 *Soc. Lond., A*, 358, 2567-2580.
1318
1319 Harrison, R. M., Tilling, R., Romero, M.S.C., Harrad, S. and Jarvis, K. (2003). A study of trace
1320 metals and polycyclic aromatic hydrocarbons in the roadside environment. *Atmospheric*
1321 *Environment*, 37, 2391-2402.
1322
1323 Harrison, R. M., Jones, A. M. and Barrowcliffe, R. (2004). Field study of the influence of
1324 meteorological factors and traffic volumes upon suspended particle mass at urban roadside sites of
1325 different geometry. *Atmospheric Environment*, 38, 6361–6369.
1326
1327 Harrison, R. M. (2009). Airborne Particulate Matter from Road Traffic: Current Status of
1328 Knowledge and Research Challenges Paper presented at the ETTA009. In 17th Transport and Air
1329 Pollution Symposium - 3rd Environment and Transport Symposium.
1330
1331 Harrison, R. M., Giorio, C., Beddows, D. C. and Dall'Osto, M. (2010). Size distribution of airborne
1332 particles controls outcomes of epidemiological studies. *Science of the Total Environment*, 409,
1333 289-293.
1334
1335 Harrison, R.M., Beddows, D.C.S. and Dall'Osto, M., (2011). PMF analysis of wide-range particle
1336 size spectra collected on a major highway. *Environmental Science and Technology*, 45, 5522-5528.
1337
1338 Harrison, R. M., Dall'Osto, M., Beddows, D. C. S., Thorpe, A. J., Bloss, W. J., Allan, J. D., Coe, H.,
1339 Dorsey, J. R., Gallagher, M., Martin, C., Whitehead, J., Williams, P. I., Jones, R. L., Langridge, J. M.,
1340 Benton, A. K., Ball, S. M., Langford, B., Hewitt, C. N., Davison, B., Martin, D., Petersson, K.,
1341 Henshaw, S. J., White, I. R., Shallcross, D. E., Barlow, J. F., Dunbar, T., Davies, F., Nemitz, E.,
1342 Phillips, G. J., Helfter, C., Di Marco, C. F. and Smith, S. (2012a). Atmospheric chemistry and
1343 physics in the atmosphere of a developed megacity (London): An overview of the REPARTEE
1344 experiment and its conclusions. *Atmospheric Chemistry and Physics*, 12, 3065-3114.
1345
1346 Harrison, R. M., Jones, A. M., Gietl, J., Yin, J. and Green, D. C. (2012b). Estimation of the
1347 contributions of brake dust, tire wear, and resuspension to nonexhaust traffic particles derived from
1348 atmospheric measurements. *Environmental Science and Technology*, 46, 6523-6529.
1349 Hassanien, M. A. and Abdel-Latif, N. M. (2008). Polycyclic aromatic hydrocarbons in road dust
1350 over Greater Cairo, Egypt. *Journal of Hazardous Materials*, 151, 247-254.
1351
1352 Hays, M. D., Cho, S.-H., Baldauf, R., Schauer, J. J. and Shafer, M. (2011). Particle size
1353 distributions of metal and non-metal elements in an urban near-highway environment. *Atmospheric*
1354 *Environment*, 45(4), 925-934.
1355
1356 He, L.-Y., Hu, M., Huang, X.-F., Zhang, Y.-H., Yu, B.-D. and Liu, D.-Q. (2006). Chemical
1357 characterization of fine particles from on-road vehicles in the Wutong tunnel in Shenzhen, China.
1358 *Chemosphere*, 62(12), 1565-1573.
1359

1360 He, L.-Y., Hu, M., Zhang, Y.-H., Huang, X.-F. and Yao, T.-T. (2008). Fine Particle Emissions from
1361 On-Road Vehicles in the Zhujiang Tunnel, China. *Environmental Science and Technology*, 42(12),
1362 4461-4466.

1363 HEI Panel on the Health Effects of Traffic-Related Air Pollution (2010). *Traffic-Related Air*
1364 *Pollution: A Critical Review of the Literature on Emissions, Exposure, and Health Effects*. HEI
1365 *Special Report*, 17, Health Effects Institute, Boston, MA.

1366

1367 Hien, T. T., Thanh, L. T., Kameda, T., Takenaka, N. and Bandow, H. (2007). Distribution
1368 characteristics of polycyclic aromatic hydrocarbons with particle size in urban aerosols at the
1369 roadside in Ho Chi Minh City, Vietnam. *Atmospheric Environment*, 41(8), 1575-1586.

1370

1371 Hildemann, L. M., Markowski, G. R., Jones, M. C. and Cass, G. R. (1991). Submicrometer aerosol
1372 mass distributions of emissions from boilers, fireplaces, automobiles, diesel trucks and meat
1373 cooking operatios. *Aerosol Science and Technology*, 14(1), 138-152.

1374

1375 Hjortenkrans, D. S. T., Bergbäck, B. G. and Häggerud, A. V. (2007). Metal Emissions from Brake
1376 Linings and Tires: Case Studies of Stockholm, Sweden 1995/1998 and 2005. *Environmental*
1377 *Science and Technology*, 41(15), 5224-5230.

1378

1379 Hopke P. K. (2003). Recent developments in receptor modelling. *Journal of Chemometrics*, 17,
1380 255-265.

1381

1382 Huang, X.-F., Chen, D.-L., Lan, Z.-J., Feng, N., He, L.-Y., Yu, G.-H. and Luan, S.-J. (2012).
1383 Characterization of organic aerosol in fine particles in a mega-city of South China: Molecular
1384 composition, seasonal variation, and size distribution. *Atmospheric Research* 114-115, 28-37.

1385

1386 Hussein, T., Johansson, C., Karlsson, H. and Hansson, H.-C. (2008). Factors affecting non-tailpipe
1387 aerosol particle emissions from paved roads: On-road measurements in Stockholm, Sweden.
1388 *Atmospheric Environment*, 42(4), 688-702.

1389

1390 Iijima, A., Sato, K., Yano, K., Tago, H., Kato, M., Kimura, H. and Furuta, N. (2007). Particle size
1391 and composition distribution analysis of automotive brake abrasion dusts for the evaluation of
1392 antimony sources of airborne particulate matter. *Atmospheric Environment*, 41(23), 4908-4919.

1393

1394 Imhof, D., Weingartner, E., Prevot, A.S.H., Ordonez, C., Kurtenbach, R., Weisen, P., Rodler, J.,
1395 Sturm, P., McCrae, I., Sjodin, A. and Baltensperger, U. (2005a). Aerosol and NO_x emissions
1396 factors and submicron particle size distributions in two road tunnels with different traffic regimes.
1397 *Atmospheric Chemistry and Physics Discussions*, 5, 5127-5166.

1398

1399 Imhof, D., Weingartner, E., Ordonez, C., Gehrig, R., Hill, M., Buchmann, B. and Baltensperger, U.
1400 (2005b). Real-world emission factors of fine and ultrafine aerosol particles for different traffic
1401 situations in Switzerland. *Environmental Science and Technology*, 39(21), 5127-5166.

1402

1403 Ingo, G. M., D'Uffizi, M., Falso, G., Bultrini, G. and Padeletti, G. (2004). Thermal and
1404 microchemical investigation of automotive brake pad wear residues. *Thermochimica Acta*, 418, 61-
1405 68.

1406

1407 Jamriska, M., Morawska, L., Thomas, S. and He, C. (2004). Diesel Bus Emissions Measured in A
1408 Tunnel Study. *Environmental Science and Technology*, 38, 6701-6709.

1409

1410

- 1411 Jayaratne, E. R., Wang, L., Heuff, D., Morawska, L. and Ferreira, L. (2009). Increase in particle
1412 number emissions from motor vehicles due to interruption of steady traffic flow. *Transportation*
1413 *Research Part D*, 14, 521-526.
1414
- 1415 Johansson, C., Norman, M. and Burman, L. (2009). Road traffic emission factors for heavy metals.
1416 *Atmospheric Environment*, 43(31), 4681-4688.
1417
- 1418 Jones, A. M. and Harrison, R. M. (2006). Estimation of emission factors of particle number and
1419 mass fractions from traffic at a site where mean vehicle speeds vary over short distances.
1420 *Atmospheric Environment*, 35, 7125-7137.
1421
- 1422 Jones, A. M., Yin, J. and Harrison, R. M. (2008). The weekday-weekend difference and the
1423 estimation of the non-vehicle contributions to the urban increment of airborne particulate matter.
1424 *Atmospheric Environment*, 42, 4467-4479.
1425
- 1426 Kam, W., Liacos, J. W., Schauer, J. J., Delfino, R. J. and Sioutas, C. (2012). Size-segregated
1427 composition of particulate matter (PM) in major roadways and surface streets. *Atmospheric*
1428 *Environment*, 55, 90-97.
1429
- 1430 Karanasiou, A., Moreno, T., Amato, F., Lumberras, J., Narros, A., Borge, R., Tobias, A., Boldo, E.,
1431 Linares, C., Pey, J., Reche, C., Alastuey, A. and Querol, X. (2011). Road dust contribution to PM
1432 levels – Evaluation of the effectiveness of street washing activities by means of Positive Matrix
1433 Factorization. *Atmospheric Environment*, 45(13), 2193-2201.
1434
- 1435 Karanasiou, A., Moreno, T., Amato, F., Tobias, A., Boldo, E., Linares, C., Lumberras, J., Borge, R.,
1436 Alastuey, A. and Querol, X. (2012). Variation of PM_{2.5} concentrations in relation to street washing
1437 activities. *Atmospheric Environment*, 54, 465- 469.
1438
- 1439 Kaunhaniemi, M., Kukkonen, J., Harkonen, J., Nikmo, J., Kangas, L., Omstedt, G., Ketzel, M.,
1440 Kousa, A., Haakana, M. and Karppinen, A. (2011). Evaluation of a road dust suspension model for
1441 predicting the concentrations of PM₁₀ in a street canyon. *Atmospheric Environment*, 45, 3646-3654.
1442
- 1443 Kelly, F.J. and Fussell, J.C. (2012). Size, source and chemical composition as determinants of
1444 toxicity attributable to ambient particulate matter. *Atmospheric Environment*, 60, 504- 526.
1445
- 1446 Kerminen, V.-M., Pakkanen, T. A., Mäkelä, T., Hillamo, R. E., Sillanpää, M., Rönkkö, T., Virtanen,
1447 A., Keskinen, J., Pirjola, L., Hussein, T. and Hameri, K. (2007). Development of particle number
1448 size distribution near a major road in Helsinki during an episodic inversion situation. *Atmospheric*
1449 *Environment*, 41(8), 1759-1767.
1450
- 1451 Ketzel, M.G., Omstedt, G., Johansson, C., Düring, I., Pohjola, M., Oettl, D., Gidhagen, L., Wahlin,
1452 P., Lohmeyer, A. and Berkowicz (2007). Estimation and validation of PM_{2.5}/PM₁₀ exhaust and
1453 non-exhaust emission factors for practical street pollution modelling. *Atmospheric Environment*, 41,
1454 9370-9385.
1455
- 1456 Keuken, M., Denier van der Gon, H. and van der Valk, K. (2010). Non-exhaust emissions of PM
1457 and the efficiency of emission reduction by road sweeping and washing in the Netherlands. *The*
1458 *Science of the Total Environment*, 408(20), 4591-4599.
1459
- 1460 Keuken, M., Henzing, B., Zandveld, P., van den Elshout, S. and Karl, M. (2012). Dispersion of
1461 particle numbers and elemental carbon from road traffic, a harbour and an airstrip in the
1462 Netherlands. *Atmospheric Environment*, 54, 320-327.

1463
1464 Kim, J. Y., Lee, J. Y., Kim, Y. P., Lee, S. B., Jin, H. C. and Bae, G. N. (2012). Seasonal
1465 characteristics of the gaseous and particulate PAHs at a roadside station in Seoul, Korea.
1466 Atmospheric Research, 116, 142-150.
1467 Kittleson, D. B., Watts, W. F. and Johnson, J.P. (2004). Nanoparticle emissions on Minnesota
1468 highways. Atmospheric Environment, 38, 9-19.
1469
1470 Klems, J. P., Pennington, M. R., Zordan, C. A. and Johnston, M. V. (2010). Ultrafine particles near
1471 a roadway intersection: Origin and apportionment of fast changes in concentration. Environmental
1472 Science and Technology, 44(20), 7903-7907.
1473
1474 Kuhlbusch, T.A.J., John, A.C. and Quass, U. (2009). Sources and source contributions to fine
1475 particles. Biomarkers, 14(1), 23- 28.
1476
1477 Kumar, P., Fennell, P., Langley, D. and Britter, R. (2008). Pseudo-simultaneous measurements for
1478 the vertical variation of coarse, fine and ultrafine particles in an urban street canyon. Atmospheric
1479 Environment, 42(18), 4304-4319.
1480
1481 Kumar, P., Pirjola, L., Ketzler, M. and Harrison, R.M. (2013). Nanoparticle emissions from 11 non-
1482 vehicle exhaust sources- A review. Atmospheric Environment, 67, 252-277.
1483
1484 Kumata, H., Yamada, J., Masuda, K., Takada, H., Sato, Y., Sakurai, T. and Fujiwara, K. (2002).
1485 Benzothiazolamines as tire-derived molecular markers: sorptive behaviour in street runoff and
1486 application to source apportioning. Environmental Science and Technology, 36(4), 702-708.
1487
1488 Kupiainen, K. J., Tervahattu, H., Räsänen, M., Mäkelä, T., Aurela, M. and Hillamo, R. (2005). Size
1489 and Composition of Airborne Particles from Pavement Wear, Tires, and Traction Sanding.
1490 Environmental Science and Technology, 39(3), 699-706.
1491
1492 Kwon, E.E. and Castaldi, M.J. (2012). Mechanistic understanding of polycyclic aromatic
1493 hydrocarbons (PAHs) from the thermal degradation of tires under various oxygen concentration
1494 atmospheres. Environmental Science and Technology, 46(23), 12921- 6.
1495
1496 Laidlaw, M. A. S., Zahran, S., Mielke, H. W., Taylor, M. P. and Filippelli, G. M. (2012). Re-
1497 suspension of lead contaminated urban soil as a dominant source of atmospheric lead in
1498 Birmingham, Chicago, Detroit and Pittsburgh, USA. Atmospheric Environment, 49, 302-310.
1499
1500 Landis, M. S., Lewis, C. W., Stevens, R. K., Keeler, G. J., Dvonch, J. T. and Tremblay, R. T.
1501 (2007). Ft. McHenry tunnel study: Source profiles and mercury emissions from diesel and gasoline
1502 powered vehicles. Atmospheric Environment, 41(38), 8711-8724.
1503
1504 Larsen, R.K. and Baker, J.E. (2003). Source apportionment of polycyclic aromatic hydrocarbons in
1505 the urban atmosphere: A comparison of three methods. Environmental Science and Technology,
1506 37(9), 1873-1881.
1507
1508 Lee, W.-J., Wang, Y.-F., Lin, T.-C., Chen, Y.-Y., Lin, W.-C., Ku, C.-C. and Cheng, J.-T. (1995).
1509 PAH characteristics in the ambient air of traffic-source. The Science of the Total Environment, 159,
1510 185-200.
1511
1512 Liacos, J. W., Kam, W., Delfino, R. J., Schauer, J. J. and Sioutas, C. (2012). Characterization of
1513 organic, metal and trace element PM_{2.5} species and derivation of freeway-based emission rates in
1514 Los Angeles, CA. Science of the total Environment, 435-436, 5283-5293.

1515
1516 Lim, L.H., Harrison, R.M. and Harrad, S. (1999). The contribution of traffic to atmospheric
1517 concentrations of polyaromatic hydrocarbons. *Environmental Science and Technology*, 33, 3538-
1518 3542.

1519 Lim, J., Yu, L.E., Kostetski, Yu. Yu., Lim, C., Ryu, J. and Kim, J. (2008). Effects of driving
1520 conditions on diesel exhaust particles. *Journal of Air and Waste Management Association*, 58,
1521 1077-1085.

1522
1523 Lin, C.-C., Chen, S.-J., Huang, K.-L., Hwang, W.-I., Chang-Chien, G.-P. and Lin, W.-Y. (2005).
1524 Characteristics of Metals in Nano/Ultrafine/Fine/Coarse Particles Collected Beside a Heavily
1525 Trafficked Road. *Environmental Science and Technology*, 39(21), 8113-8122.

1526
1527 Lin, H.-Y., Hung, C.-H., Yuan, C.-S., Chen, C.-J., Chen, W.-C., Chiang, S.-W. and Wang, C., J.-L.
1528 (2008). Characteristics and source identification of roadside suspended particles. *Transportation
1529 Research Part D: Transport and Environment*, 13(8), 531-538.

1530
1531 Lingard, J. J. N., Agus, E. L., Young, D. T., Andrews, G. E. and Tomlin, A. S. (2006). Observations
1532 of urban airborne particle number concentrations during rush-hour conditions: analysis of the
1533 number based size distributions and modal parameters. *Journal of Environmental Monitoring*, 8(12),
1534 1203-1218.

1535
1536 Liu, M., Cheng, S.B., Ou, D.N., Hou, L.J., Gao, L., Wang, L.L., Xie, Y.S., Yang, Y., Xu, S.Y.
1537 (2007). Characterization, identification of road dust PAHs in central Shanghai areas, China.
1538 *Atmospheric Environment*, 41, 8785-8795.

1539
1540 Lough, G. C., Schauer, J. J., Park, J.-S., Shafer, M. M., DeMinter, J. T. and Weinstein, J. P. (2005).
1541 Emissions of Metals Associated with Motor Vehicle Roadways. *Environmental Science and
1542 Technology*, 39(3), 826-836.

1543
1544 Lough, G. C., Christensen, C. G., Schauer, J. J., Tortorelli, J., Mani, E., Lawson, D. R., Clark, N. N.
1545 and Gabela, P. A. (2007). Development of molecular marker source profiles for emissions from on-
1546 road gasoline and diesel vehicle fleets. *Journal of the Air and Waste Management Association*,
1547 57(10), 1190-1199.

1548
1549 Luo, X. S., Yu, S., Zhu, Y. G. and Li, X. D. (2011). Trace metal contamination in urban soils of
1550 China. *The Science of the Total Environment*, 421-422, 17-30.

1551
1552 Majumdar, D., Rajaram, B., Meshram, S. and Chalapati Rao, C.V. (2012). PAHs in Road Dust:
1553 Ubiquity, Fate, and Summary of Available Data. *Critical Reviews in Environmental Science and
1554 Technology*, 42(12), 1191-1232.

1555
1556 Mancilla, Y. and Mendoza, A. (2012). A tunnel study to characterize PM_{2.5} emissions from
1557 Gasoline-powered vehicles in Monterrey, Mexico. *Atmospheric Environment*, 59, 449-460.

1558
1559 Manno, E., Varrica, D. and Dongarrà, G. (2006). Metal distribution in road dust samples collected
1560 in an urban area close to a petrochemical plant at Gela, Sicily. *Atmospheric Environment*, 40(30),
1561 5929-5941.

1562
1563 Mari, M., Harrison, R.M., Schuhmacher, M., Domingo, J.L. and Pongpiachan S. (2010). Inferences
1564 over the sources and processes affecting polycyclic aromatic hydrocarbons in the atmosphere derived
1565 from measured data. *Science of the Total Environment*, 408, 2387-2393.

1566

1567 Martuzevicius, D., Kliucininkas, L., Prasauskas, T., Krugly, E., Kauneliene, V. and Strandberg, B.
1568 (2011). Resuspension of particulate matter and PAHs from street dust. *Atmospheric Environment*,
1569 45(2), 310-317.
1570
1571 Mathissen, M., Scheer, V., Vogt, R. and Benter, T. (2011). Investigation on the potential generation
1572 of ultrafine particles from the tire–road interface. *Atmospheric Environment*, 45(34), 6172-6179.
1573
1574 Maricq, M. M. (2007). Chemical characterization of particulate emissions from diesel engines: A
1575 review. *Journal of Aerosol Science*, 38(11), 1079-1118.
1576
1577 Masiol, M., Hofer, A., Squizzato, S., Piazza, R., Rampazzo, G. and Pavoni, B. (2012). Carcinogenic
1578 and mutagenic risk associated to airborne particle-phase polycyclic aromatic hydrocarbons: A
1579 source apportionment. *Atmospheric Environment*, 60, 375- 382.
1580
1581 Mauderly, J.L. (1994). Toxicological and epidemiological evidence for health risks from inhaled
1582 engine emissions. *Environmental Health Perspectives*, 102, 165-171.
1583
1584 McLaren, R., Gertler, A.W., Wittorff, D.N., Belzer, W., Dann, T. and Singleton, D.L. (1996). Real-
1585 World Measurements of Exhaust and Evaporative Emissions in the Cassiar Tunnel Predicted by
1586 Chemical Mass Balance Modeling. *Environmental Science and Technology*, 30, 3001-3009.
1587
1588 Miguel, A. H., Kirchstetter, T. W., Harley, R. A. and Hering, S. V. (1998). On-Road Emissions of
1589 Particulate Polycyclic Aromatic Hydrocarbons and Black Carbon from Gasoline and Diesel
1590 Vehicles. *Environmental Science and Technology*, 32(4), 450-455.
1591
1592 Milani, M., Pucillo, F.P., Ballerini, M., Camatani, M., Gualtieri, M. and Martino, S. (2004). First
1593 evidence of tyre debris characterization at the nanoscale by focused ion beam. *Materials*
1594 *Characterization*, 52, 283-288.
1595
1596 Morawska, L., Jayaratne, E.R., Mengersen, K., Jamriska, M. and Thomas, S. (2002). Differences in
1597 airborne particle and gaseous concentrations in urban air between weekdays and weekends.
1598 *Atmospheric Environment*, 36, 4375-4383.
1599
1600 Nanzetta, K. M. and Holmen, B. A. (2004). Roadside Particle Number Distributions and
1601 Relationships between Number Concentrations, Meteorology, and Traffic along a Northern
1602 California Freeway. *Journal of Air and Waste Management Association*, 54, 540-554.
1603
1604 Narvaez, R. F., Hoepner, L., Chillrud, S. N., Yan, B., Garfinkel, R., Whyatt, R., Camann, D., Perera,
1605 F. P., Kinney, P. L. and Miller, R. L. (2008). Spatial and Temporal Trends of Polycyclic Aromatic
1606 Hydrocarbons and Other Traffic-Related Airborne Pollutants in New York City. *Environmental*
1607 *Science and Technology*, 42, 7330-7335.
1608
1609 Nicholson, K.W. (1988). A review of particle resuspension. *Atmospheric Environment* (1967),
1610 22(12), 2639-2651.
1611
1612 Nielsen, T. (1996). Traffic contribution of polycyclic aromatic hydrocarbons in the centre of a large
1613 city. *Atmospheric Environment*, 30(20), 3481-3490.
1614
1615 Ntziachristos, L., Ning, Z., Geller, M.D., Sheesley, R.J., Schauer, J.J. and Sioutas, C. (2007). Fine,
1616 ultrafine and nanoparticle trace element compositions near a major freeway with a high heavy-duty
1617 diesel fraction. *Atmospheric Environment*, 41, 5864-5696.
1618

1619 Oliveira, C., Pio, C., Caseiro, A., Santos, P., Nunes, T., Mao, H., Luahana, L. and Sokhi, R.S.
1620 (2010). Road traffic impact on urban atmospheric aerosol loading at Oporto, Portugal. *Atmospheric*
1621 *Environment*, 44(26), 3147-3158.
1622

1623 Oliveira, C., Martins, N., Tavares, J., Pio, C., Cerqueira, M., Matos, M., Silva, H., Oliviera, S. and
1624 Camoes, F. (2011). Size distribution of polycyclic aromatic hydrocarbons in a roadway tunnel in
1625 Lisbon, Portugal. *Chemosphere*, 83(11), 1588-1596.
1626

1627 Omar, N. Y. M. J., Abas, M. R. B., Rahman, N. A., Tahir, N. M., Rushdi, A. I. and Simoneit, B. R.
1628 T. (2007). Levels and distributions of organic source tracers in air and roadside dust particles of
1629 Kuala Lumpur, Malaysia. *Environmental Geology*, 52, 1485-1500.
1630

1631 Omstedt, G., Bringfelt, B., Johansson, C. (2005). A model for vehicle-induced non-tailpipe
1632 emissions of particles along Swedish roads. *Atmospheric Environment*, 39, 6088-6097.
1633

1634 Oanh, N. T. K., Thiansathit, W., Bond, T. C., Subramanian, R., Winijkul, E. and Paw-armart, I.
1635 (2010). Compositional characterization of PM_{2.5} emitted from in-use diesel vehicles. *Atmospheric*
1636 *Environment*, 44(1), 15-22.
1637

1638 Ondráček, J., Schwarz, J., Ždímal, V., Andělová, L., Vodička, P., Bízek, V., Tsai, C.-J., Chen, S.-C.
1639 and Smolik, J. (2011). Contribution of the road traffic to air pollution in the Prague city (busy
1640 speedway and suburban crossroads). *Atmospheric Environment*, 45(29), 5090-5100.
1641

1642 Padro-Martinez, L. T., Patton, A. P., Trull, J. B., Zamore, W., Brugge, D. and Durant, J. L. (2012).
1643 Mobile monitoring of particle number concentration and other traffic-related air pollutants in a
1644 near-highway neighbourhood over the course of a year. *Atmospheric Environment*, 61, 253-264.
1645

1646 Pant, P. and Harrison, R. M. (2012). Critical review of receptor modelling for particulate matter: A
1647 case study of India, *Atmospheric Environment*, 49, 1-12.
1648

1649 Peltier, R. E., Cromar, K. R., Ma, Y., Fan, Z.-H. and Lippmann, M. (2011). Spatial and seasonal
1650 distribution of aerosol chemical components in New York City: (2) Road dust and other tracers of
1651 traffic-generated air pollution. *Journal of Exposure Science and Environmental Epidemiology*,
1652 21(5), 484-494.
1653

1654 Perez, N., Pey, J., Cusack, M., Reche, C., Querol, X., Alastuey, A. and Viana, M. (2010).
1655 Variability of Particle Number, Black Carbon and PM₁₀, PM_{2.5} and PM₁ levels and speciation:
1656 Influence of road traffic emissions on urban air quality. *Aerosol Science and Technology*, 44(7),
1657 487-499.
1658

1659 Perrone, M. G., Larsen, B. R., Ferrero, L., Sangiorgi, G., De Gennaro, G., Udisti, R., Zangrano,
1660 R., Gambaro, A. and Bolzacchini, E. (2012). Sources of high PM_{2.5} concentrations in Milan,
1661 Northern Italy: Molecular marker data and CMB modelling. *Science of the Total Environment*, 414,
1662 343-355.
1663

1664 Pey, J., Querol, X., Alastuey, A., Rodriguez, R., Putuad, J.-P. and Van Dingenen, R. (2009).
1665 Source apportionment of urban fine and ultra-fine particle number concentration in a Western
1666 Mediterranean city. *Atmospheric Environment*, 43, 4407-4415.
1667

1668 Pey, J., Querol, X. and Alastuey, A. (2010). Discriminating the regional and urban contributions in
1669 the North-Western Mediterranean: PM levels and composition. *Atmospheric Environment*, 44(13),
1670 1587-1596.

1671 Phuleria, H. C., Geller, M. D., Fine, P. M. and Sioutas, C. (2006). Size-Resolved Emissions of
1672 Organic Tracers from Light- and Heavy-Duty Vehicles Measured in a California Roadway Tunnel.
1673 *Environmental Science and Technology*, 40(13), 4109-4118.

1674 Phuleria, H. C., Sheesley, R. J., Schauer, J. J., Fine, P. M. and Sioutas, C. (2007). Roadside
1675 measurements of size-segregated particulate organic compounds near gasoline and diesel-
1676 dominated freeways in Los Angeles, CA. *Atmospheric Environment*, 41(22), 4653-4671.

1677

1678 Prichard, H.M. and Fisher, P.C. (2012). Identification of Platinum and Palladium Particles Emitted
1679 from Vehicles and Dispersed into the Surface Environment. *Environmental Science and*
1680 *Technology*, 46(6), 3149-3154.

1681 Pulles, T., Denier van der Gon, H., Appelman, W. and Verheul, M. (2012). Emission factors for
1682 heavy metals from diesel and petrol used in European vehicles. *Atmospheric Environment*, 61, 641-
1683 651.

1684

1685 Puustinen, A., Hämeri, K., Pekkanen, J., Kulmala, M., de Hartog, J., Meliefste, K., ten Brink, H.,
1686 Kos, G., Katsouyanni, K., Karakatsani, A., Kotronarou, A., Kavouras, I., Meddings, C., Thomas, S.,
1687 Harrison, R., Ayres, J.G., van der Zee, S. and Hok, G. (2007). Spatial variation of particle number
1688 and mass over four European cities. *Atmospheric Environment*, 41(31), 6622-6636.

1689

1690 Reche, C., Querol, X., Alastuey, A., Viana, M., Pey, J., Moreno, T., Rodriguez, S., Gonzalez, Y.,
1691 Fernandez-Camacho, R., Sanchez de la Campa, A. M., de la Rosa, J., Dall'Osto, M., Prevot, A. S.
1692 H., Hueglin, C., Harrison, R.M. and Quincey, P. (2011). New considerations for PM, black carbon
1693 and particle number concentration for air quality monitoring across different European cities.
1694 *Atmospheric Chemistry and Physics*, 11, 6207-6227.

1695

1696 Reddy, C. M. and Quinn, J. G. (1997). Environmental Chemistry of Benzothiazoles Derived from
1697 Rubber. *Environmental Science and Technology*, 31, 2847-2853.

1698

1699 Rexeis, M. and Hausberger, S. (2009). Tend of vehicle emission levels until 2020 - Prognosis
1700 based on current vehicle measurements and future emission legislation. *Atmospheric Environment*,
1701 43, 4689-4698.

1702

1703 Riddle, S. G., Robert, M. A., Jakober, C. A., Hannigan, M. P. and Kleeman, M. J. (2007). Size
1704 Distribution of Trace Organic Species Emitted from Light-Duty Gasoline Vehicles. *Environmental*
1705 *Science and Technology*, 41(21), 7464-7471.

1706

1707 Rissler, J., Swietlicki, E., Bengtsson, A., Boman, C., Pagels, J., Sandstrom, T., Blomberg, A. and
1708 Londahl, J. (2012). Experimental determination of deposition of diesel exhaust particles in the
1709 human respiratory tract. *Journal of Aerosol Science*, 48, 18-33.

1710

1711 Robert, M.A., VanBergen, S., Kleeman, M.J. and Jakober, C.A. (2007a). Size and composition
1712 distributions of particulate matter emissions: Part 1- Light-duty diesel vehicles. *Journal of Air and*
1713 *Waste Management Association*, 57, 1414-1428.

1714

1715 Robert, M. A., Kleeman, M. J. and Jakober, C. A. (2007b). Size and composition distributions of
1716 particulate matter emissions: Part 2- Heavy-duty diesel vehicles. *Journal of Air and Waste*
1717 *Management Association*, 57, 1429-1438.

1718

1719 Robinson, A. L., Subramanian, R., Donahue, N. M. and Rogge, W. F. (2006). Source
1720 Apportionment of Molecular Markers and Organic Aerosol. Polycyclic Aromatic Hydrocarbons
1721 and Methodology for Data Visualization. *Environmental Science and Technology*, 40(24), 7803-
1722 7810.

- 1723 Rodriguez, S., Van Dingenen, R., Putuad, J.-P., Dell'Acqua, A., Pey, J., Querol, X., Alastuey, A.,
1724 Chenery, S., Ho, K.-F., Harrison, R., Tardivo, R., Scarnato, B. and Gemelli, V. (2007). A study on
1725 the relationship between mass concentrations, chemistry and number size distribution of urban fine
1726 aerosols in Milan, Barcelona and London. *Atmospheric Chemistry and Physics*, 35, 2217-2232.
1727
- 1728 Rogak, S. N., Green, S. I. and Pott, U. (1998). Use of Tracer Gas for Direct Calibration of
1729 Emission-Factor Measurements in a Traffic Tunnel. *Journal of the Air and Waste Management
1730 Association*, 48(6), 545-552.
1731
- 1732 Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R. and Simoneit, B. R. T. (1993).
1733 Sources of fine organic aerosol. 3. Road dust, tire debris, and organometallic brake lining dust:
1734 roads as sources and sinks. *Environmental Science and Technology*, 27(9), 1892-1904.
1735
- 1736 Rose, D., Wehner, B., Ketzel, M., Engler, C., Voigtlander, J., Tuch, T. and Wiedensohler, A. (2006).
1737 Atmospheric number size distributions of soot particles and estimation of emission factors.
1738 *Atmospheric Chemistry and Physics*, 6(4), 1021-1031.
1739
- 1740 Rosenbohm, E., Vogt, R., Scheer, V., Nielsen, O. J., Dreiseidler, A., Baumbach, G., Imhof, D.,
1741 Baltensperger, U., Fuchs, J. and Jaeschke, W. (2006). Particulate size distributions and mass
1742 measured at a motorway during the BAB II campaign. *Atmospheric Environment*, 37, 5283-5293.
1743
- 1744 Ruellan, S. and Cachier, H. (2001). Characterisation of fresh particulate vehicular exhausts near a
1745 Paris high flow road. *Atmospheric Environment*, 35(2), 453-468.
1746
- 1747 Rutter, A. P., Snyder, D. C., Schauer, J. J., Sheesley, R. J., Olson, M. R. and DeMinter, J. (2011).
1748 Contributions of resuspended soil and road dust to organic carbon in fine particulate matter in the
1749 Midwestern US. *Atmospheric Environment*, 45(2), 514-518.
1750
- 1751 Sadiktsis, I., Bergvall, C., Johansson, C. and Westerholm, R. (2012). Automobile Tires—A
1752 Potential Source of Highly Carcinogenic Dibenzopyrenes to the Environment. *Environmental
1753 Science and Technology*, 46(6), 3326-3334.
1754
- 1755 Sahu, M., Hu, S., Ryan, P. H., Le Masters, G., Grinshpun, S. A., Chow, J. C. and Biswas, P. (2011).
1756 Chemical compositions and source identification of PM_{2.5} aerosols for estimation of a diesel source
1757 surrogate. *Science of the Total Environment*, 409(13), 2642-2651.
1758
- 1759 Samara, C. and Voutsas, D. (2005). Size distribution of airborne particulate matter and associated
1760 heavy metals in the roadside environment. *Chemosphere*, 59, 1197-1206.
1761
- 1762 Sanchez-Ccoyllo, O. R., Ynoue, R. Y., Martins, L. D., Astolfo, R., Miranda, R. M., Freitas, E. D.,
1763 Borges, A.S., Fornaro, A., Freitas, H., Moreira, A. and Andrade, M. F. (2009). Vehicular particulate
1764 matter emissions in road tunnels in Sao Paulo, Brazil. *Environmental Monitoring and Assessment*,
1765 149(1-4), 241-249.
1766
- 1767 Sanders, P. G., Xu, N., Dalka, T. M. and Maricq, M. M. (2003). Airborne brake wear debirs: Size
1768 distributions, composition, and a comparison of dynamometer and vehicle tests. *Environmental
1769 Science and Technology*, 37(4), 4060-4069.
1770
- 1771 Schaap, M., Manders, A.M.M., Hendriks, E.C.J., Cnossen, J.M., Segers, A.J.S., Denier van der Gon,
1772 H.A.C., Jozwicka, M., Sauter, F., Velders, G., Matthijssen, J. and Builtjes, P.J.H. (2009). Regional
1773 modelling of PM₁₀ over the Netherlands. Technical Report 500099008. Available at
1774 <http://www.rivm.nl/bibliotheek/rapporten/500099008.pdf> [Accessed September 9, 2012].

- 1775 Schauer, J. J., Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R. and Simoneit, B. R. T.
1776 (1996). Source apportionment of airborne particulate matter using organic compounds as tracers.
1777 *Atmospheric Environment*, 30, 3837-3855.
1778
- 1779 Schauer, J. J., Lough, G. C., Shafer, M. M., Christensen, W. C., Arndt, M. F., DeMinter, J. T. and
1780 Park, J.-S. (2006). Characterization of Emissions of Metals Emitted from Motor Vehicles: Health
1781 Effects Institute.
1782
- 1783 Shen, Z., Han, Y., Cao, J., Tian, J., Zhu, C., Liu, S., Liu, P. and Wang, Y. (2010). Characteristics of
1784 traffic-related emissions: A case study in roadside ambient air over Xi'an, China. *Aerosol and Air*
1785 *Quality Research*, 10, 292-300.
1786
- 1787 Shi, J. P. and Harrison, R. M., (1999). Investigation of ultrafine particle formation during diesel
1788 exhaust dilution. *Environmental Science and Technology*, 33, 3730-3736.
1789
- 1790 Shi, J. P., Khan, A. A. and Harrison, R. M., (1999). Measurements of ultrafine particle
1791 concentration and size distribution in the urban atmosphere. *Science of the Total Environment*, 235,
1792 51-64.
1793
- 1794 Sjodin, A., Ferm, M., Bjork, A., Rahmberg, M., Gudmundsson, A., Swietlicki, E., Johansson, C.,
1795 Gustafsson, M. and Blomqvist, G. (2010). Wear particles from road traffic: a field, laboratory and
1796 modelling study. IVL Report B1830. Available at: [http://www.ivl.se/webdav/files/B-](http://www.ivl.se/webdav/files/B-rapporter/B1830.pdf)
1797 [rapporter/B1830.pdf](http://www.ivl.se/webdav/files/B-rapporter/B1830.pdf).
1798
- 1799 Smith, D. J. T. and Harrison, R. M. (1996). Concentrations, trends and vehicle source profile of
1800 polynuclear aromatic hydrocarbons in the U.K. atmosphere. *Atmospheric Environment*, 30(14),
1801 2513-2525.
1802
- 1803 Sodeman, D. A., Toner, S. M. and Prather, K. A. (2005). Determination of single particle mass
1804 spectral signatures from light-duty vehicle emissions. *Environmental Science and Technology*,
1805 39(12), 4569-4580.
1806
- 1807 Song, F. and Gao, Y. (2011). Size distributions of trace elements associated with ambient particular
1808 matter in the vicinity of a major highway in the New Jersey–New York metropolitan area.
1809 *Atmospheric Environment*, 45(37), 6714-6723.
1810
- 1811 Song, S., Wu, Y., Jiang, J., Yang, L., Cheng, Y. and Hao, J. (2012). Chemical characteristics of
1812 size-resolved PM(2.5) at a roadside environment in Beijing, China. *Environmental Pollution*, 161,
1813 215-221.
1814
- 1815 Sonntag, D. B., Bailey, C. R., Fulper, C. R. and Baldauf, R. W. (2012). Contribution of Lubricating
1816 Oil to Particulate Matter Emissions from Light-Duty Gasoline Vehicles in Kansas City.
1817 *Environmental Science and Technology*, 46(7), 4191–4199.
- 1818 Stechmann, H. and Dannecker, W. (1990). Characterization and source analysis of vehicle-
1819 generated aerosols. *Journal of Aerosol Science*, 21, Supplement 1, S287-S290.
1820
- 1821 Staehelin, J., Schlapfer, K., Burgin, T., Steinemann, U., Schneider, S., Brunner, D., Baumle, M.,
1822 Meier, M., Zahner, C., Keiser, S., Stahel, W. and Keller, C. (1995). Emission factors from road
1823 traffic from a tunnel study (Gubrist tunnel, Switzerland). Part I: concept and first results. *Science of*
1824 *the Total Environment*, 169, 141-147.
1825

1826 Sternbeck, J., Sjödin, Å. and Andréasson, K. (2002). Metal emissions from road traffic and the
1827 influence of resuspension—results from two tunnel studies. *Atmospheric Environment*, 36(30),
1828 4735-4744.
1829
1830 Sun, Y.L., Zhang, Q., Schwab, J.J., Chen, W.-N., Bae, M.-S., Hung, H.-M., Lin, Y.-C., Ng, N.L.,
1831 Jayne, J., Massoli, P., Williams, L.R. and Demerjian, K.L. (2012). Characterization of near-
1832 highway submicron aerosols in New York City with a high-resolution aerosol mass spectrometer.
1833 *Atmospheric Chemistry and Physics*, 12, 2215- 2227.
1834
1835 Sysalova, J., Sykorova, I., Havelcova, M., Szakova, J., Trejtnarova, H. and Kotlik, B. (2012).
1836 Toxicologically important trace elements and organic compounds investigated in size –fractionated
1837 urban particulate matter collected near the Prague highway. *Science of the total Environment*, 437,
1838 127- 136.
1839
1840 Tanner, P.A., Hoi-Ling, M. and Yu, P.K.N. (2008). Fingerprinting metals in street dust in Beijing,
1841 Shanghai and Hong Kong. *Environmental Science and Technology*, 42, 7111-7117.
1842
1843 Teixeira, E.C., Agudelo-Castaneda, D.M., Fachel, J.M.G., Leal, K.L., Garcia, K.O. and Wiegand, F.
1844 (2012). Source identification and seasonal variation of polycyclic aromatic hydrocarbons associated
1845 with atmospheric fine and coarse particles in the Metropolitan Area of Porto Alegre, RS, Brazil.
1846 *Atmospheric Research*, 118: 390-403.
1847
1848 Tervahattu, H., Kupiainen, K.J., Raisanen, M., Makela, T. and Hillamo, R. (2006). Generation of
1849 urban road dust from anti-skid and asphalt concrete aggregates. *Journal of Harazdous Materials*,
1850 132, 39-46.
1851
1852 Thorpe, A. J., Harrison, R. M., Boulter, P. G. and McCrae, I. S. (2007). Estimation of particle
1853 resuspension source strength on a major London Road. *Atmospheric Environment*, 41, 8007-8020.
1854
1855 Thorpe, A. and Harrison, R. M. (2008). Sources and properties of non-exhaust particulate matter
1856 from road traffic: a review. *Science of the Total Environment*, 400(1-3), 270-282.
1857
1858 Van Poppel, M., Panis, L.I., Govarts, E., Van Houtte, J. and Maenhaut, W. (2012). A comparative
1859 study of traffic related air pollution next to a motorway and a motorway flyover. *Atmospheric*
1860 *Environment*, 60, 132-141
1861
1862 Varrica, D., Bardelli, F., Dongarra, G. and Tamburro, E. (2012). Speciation of Sb in airborne
1863 particulate matter, vehicle brake linings and brake pad wear residues. *Atmospheric Environment*, 64,
1864 18-24.
1865
1866 Viana, M., Kuhlbusch, T.A.J., Querol, X., Alastuey, A., Harrison, R.M., Hopke, P.K., Winiwarter,
1867 W., Vallius, M., Szidat, S., Prevot, A.S.H., Hueglin, C., Bloemen, H., Wahlin, P., Vecchi, R.,
1868 Miranda, A.I., Kasper-Giebl, A., Maenhaut, W. And Hitzenberger, R. (2008). Source apportionment
1869 of particulate matter in Europe: A review of methods and results. *Aerosol Science*, 39, 827-849.
1870
1871 Wahlin, P., Berkowicz, R. and Palmgren, F. (2006). Characterisation of traffic-generated particulate
1872 matter in Copenhagen. *Atmospheric Environment*, 40(12), 2151-2159.
1873
1874 Wang, G., Kawamura, K., Xie, M., Hu, S., Gao, S., Cao, J., An, Z. and Wang, Z. (2009). Size
1875 distribution on n-alkanes, PAHs and hopanes and their sources in the urban, mountain and marine
1876 atmospheres over East Asia. *Atmospheric Chemistry and Physics*, 14: 8869-8882.
1877

1878 Wang, F., Ketzel, M., Ellermann, T., Wahlin, P., Jensen, S.S., Fang, D. and Massling, A. (2010).
1879 Particle number, particle mass and NO_x emission factors at a highway and an urban street in
1880 Copenhagen. *Atmospheric Chemistry and Physics*, 10, 2745-2764.
1881
1882 Wang, X., Westerdahl, D., Wu, Y., Pan, X. and Zhang, K. M. (2011). On-road emission factor
1883 distributions of individual diesel vehicles in and around Beijing, China. *Atmospheric Environment*,
1884 45(2), 503-513.
1885
1886 Watson, J.G., Chow, J.C., Lowenthal, D.H., Pritchett, L.C., Frazier, C.A., Neuroth, G.R. and
1887 Robbins, R. (1994). Differences in the carbon composition of source profiles for diesel- and
1888 gasoline- powered vehicles. *Atmospheric Environment*, 28(15), 2493-2505.
1889
1890 Watson, J. G., Zhu, T., Chow, J. C., Engelbrecht, J., Fujita, E. M. and Wilson, W. E. (2002).
1891 Receptor modeling application framework for particle source apportionment. *Chemosphere*, 49(9),
1892 1093-1136.
1893
1894 Watson, J.G. and Chow, J.C. (2007). Receptor models for source apportionment of suspended
1895 particles. In *Introduction to Environmental Forensics*, 2nd Edition, 2, B. Murphy and R. Morrison
1896 (Eds.), Academic Press, New York, NY, 279-316.
1897
1898 Wehner, B., Birmili, W., Gnauk, T. and Wiedensohler, A. (2002). Particle number size distributions
1899 in a street canyon and their transformation into the urban-air background: measurements and a
1900 simple model study. *Atmospheric Environment*, 36, 2215-2223.
1901
1902 Wei, B. and Yang, L. (2010). A review of heavy metal contaminations in urban soils, urban road
1903 dusts and agricultural soils from China. *Microchemical Journal*, 94(2), 99-107.
1904
1905 Weingartner, E., Keller, C., Stahel, W. A., Burtscher, H. and Baltensperger, U. (1997). Aerosol
1906 emission in a road tunnel. *Atmospheric Environment*, 31(3), 451-462.
1907
1908 Weijers, E. P., Khlystov, A. Y., Kos, G. P. A. and Erisman, J. W. (2004). Variability of particulate
1909 matter concentrations along roads and motorways determined by a moving measurement unit.
1910 *Atmospheric Environment*, 38, 2993-3002.
1911
1912 Westerdahl, D., Wang, X., Pan, X. and Zhang, M.K. (2009). Characterization of on-road vehicle
1913 emission factors and microenvironmental air quality in Beijing, China. *Atmospheric Environment*,
1914 43, 697-705.
1915
1916 Wik, A. and Dave, G. (2009). Occurrence and effects of tire wear particles in the environment--a
1917 critical review and an initial risk assessment. *Environmental Pollution*, 157(1), 1-11.
1918
1919 Wu, S.-P., Tao, S., Zhang, Z.-H., Lan, T. and Zuo, Q. (2007). Characterization of TSP-bound n-
1920 alkanes and polycyclic aromatic hydrocarbons at rural and urban sites of Tianjin, China.
1921 *Environmental Pollution*, 147(1), 203-210.
1922
1923 Yan, B., Zheng, M., Hu, Y., Ding, X., Sullivan, A. P., Weber, R. J., Baek, J., Edgerton, E. S. and
1924 Russell, A.G. (2009). Roadside, Urban, and Rural Comparison of Primary and Secondary Organic
1925 Molecular Markers in Ambient PM_{2.5}. *Environmental Science and Technology*, 43(12), 4287-4293.
1926
1927 Yin, J., Harrison, R. M., Chen, Q., Rutter, A. and Schauer, J. J. (2010). Source apportionment of
1928 fine particles at urban background and rural sites in the UK atmosphere. *Atmospheric Environment*,
1929 44, 841-851.

- 1930 Zhang, Y., Stedman, D. H., Bishop, G. A., Guenther, P.L. and Beaton, S. P. (1995). Worldwide On-
1931 Road Vehicle Exhaust Emissions Study by Remote Sensing. *Environmental Science*
1932 *and Technology*, 29, 2286-2294.
- 1933
- 1934 Zhu, Y., Hinds, W. C., Kim, S. and Sioutas, C. (2002a). Concentration and Size Distribution of
1935 Ultrafine Particles Near a Major Highway. *Journal and Air and Waste Management Association*, 52,
1936 1032-1042.
- 1937
- 1938 Zhu, Y., Hinds, W. C., Kim, S., Shen, S. and Sioutas, C. (2002b). Study of ultrafine particles near a
1939 major highway with heavy-duty diesel traffic. *Atmospheric Environment*, 36(27), 4323-4335.
- 1940
- 1941 Zhu, Y., Kuhn, T., Mayo, P. and Hinds, W.C. (2006). Comparison of daytime and nighttime
1942 concentration profiles and size distributions of ultrafine particles near a major highway.
1943 *Environmental Science and Technology*, 40, 2531-2536.
- 1944
- 1945 Zhu, C.-S., Chen, C.-C., Cao, J.-J, Tsai, C.-J., Chou, C. C.-K., Liu, S.-C. and Roam, G.-D. (2010).
1946 Characterization of carbon fractions for atmospheric fine particles and nanoparticles in a highway
1947 tunnel. *Atmospheric Environment*, 44, 2668-2673.
- 1948
- 1949
- 1950

1951 **TABLE LEGENDS**

1952

1953 **Table 1:** Mechanisms for non-exhaust particle emission (Rogge et al., 1993;
1954 Boulter et al., 2005; Wahlin et al., 2006; Thorpe et al. 2007; Amato et al.,
1955 2009; Harrison, 2009; Gietl et al., 2010; Barmapadimos et al., 2011;
1956 Hays et al., 2011; Denier van der Gon, 2013).

1957

1958 **Table 2:** Key tracers used for non-exhaust PM.

1959

1960 **Table 3:** Ratio of Cu to Sb characteristic of brake wear particles in ambient PM
1961 reported in the literature.

1962

1963 **Table 4:** (a) Results from mass size distribution analysis for traffic-related PM as
1964 reported in the literature (in μm); (b) Results from mass size distribution
1965 analysis for traffic-related elements as reported in the literature (in μm).
1966

1967

1968 **Table 5:** Comparison between dynamometer and tunnel/roadway measurements
1969 (Allen et al., 2001; Imhof et al., 2005a; Phuleria et al., 2006; Phuleria et
1970 al., 2007; Handler et al., 2008; He et al., 2008; El Haddad et al., 2009;
1971 Sanchez-Ccoyllo et al., 2009; Franco et al., 2013).

1972

1973 **Table 6:** Emission factors as reported in the literature (selected studies).

1974

1975 **Table 7:** Source apportionment analyses for road traffic-generated PM (selected
1976 studies).
1977

1978

1979 **FIGURE LEGEND**

1980

1981 **Figure 1:** Methods for direct road traffic emissions analyses.

1982

1983

1984

1985 **Table 1:** Mechanisms for non-exhaust particle emission (Rogge et al., 1993; Boulter et al., 2005;
 1986 Wahlin et al., 2006; Thorpe et al. 2007; Amato et al., 2009; Harrison, 2009; Gietl et al., 2010;
 1987 Barmapadimos et al., 2011; Hays et al., 2011; Denier van der Gon, 2013)

Emission type	Mechanism	Includes
Direct	Abrasion and wear and tear	Tyre, brake, clutch, road surface
	Corrosion	Vehicle, street furniture
Indirect	Re-suspension (due to tyre shear, wind and vehicle turbulence)	

1988

1989 **Table 2:** Key tracers used for non-exhaust PM

Reference	Brake wear	Tyre wear	Re-suspension
Adachi and Tainosho (2004)	Fe, Ba, Cu, Sb, Zr	Zn	-
Schauer et al. (2006)	Fe, Cu, Ba		-
Grieshop et al. (2006)	Cu, Sb, Ba and Ga		
Wahlin et al. (2006)	Cr, Fe, Cu, Zn, Zr, Mo, Sn, Sb, Ba and Pb	Al, Si, K, Ca, Ti, Mn, Fe, Zn and Sr (together with road dust)	
Tanner et al. (2008)	Cu, Cd	Zn	
Canepari et al. (2008)	Ba, Fe, Sb, Sr	-	-
Harrison (2009)	Ba, Cu	-	Al
Dongarra et al. (2009)	Cu, Mo, Sb	-	
Fabretti et al. (2009)	Cu, Zn, Sb, Sn (vehicular abrasion)	-	Rb, Sr, Mn, Fe, As
Keuken et al. (2010)	Cu	Zn	-
Bukowiecki et al. (2010)	Fe, Cu, Zn, Zr, Mo, Sn, Sb and Ba		
Pey et al. (2010)	-	-	Fe, Ca, Sb, Sn, Cu, Zn
Perez et al. (2010)	Sb, Cu, Ni, Sn (wear of brake, tire and other parts)		Fe
Amato et al. (2011a)	Fe, Cu, Zn, Cr, Sn, Sb	OC, S, Zn	Al, Ca, Fe, V
Apegyei et al. (2011)	Fe, Ti, Cu, Ba	Zn, Ca, W, K, Fe, Ti, Cr, Mo	-
Duong and Lee (2011)	Ni, Cu	Zn	
Ondracek et al. (2011)	Cu, Ba, Fe, Zn	-	
Song and Gao (2011)	Sb, Cu, Fe, Pb	Zn, Co	
Sahu et al. (2011)	Zn (brake and tyre wear)	-	
Peltier et al. (2011)	-	-	Al, Si, Ti, Fe
Harrison et al. (2012b)	Ba, Cu, Fe, Sb	Zn	Si, Al

1990

1991

1992 **Table 3:** Ratio of Cu to Sb characteristic of brake wear particles in ambient PM reported in the
 1993 literature

Reference	Location	Cu:Sb ratio
Stechmann and Dannecker (1990)	Hamburg, Germany	5.6
Adachi and Tainosho (2004)	Kobe, Japan	1.3
Lin et al. (2005)	Taiwan	9.1±1.79 (coarse fraction) 2.17±0.83 (fine fraction)
Wahlin et al. (2006)	Copenhagen, Denmark	4.4±0.3
Hjortenkrans et al. (2007)	Stockholm, Sweden	5.3
Fabretti et al. (2009)	Nice, France	7.0±1.3
Dongarra et al. (2009)	Palermo, Italy	4.9 (coarse fraction) 2.5 (fine fraction)
Amato et al. (2011b)	Barcelona, Spain	8.0 ± 1.5 (roadside) 6.8 ± 1.9 (urban)
Bukowiecki et al. (2010)	Zürich-Weststrasse, Switzerland	6.2 (coarse fraction)
Pey et al. (2010)	Barcelona, Spain	9 (coarse fraction) 10 (fine fraction)
Gietl et al. (2010)	London, UK	9.1

1994

1995

1996 **Table 4: (a)** Results from mass size distribution analysis for traffic-related PM as reported in the
 1997 literature (in μm)

Reference	Location	Site type	Component	Distribution type	Primary peak
Allen et al. (2001)	USA	Tunnel	PM ₁₀	Unimodal	0.1-0.18
Hildemann et al. (1991)	USA	Freeway	PM	Trimodal	Ultrafine (<0.2), fine (0.2-1) and coarse (1-10)
Robert et al. (2007)	USA	Dynamometer	PM _{1.8} (HDGV)	Unimodal	0.1-0.18
			PM _{1.8} (LDGV)	Unimodal	0.1-0.18

1998 **(b)** Results from mass size distribution analysis for traffic-related elements as reported in the
 1999 literature (in μm)
 2000

Reference	Location	Metal(s)	Distribution type	Primary peak	Secondary peak
Stechmann and Dannecker (1990)	Germany	Cu, Ba	Unimodal	2.5	
Funasaka et al. (1998)	Japan (tunnel)	EC	Unimodal		
		OC	Bimodal	1	2-10
Lin et al., 2005	Taiwan	Si, Al, Fe, Ca	Bimodal	3.2- 5.6	Nano size range
		Ba	Bimodal	1.0-1.8	0.18-0.32
		Pb	Bimodal	3.2- 5.6	0.32- 0.56
		Zn	Bimodal	1.0-1.8	0.32-0.56
		Mn, Cu, Cd, Sr	Bimodal		
		Cr, V	Trimodal		
Lough et al. (2005)	USA	Cu	Bimodal	3.0	0.1
Wahlin et al. (2006)	Denmark	Ba, Sb	Unimodal	3	
		Cu	Bimodal	3	0.1
Hien et al. (2007)	Vietnam	PAHs in PM _{2.5}	Bimodal	Coarse and fine	
Riddle et al. (2007)	USA	Hopanes and steranes in PM _{1.8}	Unimodal	0.10-0.18	0.32 (vehicles without catalysts)
Wang et al. (2009)	China	n-Alkanes, PAHs and Hopanes	Unimodal	0.7-1.1	
Gietl et al. (2010)	UK	Ba, Fe	Unimodal	1.2- 7.2	
		Cu, Sb	Bimodal	1.2- 7.2	0.2- 0.4
Huang et al. (2012)	China	n-Alkanes	Unimodal	0.32-0.56	
		PAHs	Unimodal	0.56-1.0	
		Hopanes	Unimodal	0.56-1.0	3.2 (in some cases)
Ondráček et al. (2011)	Czech Republic (freeway)	Fe, Cu, Mn, Zn	Unimodal	2.5	
		Si, Al, Ca and coarse mode K	Unimodal	5	
Song and Gao (2012, 2011)	USA	Al, Fe, Cu, Sb, Sc, Mn	Bimodal	3.2- 5.6	1.0-1.8
		Cd	Bimodal	0.18- 0.56	3.2- 5.6
		Pb, Zn	Trimodal	0.32-0.56	1.0-1.8, 3.2- 5.6
		Ni, V	Bimodal	0.18-0.32	3.2- 5.6

2001 **Table 5:** Comparison between dynamometer and tunnel/roadway measurements (Allen et al., 2001;
 2002 Imhof et al., 2005a; Phuleria et al., 2006; Phuleria et al., 2007; Handler et al., 2008; He et al., 2008;
 2003 El Haddad et al., 2009; Sanchez-Ccoyllo et al., 2009; Franco et al., 2013)

	Dynamometer measurements	Tunnel/Roadway measurements
Testing conditions	Precise and controlled; different test cycles and driving conditions can be analysed; can include cold-start emissions	Ambient conditions- cannot be physically controlled; cannot include cold-start emissions Better known boundary conditions and dilution effects In some cases, atmospheric conditions in tunnel may not be the same as ambient environment
Representativeness for the vehicle fleet	Low, since the tests are conducted on a sub-set of the in-use fleet and variations in engine type, vehicle age and maintenance and mixing of emissions from different vehicles etc are not accounted for. Such studies do not account for particle aging effects	High, since the measurements are made in ambient environment and for in-use mixed fleet
Emission type accounted for	Exhaust	Exhaust and non-exhaust

2004

Table 6: Emission factors as reported in the literature (selected studies)

- Particle Number

Reference	City, Country	Study type	Particle number EF
Gertler et al. (2002)	USA	Tunnel (mixed LDV and HDV)	5x 10 ¹² particles/veh/km (low HDV) 2-3 x 10 ¹³ particles/veh/km (high HDV)
Abu-Allaban et al. (2002)	USA	Tunnel	5.2- 5.4 x 10 ¹³ particles/veh/km (low HD fraction) 2.1- 3.1x 10 ¹⁴ particles/veh/km (high HD fraction)
Jamriska et al. (2004)	Australia	Tunnel (diesel powered buses)	3.11 ± 2.41 x 10 ¹⁴ particles/km
Kittleston et al. (2004)	USA	Roadside (particles > 3 nm)	1.9- 9.9 x 10 ¹⁴ particles/km (gasoline-dominant fleet)
Jones and Harrison (2006)	UK	Roadside	5.84 x 10 ¹³ particles/veh/km 6.36 x 10 ¹⁴ particles/veh/km (LDV)
Westerdahl et al. (2009)	China	Roadside- LDV (prevalent during daytime)	1.8 x 10 ¹⁵ particles/kg fuel
		Roadside- HDV (prevalent during nighttime)	1.1 x 10 ¹⁶ particles/kg fuel
Wang et al. (2009)	Denmark	Highway	(215±15) 10 ¹² particles/ veh/km
Ban-Weiss et al. (2010)	USA	Tunnel- LDV	(3.9±1.4) 10 ¹⁴ particles/kg fuel
		Tunnel-HDV	(3.3±1.3) 10 ¹⁵ particles/kg fuel
Wang et al. (2011)	China	Vehicle chase	7.1 x 10 ¹⁶ particles/kg fuel

- Particle Mass

Reference	City, Country	Study type	PM size range	EF
Weingartner et al. (1997)	Switzerland	Tunnel (mixed LDV and HDV with more LDV)	PM ₃	310 mg/km (63% vehicles running on diesel)
Gillies et al. (2001)	USA	Tunnel (mixed LDV and HDV)	PM _{2,5}	0.052 ± 0.027 g/km
			PM ₁₀	0.069 ± 0.030 g/km
Gertler et al. (2002)	USA	Tunnel (mixed LDV and HDV)	PM _{2,5}	0.062 ± 0.042 g/km

		HDV)	PM ₁₀	0.087 ± 0.054 g/km
Jamriska et al. (2004)	Australia	Tunnel (diesel powered buses)	PM _{2.5}	583± 451 mg/km
Greishop et al. (2006)	USA	Tunnel	PM _{2.5} (rush hour)	189±23 mg/kg fuel
			PM _{2.5} (mid-day)	158±29 mg/kg fuel
			PM _{2.5} (early morning-HDV)	437±76 mg/kg fuel
Handler et al. (2008)	Austria	Tunnel	TSP	129±45 mg/veh/km
			PM ₁₀	62±18 mg/veh/km
			PM _{2.5}	26±10 mg/veh/km
Sanchez-Ccoyllo et al., (2009)	Brazil		Coarse	127 mg/km
			Fine	92 mg/km
Wang et al. (2010)	Denmark	Highway	PM _{2.5}	29 mg/veh/km
Chiang and Huang (2009)	Taiwan	Freeway tunnel	PM _{2.5-10}	18±6.5 mg/veh/km
			PM _{2.5}	39±11 mg/veh/km
Cheng et al. (2010)	Hong Kong	Tunnel, roadside sites and urban site	PM _{2.5}	257±31 mg/veh/km
Bukowiecki et al. (2010)	Switzerland	Roadside	PM ₁₀ - LDV	24±8 mg/veh/km
			PM ₁₀ -HDV	498±86 mg/veh/km
Wang et al. (2011)	China	Vehicle chase	PM _{0.5}	2.35 g/kg fuel
Mancilla and Mendoza (2012)	Mexico	Tunnel	PM ₁₀	22.8±7.4 mg/veh/km
Chen et al. (2013)	China	Tunnel (mixed with higher proportion of gasoline vehicles)		687 mg/veh/km (summer) 714 mg/veh/km (winter)

- LDV and HDV vehicles (mass)

Reference	Country	Study type	PM size range	LDV EF	HDV EF
Weingartner et al. (1997)	Switzerland	Tunnel	PM ₃	8.53 ± 0.47 mg/km	383.5 ± 10.7 mg/km
Gertler et al. (2002)	USA	Tunnel (mixed LDV and HDV)	PM _{2.5}	0.014 ± 0.013 g/km	0.135 ± 0.018 g/km
			PM ₁₀	0.010 ± 0.011 g/km	0.181 ± 0.013 g/km
Abu-Allaban et al.	USA	Roadside (LD-Spark)	PM10	10-70 mg/km/vehicle	60-570 mg/km

(2003)		ignition and HD-diesel)	PM _{2.5}	10-50 mg/km/vehicle	60-480 mg/km
Chiang et al. (2012)	Taiwan	Dynamometer	PM _{2.5}	0.172 g/km (diesel)	
Oanh et al. (2010)	Bangkok	Dynamometer	PM _{2.5}	230 mg/km	176 mg/km
Jones and Harrison (2006)	UK	Roadside	PM ₁₀	0.033±0.006 g/veh/km	0.370±0.032 g/veh/km
			PM _{2.5}	0.010±0.004 g/veh/km	0.179±0.022 g/veh/km
Liacos et al. (2012)	USA	Freeway	PM _{2.5}	151189.3 mg/km/h (low HDV)	169629.5 mg/km/h (high HDV)

- Non-exhaust emissions

Reference	City, Country	Study type	PM size range	Road dust	Brake wear	Tyre wear
Abu-Allaban et al. (2003)	USA	Roadside (LD-Spark ignition and HD-diesel)	PM ₁₀ LDV	40-780 mg/km/vehicle	0-80 mg/km/vehicle	-
			PM ₁₀ HDV	230-7800 mg/km/vehicle	0-610 mg/km/vehicle	
			PM _{2.5} LDV	2-25 mg/km/vehicle	0-5 mg/km/vehicle	-
			PM _{2.5} HDV	15-300 mg/km/vehicle	0-15 mg/km/vehicle	
Bukowiecki et al. (2010)	Switzerland	Roadside	PM ₁₀	27 mg/vehicle/km	15 mg/ vehicle /km	
Amato et al. (2012)	Spain	Freeway	PM ₁₀	22.7 ± 14.2 mg/vehicle kilometer travelled	-	-

Table 7: Source apportionment analyses for road traffic-generated PM (selected studies)

Reference	Country	Method	Particle type	Sources
Lin et al. (2005)	Taiwan	PCA	Coarse particles	Fugitive dust and brake lining, diesel, fuel oil, gasoline
			Fine particles	Gasoline, diesel, brake lining/tyre wear, fuel oil
			Ultrafine and nano particles	Gasoline, diesel, industry, fuel oil
Furujsjo et al. (2007)	Stockholm, Sweden	PMF	PM ₁₀	Resuspension Vehicle derived (brake wear) Road salt Regional combustion Long range transport
Chan and Mozurkewich (2007)	Ontario, Canada	Absolute PCA	Particle number distribution with trace gas and meteorological data	Common sources across 3 sites: Nucleation particles (photochemically produced), regional pollution factor, boundary layer dynamics factor Specific sources at the three sites: Local industrial emissions (urban) Processed nucleation mode particles (polluted rural) Transported fine particles (downwind Toronto)
Amato et al. (2009)	Barcelona, Spain	PMF2 with ME	PM ₁₀ and PM _{2.5}	Road dust factor accounted for 17% PM ₁₀ , 8% PM _{2.5} and less than 2% of PM ₁ . All traffic sources contributed to 46% PM ₁₀ , 51% PM _{2.5} and 48% of PM ₁ .
Fabretti et al. (2009)	Nice, France	PMF	PM _{2.5}	Vehicle abrasion: Cu, Zn, Sb (36%) Re-suspension: Mn, Fe, As, Rb, Sr (43%) Fuel combustion: V, Ni, Co (21%)
Dreyfus et al. (2009)	Wilmington, USA	PMF	Organic carbon	Diesel factor: Alkyl fragment, benzothiazole, benzenedicarboxylic acids/3,4-dimethoxybenzaldehyde, 3,4-dimethoxybenzoic acid, hopanes Road dust/car emissions: hopanes, benzothiazole and triacontanoic acid
Mancilla and Mendoza (2012)	Monterrey, Mexico	FA	PM _{2.5}	Brake emissions Exhaust emissions
Ondracek et al. (2011)	Prague, Czech Republic			Abrasion of vehicle parts: Fe, Cu, Mn, Zn Re-suspension of road dust: Si, Al, Ca Long-range transport/regional background: Ca, K
Song and Gao (2011)	New Jersey, USA	FA	PM	Brake wear and fuel combustion: Fe, Sb, Pb, Cd (~35%) Primary fuel combustion: Cr, Ni, V, Cu (~28.3%) Tyre wear and fuel combustion: Zn, Co (~23.7%)
Yin et al. (2010)	United Kingdom	CMB	PM _{2.5}	Vegetative detritus; woodsmoke; natural gas; diesel engines (11.6%); gasoline engines (1.4%); smoking engines (9.7%); coal; dust/soil; secondary organic matter; sea salt; ammonium sulphate; ammonium nitrate




Figure 2: Methods for direct road traffic emissions analyses