Polar cap patches, GPS TEC variations, and atmospheric gravity waves

Prikryl, Paul; Gillies, Robert G.; Themens, David R.; Kunduri, Bharat S.-R.; Varney, Roger; Weygand, James M.

DOI:

[10.5194/egusphere-egu21-6775](https://doi.org/10.5194/egusphere-egu21-6775)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Prikryl, P, Gillies, RG, Themens, DR, Kunduri, BSR, Varney, R & Weygand, JM 2021, Polar cap patches, GPS TEC variations, and atmospheric gravity waves. in *EGU General Assembly 2021*, EGU21-6775, European Geosciences Union, EGU General Assembly 2021, 19/04/21. <https://doi.org/10.5194/egusphere-egu21-6775>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

EGU21-6775

<https://doi.org/10.5194/egusphere-egu21-6775>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.

Polar cap patches, GPS TEC variations, and atmospheric gravity waves

Paul Prikryl^{1,2}, Robert G. Gillies³, David R. Themens^{1,4}, Bharat S. R. Kunduri⁵, Roger Varney⁶, and James M. Weygand⁷

¹University of New Brunswick, Physics Department, Ottawa, ON, Canada

²Geomagnetic Laboratory, Natural Resources Canada, Ottawa, ON, Canada

³Department of Physics and Astronomy, University of Calgary, AB, Canada

⁴School of Engineering, University of Birmingham, Birmingham, UK

⁵Bradley Department of Electrical and Computer Engineering, Virginia Tech, Blacksburg, VA, USA

⁶Center for Geospace Studies, SRI International, Menlo Park, California, USA

⁷Earth Planetary and Space Sciences, University of California, Los Angeles, CA, USA

The southward pointing field of view of the Canadian component of the Resolute Bay Incoherent Scatter Radar (RISR-C) is well suited for observing the ionospheric signatures of flux transfer events and subsequent polar patch formation in the cusp. The fast azimuthally oriented flows and associated density depletions often show an enhanced ion temperature from Joule heating caused by the sudden change in plasma flow direction. The newly formed polar patches are then observed as they propagate through the field-of-views of both RISR-C and RISR-N. In the ionosphere, the electron density gradients imposed in the cusp, and small-scale irregularities resulting from gradient-drift instability, particularly in the trailing edges of patches, cause GPS TEC and phase variations, and sometimes amplitude scintillation. The neutral atmosphere is affected by ionospheric currents resulting in Joule heating. The pulses of ionospheric currents in the cusp launch atmospheric gravity waves (AGWs) causing traveling ionospheric disturbances, as they propagate equatorward and upward. On the other hand, the downward propagating AGW packets can impact the lower atmosphere, including the troposphere. Despite significantly reduced wave amplitudes, but subject to amplification upon over-reflection in the upper troposphere, these AGWs can trigger/release existing moist instabilities, initiating convection and latent heat release, the energy leading to intensification of storms.