The rise of non-traditional pregnancies through assisted reproductive technologies

Kaur, Amarpreet

DOI:

[10.12968/bjom.2021.29.2.82](https://doi.org/10.12968/bjom.2021.29.2.82)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kaur, A 2021, 'The rise of non-traditional pregnancies through assisted reproductive technologies', *British Journal of Midwifery*, vol. 29, no. 2, pp. 82-88. <https://doi.org/10.12968/bjom.2021.29.2.82>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This document is the Accepted Manuscript version of a Published Work that appeared in final form in the British Journal of Midwifery, copyright © MA Healthcare, after peer review and technical editing by the publisher. To access the final edited and published work see: <https://doi.org/10.12968/bjom.2021.29.2.82>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The Rise of Non-Traditional Pregnancies through Assisted Reproductive Technologies

Abstract

This article explores how the development of assisted reproductive technologies (ARTs) have perpetuated an increase in non-traditional pregnancies. It discusses what this increase means for midwifery practices and what care midwives may need to consider for such pregnancies. The discussions in this article are based on triangulated findings from a three-phase research design. The research consisted of a mixed-methods online survey of 521 citizens of the United Kingdom, semi-structured interviews with experts and professionals who could speak to the future of potential ARTs, and with people who are affected by genetic conditions. Findings reveal that applications of ARTs are anticipated to continue to expand with the potential introduction of genome editing technologies to reproductive choices for the prevention of disease. Thus, this article concludes that because of this expansion, the rise of non-traditional pregnancies is likely to continue, and that midwives may benefit from considering the implications the rise could have for midwifery care.

Keywords

Assisted Reproduction; Genetic Disease; Geriatric Mothers; High Risk Pregnancies; *In Vitro* Fertilisation; Surrogacy

Key Points

- Since the introduction of *in vitro* fertilisation (IVF) in the late 1970's, it has been used to enable a number of other applications which have led to an increase in non-traditional pregnancies.
- Current applications of IVF technologies beyond infertility include egg freezing, egg donation, preimplantation genetic testing (PGT), and surrogacy.
- Non-traditional pregnancies include those involving geriatric mothers, people with genetic conditions, surrogate mothers, and same-sex couples.
- Non-traditional pregnancies are more likely to require greater variations of typical midwifery care and/or antenatal and postnatal support.

Background

In vitro fertilisation (IVF) is when an egg is fertilised by sperm outside the human body, usually in a culture dish (Franklin 1997). This technique was developed by Sir Robert Geoffrey Edwards, a physiologist, and Dr Patrick Christopher Steptoe, an obstetrician and gynaecologist in the 1970s, as a means to treat female infertility (Mulkay 1997). The first baby born as a result of IVF, Louise Brown, was delivered via a caesarean section in July 1978 at Oldham Hospital in Greater Manchester (Brown 2018). Louise celebrated her fortieth birthday in 2018, and has two sons conceived without the use of any assisted reproductive technologies (ARTs). In 2018, the European Society of Human Reproduction and Embryology (ESHRE) reported that a global total of more than eight million babies had been born from IVF since the technology was first pioneered (ESHRE 2018 Jul).

The development of IVF technology has also enabled several other applications which extend beyond treating female infertility (Sarojini and Marwah 2015). Such

applications include circumventing male infertility, facilitating people undergoing various medical treatments, such as chemotherapy, to preserve their gametes, and women to extend their fertility through egg freezing (Edwards et al. 1999; Crawshaw et al. 2009; Inhorn 2015; van de Wiel 2015; Baldwin 2019). Additionally, other applications based on IVF have empowered people who may not have otherwise considered bearing a genetically related child to have more reproductive choices. Such people include those with a genetic condition and/or same-sex couples. These choices include preimplantation genetic testing (PGT), i.e. testing an embryo for the disease its parents' are affected by before the embryo is transferred into a uterus in the hope of establishing a pregnancy, using donated gametes, and surrogacy (Kaur and Border 2020: 2). There is potential for ARTs to expand to include editing embryos to prevent genetic disease. The development of ARTs has thus led to a rise in non-traditional pregnancies (NTPs) (Thompson 2007).

Non-traditional Pregnancies

NTPs are those which thwart conventional pregnancies, i.e. pregnancies which would otherwise be contingent on sexual reproduction, and natural biology. NTPs include those involving geriatric mothers, people with genetic conditions, surrogate mothers, and same-sex couples (Thompson 2007). Geriatric mothers are women who have their first pregnancy aged thirty-five years or older (RCOG 2011: 2). Geriatric mothers result from women being able to bypass age related infertility through IVF (Royal College of Obstetricians and Gynaecologists (RCOG) 2011), and women purposefully planning to circumvent age related fertility by freezing their eggs (Baldwin 2019). While late motherhood may be welcomed and celebrated by many women, it can generate high-risk pregnancies. This is because older women are at

higher risk of many medical complications. Examples of such complications include gestational hypertension, prolonged labour, stillbirth, and postpartum haemorrhaging (RCOG 2011).

Women with underlying health conditions, including health conditions which can result from having a genetic condition, are also often considered to be high-risk (Franklin and Roberts 2006). However, women with high-risk factors will all still require a level of routine midwifery care (Narayan and Morton 2015). Women aware of their high-risk status may require additional emotional support to alleviate their anxieties, especially if their pregnancy has been achieved following the use of ARTs (Cooper and Glazer 1999; Stevenson and Hershberger 2016). This is because IVF has a very modest average success rate of one in four, and this rate steeply declines as maternal age advances; this is illustrated in *Figure 1*.

Figure 1: Success Rate of IVF in relation to Maternal Age (HFEA 2020 Jun 30)

The physical, emotional, and financial expense of IVF can therefore make any resulting pregnancy from it even more valuable to the parents, who may have invested all they have in their desire for a child (Kaur 2020). Additionally, a review of articles on high-risk pregnancies found that women experiencing such pregnancies had worse emotional status than women with low-risk pregnancies (Rodrigues et al. 2016). Such emotional states include heightened anxiety and higher levels of stress, which may have pronounced implications antenatally, and greater intensities of depression postnatally.

For other people, ARTs may enable them to hope for a, at least partially, genetically related child if they cannot gestate a pregnancy themselves. This is true for same-sex couples, particularly gay men, and people with other biological/health limitations which could make them dependent on gamete donation and/or surrogacy to achieve their reproductive goals (Mamo 2007; Smietana et al. 2018). This could mean that antenatal and postnatal care may involve greater levels of consideration than would be typically extended. This is because in addition to the surrogate, the intended parents may wish to be part of their surrogate's care (MacCallum et al. 2003; Horsey 2010).

Furthermore, postnatally, the surrogate mother and the baby are likely to be separated, at least upon discharge from delivery settings. This would mean that community care for the mother and baby could be required at two different sites (Jadva et al. 2003; Horsey 2010). Ensuring that both parties are well may require more/different support than would traditionally be tendered. The ongoing expansion of ARTs is therefore clearly likely to impact midwifery practices to some extent, and

so, midwives should engage in discussions on how to support woman and their families who have invested in the use of ARTs to achieve their pregnancies and (hopefully) their resulting child(ren).

Methodologies

The expansion of ARTs to include a developing technology which could be used to edit human embryos before they are transferred into a uterus, was explored using a three-phase research design. All three phases of the research were designed in adherence to the British Sociological Association's (BSA) (2017) guidelines on conducting ethical research; they also received ethical approval from the University of Cambridge's Department of Sociology's Ethics Committee. The three phases of research were conducted between March 2018 and October 2019.

The first phase of the research consisted of a mixed-methods online survey of 521 citizens of the UK, aged 16-82 years, 52% of whom self-identified as female and 48% as male, on *Understandings of Genetic Editing and its Potential Uses with Human Reproduction*. Table 1 and Table 2 support this claim.

Statistics			Age Ranges			
			Frequency	Percent	Valid Percent	Cumulative Percent
What age are you?	Valid	521	16-19	48	9.2	9.2
	Missing	0	20-24	85	16.3	25.5
Minimum	16	82	25-29	78	15.0	40.5
			30-39	118	22.6	63.1
Maximum	82		40-49	91	17.5	80.6
			50-59	68	13.1	93.7
			60+	33	6.3	100.0
			Total	521	100.0	

Table 1: Respondents' Age

What gender are you?					
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Female	271	52.0	52.0	52.0
	Male	250	48.0	48.0	100.0
	Total	521	100.0	100.0	

Table 2: Respondents' Gender

The sample size was based on achieving a confidence level of 95% with a 4-5% margin of error. This meant that $\pm 5\%$ of the approximated total population that could have responded to the survey were likely to have responded in the same way as actual respondents. The approximated population was 48,043,809; this was derived from 2011 census data in the UK. The sample was nationally representative of the age and sex of citizens in the UK based on 2011 census data from the UK. Additionally, within in the sample, 36.7% of respondents self-identified as being religious (see *Table 3*) – the six main religions were represented within this percentage. The six main religions are considered to be Buddhism, Christianity, Hinduism, Islam, Judaism, and Sikhism (Garner 2004 Apr 27).

Are you religious?					
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Yes	191	36.7	36.7	36.7
	No	330	63.3	63.3	100.0
	Total	521	100.0	100.0	

Table 3: Respondents' Religiosity

Finally, 29.2% self-identified as being affected by a genetic condition (see *Table 4*).

Have you ever been affected by a genetic condition?					
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Yes	152	29.2	29.2	29.2
	No	369	70.8	70.8	100.0
	Total	521	100.0	100.0	

Table 4: Affected by Genetic Conditions

The survey was distributed via social media and email to ensure that people of all ages, from the six main world religions, and/or affected by genetic conditions were represented within the sample. This method of distribution was also considered to be the most feasible and effective way to reach the UK's wider public, i.e. its citizens, in an eco-friendly and timely manner.

The survey had four sections; the first section was on knowledge and understanding of genome editing, the second was on hypothetical practical applications relating to factors of disease, the third was on regulation and ethics, and the final section captured demographic information. Data from the survey was imported to SPSS to be cleaned and thematically analysed. SPSS is a software largely used to analyse quantitative and statistical data (IBM, 2020). Cleaning, in this context, refers to ensuring that all the data is formatted correctly; this can include modifying data so that it is formatted correctly (Schmidt 1997; Wright 2005). For example, changing 'f', 'woman' and 'girl' to 'female' and copying and pasting answers to replace 'see previous answer' or 'same as before' where appropriate.

The qualitative answers to the survey were also imported to NVivo for coding and thematic analysis. NVivo is software used to find common themes, reasons, and/or meanings in qualitative data (QSR International, 2019). Coding is a process in which data is categorised to facilitate analysis (Gibbs 2007). Findings from the survey were then used to inform the subsequent phases of research. The second phase of research consisted of semi-structured interviews with experts and professionals who could speak to the landscape of ARTs in the UK, specifically in relation to genome editing technology which could be used to edit human embryos; this possibility is referred to as human germline genome editing (hGGE).

The experts and professionals included scientists, UK government officials, and staff from the Human Fertilisation and Embryology Authority (HFEA); they were selected based on the relevance of their respective expertise. The third phase of the research comprised of interviews with twenty-one people who are affected by genetic conditions. Data from all the interviews were transcribed and also imported to NVivo for thematic analysis. Participants in the third phase were assigned pseudonyms during transcription.

The data from all three phases of the research were triangulated to produce the findings shared in this article. The research findings should be of interest to midwives because the implications of them could impact midwifery practices; thinking about these impacts and how midwives may respond to them and/or incorporate them into midwifery practices may therefore prove to be useful for continuing professional development.

Research Findings and Discussion

Triangulated findings from the primary research suggest that hGGE could be added to ARTs in the foreseeable future as a means to enable people with genetic conditions to prevent their child(ren) inheriting their condition. This finding is based on only 4.8% (25/521) of respondents to the survey being directly opposed to hGGE being legalised in the UK as a reproductive choice; 66.22%-86.56% (345-451/521) of respondents feeling that hGGE should be allowed as various factors of disease were explored; the technology being able to adopt frameworks used for existing ARTs for sound regulation (according to experts interviewed in the second phase of research), and people with genetic conditions being supportive of enabling the technology for such purposes. The following qualitative answers from participants in the third phase of research support the latter finding:

Virginia: [hGGE] should be legalised. The key reason why I hold that opinion is because I am disabled myself, I have a genetic condition, and personally I would not want to have a child or bring another person into this world with the same genetic condition as myself. I think life with this disability is a struggle in this ableist world. Frankly, I would not choose to have this condition if I had the option, therefore I would not choose it for someone else either.

(Female, 21, has Muscular Dystrophy)

George: I actually look into genome editing quite a lot because I have Charcot Marie Tooth disease, which is a genetic condition. And I feel that if it'd been possible for me to be, like fixed early on, I feel that that's almost like a necessity, like it needs to be done.

(Male, 34, has Charcot Marie Tooth Disease)

The potential addition of hGGE to ARTs would be of significance to midwives because it could result in higher levels of people with serious genetic conditions gestating a pregnancy. Such people are likely to be considered high-risk, and/or to

require greater levels of support than people without such genetic conditions (Yali and Lobel 1999). As mentioned above, even high-risk pregnancies still require a level of midwifery care (Narayan and Morton 2015). This is important to note, because to date, PGT has been relatively unsuccessful for the minority which do try to utilise the technology, with only 38% of established pregnancies resulting in a live birth (HFEA 2018 Sep 13). This means that most midwives would have been unlikely to care for a woman who has used such ARTs, but this could change if hGGE were to be introduced to ARTs, whether in the UK or abroad (Horsey 2010; Lovell-Badge 2019). Additionally, obstetricians may not have supported people with various medical complications through pregnancy, birth, and puerperium.

The concept of individuals/couples travelling abroad to access ARTs is not a new phenomenon, and if hGGE were to become available outside the UK, presuming that traveling abroad to access hGGE is reasonable (Inhorn 2015 Nov 9; Kaur 2020). 65.64% (342/521) of respondents to the survey felt that citizens of the UK should be allowed to travel abroad to access hGGE; this percentage increased if the underlying reason for accessing the technology abroad were to be for the prevention of disease. Transnational care has implications for midwifery practices, insofar as, the use of ARTs in the UK is highly regulated by the HFEA under the Human Fertilisation and Embryology (HFE) Act 1990, as amended in 2008. The same cannot be said for other countries (Jackson 2001). As such, if women were to seek ARTs abroad, the care they receive, and/or any malpractice may not be documented and/or accessible to healthcare practitioners in the UK (Bell et al. 2015; Rosemann et al. 2019; Kaur 2020). This could impact the holistic care that midwives are expected to provide for

women and their (growing) families, even those with NTPs (Berg 2005; Sittner et al. 2005).

Additionally, if/when hGGE is introduced to ARTs as a reproductive choice for the prevention of disease, this could increase the amount of live births from people who are affected by genetic conditions (Lovell-Badge 2019). Depending on how the parents' genetic condition affects them, they may need extra support in caring for their child; such situations may therefore require referral to specialist midwives and/or other support (Yali and Lobel 1999; Ogbuehi and Powell 2015: 109). In this vein, universities, Trusts, and the Nursing and Midwifery Council (NMC) may benefit from considering whether having specialist midwives or all midwives, trained in the range of ARTs and their relation to midwifery practices would be a pragmatic approach to contemplating a rise in NTPs and their outcomes; this could be done by developing e-learning modules for registered midwives and adding such content to student midwives' syllabi. This may become increasingly important to consider as ARTs are increasingly used for purposes that extend beyond infertility – such as those shared in the background section.

Furthermore, while PGT and hGGE are both based on preventing a child from being born with the genetic condition which affects its parents, this is the only condition that should be prevented (Ormond et al. 2017; Lovell-Badge 2019). This means that the resulting child still holds risks for having any other disease, including the nine diseases routinely screened for via the blood spot test (NHS 2017 Dec 21). Midwives should be prepared to explain/reiterate this to parents who may use such ARTs to conceive their child(ren), particularly if more people choose to use them. Another

prospect of hGGE is that any person born following the use of the technology, at least initially, may require additional tests and ongoing health reviews to ensure that no adverse side-effects have been encountered (Thompson 2019).

The extent to which midwives will be/are willing to be involved in such surveillance is a matter worth ruminating before decisions are made. Although ARTs are primarily concerned with conception, the impacts the use of ARTs have evidently interplay with midwifery practices. Keeping abreast of advances to ARTs, potential ARTs, and their applications should therefore be of interest to midwives. This is because decisions regarding the births and monitoring of children born from various ARTs may have direct implications for maternity documentation and care. Midwives could therefore benefit from collectively voicing their options on such matters, particularly on how they could affect midwifery practices and the care midwives are able/should be able to provide for women. This may become increasingly important as such technologies continue to advance and potentially new forms of NTPs are encountered.

While researching the potential applications of hGGE, other possible applications of ARTs were brought to attention. One such application is uterine (womb) transplants and the possibility that transwomen and transmen may try to establish a pregnancy using a combination of ARTs (Alghrani 2016). This possibility, like hGGE, is also foreseeable (Jones et al. 2019). These potential applications are envisaged due to continuing social and scientific advances (Doudna 2020; Kaur and Border 2020), and are likely to continue to develop. How midwives will/could respond to these within clinical settings, is a topic which could benefit from further discussion and/or

research. Such NTPs are likely to challenge routine clinical observations, could potentially raise issues in maternity related documentation (Horsey 2010; Sarojini and Marwah 2015), and will have repercussions antenatally and postnatally.

Conclusion

The expansion of ARTs and their various implications are likely to increasingly impact midwifery care. This argument is based on existing ARTs, such as egg freezing and surrogacy, and potential ARTs, such as hGGE, conceivably leading to a rise in NTPs which may require greater and/or alternate levels of care. Potential NTPs include higher numbers of geriatric mothers, high-risk women, surrogate mothers, and trans-mothers (Sittner et al. 2005; Horsey 2010; van de Wiel 2015; Alghrani 2016; Baldwin 2019). All these NTPs could be exacerbated by the introduction of hGGE to ARTs due to the reproductive choices the technology could enable. To this end, this article concludes that because advances in and applications of ARTs are likely to generate an ongoing rise of NTPs, midwives could benefit from contemplating this prospect within their continuing professional development and prepare for this foreseeable eventuality.

Reflective Questions

1. How does the mode/means of conception affect the provision of midwifery care?
2. Will the expansion of ARTs require additional documentation?
3. What implications do surrogate pregnancies pose on antenatal and postnatal midwifery care?
4. What support is available for parents with genetic conditions?
5. Are all non-traditional pregnancies high-risk pregnancies?

References

- Alghrani A. 2016. Uterus transplantation: does procreative liberty encompass a right to gestate? *J Law Biosci.* 3(3):636–641. doi:10.1093/jlb/lsw048.
- Baldwin K. 2019. *Egg Freezing, Fertility and Reproductive Choice: Negotiating Responsibility, Hope and Modern Motherhood.* Emerald Publishing Limited.
- Bell D, Holliday R, Ormond M, Mainil T. 2015. Transnational Healthcare, Cross-Border Perspectives. *Social Science & Medicine.* 124:284–289. doi:10.1016/j.socscimed.2014.11.014.
- Berg M. 2005. A midwifery model of care for childbearing women at high risk: genuine caring in caring for the genuine. *J Perinat Educ.* 14(1):9–21. doi:10.1624/105812405X23577.
- British Sociological Association. 2017. BSA Statement of Ethical Practice 2017. [accessed 2020 Jan 15]. https://www.britsoc.co.uk/media/24310/bsa_statement_of_ethical_practice.pdf.
- Brown. 2018. *Louise Brown: 40 Years of IVF.* Place of publication not identified: Bristol Books CIC.
- Cooper S, Glazer ES. 1999. *Choosing Assisted Reproduction: Social, Emotional & Ethical Considerations.* Perspectives Pr.
- Crawshaw MA, Glaser AW, Hale JP, Sloper P. 2009. Male and female experiences of having fertility matters raised alongside a cancer diagnosis during the teenage and young adult years. *Eur J Cancer Care (Engl).* 18(4):381–390. doi:10.1111/j.1365-2354.2008.01003.x.
- Doudna JA. 2020. The Promise and Challenge of Therapeutic Genome Editing. *Nature.* 578(7794):229–236. doi:10.1038/s41586-020-1978-5.
- Edwards J, Franklin S, Hirsch E, Price F. 1999. *Technologies of Procreation: Kinship in the Age of Assisted Conception.* 2nd Edition. New York: Routledge.
- ESHRE. 2018 Jul. More than 8 Million Babies Born from IVF Since the World's First in 1978. *ScienceDaily.* [accessed 2020 Apr 15]. <https://www.sciencedaily.com/releases/2018/07/180703084127.htm>.
- Franklin S. 1997. *Embodied Progress: A Cultural Account of Assisted Conception.* 1 edition. London ; New York: Routledge.
- Franklin S, Roberts C. 2006. *Born and Made: An Ethnography of Preimplantation Genetic Diagnosis.* Princeton University Press.
- Garner R. 2004 Apr 27. Schools to Teach UK's Six Main Religions. *The Independent.* [accessed 2020 Feb 4]. <http://www.independent.co.uk/news/education/education-news/schools-to-teach-uks-six-main-religions-561387.html>.

Gibbs G. 2007. *Analyzing Qualitative Data*. 1 edition. London: SAGE Publications.

HFEA. 2018 Sep 13. Pre-implantation Genetic Diagnosis (PGD). Human Fertilisation and Embryology Authority. [accessed 2020 Jan 15].
<https://www.hfea.gov.uk/treatments/embryo-testing-and-treatments-for-disease/pre-implantation-genetic-diagnosis-pgd/>.

HFEA. 2020 Jun 30. Fertility Treatment 2018: Trends and Figures. Human Fertilisation and Embryology Authority. [accessed 2020 Jul 1].
https://www.hfea.gov.uk/about-us/publications/research-and-data/fertility-treatment-2018-trends-and-figures/?utm_source=Twitter&utm_medium=Tweet&utm_campaign=fertility-treatment-2018-trends-and-figures_30/06/2020.

Horsey K. 2010. Challenging Presumptions: Legal Parenthood and Surrogacy Arrangements. *Child & Fam L Q*. 22(4):449–474.

Inhorn. 2015. *Cosmopolitan Conceptions: IVF Sojourns in Global Dubai*. United States: Duke University Press.

Inhorn. 2015 Nov 9. The Rise of Reprotourism. [accessed 2020 Jan 15].
<https://www.foreignaffairs.com/articles/2015-11-09/rise-reprotourism>.

Jackson E. 2001. *Regulating Reproduction: Law, Technology and Autonomy*. 01 edition. Oxford ; Portland, Or: Hart Publishing.

Jadva V, Murray C, Lycett E, MacCallum F, Golombok S. 2003. Surrogacy: the experiences of surrogate mothers. *Hum Reprod*. 18(10):2196–2204.
doi:10.1093/humrep/deg397.

Jones BP, Williams NJ, Saso S, Thum M-Y, Quiroga I, Yazbek J, Wilkinson S, Ghaem-Maghami S, Thomas P, Smith JR. 2019. Uterine transplantation in transgender women. *BJOG: An International Journal of Obstetrics & Gynaecology*. 126(2):152–156. doi:10.1111/1471-0528.15438.

Kaur A. 2020. Could Seeking Human Germline Genome Editing Force Journeys of Transnational Care? *Multidisciplinary Journal of Gender Studies*. 9(2):184–209.
doi:10.17583/generos.2020.5077.

Kaur A, Border P. 2020. Human Germline Genome Editing. Parliamentary Office of Science and Technology POSTnote Report No.: 611.

Lovell-Badge R. 2019. CRISPR Babies: A View from the Centre of the Storm. *Development*. 146(3). doi:10.1242/dev.175778. [accessed 2020 Feb 8].
<https://dev.biologists.org/content/146/3/dev175778>.

MacCallum F, Lycett E, Murray C, Jadva V, Golombok S. 2003. Surrogacy: The experience of commissioning couples. *Hum Reprod*. 18(6):1334–1342.
doi:10.1093/humrep/deg253.

Mamo L. 2007. *Queering Reproduction: Achieving Pregnancy in the Age of Technoscience*. Durham, N.C: Duke University Press.

- Mulkay M. 1997. *The Embryo Research Debate: Science and the Politics of Reproduction*. New York: Cambridge University Press.
- Narayan H, Morton A. 2015. Role of the midwife in the care of high-risk pregnancies. In: *Compendium for the Antenatal Care of High-Risk Pregnancies*. Oxford University Press. [accessed 2020 Jun 29].
<https://oxfordmedicine.com/view/10.1093/med/9780199673643.001.0001/med-9780199673643-appendix-0030>.
- NHS. 2017 Dec 21. Newborn Blood Spot Test. nhs.uk. [accessed 2020 Jun 30].
<https://www.nhs.uk/conditions/pregnancy-and-baby/newborn-blood-spot-test/>.
- Ogbuehi O, Powell J. 2015. Parenthood. In: Lewis L, editor. *Fundamentals of Midwifery: A Textbook for Students*. Wiley-Blackwell. p. 90–119.
- Ormond KE, Mortlock DP, Scholes DT, Bombard Y, Brody LC, Faucett WA, Garrison NA, Hercher L, Isasi R, Middleton A, et al. 2017. Human Germline Genome Editing. *The American Journal of Human Genetics*. 101(2):167–176.
doi:10.1016/j.ajhg.2017.06.012.
- Rodrigues PB, Zambaldi CF, Cantilino A, Sougey EB. 2016. Special features of high-risk pregnancies as factors in development of mental distress: a review. *Trends Psychiatry Psychother*. 38(3):136–140. doi:10.1590/2237-6089-2015-0067.
- Rosemann A, Balen A, Nerlich B, Hauskeller C, Sleeboom-Faulkner M, Hartley S, Zhang X, Lee N. 2019. Heritable Genome Editing in a Global Context: National and International Policy Challenges. *Hastings Cent Rep*. 49(3):30–42.
doi:10.1002/hast.1006.
- Royal College of Obstetricians and Gynaecologists. 2011. *Reproductive Ageing. Scientific Impact Report No.: 24*.
https://www.rcog.org.uk/globalassets/documents/guidelines/scientific-impact-papers/sip_24.pdf.
- Sarojini N, Marwah V. 2015. *Reconfiguring Reproduction Feminist Health Perspectives on Assisted Reproductive Technologies*. 1 edition. New Delhi: Zubaan Books.
- Schmidt WC. 1997. World-Wide Web Survey Research: Benefits, Potential Problems, and Solutions. *Behavior Research Methods, Instruments, & Computers*. 29(2):274–279. doi:10.3758/BF03204826.
- Sittner BJ, DeFrain J, Hudson DB. 2005. Effects of high-risk pregnancies on families. *MCN Am J Matern Child Nurs*. 30(2):121–126. doi:10.1097/00005721-200503000-00010.
- Smietana M, Thompson C, Twine FW. 2018. Making and breaking families – reading queer reproductions, stratified reproduction and reproductive justice together. *RBMSociety*. 7:112–130. doi:10.1016/j.rbms.2018.11.001.

Stevenson EL, Hershberger PF, editors. 2016. Contemporary Fertility and Assisted Reproductive Technology (Art): Advancing Global Discussion in Theory, Practice, Policy, and Research. 1st edition. New York: Springer Publishing.

Thompson C. 2007. Making Parents: The Ontological Choreography of Reproductive Technologies. New Ed edition. Cambridge, Mass.: MIT Press.

Thompson C. 2019. How Should “CRISPRed” Babies Be Monitored Over Their Life Course to Promote Health Equity? *AMA Journal of Ethics*. 21(12):1036–1041. doi:10.1001/amajethics.2019.1036.

van de Wiel L. 2015. Frozen in anticipation: Eggs for later. *Women’s Studies International Forum*. 53:119–128. doi:10.1016/j.wsif.2014.10.019.

Wright KB. 2005. Researching Internet-Based Populations: Advantages and Disadvantages of Online Survey Research, Online Questionnaire Authoring Software Packages, and Web Survey Services. *J Comput Mediat Commun*. 10(3). doi:10.1111/j.1083-6101.2005.tb00259.x. [accessed 2020 Feb 23]. <https://academic.oup.com/jcmc/article/10/3/JCMC1034/4614509>.

Yali AM, Lobel M. 1999. Coping and distress in pregnancy: an investigation of medically high risk women. *J Psychosom Obstet Gynaecol*. 20(1):39–52. doi:10.3109/01674829909075575.