

# Effect of exercise on acute senescent lymphocyte counts

Veiga Sardeli, Amanda; Mori, Marcelo Alves; Lord, Janet

DOI:

[10.1159/000520528](https://doi.org/10.1159/000520528)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Veiga Sardeli, A, Mori, MA & Lord, J 2022, 'Effect of exercise on acute senescent lymphocyte counts: a systematic review and meta-analysis', *Gerontology*. <https://doi.org/10.1159/000520528>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

This is the accepted manuscript version of an article published by S. Karger AG in *Gerontology*, 2022, DOI: 10.1159/000520528, available on <https://www.karger.com/Article/FullText/520528>.

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Gerontology

<b>Manuscript:</b>	GER-2021-8-38/R1 RESUBMISSION
<b>Title:</b>	Effect of exercise on acute senescent lymphocyte counts: a systematic review and meta-analysis.
<b>Authors(s):</b>	Amanda Veiga Sardeli (Corresponding Author), Marcelo A. Mori (Co-author), Janet M. Lord (Co-author)
<b>Keywords:</b>	Aging, Cellular senescence, Exercise, Immunology, Immunosenescence
<b>Type:</b>	Meta-Analysis

Dear Monika Lechleitner,

**Re: GER-2021-8-38 - Effect of exercise on acute senescent lymphocyte counts: a systematic review and meta-analysis.**

Thank you for passing on the reviewers comments and for giving us the opportunity to revise the manuscript. We would like to thank the reviewers and section editor for their comments which helped us to improve the quality of our manuscript. We have revised our manuscript with all changes marked in red and our detailed response to each comment is described below.

We look forward to your response.

Yours sincerely

**Section Editor comments:**

*This review and meta-analysis about the role of exercise on acute senescent lymphocyte counts is of clinical interest.*

*According to the statements of the reviewers there remain some minor points of concerns:*

*- the authors have included several immunosenescence markers, however, p16 and p 21 were not included.*

Answer: We did not exclude studies using these classic markers, however, the only two studies testing the effects of exercise on SA- $\beta$ -gal (Wu et al., 2018) and p16<sup>INK4a</sup> (Yang et al., 2018) did so in muscle and endothelial progenitor cells, respectively, which was out of the scope of our analysis. In the immune field the markers of senescence used tend to be different to non-immune cells and focus on the cell membrane phenotype, such as expression of CD57. We now include a comment on this point in the methods section (page 6, L 149).

*- did the authors consider the differences in the mode of exercise ? (such as bicycle versus treadmill)*

Answer: Originally we did include this variable but the results showed no difference between exercise type (treadmill vs cycling) and we decided not to show the data as the number of studies in the different categories were very unbalanced, ranging from 2 to 40 studies. We now mention this finding in the discussion section (page 13, L 347), but do not show the data, we hope this is acceptable.

*- units should be added to the data*

Answer: As stated in the methods (page 9, L 216) “We analysed the absolute cell count as the outcome measure, considering the standardized mean difference (SMD) and 95% of confidence interval between baseline levels and post exercise time-points since the units of measure were not consistent across studies”. This allowed us to overcome the issue of different units being used in different studies and instead bases the analysis on effect size.

- in the introduction section the purpose of the review should be more clearly defined

Answer: We have revised the paragraph to clarify the aim of the study and hope that this is now satisfactory (page 6, L 149).

-----  
Reviewer 1:

*My only criticism is the very generous use of the term senescence. In the context of T cell differentiation, senescence is an ill-defined term and CMV-specific T cells in a 40 yo as included in this study does not necessarily have cellular senescence and it is not clear where effector functions of T cells ends and SASP start. Although transient loss of CD28 in effector T cells is normal in an immune response and not senescent (see for example the studies by Rafi Ahmed). the authors should use the introduction to discuss this issue and give a clear operative definition for the purpose of this paper.*

Answer: This is a good point and the field of immunosenescence does have distinct features from senescence in non-immune cells. We have revised the introduction in order to clarify these issues in more detail (page 4, L 78).

*Also, one of the limitations of the study that should be mentioned is that senescent cells in this study are a mixed bag. CD57 TEMRAs (representing the cell type that is closest to senescence) are not specifically identified in the published papers. Moreover, TEMRAs and CD8 EM that may be negative for CD27 or CD28 are quite different differentiation stages.*

Answer: This is a valid comment, and we were of course aware of this limitation. We have now added a comment on this point to the limitations section in the discussion (page 14, L 379).

*A minor issue is that data are given without units. I suppose that the included papers provide absolute numbers and not percentages and the unit is per ul.*

Answer: This issue was also raised by the section editor and our response is shown above.

-----  
Reviewer 2:

*Authors have included several immunosenescence markers, however, were classic senescence markers like p16, p19, p21 also looked at? For e.g. studies have shown that p16 and p21 expression was higher in CD28+ CD57+ senescent T cell populations.*

Answer: This point was also raised by the section editor and our response is given above.

*Was the mode of aerobic exercise – bicycle vs. treadmill – normalized in anyway? In table 1, Azali Alamdari et. al., and Turner use a treadmill in their study vs. other authors that use a*

*bicycle. Therefore, what was the rationale/parameter for clubbing both these modes under “aerobic exercise” for the meta-analysis?*

Answer: Please see the comment above to the section editor.

*In table 1, how does intensity “until exhaustion” correspond to a VO2max value?*

Answer: The studies classified as “until exhaustion” analysed the frequency of senescent lymphocytes immediately after a maximum test, which could be considered as 100% of effort, thus equivalent to VO2max intensity.

### **References:**

Wu, J., Saovieng, S., Cheng, I. S., Liu, T., Hong, S., Lin, C. Y., Su, I. C., Huang, C. Y., & Kuo, C. H. (2019). Ginsenoside Rg1 supplementation clears senescence-associated beta-galactosidase in exercising human skeletal muscle. *Journal of Ginseng Research*, **43**, 580–588.

Yang, C., Jiao, Y., Wei, B., Yang, Z., Wu, J.-F., Jensen, J., Jean, W.-H., Huang, C.-Y., & Kuo, C.-H. (2018). Aged cells in human skeletal muscle after resistance exercise. *Aging (Albany NY)*, **10**, 1356–1365.

1 ***Meta-Analysis***

2 **Effect of exercise on acute senescent lymphocyte counts: a systematic review and meta-analysis.**

3  
4 Amanda V. Sardeli <sup>a,b\*</sup>, Marcelo A. Mori <sup>c, d, e</sup>, Janet M. Lord <sup>a, f</sup>

5  
6 <sup>a</sup> MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, Institute of Inflammation and  
7 Ageing, University of Birmingham, Birmingham, UK.

8 <sup>b</sup> Laboratory of Exercise Physiology (FISEX), University of Campinas, Campinas, Brazil.

9 <sup>c</sup> Department of Biochemistry and Tissue Biology, Institute of Biology, University of Campinas,  
10 Campinas, Brazil.

11 <sup>d</sup> Experimental Medicine Research Cluster, University of Campinas, Campinas, Brazil.

12 <sup>e</sup> Obesity and Comorbidities Research Center, University of Campinas, Campinas, Brazil.

13 <sup>f</sup> NIHR Birmingham Biomedical Research Centre, University Hospital Birmingham and University of  
14 Birmingham, Birmingham, UK.

15  
16  
17 Short Title: Effect of exercise on acute senescent lymphocyte counts

18  
19 Corresponding Author:

20 Amanda Veiga Sardeli

21 Institute of Inflammation and Ageing

22 Queen Elizabeth Hospital

23 Mindelsohn Way

24 Birmingham, West Midlands, B15 2WB, UK

25 Tel: +44 (0)121 414 3344

26 E-mail: amandavsardeli@gmail.com

27

28 Number of Tables: 2.

29 Number of Figures: 5.

30 Word count: 4832.

31 Keywords: Exercise, Ageing, Cellular senescence, Immunosenescence, Immunology.

32 **Abstract**

33 **Background:** Highly differentiated, senescent lymphocytes are pro-inflammatory and contribute to  
34 age-related systemic inflammation, inflammageing. There are several reports of acute changes in  
35 senescent lymphocyte counts post-exercise which potentially has consequences for systemic  
36 inflammation. However, there is little consensus since the studies differ with respect to participants,  
37 exercise protocols, cellular markers assessed, and the time point of assessment post-exercise.

38 **Objective:** We performed a systematic review and meta-analysis to assess the impact of exercise on  
39 senescent lymphocyte counts in blood immediately, 1h and 2h post exercise.

40 **Methods:** The search was performed in PubMed (MEDLINE), Web of Science, Embase, Scopus and  
41 Cochrane, on January 11, 2021. The 13 studies selected tested aerobic exercise effects, mainly in  
42 young men. They assessed the counts of lymphocytes (CD4 T cells, CD8 T cells, NK cells), with the  
43 following immune cell marker combinations: KLRG1+, CD57+ (only NK cells), EMRA T cells  
44 (CD45RA+CCR7-CD28-CD27-), CD28-CD27-, KLRG1+CD28- and CD28-. Independent extraction of  
45 articles by 2 researchers.

46 **Results:** Standardized mean difference (SMD) and 95% confidence interval between baseline and  
47 post-exercise showed significant increases (SMD > 0.9, p<0.003) in all types of lymphocyte counts  
48 immediately post exercise. At 1h post exercise senescent CD4 T cells returned to baseline values  
49 ( $p=0.74$ ), CD8 T cells were reduced ( $-0.26 [-0.41; -0.11]$ ,  $p=0.001$ ), and senescent NK cells were raised  
50 ( $0.62 [0.14; 1.10]$ ,  $p=0.01$ ) above baseline. By 2 hours post exercise, senescent CD4 T cells were  
51 reduced ( $-0.94 [-1.40; -0.48]$ ,  $p<0.001$ ), CD8 T cells remained below baseline ( $-0.53 [-1.04; -0.009]$ ,  
52  $p=0.04$ ), and NK cells had returned to baseline values ( $-0.29 [-0.64; 0.07]$ ,  $p=0.11$ ). The main  
53 determinants of heterogeneity between studies were cytomegalovirus (CMV) serostatus and the  
54 characteristics of exercise protocols. CMV+ individuals had a higher immediate lymphocytosis and 1h  
55 post lymphopenia than CMV- individuals. Exercise performed at higher intensities and shorter  
56 durations led to higher magnitude of change in senescent lymphocyte counts at all time-points.

57 **Conclusion:** The differing effects of exercise on senescent NK cells and CD4 and CD8 T cells suggest  
58 differing susceptibility to factors modulating lymphocyte extravasation such as adrenaline and  
59 exercise intensity.

60

61



## 62 **Introduction**

63 Immunosenescence, the gradual remodelling of the immune system, is an integral component of the  
64 ageing process [1]. Advanced age impairs innate immune responses, contributes to chronic low-  
65 grade inflammation (inflammageing) and reduces immunity, increasing the risk of infections,  
66 autoimmunity and overall poor health in the older adult [2,3]. Among the features associated with  
67 adaptive immunosenescence are the atrophy of the thymus, which reduces naïve T cell output, and  
68 the subsequent increased number of highly differentiated, senescent T cells in the circulation [3,4].  
69 Senescent cells are one of the causes of detrimental effects to the body during ageing, contributing  
70 to chronic diseases, such as idiopathic pulmonary fibrosis, diabetes, and osteoarthritis [5]. It has been  
71 shown recently that mice with high levels of senescent T cells, due to dysfunctional mitochondria,  
72 enter premature senescence and a broad range of age-related diseases [6]. Immunosenescence,  
73 especially T cell senescence, may therefore be a major contributor to the ageing process.

74           Senescent cells undergo a state of cell quiescence with permanent cell cycle arrest induced  
75 by different sources of stress and damage to the cell. These cells produce a senescence-associated  
76 secretory phenotype (SASP), which is composed of pro-inflammatory cytokines, chemokines, growth  
77 factors and proteases. Cells releasing SASP alter the tissue microenvironment, affect neighboring  
78 cells, and are thus deleterious [7,8]. **In the immune system there are some subtle differences. For  
79 example, T cells can have a functionally exhausted phenotype resulting from chronic stimulation,  
80 which is distinct from a senescent phenotype resulting from ageing or chronic infection. These  
81 phenotypes can be differentiated by cell surface markers [9]. We have therefore used the markers  
82 identified as relating to senescent T cells such as loss of CD28 and CD27 and expression of KLRG1 and  
83 CD57. Importantly, senescent T cells also produce a SASP that is highly pro-inflammatory and similar  
84 in content to that of non-immune senescent cells [10], therefore they are likely to contribute to  
85 inflammageing and tissue compromise during ageing.**

86           The immunomodulatory effects of exercise have been widely explored and could be  
87 associated with the reduction in senescent cell counts [10], for example obese mice provided with an  
88 exercise wheel had reduced numbers of senescent cells in their adipose tissue [11]. Exercise has been  
89 reported to have a range of immune enhancing effects including reducing chronic low-grade  
90 inflammation [12], improving responses to vaccination [13], reducing the risk of infection [14,15],  
91 improving the immune response against viruses and bacteria and reducing the burden of latent viral  
92 infections [16–18]. Among the main physiological mechanisms mediating the immunomodulatory

93 benefits of exercise are the reduction in body fat and the release of anti-inflammatory cytokines,  
94 such as interleukin-6 (IL-6) and IL-1RA, by the exercising muscle [12,19].

95           Recently, Duggal *et al.* [20] have reported the benefits of sustained physical activity in to old  
96 age on adaptive immune phenotype and immunosenescence. They reported that thymic health, as  
97 measured by the frequency of naïve T cells and recent thymic emigrants (RTE), was better preserved  
98 in older exercisers (cyclists) compared to inactive elders. Older cyclists also had significantly higher  
99 serum levels of the thymoprotective cytokine interleukin-7 (IL-7), higher B regulatory cell frequency,  
100 lower IL-6 and reduced Th17 polarization, all markers of an aged immune system. However, they also  
101 reported that the age-related increase in senescent T cells was not prevented in the cyclists [20].

102           Despite the chronic benefits of exercise being well established, whether acute exercise  
103 increases susceptibility to infection or confers immune protection is still a matter of debate [21].  
104 However, an increase in lymphocyte counts in the blood (lymphocytosis), followed by a decrease  
105 (lymphocytopenia) post exercise has generally been reported [22]. Lymphocytes are proposed to  
106 migrate from the marginal pool, the spleen and lymph nodes in to the blood, as well as increased  
107 release from the bone marrow to produce the lymphocytosis. This migration is mediated by exercise-  
108 induced shear stress on blood vessels and catecholamines, as well as cortisol and to some extent  
109 cytokines such as IL-6 [23–25].

110           What is less clear is the impact of exercise on specific immune cell types and their  
111 differentiation state, notably senescent immune cells. This is important bearing in mind their pro-  
112 inflammatory nature and potential role in driving inflammaging and the aged phenotype [26].  
113 Studies investigating senescent lymphocyte counts in circulation post-exercise have shown a variety  
114 of responses [27–31], including a reduction on leukocyte counts [32]. However, senescent, or highly  
115 differentiated lymphocytes appear to be more likely to increase in blood with exercise than  
116 lymphocytes in earlier stages of differentiation. This could be beneficial in leading to their  
117 subsequent removal by NK cells or CD8 T cells which can detect senescent cells and kill them by  
118 apoptosis [22,33].

119           Another important confounding factor in the various exercise intervention studies, is  
120 infection by cytomegalovirus (CMV) that increases with age and has deleterious effects on  
121 lymphocyte immunity, accelerating immunosenescence [22,34,35]. The higher baseline cell counts of  
122 senescent lymphocytes in CMV+ individuals could lead to higher magnitude of change in these  
123 individuals after exercise, and thus the CMV serostatus might be an important confounding factor  
124 between studies [27,28]. Other factors that may cause different results between studies are: the

125 comparison between absolute cell counts and the frequency of cells in the circulation; the different  
126 types of lymphocytes assessed; the membrane markers used to identify cell senescence;  
127 characteristics of the study population (age, sex and physical activity level) and the exercise protocols  
128 used (type of exercise, volume and intensity).

129 To derive a consensus from the literature it is important to isolate the variety of confounding  
130 factors among the studies and to run a pooled effects meta-analysis. Thus, we aimed to identify the  
131 impact of acute exercise on the frequency of senescent T cells and NK cells, taking in to account  
132 variables such as CMV serostatus, age, training status and specifics of the exercise protocols.

## 133 **Methods**

134 This systematic review and meta-analysis was registered on PROSPERO under the number  
135 CRD42021267078, that can be assessed at <https://www.crd.york.ac.uk/prospero/>, and it was  
136 reported in accordance with the recommendations of Preferred Reporting Items for Systematic  
137 Reviews and Meta Analyses (PRISMA) statement [36].

### 138 *Search strategy*

139 On January 11, 2021 the search was updated at PubMed (Medline), Web of Science, Embase, Scopus  
140 and Cochrane. It combined the synonyms of “senescent markers” and “exercise” according to each  
141 data base descriptor and field of search as detailed in the Supplementary material.

### 142 *Eligibility criteria*

143 Figure 1 shows the study selection process, completed by two independent reviewers. We included  
144 studies: (1) of acute interventional exercise; (2) with no associated intervention, i.e. exercise only  
145 group; (3) in humans from both sexes; (4) comparing resting and immediately, 1h and 2h post  
146 exercise condition; (5) assessing any bona fide markers of immunosenescence; (6) assessing CD4+ or  
147 CD8+ T cells, or NK lymphocytes; (7) written in English.

### 148 *Immunosenescence cell markers*

149 Markers of senescence traditionally used for non-immune cells, such as p16<sup>ink4a</sup> and SA-βGal have not  
150 been used in studies of immunosenescence which focus on cell membrane markers. We therefore  
151 selected several broadly accepted markers of immunosenescence to use in this study and the  
152 characteristics of each of them are described below.

153 **CD57<sup>+</sup>**. CD57<sup>+</sup> NK cells have been attributed a senescent-like phenotype due to their short telomeres  
154 and inability to proliferate [1,37,38].

155 **CD28<sup>-</sup> CD27<sup>-</sup>**. CD27 and CD28 are costimulatory receptors and T cells lacking CD27 and CD28 are  
156 thought to be fully differentiated T cells exhibiting shorter telomere length [39]. When the  
157 expression of CD27 and CD28 is lost, there is no evidence of subsequent re-expression and the  
158 downregulation of these molecules are linked to dysfunctional T cells with a SASP secretome [40,41].

159 **KLRG1<sup>+</sup>**. T-lymphocytes expressing KLRG1 have impaired capacity to proliferate, yet  
160 maintain immediate effector cell capabilities such as the recognition and killing of target cells [42].

161 **EMRA (CD45RA<sup>+</sup>CCR7<sup>-</sup>CD28<sup>-</sup>CD27<sup>-</sup>)**. EMRA, for terminally differentiated effector memory cells re-  
162 expressing CD45RA, have the key features of cell senescence, with low proliferation response and a  
163 highly inflammatory phenotype [10]. They also have high levels of DNA damage and loss of  
164 telomerase activity [43]. However, due to their ability to proliferate under specific conditions their  
165 phenotype is distinct from non-immune senescent cells which cannot proliferate [44].

#### 166 *Exclusion criteria*

167 We excluded studies that: (1) did not have original data or did not undergo peer-review such as  
168 reviews, commentaries, editorials, letter to the editors, case reports or conference abstracts; (2)  
169 assessed other senescence markers such as telomere shortening, or telomerase activity; (3) had not  
170 tested exercise effects; (4) had not assessed immunosenescence in humans; (5) assessed senescence  
171 in other cells, besides lymphocytes and NK cells; (6) were not written in English.

#### 172 **Data collection and data items**

173 Data collection was performed by two independent researchers. The means and a measure of  
174 dispersion of the senescent cell counts were extracted for each subgroup within studies. Mean,  
175 standard deviation (SD) and sample number (n) were used for the meta-analyses. Standard error (SE)  
176 was converted to SD by the equation  $SD = SE \times (\sqrt{n})$ , if SD was not provided in the original study.

177 For subgroup analysis we extracted information about participants sex, age, level of training,  
178 health condition and CMV serostatus, type of lymphocytes assessed, membrane markers used, unit  
179 of measurement, and the characteristics of the exercise bout such as intensity, volume, and type of  
180 equipment.

181 The sample of studies was classified as young, middle aged and old according to the mean  
182 age reported (young [ $<30$ yrs], middle aged [30-40yrs] and older adults [ $>50$ yrs]).

183 The participants were considered “trained” when the studies classified them as elite athletes,  
184 trained, physically active, cyclists or when the  $VO_2$ max was above the 50% percentile according to  
185 their age [45]; they were considered “untrained” when the studies classified them as untrained, or  
186 doing no regular physical activity or sedentary. The studies that did not report the participant’s  
187 physical activity level or reported a too wide range of physical activity level among their participants  
188 were excluded from this subgroup analysis. Individuals undergoing exercise chronic intervention  
189 were considered trained [32,46]; while the individuals undergoing non-exercise intervention in Wang  
190 et. al. [32] were classified as untrained and the individuals undergoing non-exercise intervention in  
191 Azali Alamdari et al. [46] were excluded for training status analysis, since they were athletes at  
192 baseline.

193 Regarding health status, only Curran et al. [47] have included individuals with type I diabetes,  
194 while the other studies only included healthy participants.

195 The exercise intensity was classified according to the percentage of  $VO_2$ max described by the  
196 American college of Sports Medicine [48], in which 46-63% is moderate, 64-90% is vigorous and  $>91\%$   
197 is near maximum. The intensity reported on Ingram et al. study [49], in watts was estimated as 73.7%  
198 of maximum according to data from participants of a similar age. Another study tested different  
199 protocols according to their lactate threshold (5% under LT, 5% above LT and 15% above LT) in the  
200 same individuals and each of them were included in the meta-analysis as a separate study [27].The  
201 intensity was also converted to percentage according to Farina *et al.* [50], in which 5%  $<LT$  was  
202 considered 61.1%, 5% $>LT$  was considered 71.1%, and 15%  $>LT$  was considered 81.1%. The studies  
203 reporting percentage of estimated maximum power or percentage of ventilatory threshold work  
204 rate, instead of  $VO_2$ max, were classified for subgroup analysis as these markers were proportionally  
205 equivalent.

206 The studies applying incremental maximum effort tests and other protocols expected to last  
207 less than 20 min were considered short, the ones applying 30 min duration were considered  
208 moderate and above this they were considered long duration.

209 Only Azali Alamdari et al.[46] had a control group, and thus, the change of control group was  
210 subtracted from the exercise change to increase the robustness of the analysis. Although Turner et  
211 al. [51] had also reported a control group, they did not present the effects of the control period on

212 CD28-CD27- markers, in this way the control group was not considered for analysis. Two studies  
213 presented acute exercise effects before and after a variety of chronic interventions [32,46] and thus  
214 we included only their post intervention session to avoid sample overlapping in the analysis.

## 215 **Statistical analysis**

216 We analysed the absolute cell count as the outcome measure, considering the standardized mean  
217 difference (SMD) and 95% of confidence interval between baseline levels and post exercise time-  
218 points since the units of measure were not consistent across studies.

219 The 3 main meta-analyses, for each time point (immediately, 1h and 2h post exercise) and  
220 the subgroup analyses were performed using Comprehensive Meta-Analysis software, version  
221 3.3.070. When there was statistical significance for heterogeneity, randomized effect models were  
222 selected and when there was no significant heterogeneity, fixed effects were applied. The  
223 inconsistency between studies was reported as a percentage ( $I^2$ ), based on difference between  
224 expected heterogeneity (df) and true heterogeneity (Q-value).

225 For subgroup analysis we tested the influence of the following confounding factors: sex (men  
226 and women); age (young [ $<30$  yrs], middle aged [ $30-40$  yrs] and older adults [ $>50$  yrs]); type of  
227 lymphocytes (CD4+, CD8+ and NK); type of senescence marker (KLRG1+, CD57+, EMRA  
228 [CD45RA+CCR7-CD28-CD27-], CD28-CD27-, KLRG1+CD28- and CD28-), level of training (trained and  
229 untrained); health condition (healthy and diseased); CMV serostatus (CMV+ and CMV-); exercise  
230 intensity (moderate, vigorous, near maximum); and exercise volume (short, moderate and long). Q  
231 tests were applied to group comparisons, considering 95% confidence.

232 Egger's tests were performed to check the risk of publication bias in each meta-analysis [52].

## 233 **Results**

234 We included thirteen studies [27–29,32,46,47,49,51,53–57] testing acute aerobic exercise effects on  
235 senescent T lymphocytes and NK cell counts (shown in Figure 1). It is noteworthy that some studies  
236 had to be excluded due to the absence of specific description of absolute senescent lymphocyte  
237 counts [30,58–64]. Most studies included, reported their results among different subgroups of  
238 individuals with different sex, ages, CMV serostatus, types of exercise protocols and time points of  
239 analysis that were analyzed as a sub-study.

240 **\*\*\*please insert Figure 1 here\*\*\***

241 *Study characteristics*

242 Table 1 shows the characteristics of the studies included. Only Curran et al.[47] included a type I  
243 diabetes group, while the other studies only included healthy participants. While twelve studies  
244 tested exercise effects on males, just one tested exercise effects on participants from both sexes  
245 [27], and thus comparisons between men and women were not possible in subgroup analysis. One  
246 study included middle aged [28], two included older adults [28,29] and all of them (thirteen) tested  
247 young adults. Our analysis reported the effect of exercise on T CD4+, T CD8+ and NK cell counts. All  
248 studies tested the effects of aerobic exercise, the majority of them used bicycle, and a few used  
249 treadmill [46,51].

250 **\*\*\*please insert Table 1 here\*\*\***

251 *Syntheses of the results*

252 **Lymphocyte counts immediately post exercise.** Figure 2 shows there were significant increases on  
253 senescent CD4 T cells (SMD 0.96 [0.67; 1.25],  $p<0.001$ ), CD8 T cells (SMD 1.26 [0.93; 1.59],  $p<0.001$ )  
254 and NK cells counts immediately post exercise (SMD 1.04 [0.35; 1.72],  $p=0.003$ ). However, all those  
255 analyses were heterogeneous, reinforcing the need for further subgroup analyses. Furthermore, the  
256 analysis of senescent CD4 T cells and CD8 T cell counts had significant risk of bias, evidencing that  
257 studies with low precision conducted the main effects.

258 Table 2 shows no effect of age ( $p= 0.46$ ) or training status ( $p=0.35$ ) on outcomes immediately  
259 post exercise. On the other hand, the intensity and duration of exercise protocols and CMV status  
260 influenced the post exercise senescent lymphocyte counts (Table 2). Specifically, the higher  
261 magnitude of increase in senescent lymphocytes were seen in the maximum intensity and short  
262 duration protocols (SMD 1.81 [1.45; 2.1],  $p<0.001$ ) compared to the others (SMD  $<0.85$ ,  $p<0.05$ ).  
263 There was a trend to higher senescent lymphocyte counts in CMV positive participants compared to  
264 CMV- ( $p=0.09$ ). The CMV status analysis for each subgroup of T lymphocyte showed higher increase  
265 in senescent CD8+ T cells for CMV+ (SMD 1.60 [ 0.73; 2.46],  $p<0.001$ ) compared to CMV- (SMD 0.58  
266 [0.33; 0.83],  $p<0.001$ ), with no difference for senescent CD4+ T cells regarding CMV status (SMD  
267 CMV+: 0.42 [0.02; 0.82],  $p=0.038$  and CMV-: 0.50 [0.18; 0.82],  $p=0.002$ ).

268 **\*Please, insert Figure 2 here\***

269 **Lymphocyte counts one hour post exercise.** Figure 3 shows senescent CD8+ T cell counts were lower  
270 compared to baseline levels (SMD -0.28 [-0.44; -0.13],  $p<0.001$ ), while CD4+ T cell counts returned to  
271 baseline levels (SMD -0.13 [-0.37; 0.11],  $p=0.28$ ) and NK cells were still above baseline values (SMD

272 0.62 [0.14; 1.09],  $p=0.11$ ). These analyses were homogeneous ( $p>0.53$ ,  $I^2=0\%$ ), confirming that each  
273 of these senescent cells have very consistent response 1h post exercise across the different studies.

274 Table 2 shows there was a significant reduction in the senescent lymphocyte count only in  
275 CMV+ and not CMV- individuals, with significant difference between groups. Regarding each  
276 subgroup of T lymphocyte there was no significant reduction for senescent CD4+ T or CD8+ cells in  
277 CMV- (SMD CD4+: 0 [-0.30; 0.31],  $p=0.97$  and CD8+: -0.13 [-0.36; 0.09],  $p=0.25$ ) while there was a  
278 trend of senescent CD4+ reduction in CMV+ individuals (SMD -0.35 [-0.74; 0.04],  $p=0.075$ ), and  
279 reduction of senescent CD8+ T cells in CMV+ (SMD -0.46 [-0.75; -0.18]  $p=0.001$ ). Only vigorous  
280 intensity and long duration exercise protocols led to significant reduction of senescent lymphocytes  
281 (SMD -0.5 [-0.8; -0.2],  $p<0.001$ ) while the other intensities and durations did not vary significantly  
282 ( $p>0.16$ ).

283 **\*Please, insert Figure 3 here\***

284 **Lymphocyte counts two hours post exercise.** Figure 4 shows senescent CD4 T cells were reduced  
285 (SMD -0.94 [-1.40; -0.48],  $p<0.001$ ), CD8 T cells remained below baseline (SMD -0.53 [-1.04; -0.009],  
286  $p=0.04$ ), and NK cells had returned to baseline values (SMD -0.29 [-0.64; 0.07],  $p=0.11$ ). There was  
287 significant risk of publication bias for the analysis of senescent CD4 T cells (Egger test  $p$ -value  $<0.001$ ),  
288 evidencing that studies with low precision conducted the main effects in this analysis.

289 All these three meta-analyses were heterogeneous, however, due to the low number of  
290 subgroups in these analyses, only training status, intensity and volume of exercise protocols were  
291 analyzed. No difference between trained and untrained individuals was noticed ( $p=0.81$ ) and only  
292 maximum intensity and short duration protocols reduced senescent cell counts (SMD -0.7 [-1; -0.4],  
293  $p<0.001$ ), however, it is noteworthy there was very low number of studies in the other categories  
294 (Table 2).

295 **\*Please, insert figure 4 here\***

## 296 **Discussion**

297 The main findings of the present meta-analysis were the significant increase in senescent CD8+, CD4+  
298 and NK cell counts immediately post exercise followed by a reduction in senescent CD8+ T cells at 1h  
299 and 2h post exercise, a reduction in senescent CD4+ T cells at 2h post exercise and maintenance of  
300 increased NK senescent cells at 1h post exercise with a return to baseline at 2h post exercise (Figure  
301 5). Although there is no consensus about the exact role of these redistributions of senescent  
302 lymphocytes post exercise, it has been proposed that senescent lymphocytes are preferentially



303 recruited for immune surveillance and removal by NK and CD8+ T cells, resulting in an exercise-  
304 induced senolytic effect [22].

305 In fact, it is known that T-cells with high cytotoxic capabilities and tissue migration potential,  
306 which are characteristics of highly differentiated lymphocytes, are preferentially mobilised by acute  
307 stress and exercise [65]. These lymphocytes could be recruited due to their high  $\beta_2$ -adrenergic  
308 receptor expression [66] even though they have impaired replicability and co-stimulatory potential.  
309 Furthermore, in mice, NK cells are the main mediator of the antitumor effects of exercise. These  
310 effects depend on the mobilisation of these cells [67], which are the most responsive lymphocyte  
311 subset to acute exercise due to their high  $\beta$ -adrenergic receptor expression [68]. Mobilisation of the  
312 senescent, less functional form of these cells could be beneficial if they are then removed, improving  
313 the overall quality of the lymphocyte pool.

314 Following their mobilisation it is possible that T-lymphocytes egressing to the peripheral  
315 tissues may experience a pro-apoptotic environment [69], as Kruger et. al. [70] showed the number  
316 of highly differentiated CD3+ T cells remained reduced 3h and 24h post exercise. Another possibility  
317 could be the return of senescent cells to lymph nodes but most of these cells lack CCR7, a secondary  
318 lymphoid organ-homing marker, this is unlikely.

319 In theory, when senescent T-lymphocytes undergo apoptosis, a subsequent feedback loop  
320 could increase the output of naïve T-lymphocytes from the thymus, restoring the peripheral T-  
321 lymphocyte pool [22,59]. In fact, naïve lymphocytes counts are increasing 1h post exercise [30,59].  
322 Furthermore, older adults involved in regular exercise have higher serum levels of the  
323 thymoprotective IL-7 and higher frequency of RTE than sedentary controls [20], which could be  
324 stimulated by senescent lymphocyte clearance post each exercise bout.

325 In an opposite way, exercise-induced hematopoiesis [25,71], could also affect the thymic  
326 feedback loop, increasing the stimuli for senescent lymphocyte removal. Cross-sectional studies  
327 showed physically active individuals have lower markers of senescent T lymphocytes [10,20] and  
328 master athletes have longer lymphocyte telomere length than untrained controls [72]. Nevertheless,  
329 it is noteworthy that highly differentiated, senescent lymphocytes are less sensitive to apoptotic  
330 signals [73,74], and the exercise effects on senescent cell apoptosis is still unknown.

331 Most studies tested the influence of CMV serostatus on exercise responses. CMV reactivation  
332 can be triggered through catecholamine-responsive elements [75] and stress hormone levels, which  
333 are known to correlate strongly with CMV reactivation in astronauts before and after spaceflight

334 [76]. Thus, it is believed that CMV<sup>+</sup> individuals have reduced sensitivity to  $\beta$ -adrenergic stimulation  
335 and decreased  $\beta_2$ -adrenergic receptor expression to prevent CMV reactivation [58]. However, a  
336 reduced  $\beta$ -adrenergic sensitivity of T cells in CMV<sup>+</sup> individuals is not supported by our analysis, and in  
337 fact we saw a larger magnitude of changes in CMV<sup>+</sup> individuals. Thus, we believe the expected higher  
338 baseline senescent lymphocyte counts in CMV<sup>+</sup> individuals [28,77], especially for senescent CD8<sup>+</sup> T  
339 cells, explains the higher magnitude of change with exercise in this population.

340         There was a greater magnitude of increase in senescent lymphocyte counts immediately post  
341 maximum intensity and short duration protocols and greater magnitude of reduction 2h post  
342 exercise compared to lower intensities and longer duration protocols. These differences could be  
343 explained by higher sympathetic activation and sustained release of epinephrine reported in higher  
344 intensities protocols [70,78,79]. However, there is also evidence that cortisol affects lymphocyte  
345 counts during exercise [70,80–83]. Exercise of high intensity leads to greater release of cortisol in to  
346 the blood and for a longer time and may explain the reduced cell counts at later time points since  
347 cortisol induces apoptosis in lymphocytes [84]. *We also considered the type of exercise and whether  
348 this may make a difference to the senescent cell response. However, we found no difference at any  
349 of the time points between treadmill and cycling protocols (data not shown), though this is with the  
350 caveat that the number of studies per subgroup category varied greatly.*

351         It is unlikely that IL-6 released by muscle cells during exercise [85], explains the difference  
352 between exercise protocols. It is known, that IL-6 attracts lymphocytes to the circulation together  
353 with  $\beta$ -adrenergic signaling during exercise [67], however, there is a higher release of IL-6 within  
354 exercise protocols with higher energetic demand, such as the higher volume and during regimens  
355 [85–87] which do not agree with our findings.

356         Finally, exercise hypoxia may explain at least part of the changes in T lymphocytes and NK  
357 counts with exercise, possibly mediated by the same neuroendocrinological factors released by other  
358 stress conditions (i.e.: catecholamines, cortisol) [57,88].

### 359 *Limitations*

360         The first limitation of this study was that most studies included young individuals. The unbalanced  
361 subgroup analysis suggested there is higher magnitude of change on senescent counts in young than  
362 older or middle-aged individuals immediately and 1h post exercise. Whether it is a true effect is  
363 unclear, it could be explained by reduced  $\beta_2$ -adrenergic receptor sensitivity with ageing [89], which in

364 turn increases the threshold for catecholamine-induced lymphocyte recruitment. In this way, it is  
365 important to confirm these results with a larger sample of older adults.

366 Comparisons between men and women were also not possible due to the lack of studies in  
367 women. An exploratory analysis showed immediately post exercise there was a large ( $p < 0.001$ )  
368 increase of senescent lymphocytes in men (1.23 [0.99; 1.47],  $p < 0.001$ ,  $k = 44$ ) compared to studies  
369 with mixed sex samples (0.48 [0.19; 0.78],  $p < 0.001$ ,  $k = 6$ ), while there was no difference between  
370 these subgroups 1h post. Future studies should test to what extent the results in men are also  
371 applied to women.

372 Another limitation was the lack of a control group, i.e. without exercise, within the original  
373 studies which precluded a proper risk of bias assessment. In the other hand, the comparison of the  
374 same participants along time removes the between subjects' effects, which in turn contributes to the  
375 isolation of exercise effects. Furthermore, we explored possible influences of the confounding factors  
376 in subgroup analysis. At last, it is noteworthy that only two studies investigated exercise effects on  
377 senescent NK cell counts, with is a limitation of the literature and future studies should fill these gaps  
378 to strengthen knowledge in the field.

379 Lastly, one additional issue was the use of markers to identify the different stages of T cell  
380 differentiation and their relation to T cell senescence. Thus, no studies enumerated CD57 TEMRA  
381 cells, the ones that are closest to a senescent phenotype.

### 382 *Conclusions*

383 Senescent lymphocyte counts change significantly in the acute response to aerobic exercise.  
384 However, a complex picture has emerged where senescent CD8+ cells had a higher immediate  
385 lymphocytosis and subsequent lymphopenia (1h and 2h post), senescent CD4+ T cells followed a  
386 similar profile but with lower magnitude of change, and senescent NK cells increased but had a  
387 delayed return to baseline levels. There was higher magnitude of lymphocytosis and  
388 lymphocytopenia for CMV+ individuals and near maximum intensity and short duration protocols.  
389 The differing effects of exercise on senescent NK cells and CD4+ and CD8+ T cells suggest differing  
390 susceptibility to factors modulating lymphocyte extravasation such as adrenaline that is also  
391 regulated by exercise intensity. More studies are needed for understanding exercise effects on  
392 senescent NK cells, in older adults and in women.

393

394 **Statements**

395 **Acknowledgement**

396 The authors thanks Diego Nacarato Pereira da Silva for the contribution in the selection of studies  
397 and data acquisition.

398 **Statement of Ethics**

399 An ethics statement is not applicable because this study is based exclusively on published literature.

400 **Conflict of Interest Statement**

401 The authors have no conflicts of interest to declare.

402 **Funding Sources**

403 This work was supported by the Newton International Fellowship generously awarded to Amanda  
404 Veiga Sardeli by the Academy of Medical Sciences through the UK Government's Newton Fund  
405 Programme [NIFR7\1031]. JML is supported by the NIHR Birmingham Biomedical Research Centre.  
406 The views expressed here are those of the authors and not necessarily those of the NIHR or the  
407 Department of Health and Social Care.

408 **Author Contributions**

409 All three authors have given substantial contributions to the conception and the design of the  
410 manuscript; AVS did the studies selection, data collection and analysis. AVS, MAM and JML  
411 interpreted the data. AVS did the first draft while MAM and JML reviewed it critically for important  
412 intellectual content. All authors read and approved the final version of the manuscript.

413 **Data Availability Statement**

The data in this study was obtained from the previous studies where specific restrictions for public sharing their data may apply according to each journal politics. Such dataset may be requested by the corresponding author e-mail.

## References

- 414 1. Alpert A, Pickman Y, Leipold M, Rosenberg-Hasson Y, Ji X, Gaujoux R, et al. A clinically meaningful  
415 metric of immune age derived from high-dimensional longitudinal monitoring. *Nat Med*. Nature  
416 Publishing Group; 2019;25:487–95.
- 417 2. Franceschi C, Campisi J. Chronic inflammation (Inflammaging) and its potential contribution to age-  
418 associated diseases [Internet]. *Journals Gerontol - Ser A Biol Sci Med Sci*. Oxford University Press;  
419 2014. p. S4–9.
- 420 3. Hazeldine J, Lord J, Hampson P. Immunosenescence and inflammaging: A contributory factor in the  
421 poor outcome of the geriatric trauma patient. *Ageing Res Rev*. *Ageing Res Rev*; 2015;24:349–57.
- 422 4. Majumdar S, Nandi D. Thymic Atrophy: Experimental Studies and Therapeutic Interventions  
423 [Internet]. *Scand J Immunol*. Blackwell Publishing Ltd; 2018. p. 4–14.
- 424 5. Muñoz-Espín D, Serrano M. Cellular senescence: from physiology to pathology. *Nat Publ Gr*. 2014;
- 425 6. Vasileiou PVS, Evangelou K, Vlasis K, Fildis G, Panayiotidis MI, Chronopoulos E, et al.  
426 Mitochondrial Homeostasis and Cellular Senescence. *Cells*. Multidisciplinary Digital Publishing  
427 Institute (MDPI); 2019;8:686.
- 428 7. Coppé JP, Patil CK, Rodier F, Sun Y, Muñoz DP, Goldstein J, et al. Senescence-associated secretory  
429 phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor.  
430 *PLoS Biol*. *PLoS Biol*; 2008;6.
- 431 8. Campisi J. Aging, cellular senescence, and cancer [Internet]. *Annu Rev Physiol*. *Annu Rev Physiol*;  
432 2013. p. 685–705.
- 433 9. Huff W, Kwon J, Henriquez M, Fetcko K, Dey M. The Evolving Role of CD8 + CD28 -  
434 Immunosenescent T Cells in Cancer Immunology. *Int J Mol Sci*. *Int J Mol Sci*; 2019;20.
- 435 10. Chen X, Yi Z, Wong GT, Hasan KMM, Kwan JS, Ma AC, et al. Is exercise a senolytic medicine? A  
436 systematic review. *Ageing Cell*. Blackwell Publishing Ltd; 2020;
- 437 11. Schafer MJ, White TA, Iijima K, Haak AJ, Ligresti G, Atkinson EJ, et al. Cellular senescence  
438 mediates fibrotic pulmonary disease. *Nat Commun*. Nature Publishing Group; 2017;8:14532.
- 439 12. Gleeson M, Bishop N, Walsh N. *Exercise Immunology*. New York: Routledge/Taylor & Francis  
440 Group; 2013.

- 441 13. Pascoe AR, Fiatarone Singh MA, Edwards KM. The effects of exercise on vaccination responses: A  
442 review of chronic and acute exercise interventions in humans [Internet]. *Brain Behav Immun*.  
443 Academic Press Inc.; 2014. p. 33–41.
- 444 14. Pape K, Ryttergaard L, Rotevatn TA, Nielsen BJ, Torp-Pedersen C, Overgaard C, et al. Leisure-Time  
445 Physical Activity and the Risk of Suspected Bacterial Infections. *Med Sci Sports Exerc*. Lippincott  
446 Williams and Wilkins; 2016;48:1737–44.
- 447 15. Nieman DC, Wentz LM. The compelling link between physical activity and the body’s defense  
448 system [Internet]. *J Sport Heal Sci*. Elsevier B.V.; 2019. p. 201–17.
- 449 16. Simpson RJ, Hussain M, Baker F, Bigley AB, Peek MK, Stowe RP. Cardiorespiratory fitness is  
450 associated with better control of latent herpesvirus infections in a large ethnically diverse community  
451 sample: Evidence from the Texas City Stress and Health Study. *Brain Behav Immun* [Internet]. Elsevier  
452 BV; 2017;66:e35.
- 453 17. Agha NH, Mehta SK, Rooney B V., Laughlin MS, Markofski MM, Pierson DL, et al. Exercise as a  
454 countermeasure for latent viral reactivation during long duration space flight. *FASEB J*. John Wiley  
455 and Sons Inc.; 2020;34:2869–81.
- 456 18. Grande AJ, Keogh J, Silva V, Scott AM. Exercise versus no exercise for the occurrence, severity,  
457 and duration of acute respiratory infections [Internet]. *Cochrane Database Syst Rev*. John Wiley and  
458 Sons Ltd; 2020.
- 459 19. Duggal NA, Niemi G, Harridge SDR, Simpson RJ, Lord JM. Can physical activity ameliorate  
460 immunosenescence and thereby reduce age-related multi-morbidity? [Internet]. *Nat Rev Immunol*.  
461 Nature Publishing Group; 2019. p. 563–72.
- 462 20. Duggal NA, Pollock RD, Lazarus NR, Harridge S, Lord JM. Major features of immunesenescence,  
463 including reduced thymic output, are ameliorated by high levels of physical activity in adulthood.  
464 *Aging Cell* [Internet]. Blackwell Publishing Ltd; 2018;17.
- 465 21. Simpson RJ, Campbell JP, Gleeson M, Krüger K, Nieman DC, Pyne DB, et al. Can exercise affect  
466 immune function to increase susceptibility to infection? *Exerc Immunol Rev* [Internet]. 2020;26:8–22.
- 467 22. Simpson RJ. Aging, persistent viral infections, and immunosenescence: Can exercise “make  
468 space”? *Exerc Sport Sci Rev*. *Exerc Sport Sci Rev*; 2011;39:23–33.
- 469 23. Agha NH, Baker FL, Kunz HE, Graff R, Azadan R, Dolan C, et al. Vigorous exercise mobilizes CD34+

470 hematopoietic stem cells to peripheral blood via the  $\beta$  2 -adrenergic receptor. *Brain Behav Immun.*  
471 Academic Press Inc.; 2018;68:66–75.

472 24. Pedersen BK. Physical activity and muscle–brain crosstalk [Internet]. *Nat Rev Endocrinol.* Nature  
473 Publishing Group; 2019. p. 383–92.

474 25. McCarthy DA, Macdonald I, Grant M, Marbut M, Watling M, Nicholson S, et al. Studies on the  
475 immediate and delayed leucocytosis elicited by brief (30-min) strenuous exercise. *Eur J Appl Physiol*  
476 *Occup Physiol.* Springer-Verlag; 1992;64:513–7.

477 26. Desdín-Micó G, Soto-Heredero G, Aranda JF, Oller J, Carrasco E, Gabandé-Rodríguez E, et al. T  
478 cells with dysfunctional mitochondria induce multimorbidity and premature senescence. *Science (80-*  
479 *).* American Association for the Advancement of Science; 2020;368:1371–6.

480 27. LaVoy EC, Hussain M, Reed J, Kunz H, Pistillo M, Bigley AB, et al. T-cell redeployment and  
481 intracellular cytokine expression following exercise: effects of exercise intensity and cytomegalovirus  
482 infection. *Physiol Rep.* American Physiological Society; 2017;5.

483 28. Spielmann G, Bollard CM, Bigley AB, Hanley PJ, Blaney JW, LaVoy ECP, et al. The effects of age and  
484 latent cytomegalovirus infection on the redeployment of CD8+ T cell subsets in response to acute  
485 exercise in humans. *Brain Behav Immun.* Academic Press Inc.; 2014;39:142–51.

486 29. Ross M, Ingram L, Taylor G, Malone E, Simpson RJ, West D, et al. Older men display elevated  
487 levels of senescence-associated exercise-responsive CD28null angiogenic T cells compared with  
488 younger men. *Physiol Rep.* American Physiological Society; 2018;6.

489 30. Simpson RJ, Cosgrove C, Ingram LA, Florida-James GD, Whyte GP, Pircher H, et al. Senescent T-  
490 lymphocytes are mobilised into the peripheral blood compartment in young and older humans after  
491 exhaustive exercise. *Brain Behav Immun.* Brain Behav Immun; 2008;22:544–51.

492 31. Hanson ED, Sakkal S, Que S, Cho E, Spielmann G, Kadife E, et al. Natural killer cell mobilization and  
493 egress following acute exercise in men with prostate cancer. *Exp Physiol.* Blackwell Publishing Ltd;  
494 2020;105:1524–39.

495 32. Wang JS, Chen WL, Weng TP. Hypoxic exercise training reduces senescent T-lymphocyte subsets  
496 in blood. *Brain Behav Immun* [Internet]. Academic Press; 2011;25:270–8.

497 33. Ovadya Y, Landsberger T, Leins H, Vadai E, Gal H, Biran A, et al. Impaired immune surveillance  
498 accelerates accumulation of senescent cells and aging. *Nat Commun* 2018 91. Nature Publishing

499 Group; 2018;9:1–15.

500 34. Koch S, Larbi A, Özcelik D, Solana R, Gouttefangeas C, Attig S, et al. Cytomegalovirus infection: A  
501 driving force in human T cell immunosenescence. *Ann N Y Acad Sci.* Blackwell Publishing Inc.; 2007. p.  
502 23–35.

503 35. Pawelec G, Larbi A, Derhovanessian E. Senescence of the Human Immune System. *J Comp Pathol.*  
504 W.B. Saunders Ltd; 2010;142.

505 36. Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, et al. Preferred reporting items for  
506 systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* 2009.

507 37. Brenchley JM, Karandikar NJ, Betts MR, Ambrozak DR, Hill BJ, Crotty LE, et al. Expression of CD57  
508 defines replicative senescence and antigen-induced apoptotic death of CD8+ T cells. *Blood.* *Blood*;  
509 2003;101:2711–20.

510 38. Lopez-Vergès S, Milush JM, Pandey S, York VA, Arakawa-Hoyt J, Pircher H, et al. CD57 defines a  
511 functionally distinct population of mature NK cells in the human CD56dimCD16+ NK-cell subset.  
512 *Blood.* *American Society of Hematology*; 2010;116:3865–74.

513 39. Appay V, Van Lier RAW, Sallusto F, Roederer M. Phenotype and function of human T lymphocyte  
514 subsets: Consensus and issues [Internet]. *Cytom Part A. Cytometry A*; 2008. p. 975–83.

515 40. Henson SM, Akbar AN. KLRG1-more than a marker for T cell senescence [Internet]. *Age (Omaha).*  
516 *Age (Dordr)*; 2009. p. 285–91.

517 41. Plunkett FJ, Franzese O, Finney HM, Fletcher JM, Belaramani LL, Salmon M, et al. The Loss of  
518 Telomerase Activity in Highly Differentiated CD8 + CD28 – CD27 – T Cells Is Associated with  
519 Decreased Akt (Ser 473 ) Phosphorylation . *J Immunol.* *The American Association of Immunologists*;  
520 2007;178:7710–9.

521 42. Voehringer D, Koschella M, Pircher H. Lack of proliferative capacity of human effector and  
522 memory T cells expressing killer cell lectinlike receptor G1 (KLRG1). *Blood.* *Blood*; 2002;100:3698–  
523 702.

524 43. Callender LA, Carroll EC, Beal RWJ, Chambers ES, Nourshargh S, Akbar AN, et al. Human CD8 +  
525 EMRA T cells display a senescence-associated secretory phenotype regulated by p38 MAPK. *Aging*  
526 *Cell.* *Blackwell Publishing Ltd*; 2018;17.



- 527 44. Zhou D, Borsa M, Simon AK. Hallmarks and detection techniques of cellular senescence and  
528 cellular ageing in immune cells [Internet]. *Aging Cell*. Blackwell Publishing Ltd; 2021. p. e13316.
- 529 45. Kaminsky LA, Arena R, Myers J. Reference standards for cardiorespiratory fitness measured with  
530 cardiopulmonary exercise testing data from the fitness registry and the importance of exercise  
531 national database. *Mayo Clin Proc*. Elsevier Ltd; 2015;90:1515–23.
- 532 46. Azali Alamdari K, Bashiri J. Effects of hypobaric Endurance Training on Graded Exercise Induced  
533 Lymphocyte Mobilization, Senescence and Their Surface Thiol Levels in Elite Male Athletes. *Int J Appl*  
534 *Exerc Physiol*. International Society of Communication and Development Between Universities  
535 (ISCDBU); 2018;7:48–55.
- 536 47. Curran M, Campbell JP, Powell E, Chikhliya A, Narendran P. The mobilisation of early mature  
537 CD56dim-CD16bright NK cells is blunted following a single bout of vigorous intensity exercise in Type  
538 1 Diabetes. *Exerc Immunol Rev* [Internet]. 2020;26:116–31.
- 539 48. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. Quantity and  
540 quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and  
541 neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Med Sci Sports*  
542 *Exerc*. *Med Sci Sports Exerc*; 2011;43:1334–59.
- 543 49. Ingram LA, Simpson RJ, Malone E, Florida-James GD. Sleep disruption and its effect on  
544 lymphocyte redeployment following an acute bout of exercise. *Brain Behav Immun*. Academic Press  
545 Inc.; 2015;47:100–8.
- 546 50. Farina D, Ferguson RA, Macaluso A, De Vito G. Correlation of average muscle fiber conduction  
547 velocity measured during cycling exercise with myosin heavy chain composition, lactate threshold,  
548 and VO<sub>2</sub>max. *J Electromyogr Kinesiol*. *J Electromyogr Kinesiol*; 2007;17:393–400.
- 549 51. Turner JE, Aldred S, Witard OC, Drayson MT, Moss PM, Bosch JA. Latent Cytomegalovirus  
550 infection amplifies CD8 T-lymphocyte mobilisation and egress in response to exercise. *Brain Behav*  
551 *Immun*. *Brain Behav Immun*; 2010;24:1362–70.
- 552 52. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple , graphical  
553 test measures of funnel plot asymmetry. *BMJ*. 1997;315:629–34.
- 554 53. Curran M, Campbell J, Drayson M, Andrews R, Narendran P. Type 1 diabetes impairs the  
555 mobilisation of highly-differentiated CD8+T cells during a single bout of acute exercise. *Exerc*

556 Immunol Rev [Internet]. 2019;25:64–82.

557 54. Krzywkowski K, Petersen EW, Ostrowski K, Kristensen JH, Boza J, Pedersen BK. Effect of glutamine  
558 supplementation on exercise-induced changes in lymphocyte function. *Am J Physiol - Cell Physiol*.  
559 American Physiological Society; 2001;281.

560 55. Lavoy EC, Bigley AB, Spielmann G, Rector JL, Morrison MR, O'Connor DP, et al. CMV amplifies T-  
561 cell redeployment to acute exercise independently of HSV-1 serostatus. *Med Sci Sports Exerc*. *Med*  
562 *Sci Sports Exerc*; 2014;46:257–67.

563 56. Bigley AB, Lowder TW, Spielmann G, Rector JL, Pircher H, Woods JA, et al. NK-cells have an  
564 impaired response to acute exercise and a lower expression of the inhibitory receptors KLRG1 and  
565 CD158a in humans with latent cytomegalovirus infection. *Brain Behav Immun* [Internet]. Academic  
566 Press; 2012;26:177–86.

567 57. Wang JS, Wu CK. Systemic hypoxia affects exercise-mediated antitumor cytotoxicity of natural  
568 killer cells. *J Appl Physiol*. *J Appl Physiol* (1985); 2009;107:1817–24.

569 58. Bigley AB, Rezvani K, Pistillo M, Reed J, Agha N, Kunz H, et al. Acute exercise preferentially  
570 redeploys NK-cells with a highly-differentiated phenotype and augments cytotoxicity against  
571 lymphoma and multiple myeloma target cells. Part II: Impact of latent cytomegalovirus infection and  
572 catecholamine sensitivity. *Brain Behav Immun*. Academic Press Inc.; 2015;49:59–65.

573 59. Brown FF, Bigley AB, Sherry C, Neal CM, Witard OC, Simpson RJ, et al. Training status and sex  
574 influence on senescent T-lymphocyte redistribution in response to acute maximal exercise. *Brain*  
575 *Behav Immun*. Academic Press Inc.; 2014;39:152–9.

576 60. Bruunsgaard H, Jensen M, Schjerling P, Halkjaer-Kristensen J, Ogawa K, Skinhøj P, et al. Exercise  
577 induces recruitment of lymphocytes with an activated phenotype and short telomeres in young and  
578 elderly humans. *Life Sci*. *Life Sci*; 1999;65:2623–33.

579 61. JS W, TP W. Hypoxic exercise training promotes antitumour cytotoxicity of natural killer cells in  
580 young men. *Clin Sci (Lond)*. *Clin Sci (Lond)*; 2011;121:343–53.

581 62. JP C, NE R, VE B, M T, JJ van Z, MT D, et al. Acute exercise mobilises CD8+ T lymphocytes  
582 exhibiting an effector-memory phenotype. *Brain Behav Immun*. *Brain Behav Immun*; 2009;23:767–  
583 75.

584 63. Minuzzi L, Rama L, Bishop N, Rosado F, Martinho A, Paiva A, et al. Lifelong training improves anti-

585 inflammatory environment and maintains the number of regulatory T cells in masters athletes. *Eur J*  
586 *Appl Physiol. Eur J Appl Physiol*; 2017;117:1131–40.

587 64. Simpson RJ, Cosgrove C, Chee MM, McFarlin BK, Bartlett DB, Spielmann G, et al. Senescent  
588 phenotypes and telomere lengths of peripheral blood T-cells mobilized by acute exercise in humans.  
589 *Exerc Immunol Rev [Internet]*. 2010;16:40–55.

590 65. Anane LH, Edwards KM, Burns VE, Drayson MT, Riddell NE, van Zanten JJCSV, et al. Mobilization  
591 of  $\gamma\delta$  T lymphocytes in response to psychological stress, exercise, and  $\beta$ -agonist infusion. *Brain Behav*  
592 *Immun. Brain Behav Immun*; 2009;23:823–9.

593 66. Fan X, Wang Y.  $\beta$ 2 Adrenergic receptor on T lymphocytes and its clinical implications. *Prog Nat*  
594 *Sci. Science Press*; 2009. p. 17–23.

595 67. Pedersen L, Idorn M, Olofsson GH, Lauenborg B, Nookaew I, Hansen RH, et al. Voluntary running  
596 suppresses tumor growth through epinephrine- and IL-6-dependent NK cell mobilization and  
597 redistribution. *Cell Metab. Cell Press*; 2016;23:554–62.

598 68. Dimitrov S, Lange T, Born J. Selective Mobilization of Cytotoxic Leukocytes by Epinephrine. *J*  
599 *Immunol. The American Association of Immunologists*; 2010;184:503–11.

600 69. Krüger K, Lechtermann A, Fobker M, Völker K, Mooren FC. Exercise-induced redistribution of T  
601 lymphocytes is regulated by adrenergic mechanisms. *Brain Behav Immun [Internet]*. Academic Press;  
602 2008;22:324–38.

603 70. Krüger K, Alack K, Ringseis R, Mink L, Pfeifer E, Schinle M, et al. Apoptosis of T-Cell Subsets after  
604 Acute High-Intensity Interval Exercise. *Med Sci Sports Exerc. Lippincott Williams and Wilkins*;  
605 2016;48:2021–9.

606 71. Bujko K, Cymer M, Adamiak M, Ratajczak MZ. An Overview of Novel Unconventional Mechanisms  
607 of Hematopoietic Development and Regulators of Hematopoiesis – a Roadmap for Future  
608 Investigations. *Stem Cell Rev Reports. Springer*; 2019;15:785–94.

609 72. Abrahin O, Cortinhas-Alves EA, Vieira RP, Guerreiro JF. Elite athletes have longer telomeres than  
610 sedentary subjects: A meta-analysis. *Exp Gerontol [Internet]*. Elsevier Inc.; 2019;119:138–45.

611 73. Schirmer M, Vallejo AN, Weyand CM, Goronzy JJ. Resistance to apoptosis and elevated  
612 expression of Bcl-2 in clonally expanded CD4+CD28- T cells from rheumatoid arthritis patients. *J*  
613 *Immunol [Internet]*. 1998;161:1018–25.

- 614 74. Vallejo AN, Schirmer M, Weyand CM, Goronzy JJ. Clonality and Longevity of CD4 + CD28 null T  
615 Cells Are Associated with Defects in Apoptotic Pathways . J Immunol. The American Association of  
616 Immunologists; 2000;165:6301–7.
- 617 75. Prösch S, Wendt CEC, Reinke P, Priemer C, Oppert M, Krüger DH, et al. A novel link between  
618 stress and human cytomegalovirus (HCMV) infection: Sympathetic hyperactivity stimulates HCMV  
619 activation. Virology [Internet]. Academic Press Inc.; 2000;272:357–65.
- 620 76. Mehta SK, Stowe RP, Feiveson AH, Tyring SK, Pierson DL. Reactivation and Shedding of  
621 Cytomegalovirus in Astronauts during Spaceflight. J Infect Dis. Oxford Academic; 2000;182:1761–4.
- 622 77. Spielmann G, McFarlin BK, O’Connor DP, Smith PJW, Pircher H, Simpson RJ. Aerobic fitness is  
623 associated with lower proportions of senescent blood T-cells in man. Brain Behav Immun. Brain  
624 Behav Immun; 2011;25:1521–9.
- 625 78. Fisher JP, Young CN, Fadel PJ. Autonomic adjustments to exercise in humans. Compr Physiol.  
626 Wiley-Blackwell Publishing Ltd; 2015;5:475–512.
- 627 79. Fritsche A, Stumvoll M, Grub M, Sieslack S, Renn W, Schmölling RM, et al. Effect of hypoglycemia  
628 on  $\beta$ -adrenergic sensitivity in normal and type 1 diabetic subjects. Diabetes Care. American Diabetes  
629 Association Inc.; 1998;21:1505–10.
- 630 80. Liston A, Gray DHD. Homeostatic control of regulatory T cell diversity [Internet]. Nat Rev  
631 Immunol. Nat Rev Immunol; 2014. p. 154–65.
- 632 81. Krüger K, Mooren FC. T cell homing and exercise. Exerc Immunol Rev [Internet]. 2007;13:37–54.
- 633 82. Dimitrov S, Benedict C, Heutling D, Westermann J, Born J, Lange T. Cortisol and epinephrine  
634 control opposing circadian rhythms in T cell subsets. Blood. Blood; 2009;113:5134–43.
- 635 83. Benschop RJ, Oostveen FG, Heijnen CJ, Ballieux RE.  $\beta$  2-Adrenergic stimulation causes  
636 detachment of natural killer cells from cultured endothelium. Eur J Immunol. Eur J Immunol;  
637 1993;23:3242–7.
- 638 84. Navalta JW, Sedlock DA, Park KS. Effect of exercise intensity on exercise-induced lymphocyte  
639 apoptosis. Int J Sports Med. Int J Sports Med; 2007;28:539–42.
- 640 85. Pizza FX, Mitchell JB, Davis BH, Starling RD, Holtz RW, Bigelow N. Exercise-induced muscle  
641 damage: effect on circulating leukocyte and lymphocyte subsets. Med Sci Sport Exerc [Internet].

642 1995;27:363–70.

643 86. Pedersen BK. Muscle as a secretory organ. *Compr Physiol. Compr Physiol*; 2013;3:1337–62.

644 87. Pedersen BK, Febbraio MA. Muscles, exercise and obesity: Skeletal muscle as a secretory organ  
645 [Internet]. *Nat Rev Endocrinol. Nature Publishing Group*; 2012. p. 457–65.

646 88. Akbar AN. The convergence of senescence and nutrient sensing during lymphocyte ageing  
647 [Internet]. *Clin Exp Immunol. Blackwell Publishing Ltd*; 2017. p. 4–5.

648 89. Feldman RD, Limbird LE, Nadeau J, Robertson D, Wood AJJ. Alterations in Leukocyte  $\beta$ -Receptor  
649 Affinity with Aging. *N Engl J Med. Massachusetts Medical Society*; 1984;310:815–9.

650

## Figure Legends

651 Fig. 1. Flowchart of study selection.

652 Fig. 2. Forest plot of standardized mean difference (SMD) and 95% confidence interval for the overall  
653 effect immediately post exercise compared to baseline values. CMV: Cytomegalovirus; H-AT: hypoxic-  
654 absolute exercise; HC: hypobaric control; H-C: hypoxic resting; HE: hypobaric exercise; HI: High  
655 intensity; H-RT: hypoxic-relative exercise; HSV1: herpes simplex virus 1; LL: Lower limit of 95%  
656 confidence interval; LT: Lactate threshold; MI: Moderate intensity. NC: normobaric control; N-C:  
657 normoxic resting; NE: normobaric exercise; N-T: normoxic exercise; TD1: Type 1 diabetes; TR: Trained;  
658 UL: Upper limit of 95% confidence interval; UN: Untrained.

659 Fig. 3. . Forest plot of standardized mean difference (SMD) and 95% confidence interval for the  
660 overall effect 1h post exercise compared to baseline values. CMV: Cytomegalovirus; H-AT: hypoxic-  
661 absolute exercise; HC: hypobaric control; H-C: hypoxic resting; HE: hypobaric exercise; HI: High  
662 intensity; H-RT: hypoxic-relative exercise; HSV1: herpes simplex virus 1; LL: Lower limit of 95%  
663 confidence interval; LT: Lactate threshold; MI: Moderate intensity. NC: normobaric control; N-C:  
664 normoxic resting; NE: normobaric exercise; N-T: normoxic exercise; TD1: Type 1 diabetes; TR: Trained;  
665 UL: Upper limit of 95% confidence interval; UN: Untrained.

666 Fig. 4. Forest plot of standardized mean difference (SMD) and 95% confidence interval for the overall  
667 effect 2h post exercise compared to baseline values. CMV: Cytomegalovirus; H-AT: hypoxic-absolute  
668 exercise; HC: hypobaric control; H-C: hypoxic resting; HE: hypobaric exercise; HI: High intensity; H-RT:  
669 hypoxic-relative exercise; HSV1: herpes simplex virus 1; LL: Lower limit of 95% confidence interval; LT:  
670 Lactate threshold; MI: Moderate intensity. NC: normobaric control; N-C: normoxic resting; NE:  
671 normobaric exercise; N-T: normoxic exercise; TD1: Type 1 diabetes; TR: Trained; UL: Upper limit of 95%  
672 confidence interval; UN: Untrained.

673 Fig 5. The figure summarizes the lymphocytes count fold change from baseline to each time point for  
674 the senescent cells: CD8 T cells (in blue), CD4 T cell (in dark pink) and NK cells (in light pink and black  
675 centre). The position of the cells represents the SMD of each meta-analysis at each time-point post  
676 exercise.

677

## ***Meta-Analysis***

### **Effect of exercise on acute senescent lymphocyte counts: a systematic review and meta-analysis.**

Amanda V. Sardeli <sup>a,b\*</sup>, Marcelo A. Mori <sup>c, d, e</sup>, Janet M. Lord <sup>a, f</sup>

<sup>a</sup> MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, UK.

<sup>b</sup> Laboratory of Exercise Physiology (FISEX), University of Campinas, Campinas, Brazil.

<sup>c</sup> Department of Biochemistry and Tissue Biology, Institute of Biology, University of Campinas, Campinas, Brazil.

<sup>d</sup> Experimental Medicine Research Cluster, University of Campinas, Campinas, Brazil.

<sup>e</sup> Obesity and Comorbidities Research Center, University of Campinas, Campinas, Brazil.

<sup>f</sup> NIHR Birmingham Biomedical Research Centre, University Hospital Birmingham and University of Birmingham, Birmingham, UK.

Short Title: Effect of exercise on acute senescent lymphocyte counts

Corresponding Author:

Amanda Veiga Sardeli

Institute of Inflammation and Ageing

Queen Elizabeth Hospital

Mindelsohn Way

Birmingham, West Midlands, B15 2WB, UK

Tel: +44 (0)121 414 3344

E-mail: amandavsardeli@gmail.com

Number of Tables: 2.

Number of Figures: 5.

Word count: 4832.

Keywords: Exercise, Ageing, Cellular senescence, Immunosenescence, Immunology.



## Abstract

**Background:** Highly differentiated, senescent lymphocytes are pro-inflammatory and contribute to age-related systemic inflammation, inflammageing. There are several reports of acute changes in senescent lymphocyte counts post-exercise which potentially has consequences for systemic inflammation. However, there is little consensus since the studies differ with respect to participants, exercise protocols, cellular markers assessed, and the time point of assessment post-exercise.

**Objective:** We performed a systematic review and meta-analysis to assess the impact of exercise on senescent lymphocyte counts in blood immediately, 1h and 2h post exercise.

**Methods:** The search was performed in PubMed (MEDLINE), Web of Science, Embase, Scopus and Cochrane, on January 11, 2021. The 13 studies selected tested aerobic exercise effects, mainly in young men. They assessed the counts of lymphocytes (CD4 T cells, CD8 T cells, NK cells), with the following immune cell marker combinations: KLRG1+, CD57+ (only NK cells), EMRA T cells (CD45RA+CCR7-CD28-CD27-), CD28-CD27-, KLRG1+CD28- and CD28-. Independent extraction of articles by 2 researchers.

**Results:** Standardized mean difference (SMD) and 95% confidence interval between baseline and post-exercise showed significant increases (SMD > 0.9,  $p < 0.003$ ) in all types of lymphocyte counts immediately post exercise. At 1h post exercise senescent CD4 T cells returned to baseline values ( $p = 0.74$ ), CD8 T cells were reduced ( $-0.26 [-0.41; -0.11]$ ,  $p = 0.001$ ), and senescent NK cells were raised ( $0.62 [0.14; 1.10]$ ,  $p = 0.01$ ) above baseline. By 2 hours post exercise, senescent CD4 T cells were reduced ( $-0.94 [-1.40; -0.48]$ ,  $p < 0.001$ ), CD8 T cells remained below baseline ( $-0.53 [-1.04; -0.009]$ ,  $p = 0.04$ ), and NK cells had returned to baseline values ( $-0.29 [-0.64; 0.07]$ ,  $p = 0.11$ ). The main determinants of heterogeneity between studies were cytomegalovirus (CMV) serostatus and the characteristics of exercise protocols. CMV+ individuals had a higher immediate lymphocytosis and 1h post lymphopenia than CMV- individuals. Exercise performed at higher intensities and shorter durations led to higher magnitude of change in senescent lymphocyte counts at all time-points.

**Conclusion:** The differing effects of exercise on senescent NK cells and CD4 and CD8 T cells suggest differing susceptibility to factors modulating lymphocyte extravasation such as adrenaline and exercise intensity.

## Introduction

Immunosenescence, the gradual remodelling of the immune system, is an integral component of the ageing process [1]. Advanced age impairs innate immune responses, contributes to chronic low-grade inflammation (inflammageing) and reduces immunity, increasing the risk of infections, autoimmunity and overall poor health in the older adult [2,3]. Among the features associated with adaptive immunosenescence are the atrophy of the thymus, which reduces naïve T cell output, and the subsequent increased number of highly differentiated, senescent T cells in the circulation [3,4]. Senescent cells are one of the causes of detrimental effects to the body during ageing, contributing to chronic diseases, such as idiopathic pulmonary fibrosis, diabetes, and osteoarthritis [5]. It has been shown recently that mice with high levels of senescent T cells, due to dysfunctional mitochondria, enter premature senescence and a broad range of age-related diseases [6]. Immunosenescence, especially T cell senescence, may therefore be a major contributor to the ageing process.

Senescent cells undergo a state of cell quiescence with permanent cell cycle arrest induced by different sources of stress and damage to the cell. These cells produce a senescence-associated secretory phenotype (SASP), which is composed of pro-inflammatory cytokines, chemokines, growth factors and proteases. Cells releasing SASP alter the tissue microenvironment, affect neighboring cells, and are thus deleterious [7,8]. In the immune system there are some subtle differences. For example, T cells can have a functionally exhausted phenotype resulting from chronic stimulation, which is distinct from a senescent phenotype resulting from ageing or chronic infection. These phenotypes can be differentiated by cell surface markers [9]. We have therefore used the markers identified as relating to senescent T cells such as loss of CD28 and CD27 and expression of KLRG1 and CD57. Importantly, senescent T cells also produce a SASP that is highly pro-inflammatory and similar in content to that of non-immune senescent cells [10], therefore they are likely to contribute to inflammageing and tissue compromise during ageing.

The immunomodulatory effects of exercise have been widely explored and could be associated with the reduction in senescent cell counts [10], for example obese mice provided with an exercise wheel had reduced numbers of senescent cells in their adipose tissue [11]. Exercise has been reported to have a range of immune enhancing effects including reducing chronic low-grade inflammation [12], improving responses to vaccination [13], reducing the risk of infection [14,15], improving the immune response against viruses and bacteria and reducing the burden of latent viral infections [16–18]. Among the main physiological mechanisms mediating the immunomodulatory

benefits of exercise are the reduction in body fat and the release of anti-inflammatory cytokines, such as interleukin-6 (IL-6) and IL-1RA, by the exercising muscle [12,19].

Recently, Duggal *et al.* [20] have reported the benefits of sustained physical activity in to old age on adaptive immune phenotype and immunosenescence. They reported that thymic health, as measured by the frequency of naïve T cells and recent thymic emigrants (RTE), was better preserved in older exercisers (cyclists) compared to inactive elders. Older cyclists also had significantly higher serum levels of the thymoprotective cytokine interleukin-7 (IL-7), higher B regulatory cell frequency, lower IL-6 and reduced Th17 polarization, all markers of an aged immune system. However, they also reported that the age-related increase in senescent T cells was not prevented in the cyclists [20].

Despite the chronic benefits of exercise being well established, whether acute exercise increases susceptibility to infection or confers immune protection is still a matter of debate [21]. However, an increase in lymphocyte counts in the blood (lymphocytosis), followed by a decrease (lymphocytopenia) post exercise has generally been reported [22]. Lymphocytes are proposed to migrate from the marginal pool, the spleen and lymph nodes in to the blood, as well as increased release from the bone marrow to produce the lymphocytosis. This migration is mediated by exercise-induced shear stress on blood vessels and catecholamines, as well as cortisol and to some extent cytokines such as IL-6 [23–25].

What is less clear is the impact of exercise on specific immune cell types and their differentiation state, notably senescent immune cells. This is important bearing in mind their pro-inflammatory nature and potential role in driving inflammaging and the aged phenotype [26]. Studies investigating senescent lymphocyte counts in circulation post-exercise have shown a variety of responses [27–31], including a reduction on leukocyte counts [32]. However, senescent, or highly differentiated lymphocytes appear to be more likely to increase in blood with exercise than lymphocytes in earlier stages of differentiation. This could be beneficial in leading to their subsequent removal by NK cells or CD8 T cells which can detect senescent cells and kill them by apoptosis [22,33].

Another important confounding factor in the various exercise intervention studies, is infection by cytomegalovirus (CMV) that increases with age and has deleterious effects on lymphocyte immunity, accelerating immunosenescence [22,34,35]. The higher baseline cell counts of senescent lymphocytes in CMV+ individuals could lead to higher magnitude of change in these individuals after exercise, and thus the CMV serostatus might be an important confounding factor between studies [27,28]. Other factors that may cause different results between studies are: the

comparison between absolute cell counts and the frequency of cells in the circulation; the different types of lymphocytes assessed; the membrane markers used to identify cell senescence; characteristics of the study population (age, sex and physical activity level) and the exercise protocols used (type of exercise, volume and intensity).

To derive a consensus from the literature it is important to isolate the variety of confounding factors among the studies and to run a pooled effects meta-analysis. Thus, we aimed to identify the impact of acute exercise on the frequency of senescent T cells and NK cells, taking in to account variables such as CMV serostatus, age, training status and specifics of the exercise protocols.

## **Methods**

This systematic review and meta-analysis was registered on PROSPERO under the number CRD42021267078, that can be assessed at <https://www.crd.york.ac.uk/prospero/>, and it was reported in accordance with the recommendations of Preferred Reporting Items for Systematic Reviews and Meta Analyses (PRISMA) statement [36].

### *Search strategy*

On January 11, 2021 the search was updated at PubMed (Medline), Web of Science, Embase, Scopus and Cochrane. It combined the synonyms of “senescent markers” and “exercise” according to each data base descriptor and field of search as detailed in the Supplementary material.

### *Eligibility criteria*

Figure 1 shows the study selection process, completed by two independent reviewers. We included studies: (1) of acute interventional exercise; (2) with no associated intervention, i.e. exercise only group; (3) in humans from both sexes; (4) comparing resting and immediately, 1h and 2h post exercise condition; (5) assessing any bona fide markers of immunosenescence; (6) assessing CD4+ or CD8+ T cells, or NK lymphocytes; (7) written in English.

### *Immunosenescence cell markers*

Markers of senescence traditionally used for non-immune cells, such as p16<sup>ink4a</sup> and SA-βGal have not been used in studies of immunosenescence which focus on cell membrane markers. We therefore selected several broadly accepted markers of immunosenescence to use in this study and the characteristics of each of them are described below.

**CD57<sup>+</sup>.** CD57<sup>+</sup> NK cells have been attributed a senescent-like phenotype due to their short telomeres and inability to proliferate [1,37,38].

**CD28<sup>-</sup> CD27<sup>-</sup>.** CD27 and CD28 are costimulatory receptors and T cells lacking CD27 and CD28 are thought to be fully differentiated T cells exhibiting shorter telomere length [39]. When the expression of CD27 and CD28 is lost, there is no evidence of subsequent re-expression and the downregulation of these molecules are linked to dysfunctional T cells with a SASP secretome [40,41].

**KLRG1<sup>+</sup>.** T-lymphocytes expressing KLRG1 have impaired capacity to proliferate, yet maintain immediate effector cell capabilities such as the recognition and killing of target cells [42].

**EMRA (CD45RA<sup>+</sup>CCR7<sup>-</sup>CD28<sup>-</sup>CD27<sup>-</sup>).** EMRA, for terminally differentiated effector memory cells re-expressing CD45RA, have the key features of cell senescence, with low proliferation response and a highly inflammatory phenotype [10]. They also have high levels of DNA damage and loss of telomerase activity [43]. However, due to their ability to proliferate under specific conditions their phenotype is distinct from non-immune senescent cells which cannot proliferate [44].

#### *Exclusion criteria*

We excluded studies that: (1) did not have original data or did not undergo peer-review such as reviews, commentaries, editorials, letter to the editors, case reports or conference abstracts; (2) assessed other senescence markers such as telomere shortening, or telomerase activity; (3) had not tested exercise effects; (4) had not assessed immunosenescence in humans; (5) assessed senescence in other cells, besides lymphocytes and NK cells; (6) were not written in English.

#### **Data collection and data items**

Data collection was performed by two independent researchers. The means and a measure of dispersion of the senescent cell counts were extracted for each subgroup within studies. Mean, standard deviation (SD) and sample number (n) were used for the meta-analyses. Standard error (SE) was converted to SD by the equation  $SD = SE \times (\sqrt{n})$ , if SD was not provided in the original study.

For subgroup analysis we extracted information about participants sex, age, level of training, health condition and CMV serostatus, type of lymphocytes assessed, membrane markers used, unit of measurement, and the characteristics of the exercise bout such as intensity, volume, and type of equipment.

The sample of studies was classified as young, middle aged and old according to the mean age reported (young [ $<30$  yrs], middle aged [30-40 yrs] and older adults [ $>50$  yrs]).

The participants were considered “trained” when the studies classified them as elite athletes, trained, physically active, cyclists or when the  $VO_2$ max was above the 50% percentile according to their age [45]; they were considered “untrained” when the studies classified them as untrained, or doing no regular physical activity or sedentary. The studies that did not report the participant’s physical activity level or reported a too wide range of physical activity level among their participants were excluded from this subgroup analysis. Individuals undergoing exercise chronic intervention were considered trained [32,46]; while the individuals undergoing non-exercise intervention in Wang et al. [32] were classified as untrained and the individuals undergoing non-exercise intervention in Azali Alamdari et al. [46] were excluded for training status analysis, since they were athletes at baseline.

Regarding health status, only Curran et al. [47] have included individuals with type I diabetes, while the other studies only included healthy participants.

The exercise intensity was classified according to the percentage of  $VO_2$ max described by the American college of Sports Medicine [48], in which 46-63% is moderate, 64-90% is vigorous and  $>91\%$  is near maximum. The intensity reported on Ingram et al. study [49], in watts was estimated as 73.7% of maximum according to data from participants of a similar age. Another study tested different protocols according to their lactate threshold (5% under LT, 5% above LT and 15% above LT) in the same individuals and each of them were included in the meta-analysis as a separate study [27]. The intensity was also converted to percentage according to Farina *et al.* [50], in which 5%  $<LT$  was considered 61.1%, 5%  $>LT$  was considered 71.1%, and 15%  $>LT$  was considered 81.1%. The studies reporting percentage of estimated maximum power or percentage of ventilatory threshold work rate, instead of  $VO_2$ max, were classified for subgroup analysis as these markers were proportionally equivalent.

The studies applying incremental maximum effort tests and other protocols expected to last less than 20 min were considered short, the ones applying 30 min duration were considered moderate and above this they were considered long duration.

Only Azali Alamdari et al. [46] had a control group, and thus, the change of control group was subtracted from the exercise change to increase the robustness of the analysis. Although Turner et al. [51] had also reported a control group, they did not present the effects of the control period on

CD28-CD27- markers, in this way the control group was not considered for analysis. Two studies presented acute exercise effects before and after a variety of chronic interventions [32,46] and thus we included only their post intervention session to avoid sample overlapping in the analysis.

### **Statistical analysis**

We analysed the absolute cell count as the outcome measure, considering the standardized mean difference (SMD) and 95% of confidence interval between baseline levels and post exercise time-points since the units of measure were not consistent across studies.

The 3 main meta-analyses, for each time point (immediately, 1h and 2h post exercise) and the subgroup analyses were performed using Comprehensive Meta-Analysis software, version 3.3.070. When there was statistical significance for heterogeneity, randomized effect models were selected and when there was no significant heterogeneity, fixed effects were applied. The inconsistency between studies was reported as a percentage ( $I^2$ ), based on difference between expected heterogeneity (df) and true heterogeneity (Q-value).

For subgroup analysis we tested the influence of the following confounding factors: sex (men and women); age (young [ $<30$  yrs], middle aged [ $30-40$  yrs] and older adults [ $>50$  yrs]); type of lymphocytes (CD4+, CD8+ and NK); type of senescence marker (KLRG1+, CD57+, EMRA [CD45RA+CCR7-CD28-CD27-], CD28-CD27-, KLRG1+CD28- and CD28-), level of training (trained and untrained); health condition (healthy and diseased); CMV serostatus (CMV+ and CMV-); exercise intensity (moderate, vigorous, near maximum); and exercise volume (short, moderate and long). Q tests were applied to group comparisons, considering 95% confidence.

Egger's tests were performed to check the risk of publication bias in each meta-analysis [52].

### **Results**

We included thirteen studies [27–29,32,46,47,49,51,53–57] testing acute aerobic exercise effects on senescent T lymphocytes and NK cell counts (shown in Figure 1). It is noteworthy that some studies had to be excluded due to the absence of specific description of absolute senescent lymphocyte counts [30,58–64]. Most studies included, reported their results among different subgroups of individuals with different sex, ages, CMV serostatus, types of exercise protocols and time points of analysis that were analyzed as a sub-study.

**\*\*\*please insert Figure 1 here\*\*\***

*Study characteristics*

Table 1 shows the characteristics of the studies included. Only Curran et al.[47] included a type I diabetes group, while the other studies only included healthy participants. While twelve studies tested exercise effects on males, just one tested exercise effects on participants from both sexes [27], and thus comparisons between men and women were not possible in subgroup analysis. One study included middle aged [28], two included older adults [28,29] and all of them (thirteen) tested young adults. Our analysis reported the effect of exercise on T CD4+, T CD8+ and NK cell counts. All studies tested the effects of aerobic exercise, the majority of them used bicycle, and a few used treadmill [46,51].

**\*\*\*please insert Table 1 here\*\*\***

### *Syntheses of the results*

**Lymphocyte counts immediately post exercise.** Figure 2 shows there were significant increases on senescent CD4 T cells (SMD 0.96 [0.67; 1.25],  $p<0.001$ ), CD8 T cells (SMD 1.26 [0.93; 1.59],  $p<0.001$ ) and NK cells counts immediately post exercise (SMD 1.04 [0.35; 1.72],  $p=0.003$ ). However, all those analyses were heterogeneous, reinforcing the need for further subgroup analyses. Furthermore, the analysis of senescent CD4 T cells and CD8 T cell counts had significant risk of bias, evidencing that studies with low precision conducted the main effects.

Table 2 shows no effect of age ( $p= 0.46$ ) or training status ( $p=0.35$ ) on outcomes immediately post exercise. On the other hand, the intensity and duration of exercise protocols and CMV status influenced the post exercise senescent lymphocyte counts (Table 2). Specifically, the higher magnitude of increase in senescent lymphocytes were seen in the maximum intensity and short duration protocols (SMD 1.81 [1.45; 2.1],  $p<0.001$ ) compared to the others (SMD  $<0.85$ ,  $p<0.05$ ). There was a trend to higher senescent lymphocyte counts in CMV positive participants compared to CMV- ( $p=0.09$ ). The CMV status analysis for each subgroup of T lymphocyte showed higher increase in senescent CD8+ T cells for CMV+ (SMD 1.60 [ 0.73; 2.46],  $p<0.001$ ) compared to CMV- (SMD 0.58 [0.33; 0.83],  $p<0.001$ ), with no difference for senescent CD4+ T cells regarding CMV status (SMD CMV+: 0.42 [0.02; 0.82],  $p=0.038$  and CMV-: 0.50 [0.18; 0.82],  $p=0.002$ ).

**\*Please, insert Figure 2 here\***

**Lymphocyte counts one hour post exercise.** Figure 3 shows senescent CD8+ T cell counts were lower compared to baseline levels (SMD -0.28 [-0.44; -0.13],  $p<0.001$ ), while CD4+ T cell counts returned to baseline levels (SMD -0.13 [-0.37; 0.11],  $p=0.28$ ) and NK cells were still above baseline values (SMD



0.62 [0.14; 1.09],  $p=0.11$ ). These analyses were homogeneous ( $p>0.53$ ,  $I^2=0\%$ ), confirming that each of these senescent cells have very consistent response 1h post exercise across the different studies.

Table 2 shows there was a significant reduction in the senescent lymphocyte count only in CMV+ and not CMV- individuals, with significant difference between groups. Regarding each subgroup of T lymphocyte there was no significant reduction for senescent CD4+ T or CD8+ cells in CMV- (SMD CD4+: 0 [-0.30; 0.31],  $p=0.97$  and CD8+: -0.13 [-0.36; 0.09],  $p=0.25$ ) while there was a trend of senescent CD4+ reduction in CMV+ individuals (SMD -0.35 [-0.74; 0.04],  $p=0.075$ ), and reduction of senescent CD8+ T cells in CMV+ (SMD -0.46 [-0.75; -0.18]  $p=0.001$ ). Only vigorous intensity and long duration exercise protocols led to significant reduction of senescent lymphocytes (SMD -0.5 [-0.8; -0.2],  $p<0.001$ ) while the other intensities and durations did not vary significantly ( $p>0.16$ ).

**\*Please, insert Figure 3 here\***

**Lymphocyte counts two hours post exercise.** Figure 4 shows senescent CD4 T cells were reduced (SMD -0.94 [-1.40; -0.48],  $p<0.001$ ), CD8 T cells remained below baseline (SMD -0.53 [-1.04; -0.009],  $p=0.04$ ), and NK cells had returned to baseline values (SMD -0.29 [-0.64; 0.07],  $p=0.11$ ). There was significant risk of publication bias for the analysis of senescent CD4 T cells (Egger test  $p$ -value  $<0.001$ ), evidencing that studies with low precision conducted the main effects in this analysis.

All these three meta-analyses were heterogeneous, however, due to the low number of subgroups in these analyses, only training status, intensity and volume of exercise protocols were analyzed. No difference between trained and untrained individuals was noticed ( $p=0.81$ ) and only maximum intensity and short duration protocols reduced senescent cell counts (SMD -0.7 [-1; -0.4],  $p<0.001$ ), however, it is noteworthy there was very low number of studies in the other categories (Table 2).

**\*Please, insert figure 4 here\***

## Discussion

The main findings of the present meta-analysis were the significant increase in senescent CD8+, CD4+ and NK cell counts immediately post exercise followed by a reduction in senescent CD8+ T cells at 1h and 2h post exercise, a reduction in senescent CD4+ T cells at 2h post exercise and maintenance of increased NK senescent cells at 1h post exercise with a return to baseline at 2h post exercise (Figure 5). Although there is no consensus about the exact role of these redistributions of senescent lymphocytes post exercise, it has been proposed that senescent lymphocytes are preferentially

recruited for immune surveillance and removal by NK and CD8+ T cells, resulting in an exercise-induced senolytic effect [22].

In fact, it is known that T-cells with high cytotoxic capabilities and tissue migration potential, which are characteristics of highly differentiated lymphocytes, are preferentially mobilised by acute stress and exercise [65]. These lymphocytes could be recruited due to their high  $\beta_2$ -adrenergic receptor expression [66] even though they have impaired replicability and co-stimulatory potential. Furthermore, in mice, NK cells are the main mediator of the antitumor effects of exercise. These effects depend on the mobilisation of these cells [67], which are the most responsive lymphocyte subset to acute exercise due to their high  $\beta$ -adrenergic receptor expression [68]. Mobilisation of the senescent, less functional form of these cells could be beneficial if they are then removed, improving the overall quality of the lymphocyte pool.

Following their mobilisation it is possible that T-lymphocytes egressing to the peripheral tissues may experience a pro-apoptotic environment [69], as Kruger et. al. [70] showed the number of highly differentiated CD3+ T cells remained reduced 3h and 24h post exercise. Another possibility could be the return of senescent cells to lymph nodes but most of these cells lack CCR7, a secondary lymphoid organ-homing marker, this is unlikely.

In theory, when senescent T-lymphocytes undergo apoptosis, a subsequent feedback loop could increase the output of naïve T-lymphocytes from the thymus, restoring the peripheral T-lymphocyte pool [22,59]. In fact, naïve lymphocytes counts are increasing 1h post exercise [30,59]. Furthermore, older adults involved in regular exercise have higher serum levels of the thymoprotective IL-7 and higher frequency of RTE than sedentary controls [20], which could be stimulated by senescent lymphocyte clearance post each exercise bout.

In an opposite way, exercise-induced hematopoiesis [25,71], could also affect the thymic feedback loop, increasing the stimuli for senescent lymphocyte removal. Cross-sectional studies showed physically active individuals have lower markers of senescent T lymphocytes [10,20] and master athletes have longer lymphocyte telomere length than untrained controls [72]. Nevertheless, it is noteworthy that highly differentiated, senescent lymphocytes are less sensitive to apoptotic signals [73,74], and the exercise effects on senescent cell apoptosis is still unknown.

Most studies tested the influence of CMV serostatus on exercise responses. CMV reactivation can be triggered through catecholamine-responsive elements [75] and stress hormone levels, which are known to correlate strongly with CMV reactivation in astronauts before and after spaceflight

[76]. Thus, it is believed that CMV<sup>+</sup> individuals have reduced sensitivity to  $\beta$ -adrenergic stimulation and decreased  $\beta_2$ -adrenergic receptor expression to prevent CMV reactivation [58]. However, a reduced  $\beta$ -adrenergic sensitivity of T cells in CMV<sup>+</sup> individuals is not supported by our analysis, and in fact we saw a larger magnitude of changes in CMV<sup>+</sup> individuals. Thus, we believe the expected higher baseline senescent lymphocyte counts in CMV<sup>+</sup> individuals [28,77], especially for senescent CD8<sup>+</sup> T cells, explains the higher magnitude of change with exercise in this population.

There was a greater magnitude of increase in senescent lymphocyte counts immediately post maximum intensity and short duration protocols and greater magnitude of reduction 2h post exercise compared to lower intensities and longer duration protocols. These differences could be explained by higher sympathetic activation and sustained release of epinephrine reported in higher intensities protocols [70,78,79]. However, there is also evidence that cortisol affects lymphocyte counts during exercise [70,80–83]. Exercise of high intensity leads to greater release of cortisol in to the blood and for a longer time and may explain the reduced cell counts at later time points since cortisol induces apoptosis in lymphocytes [84]. We also considered the type of exercise and whether this may make a difference to the senescent cell response. However, we found no difference at any of the time points between treadmill and cycling protocols (data not shown), though this is with the caveat that the number of studies per subgroup category varied greatly.

It is unlikely that IL-6 released by muscle cells during exercise [85], explains the difference between exercise protocols. It is known, that IL-6 attracts lymphocytes to the circulation together with  $\beta$ -adrenergic signaling during exercise [67], however, there is a higher release of IL-6 within exercise protocols with higher energetic demand, such as the higher volume and during regimens [85–87] which do not agree with our findings.

Finally, exercise hypoxia may explain at least part of the changes in T lymphocytes and NK counts with exercise, possibly mediated by the same neuroendocrinological factors released by other stress conditions (i.e.: catecholamines, cortisol) [57,88].

### *Limitations*

The first limitation of this study was that most studies included young individuals. The unbalanced subgroup analysis suggested there is higher magnitude of change on senescent counts in young than older or middle-aged individuals immediately and 1h post exercise. Whether it is a true effect is unclear, it could be explained by reduced  $\beta_2$ -adrenergic receptor sensitivity with ageing [89], which in

turn increases the threshold for catecholamine-induced lymphocyte recruitment. In this way, it is important to confirm these results with a larger sample of older adults.

Comparisons between men and women were also not possible due to the lack of studies in women. An exploratory analysis showed immediately post exercise there was a large ( $p < 0.001$ ) increase of senescent lymphocytes in men (1.23 [0.99; 1.47],  $p < 0.001$ ,  $k=44$ ) compared to studies with mixed sex samples (0.48 [0.19; 0.78],  $p < 0.001$ ,  $k=6$ ), while there was no difference between these subgroups 1h post. Future studies should test to what extent the results in men are also applied to women.

Another limitation was the lack of a control group, i.e. without exercise, within the original studies which precluded a proper risk of bias assessment. In the other hand, the comparison of the same participants along time removes the between subjects' effects, which in turn contributes to the isolation of exercise effects. Furthermore, we explored possible influences of the confounding factors in subgroup analysis. At last, it is noteworthy that only two studies investigated exercise effects on senescent NK cell counts, with is a limitation of the literature and future studies should fill these gaps to strengthen knowledge in the field.

Lastly, one additional issue was the use of markers to identify the different stages of T cell differentiation and their relation to T cell senescence. Thus, no studies enumerated CD57 TEMRA cells, the ones that are closest to a senescent phenotype.

### *Conclusions*

Senescent lymphocyte counts change significantly in the acute response to aerobic exercise. However, a complex picture has emerged where senescent CD8+ cells had a higher immediate lymphocytosis and subsequent lymphopenia (1h and 2h post), senescent CD4+ T cells followed a similar profile but with lower magnitude of change, and senescent NK cells increased but had a delayed return to baseline levels. There was higher magnitude of lymphocytosis and lymphocytopenia for CMV+ individuals and near maximum intensity and short duration protocols. The differing effects of exercise on senescent NK cells and CD4+ and CD8+ T cells suggest differing susceptibility to factors modulating lymphocyte extravasation such as adrenaline that is also regulated by exercise intensity. More studies are needed for understanding exercise effects on senescent NK cells, in older adults and in women.

## **Statements**

### **Acknowledgement**

The authors thanks Diego Nacarato Pereira da Silva for the contribution in the selection of studies and data acquisition.

### **Statement of Ethics**

An ethics statement is not applicable because this study is based exclusively on published literature.

### **Conflict of Interest Statement**

The authors have no conflicts of interest to declare.

### **Funding Sources**

This work was supported by the Newton International Fellowship generously awarded to Amanda Veiga Sardeli by the Academy of Medical Sciences through the UK Government's Newton Fund Programme [NIFR7\1031]. JML is supported by the NIHR Birmingham Biomedical Research Centre. The views expressed here are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

### **Author Contributions**

All three authors have given substantial contributions to the conception and the design of the manuscript; AVS did the studies selection, data collection and analysis. AVS, MAM and JML interpreted the data. AVS did the first draft while MAM and JML reviewed it critically for important intellectual content. All authors read and approved the final version of the manuscript.

### **Data Availability Statement**

The data in this study was obtained from the previous studies where specific restrictions for public sharing their data may apply according to each journal politics. Such dataset may be requested by the corresponding author e-mail.

## References

1. Alpert A, Pickman Y, Leipold M, Rosenberg-Hasson Y, Ji X, Gaujoux R, et al. A clinically meaningful metric of immune age derived from high-dimensional longitudinal monitoring. *Nat Med*. Nature Publishing Group; 2019;25:487–95.
2. Franceschi C, Campisi J. Chronic inflammation (Inflammaging) and its potential contribution to age-associated diseases [Internet]. *Journals Gerontol - Ser A Biol Sci Med Sci*. Oxford University Press; 2014. p. S4–9.
3. Hazeldine J, Lord J, Hampson P. Immunesenescence and inflammaging: A contributory factor in the poor outcome of the geriatric trauma patient. *Ageing Res Rev*. *Ageing Res Rev*; 2015;24:349–57.
4. Majumdar S, Nandi D. Thymic Atrophy: Experimental Studies and Therapeutic Interventions [Internet]. *Scand J Immunol*. Blackwell Publishing Ltd; 2018. p. 4–14.
5. Muñoz-Espín D, Serrano M. Cellular senescence: from physiology to pathology. *Nat Publ Gr*. 2014;
6. Vasileiou PVS, Evangelou K, Vlasis K, Fildis G, Panayiotidis MI, Chronopoulos E, et al. Mitochondrial Homeostasis and Cellular Senescence. *Cells*. Multidisciplinary Digital Publishing Institute (MDPI); 2019;8:686.
7. Coppé JP, Patil CK, Rodier F, Sun Y, Muñoz DP, Goldstein J, et al. Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS Biol*. *PLoS Biol*; 2008;6.
8. Campisi J. Aging, cellular senescence, and cancer [Internet]. *Annu Rev Physiol*. *Annu Rev Physiol*; 2013. p. 685–705.
9. Huff W, Kwon J, Henriquez M, Fetcko K, Dey M. The Evolving Role of CD8 + CD28 - Immunosenescent T Cells in Cancer Immunology. *Int J Mol Sci*. *Int J Mol Sci*; 2019;20.
10. Chen X, Yi Z, Wong GT, Hasan KMM, Kwan JS, Ma AC, et al. Is exercise a senolytic medicine? A systematic review. *Ageing Cell*. Blackwell Publishing Ltd; 2020;
11. Schafer MJ, White TA, Iijima K, Haak AJ, Ligresti G, Atkinson EJ, et al. Cellular senescence mediates fibrotic pulmonary disease. *Nat Commun*. Nature Publishing Group; 2017;8:14532.
12. Gleeson M, Bishop N, Walsh N. *Exercise Immunology*. New York: Routledge/Taylor & Francis Group; 2013.

13. Pascoe AR, Fiatarone Singh MA, Edwards KM. The effects of exercise on vaccination responses: A review of chronic and acute exercise interventions in humans [Internet]. *Brain Behav Immun*. Academic Press Inc.; 2014. p. 33–41.
14. Pape K, Ryttergaard L, Rotevatn TA, Nielsen BJ, Torp-Pedersen C, Overgaard C, et al. Leisure-Time Physical Activity and the Risk of Suspected Bacterial Infections. *Med Sci Sports Exerc*. Lippincott Williams and Wilkins; 2016;48:1737–44.
15. Nieman DC, Wentz LM. The compelling link between physical activity and the body’s defense system [Internet]. *J Sport Heal Sci*. Elsevier B.V.; 2019. p. 201–17.
16. Simpson RJ, Hussain M, Baker F, Bigley AB, Peek MK, Stowe RP. Cardiorespiratory fitness is associated with better control of latent herpesvirus infections in a large ethnically diverse community sample: Evidence from the Texas City Stress and Health Study. *Brain Behav Immun* [Internet]. Elsevier BV; 2017;66:e35.
17. Agha NH, Mehta SK, Rooney B V., Laughlin MS, Markofski MM, Pierson DL, et al. Exercise as a countermeasure for latent viral reactivation during long duration space flight. *FASEB J*. John Wiley and Sons Inc.; 2020;34:2869–81.
18. Grande AJ, Keogh J, Silva V, Scott AM. Exercise versus no exercise for the occurrence, severity, and duration of acute respiratory infections [Internet]. *Cochrane Database Syst Rev*. John Wiley and Sons Ltd; 2020.
19. Duggal NA, Niemi G, Harridge SDR, Simpson RJ, Lord JM. Can physical activity ameliorate immunosenescence and thereby reduce age-related multi-morbidity? [Internet]. *Nat Rev Immunol*. Nature Publishing Group; 2019. p. 563–72.
20. Duggal NA, Pollock RD, Lazarus NR, Harridge S, Lord JM. Major features of immunosenescence, including reduced thymic output, are ameliorated by high levels of physical activity in adulthood. *Aging Cell* [Internet]. Blackwell Publishing Ltd; 2018;17.
21. Simpson RJ, Campbell JP, Gleeson M, Krüger K, Nieman DC, Pyne DB, et al. Can exercise affect immune function to increase susceptibility to infection? *Exerc Immunol Rev* [Internet]. 2020;26:8–22.
22. Simpson RJ. Aging, persistent viral infections, and immunosenescence: Can exercise “make space”? *Exerc Sport Sci Rev*. *Exerc Sport Sci Rev*; 2011;39:23–33.
23. Agha NH, Baker FL, Kunz HE, Graff R, Azadan R, Dolan C, et al. Vigorous exercise mobilizes CD34+

hematopoietic stem cells to peripheral blood via the  $\beta$  2 -adrenergic receptor. *Brain Behav Immun*. Academic Press Inc.; 2018;68:66–75.

24. Pedersen BK. Physical activity and muscle–brain crosstalk [Internet]. *Nat Rev Endocrinol*. Nature Publishing Group; 2019. p. 383–92.

25. McCarthy DA, Macdonald I, Grant M, Marbut M, Watling M, Nicholson S, et al. Studies on the immediate and delayed leucocytosis elicited by brief (30-min) strenuous exercise. *Eur J Appl Physiol Occup Physiol*. Springer-Verlag; 1992;64:513–7.

26. Desdín-Micó G, Soto-Heredero G, Aranda JF, Oller J, Carrasco E, Gabandé-Rodríguez E, et al. T cells with dysfunctional mitochondria induce multimorbidity and premature senescence. *Science* (80-). American Association for the Advancement of Science; 2020;368:1371–6.

27. LaVoy EC, Hussain M, Reed J, Kunz H, Pistillo M, Bigley AB, et al. T-cell redeployment and intracellular cytokine expression following exercise: effects of exercise intensity and cytomegalovirus infection. *Physiol Rep*. American Physiological Society; 2017;5.

28. Spielmann G, Bollard CM, Bigley AB, Hanley PJ, Blaney JW, LaVoy ECP, et al. The effects of age and latent cytomegalovirus infection on the redeployment of CD8+ T cell subsets in response to acute exercise in humans. *Brain Behav Immun*. Academic Press Inc.; 2014;39:142–51.

29. Ross M, Ingram L, Taylor G, Malone E, Simpson RJ, West D, et al. Older men display elevated levels of senescence-associated exercise-responsive CD28null angiogenic T cells compared with younger men. *Physiol Rep*. American Physiological Society; 2018;6.

30. Simpson RJ, Cosgrove C, Ingram LA, Florida-James GD, Whyte GP, Pircher H, et al. Senescent T-lymphocytes are mobilised into the peripheral blood compartment in young and older humans after exhaustive exercise. *Brain Behav Immun*. *Brain Behav Immun*; 2008;22:544–51.

31. Hanson ED, Sakkal S, Que S, Cho E, Spielmann G, Kadife E, et al. Natural killer cell mobilization and egress following acute exercise in men with prostate cancer. *Exp Physiol*. Blackwell Publishing Ltd; 2020;105:1524–39.

32. Wang JS, Chen WL, Weng TP. Hypoxic exercise training reduces senescent T-lymphocyte subsets in blood. *Brain Behav Immun* [Internet]. Academic Press; 2011;25:270–8.

33. Ovadya Y, Landsberger T, Leins H, Vadai E, Gal H, Biran A, et al. Impaired immune surveillance accelerates accumulation of senescent cells and aging. *Nat Commun* 2018 91. Nature Publishing



Group; 2018;9:1–15.

34. Koch S, Larbi A, Özcelik D, Solana R, Gouttefangeas C, Attig S, et al. Cytomegalovirus infection: A driving force in human T cell immunosenescence. *Ann N Y Acad Sci*. Blackwell Publishing Inc.; 2007. p. 23–35.

35. Pawelec G, Larbi A, Derhovanessian E. Senescence of the Human Immune System. *J Comp Pathol*. W.B. Saunders Ltd; 2010;142.

36. Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, et al. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med*. 2009.

37. Brenchley JM, Karandikar NJ, Betts MR, Ambrozak DR, Hill BJ, Crotty LE, et al. Expression of CD57 defines replicative senescence and antigen-induced apoptotic death of CD8<sup>+</sup> T cells. *Blood*. *Blood*; 2003;101:2711–20.

38. Lopez-Vergès S, Milush JM, Pandey S, York VA, Arakawa-Hoyt J, Pircher H, et al. CD57 defines a functionally distinct population of mature NK cells in the human CD56dimCD16<sup>+</sup> NK-cell subset. *Blood*. *American Society of Hematology*; 2010;116:3865–74.

39. Appay V, Van Lier RAW, Sallusto F, Roederer M. Phenotype and function of human T lymphocyte subsets: Consensus and issues [Internet]. *Cytom Part A. Cytometry A*; 2008. p. 975–83.

40. Henson SM, Akbar AN. KLRG1-more than a marker for T cell senescence [Internet]. *Age (Omaha)*. *Age (Dordr)*; 2009. p. 285–91.

41. Plunkett FJ, Franzese O, Finney HM, Fletcher JM, Belaramani LL, Salmon M, et al. The Loss of Telomerase Activity in Highly Differentiated CD8<sup>+</sup> CD28<sup>-</sup> CD27<sup>-</sup> T Cells Is Associated with Decreased Akt (Ser 473 ) Phosphorylation . *J Immunol*. *The American Association of Immunologists*; 2007;178:7710–9.

42. Voehringer D, Koschella M, Pircher H. Lack of proliferative capacity of human effector and memory T cells expressing killer cell lectinlike receptor G1 (KLRG1). *Blood*. *Blood*; 2002;100:3698–702.

43. Callender LA, Carroll EC, Beal RWJ, Chambers ES, Nourshargh S, Akbar AN, et al. Human CD8<sup>+</sup> EMRA T cells display a senescence-associated secretory phenotype regulated by p38 MAPK. *Aging Cell*. *Blackwell Publishing Ltd*; 2018;17.

44. Zhou D, Borsa M, Simon AK. Hallmarks and detection techniques of cellular senescence and cellular ageing in immune cells [Internet]. *Aging Cell*. Blackwell Publishing Ltd; 2021. p. e13316.
45. Kaminsky LA, Arena R, Myers J. Reference standards for cardiorespiratory fitness measured with cardiopulmonary exercise testing data from the fitness registry and the importance of exercise national database. *Mayo Clin Proc*. Elsevier Ltd; 2015;90:1515–23.
46. Azali Alamdari K, Bashiri J. Effects of hypobaric Endurance Training on Graded Exercise Induced Lymphocyte Mobilization, Senescence and Their Surface Thiol Levels in Elite Male Athletes. *Int J Appl Exerc Physiol*. International Society of Communication and Development Between Universities (ISCDBU); 2018;7:48–55.
47. Curran M, Campbell JP, Powell E, Chikhliya A, Narendran P. The mobilisation of early mature CD56dim-CD16bright NK cells is blunted following a single bout of vigorous intensity exercise in Type 1 Diabetes. *Exerc Immunol Rev* [Internet]. 2020;26:116–31.
48. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Med Sci Sports Exerc*. *Med Sci Sports Exerc*; 2011;43:1334–59.
49. Ingram LA, Simpson RJ, Malone E, Florida-James GD. Sleep disruption and its effect on lymphocyte redeployment following an acute bout of exercise. *Brain Behav Immun*. Academic Press Inc.; 2015;47:100–8.
50. Farina D, Ferguson RA, Macaluso A, De Vito G. Correlation of average muscle fiber conduction velocity measured during cycling exercise with myosin heavy chain composition, lactate threshold, and VO<sub>2</sub>max. *J Electromyogr Kinesiol*. *J Electromyogr Kinesiol*; 2007;17:393–400.
51. Turner JE, Aldred S, Witard OC, Drayson MT, Moss PM, Bosch JA. Latent Cytomegalovirus infection amplifies CD8 T-lymphocyte mobilisation and egress in response to exercise. *Brain Behav Immun*. *Brain Behav Immun*; 2010;24:1362–70.
52. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple , graphical test measures of funnel plot asymmetry. *BMJ*. 1997;315:629–34.
53. Curran M, Campbell J, Drayson M, Andrews R, Narendran P. Type 1 diabetes impairs the mobilisation of highly-differentiated CD8+T cells during a single bout of acute exercise. *Exerc*

Immunol Rev [Internet]. 2019;25:64–82.

54. Krzywkowski K, Petersen EW, Ostrowski K, Kristensen JH, Boza J, Pedersen BK. Effect of glutamine supplementation on exercise-induced changes in lymphocyte function. *Am J Physiol - Cell Physiol*. American Physiological Society; 2001;281.

55. Lavoy EC, Bigley AB, Spielmann G, Rector JL, Morrison MR, O'Connor DP, et al. CMV amplifies T-cell redeployment to acute exercise independently of HSV-1 serostatus. *Med Sci Sports Exerc*. *Med Sci Sports Exerc*; 2014;46:257–67.

56. Bigley AB, Lowder TW, Spielmann G, Rector JL, Pircher H, Woods JA, et al. NK-cells have an impaired response to acute exercise and a lower expression of the inhibitory receptors KLRG1 and CD158a in humans with latent cytomegalovirus infection. *Brain Behav Immun* [Internet]. Academic Press; 2012;26:177–86.

57. Wang JS, Wu CK. Systemic hypoxia affects exercise-mediated antitumor cytotoxicity of natural killer cells. *J Appl Physiol*. *J Appl Physiol* (1985); 2009;107:1817–24.

58. Bigley AB, Rezvani K, Pistillo M, Reed J, Agha N, Kunz H, et al. Acute exercise preferentially redeploys NK-cells with a highly-differentiated phenotype and augments cytotoxicity against lymphoma and multiple myeloma target cells. Part II: Impact of latent cytomegalovirus infection and catecholamine sensitivity. *Brain Behav Immun*. Academic Press Inc.; 2015;49:59–65.

59. Brown FF, Bigley AB, Sherry C, Neal CM, Witard OC, Simpson RJ, et al. Training status and sex influence on senescent T-lymphocyte redistribution in response to acute maximal exercise. *Brain Behav Immun*. Academic Press Inc.; 2014;39:152–9.

60. Bruunsgaard H, Jensen M, Schjerling P, Halkjaer-Kristensen J, Ogawa K, Skinhøj P, et al. Exercise induces recruitment of lymphocytes with an activated phenotype and short telomeres in young and elderly humans. *Life Sci*. *Life Sci*; 1999;65:2623–33.

61. JS W, TP W. Hypoxic exercise training promotes antitumour cytotoxicity of natural killer cells in young men. *Clin Sci (Lond)*. *Clin Sci (Lond)*; 2011;121:343–53.

62. JP C, NE R, VE B, M T, JJ van Z, MT D, et al. Acute exercise mobilises CD8+ T lymphocytes exhibiting an effector-memory phenotype. *Brain Behav Immun*. *Brain Behav Immun*; 2009;23:767–75.

63. Minuzzi L, Rama L, Bishop N, Rosado F, Martinho A, Paiva A, et al. Lifelong training improves anti-

inflammatory environment and maintains the number of regulatory T cells in masters athletes. *Eur J Appl Physiol*. *Eur J Appl Physiol*; 2017;117:1131–40.

64. Simpson RJ, Cosgrove C, Chee MM, McFarlin BK, Bartlett DB, Spielmann G, et al. Senescent phenotypes and telomere lengths of peripheral blood T-cells mobilized by acute exercise in humans. *Exerc Immunol Rev* [Internet]. 2010;16:40–55.

65. Anane LH, Edwards KM, Burns VE, Drayson MT, Riddell NE, van Zanten JJCSV, et al. Mobilization of  $\gamma\delta$  T lymphocytes in response to psychological stress, exercise, and  $\beta$ -agonist infusion. *Brain Behav Immun*. *Brain Behav Immun*; 2009;23:823–9.

66. Fan X, Wang Y.  $\beta$ 2 Adrenergic receptor on T lymphocytes and its clinical implications. *Prog Nat Sci*. Science Press; 2009. p. 17–23.

67. Pedersen L, Idorn M, Olofsson GH, Lauenborg B, Nookaew I, Hansen RH, et al. Voluntary running suppresses tumor growth through epinephrine- and IL-6-dependent NK cell mobilization and redistribution. *Cell Metab*. *Cell Press*; 2016;23:554–62.

68. Dimitrov S, Lange T, Born J. Selective Mobilization of Cytotoxic Leukocytes by Epinephrine. *J Immunol*. The American Association of Immunologists; 2010;184:503–11.

69. Krüger K, Lechtermann A, Fobker M, Völker K, Mooren FC. Exercise-induced redistribution of T lymphocytes is regulated by adrenergic mechanisms. *Brain Behav Immun* [Internet]. Academic Press; 2008;22:324–38.

70. Krüger K, Alack K, Ringseis R, Mink L, Pfeifer E, Schinle M, et al. Apoptosis of T-Cell Subsets after Acute High-Intensity Interval Exercise. *Med Sci Sports Exerc*. Lippincott Williams and Wilkins; 2016;48:2021–9.

71. Bujko K, Cymer M, Adamiak M, Ratajczak MZ. An Overview of Novel Unconventional Mechanisms of Hematopoietic Development and Regulators of Hematopoiesis – a Roadmap for Future Investigations. *Stem Cell Rev Reports*. Springer; 2019;15:785–94.

72. Abrahin O, Cortinhas-Alves EA, Vieira RP, Guerreiro JF. Elite athletes have longer telomeres than sedentary subjects: A meta-analysis. *Exp Gerontol* [Internet]. Elsevier Inc.; 2019;119:138–45.

73. Schirmer M, Vallejo AN, Weyand CM, Goronzy JJ. Resistance to apoptosis and elevated expression of Bcl-2 in clonally expanded CD4+CD28- T cells from rheumatoid arthritis patients. *J Immunol* [Internet]. 1998;161:1018–25.

74. Vallejo AN, Schirmer M, Weyand CM, Goronzy JJ. Clonality and Longevity of CD4 + CD28 null T Cells Are Associated with Defects in Apoptotic Pathways . J Immunol. The American Association of Immunologists; 2000;165:6301–7.
75. Prösch S, Wendt CEC, Reinke P, Priemer C, Oppert M, Krüger DH, et al. A novel link between stress and human cytomegalovirus (HCMV) infection: Sympathetic hyperactivity stimulates HCMV activation. Virology [Internet]. Academic Press Inc.; 2000;272:357–65.
76. Mehta SK, Stowe RP, Feiveson AH, Tyring SK, Pierson DL. Reactivation and Shedding of Cytomegalovirus in Astronauts during Spaceflight. J Infect Dis. Oxford Academic; 2000;182:1761–4.
77. Spielmann G, McFarlin BK, O'Connor DP, Smith PJW, Pircher H, Simpson RJ. Aerobic fitness is associated with lower proportions of senescent blood T-cells in man. Brain Behav Immun. Brain Behav Immun; 2011;25:1521–9.
78. Fisher JP, Young CN, Fadel PJ. Autonomic adjustments to exercise in humans. Compr Physiol. Wiley-Blackwell Publishing Ltd; 2015;5:475–512.
79. Fritsche A, Stumvoll M, Grub M, Sieslack S, Renn W, Schmölling RM, et al. Effect of hypoglycemia on  $\beta$ -adrenergic sensitivity in normal and type 1 diabetic subjects. Diabetes Care. American Diabetes Association Inc.; 1998;21:1505–10.
80. Liston A, Gray DHD. Homeostatic control of regulatory T cell diversity [Internet]. Nat Rev Immunol. Nat Rev Immunol; 2014. p. 154–65.
81. Krüger K, Mooren FC. T cell homing and exercise. Exerc Immunol Rev [Internet]. 2007;13:37–54.
82. Dimitrov S, Benedict C, Heutling D, Westermann J, Born J, Lange T. Cortisol and epinephrine control opposing circadian rhythms in T cell subsets. Blood. Blood; 2009;113:5134–43.
83. Benschop RJ, Oostveen FG, Heijnen CJ, Ballieux RE.  $\beta$  2-Adrenergic stimulation causes detachment of natural killer cells from cultured endothelium. Eur J Immunol. Eur J Immunol; 1993;23:3242–7.
84. Navalta JW, Sedlock DA, Park KS. Effect of exercise intensity on exercise-induced lymphocyte apoptosis. Int J Sports Med. Int J Sports Med; 2007;28:539–42.
85. Pizza FX, Mitchell JB, Davis BH, Starling RD, Holtz RW, Bigelow N. Exercise-induced muscle damage: effect on circulating leukocyte and lymphocyte subsets. Med Sci Sport Exerc [Internet].

1995;27:363–70.

86. Pedersen BK. Muscle as a secretory organ. *Compr Physiol. Compr Physiol*; 2013;3:1337–62.

87. Pedersen BK, Febbraio MA. Muscles, exercise and obesity: Skeletal muscle as a secretory organ [Internet]. *Nat Rev Endocrinol. Nature Publishing Group*; 2012. p. 457–65.

88. Akbar AN. The convergence of senescence and nutrient sensing during lymphocyte ageing [Internet]. *Clin Exp Immunol. Blackwell Publishing Ltd*; 2017. p. 4–5.

89. Feldman RD, Limbird LE, Nadeau J, Robertson D, Wood AJJ. Alterations in Leukocyte  $\beta$ -Receptor Affinity with Aging. *N Engl J Med. Massachusetts Medical Society*; 1984;310:815–9.

## Figure Legends

Fig. 1. Flowchart of study selection.

Fig. 2. Forest plot of standardized mean difference (SMD) and 95% confidence interval for the overall effect immediately post exercise compared to baseline values. CMV: Cytomegalovirus; H-AT: hypoxic-absolute exercise; HC: hypobaric control; H-C: hypoxic resting; HE: hypobaric exercise; HI: High intensity; H-RT: hypoxic-relative exercise; HSV1: herpes simplex virus 1; LL: Lower limit of 95% confidence interval; LT: Lactate threshold; MI: Moderate intensity. NC: normobaric control; N-C: normoxic resting; NE: normobaric exercise; N-T: normoxic exercise; TD1: Type 1 diabetes; TR: Trained; UL: Upper limit of 95% confidence interval; UN: Untrained.

Fig. 3. . Forest plot of standardized mean difference (SMD) and 95% confidence interval for the overall effect 1h post exercise compared to baseline values. CMV: Cytomegalovirus; H-AT: hypoxic-absolute exercise; HC: hypobaric control; H-C: hypoxic resting; HE: hypobaric exercise; HI: High intensity; H-RT: hypoxic-relative exercise; HSV1: herpes simplex virus 1; LL: Lower limit of 95% confidence interval; LT: Lactate threshold; MI: Moderate intensity. NC: normobaric control; N-C: normoxic resting; NE: normobaric exercise; N-T: normoxic exercise; TD1: Type 1 diabetes; TR: Trained; UL: Upper limit of 95% confidence interval; UN: Untrained.

Fig. 4. Forest plot of standardized mean difference (SMD) and 95% confidence interval for the overall effect 2h post exercise compared to baseline values. CMV: Cytomegalovirus; H-AT: hypoxic-absolute exercise; HC: hypobaric control; H-C: hypoxic resting; HE: hypobaric exercise; HI: High intensity; H-RT: hypoxic-relative exercise; HSV1: herpes simplex virus 1; LL: Lower limit of 95% confidence interval; LT: Lactate threshold; MI: Moderate intensity. NC: normobaric control; N-C: normoxic resting; NE: normobaric exercise; N-T: normoxic exercise; TD1: Type 1 diabetes; TR: Trained; UL: Upper limit of 95% confidence interval; UN: Untrained.

Fig 5. The figure summarizes the lymphocytes count fold change from baseline to each time point for the senescent cells: CD8 T cells (in blue), CD4 T cell (in dark pink) and NK cells (in light pink and black centre). The position of the cells represents the SMD of each meta-analysis at each time-point post exercise.

**Table 1.** Characteristics of the studies included.

First author, year (subgroup)	Time points	Sex	Population characteristics			Exercise protocols					
			Mean age $\pm$ SD, or range	Training status	CMV serostatus	Equipment	Intensity	Duration	Intensity & duration	Cells analysed	Senescent marker
Azali Alamdari, 2018 [46] (post HC, HE, NC, NE)	0-15" & 1'	M	4 Groups mean: 21.8 $\pm$ 1.58 <sup>†</sup>	TR & NR*	NA	Treadmill	Until exhaustion	Incremental	Maximum & short	CD4+ & CD8+	KLRG1+
Krzywkowski, 2001 [54]	0-15" & 2'	M	37 (25-48)	TR	NA	Bicycle	75% VO <sub>2</sub> max	120min	Vigorous & long	CD4+ & CD8+	CD28-
Lavoy, 2017 [27]	0-15" & 1'	B	30.9 $\pm$ 5.0	TR	CMV- & CMV+	Bicycle	-5% BLT, +5% BLT & +15% BLT	30min	Moderate & moderate	CD4+	CD28- CD27-
Ross, 2018 [29]	0-15" & 1'	M	60-75 & 18-25	TR	CMV-	Bicycle	70% VO <sub>2</sub> peak	30min	Vigorous & Moderate	CD4+ & CD8+	CD28-
Wang, 2011 [32] (post H-AT, H-C, H-RT, N-C, N-T)	0-15" & 2'	M	5 groups mean: 22.46 $\pm$ 0.6 <sup>†</sup>	UN & TR	NA	Bicycle	Until exhaustion	Incremental	Maximum & short	CD4+ & CD8+	KLRG1+
Bigley, 2012 [56]	0-15" & 1'	M	2 Groups mean: 28.55 $\pm$ 5.35	NR	CMV- & CMV+	Bicycle	85% EMP	30min	Vigorous & Moderate	CD8+	KLRG1+
Curran, 2019 [53] (Control & TD1)	0-15" & 1'	M	2 Groups mean: 31 $\pm$ 7.15	NR	NA	Bicycle	80% VO <sub>2</sub> max	30min	Vigorous & Moderate	CD8+	EMRA
Lavoy, 2014 [55] (HSV1+ & HSV1-)	0-15" & 1'	M	4 Groups mean: 38.72 $\pm$ 15.22	TR	CMV- & CMV+	Bicycle	80-85% EMP	30min	Vigorous & Moderate	CD8+	KLRG1+CD28-
Spielmann, 2014 [28]	0-15" & 1'	M	2 Groups mean (Older): 55.35 $\pm$ 4.1 2 Groups mean (Younger): 28.5 $\pm$ 4.9	TR	CMV- & CMV+	Bicycle	80-85% PP	30min	Vigorous & Moderate	CD8+	KLRG1+CD28-
Turner, 2010 [51]	0-15" & 1'	M	35 $\pm$ 14	TR	CMV- & CMV+	Treadmill	80% VO <sub>2</sub> max	60min	Vigorous & long	CD8+	CD28- CD27-
Ingram, 2015 [49] (Disrupted)	0-15" & 1'	M	27 $\pm$ 8	TR	NA	Bicycle	265 $\pm$ 27 Watts	55:12 min	Vigorous & long	CD8+	KLRG1+



& Undisrupted sleep)


Curran, 2020 [47]	0-15" & 1'	M	2 Groups mean: 31 ± 7.15	NR	NA	Bicycle	80% VO <sub>2</sub> max	30min	Vigorous & Moderate	NK	CD57+
Wang, 2009 [57] (HI & MI)	0-15" & 2'	M	24.2 ± 1.2	UN	NA	Bicycle	Until exhaustion & 50% PD VO <sub>2</sub> max	Incremental & 30min	Maximum & short; Moderate & moderate	NK	CD57+

**Legend:** BLT: blood lactate threshold; C: Control; CMV: cytomegalovirus; EMP: estimated max power; EMRA: CD45RA+CCR7-CD28-CD27-; H-AT: hypoxic-absolute exercise; HC: hypobaric control; H-C: hypoxic resting; HE: hypobaric exercise; HI: High intensity; H-RT: hypoxic-relative exercise; HSV1: herpes simplex virus 1; MI: Moderate intensity. NA: not applicable (when did not use just one subgroup); NC: normobaric control; N-C: normoxic resting; NE: normobaric exercise; NR: not reported; ; NR\*: groups undergoing control period (NC and HC) were not analysed for training status, since they were trained at baseline and it was not clear how much untrained they became after intervention; N-T: normoxic exercise; PP: peak power; TD1: Type 1 diabetes; TR: Trained; UN: Untrained; USA: United States of America; VTWR: ventilatory threshold work rate †Standard error.


**Table 2.** Subgroups comparison for the effects of exercise on aged immune cells.

Time-point	Subgroups	Categories	K	SMD	LL	UL	p-value	Sample	p-diff
Immediately post exercise	<b>Training status</b>	TR	36	1.13	0.86	1.4	<b>&lt;0.001</b>	345	0.35
		UN	8	1.38	0.92	1.84	<b>&lt;0.001</b>	96	
	<b>CMV status</b>	CMV-	13	0.55	0.35	0.75	<b>&lt;0.001</b>	120	0.09
		CMV+	9	1.09	0.55	1.63	<b>&lt;0.001</b>	81	
	<b>Age</b>	Middle-aged	2	1.2	0.52	1.88	<b>0.001</b>	16	0.46
		Older adults	4	0.74	0.1	1.38	<b>0.02</b>	36	
		Young adults	44	1.14	0.91	1.37	<b>&lt;0.001</b>	449	
	<b>Intensity &amp; duration</b>	Vigorous & long <sup>b</sup>	6	0.4	0.14	0.66	<b>0.002</b>	66	<b>&lt;0.001</b>
		Moderate & moderate	3	0.38	0.02	0.73	<b>0.04</b>	33	
		Vigorous & Moderate <sup>ab</sup>	22	0.85	0.61	1.1	<b>&lt;0.001</b>	196	
Maximum & short <sup>a</sup>		19	1.81	1.45	2.16	<b>&lt;0.001</b>	206		
1h post exercise	<b>Age</b>	Middle-aged	2	-0.3	-0.8	0.17	0.2	16	0.81
		Older adults	4	-0.2	-0.5	0.13	0.23	36	
		Young adults	22	-0.2	-0.3	-0	<b>0.02</b>	209	
	<b>CMV status</b>	CMV-	13	-0.1	-0.3	0.1	0.37	120	0.02
		CMV+	9	-0.4	-0.7	-0.2	<b>&lt;0.001</b>	81	
	<b>Intensity &amp; duration</b>	Vigorous & long	4	-0.5	-0.8	-0.2	<b>&lt;0.001</b>	48	0.09
		Moderate & moderate	2	-0.3	-0.8	0.16	0.18	17	
	Vigorous & Moderate <sup>c</sup>	22	-0.1	-0.3	0.04	0.16	196		
2h post exercise	<b>Training status</b>	TR	14	-0.6	-1	-0.3	<b>&lt;0.001</b>	146	0.81
		UN	8	-0.6	-0.9	-0.4	<b>&lt;0.001</b>	96	
	<b>Intensity &amp; duration</b>	Vigorous & long	2	-0.4	-0.9	0.11	0.12	18	0.21
		Moderate & moderate	1	-0.2	-0.7	0.29	0.41	16	
		Maximum & short	19	-0.7	-1	-0.4	<b>&lt;0.001</b>	208	

**Legend:** SMD: Standardized mean difference; K: number of study groups; LL: Lower limit of 95% confidence interval; UL: Upper limit of 95% confidence interval; p-value: p-value for significance (<0.05) change of senescent cell counts within categories of subgroup; p-diff: p-value for significance (<0.05) change of senescent cell counts between categories of subgroup; TR: trained individuals; UN: untrained individuals; <sup>a</sup>: different of Moderate & moderate; <sup>b</sup>: Different of Maximum & short; <sup>c</sup>: different of Vigorous & long.




## 2.a



Heterogeneity tests:  $Q=39.18$ ,  $df=17$ ,  $p=0.001$ ,  $I^2=56.62\%$ ;  
Hypothesis test:  $Z=6.59$ ,  $p<0.001$ ; Egger test:  $p<0.001$ .

Δ Immediately post exercise


## 2.b



Heterogeneity tests:  $Q=111.94$ ,  $df=27$ ,  $p<0.001$ ,  $I^2=75.88\%$ ;  
Hypothesis test:  $Z=7.38$ ,  $p<0.001$ ; Egger test:  $p<0.001$ .

Δ Immediately post exercise


## 2.c



### 3.a

First author, year	Subgroup	SMD	LL	UL	p-Value	Total	Weight
Lavoy, 2017 [27]	CMV-/ +15%LT	-0.124	-0.820	0.571	0.726	8	11.96
Lavoy, 2017 [27]	CMV-/ +5%LT	-0.163	-0.860	0.535	0.647	8	11.89
Lavoy, 2017 [27]	CMV-/ -5%LT	-0.272	-0.978	0.433	0.449	8	11.62
Lavoy, 2017 [27]	CMV+/ +15%LT	-0.499	-1.192	0.194	0.158	9	12.06
Lavoy, 2017 [27]	CMV+/ +5%LT	-0.189	-0.848	0.470	0.573	9	13.32
Lavoy, 2017 [27]	CMV+/ -5%LT	-0.390	-1.067	0.288	0.260	9	12.60
Ross, 2018 [29]	Older	0.000	-0.620	0.620	1.000	10	15.06
Ross, 2018 [29]	Young	0.599	-0.111	1.308	0.098	9	11.50
<b>Summarized fixed effects</b>		<b>-0.132</b>	<b>-0.372</b>	<b>0.109</b>	<b>0.284</b>	<b>70</b>	<b>100</b>

SMD and 95% CI for senescent CD4 T cells




Heterogeneity tests:  $Q=6.07$ ,  $df=7$ ,  $p=0.53$ ,  $I^2=0\%$ ;  
Hypothesis test:  $Z=-1.07$ ,  $p=0.28$ ; Egger test:  $p=0.90$ .

### 3.b


First author, year	Subgroup	SMD	LL	UL	p-Value	Total	Weight
Bigley, 2012 [56]	CMV-	0.140	-0.429	0.709	0.629	12	7.38
Bigley, 2012 [56]	CMV+	-0.219	-0.920	0.482	0.540	8	4.85
Curran, 2019 [53]	Control	-0.311	-0.916	0.294	0.314	11	6.52
Curran, 2019 [53]	TD1	-0.325	-0.995	0.345	0.342	9	5.31
Ingram, 2015 [49]	disrupted sleep	-0.450	-1.100	0.201	0.175	10	5.64
Ingram, 2015 [49]	undisrupted sleep	-0.361	-1.001	0.278	0.268	10	5.83
Lavoy, 2014 [55]	CMV+HSV1-	-0.569	-1.316	0.178	0.135	8	4.28
Lavoy, 2014 [55]	CMV+HSV1+	-0.333	-1.045	0.379	0.359	8	4.71
Lavoy, 2014 [55]	CMV-HSV1-	-0.132	-0.828	0.564	0.709	8	4.92
Lavoy, 2014 [55]	CMV-HSV1+	-0.038	-0.731	0.655	0.914	8	4.96
Ross, 2018 [29]	Older	-0.250	-0.880	0.379	0.436	10	6.02
Ross, 2018 [29]	Young	-0.156	-0.813	0.501	0.642	9	5.52
Spielmann, 2014 [28]	CMV-/older	-0.183	-0.882	0.516	0.608	8	4.89
Spielmann, 2014 [28]	CMV-/young	-0.165	-0.863	0.533	0.643	8	4.90
Spielmann, 2014 [28]	CMV+/older	-0.433	-1.157	0.292	0.242	8	4.54
Spielmann, 2014 [28]	CMV+/young	-0.313	-1.023	0.397	0.387	8	4.74
Turner, 2010 [51]	CMV-	-0.274	-0.808	0.259	0.314	14	8.38
Turner, 2010 [51]	CMV+	-0.793	-1.394	-0.193	0.010	14	6.61
<b>Summarized fixed effects</b>		<b>-0.283</b>	<b>-0.438</b>	<b>-0.129</b>	<b>0.000</b>	<b>171</b>	<b>100</b>

SMD and 95% CI for senescent CD8 T cells





### 3.c




Heterogeneity tests:  $Q=0.03$ ,  $df=1$ ,  $p=0.87$ ,  $I^2=0\%$ ;  
Hypothesis test:  $Z=2.53$ ,  $p=0.011$ ; Egger test: NA.

Δ 1h post exercise

## 4.a


First author, year	Subgroup	SMD	LL	UL	p-Value	Total	Weight
Azali Alamdari, 2018 [46]	HC	-1.064	-1.804	-0.324	0.005	11	10.40
Azali Alamdari, 2018 [46]	HE	-0.764	-1.407	-0.121	0.020	12	11.37
Azali Alamdari, 2018 [46]	NC	-0.687	-1.344	-0.030	0.040	11	11.23
Azali Alamdari, 2018 [46]	NE	-1.108	-1.826	-0.389	0.003	12	10.61
Krzywkowski, 2001 [54]		0.000	-0.653	0.653	1.000	9	11.27
Wang, 2011 [32]	H-AT	-1.461	-2.352	-0.570	0.001	10	8.97
Wang, 2011 [32]	H-C	-1.739	-2.722	-0.757	0.001	10	8.20
Wang, 2011 [32]	H-RT	-1.917	-2.961	-0.873	0.000	10	7.71
Wang, 2011 [32]	N-C	-1.579	-2.508	-0.650	0.001	10	8.64
Wang, 2011 [32]	N-T	-0.073	-0.693	0.548	0.818	10	11.60
<b>Summarized random effects</b>		<b>-0.958</b>	<b>-1.361</b>	<b>-0.556</b>	<b>0.000</b>	<b>105</b>	<b>100</b>



Heterogeneity tests:  $Q=24.97$ ,  $df=9$ ,  $p=0.003$ ,  $I^2=63.96\%$ ;  
Hypothesis test:  $Z=-4.67$ ,  $p<0.001$ ; Egger test:  $p<0.001$ .


## 4.b



First author, year	Subgroup	SMD	LL	UL	p-Value	Total	Weight
Azali Alamdari, 2018 [46]	HC	-0.605	-1.248	0.038	0.065	11	10.67
Azali Alamdari, 2018 [46]	HE	-0.098	-0.665	0.469	0.735	12	11.40
Azali Alamdari, 2018 [46]	NC	-0.714	-1.376	-0.052	0.034	11	10.49
Azali Alamdari, 2018 [46]	NE	-0.641	-1.262	-0.020	0.043	12	10.88
Krzywkowski, 2001 [54]		-1.000	-1.800	-0.200	0.014	9	9.22
Wang, 2011 [32]	H-AT	-0.459	-1.111	0.192	0.167	10	10.59
Wang, 2011 [32]	H-C	-0.983	-1.738	-0.228	0.011	10	9.62
Wang, 2011 [32]	H-RT	-0.805	-1.518	-0.092	0.027	10	10.01
Wang, 2011 [32]	N-C	-0.916	-1.655	-0.178	0.015	10	9.77
Wang, 2011 [32]	N-T	1.875	0.846	2.904	0.000	10	7.36
<b>Summarized random effects</b>		<b>-0.488</b>	<b>-0.880</b>	<b>-0.096</b>	<b>0.015</b>	<b>105</b>	<b>100</b>



Heterogeneity tests:  $Q=28.05$ ,  $df=9$ ,  $p=0.001$ ,  $I^2=67.92\%$ ;  
 Hypothesis test:  $Z=-2.44$ ,  $p=0.015$ ; Egger test:  $p=0.29$ .

### 4.c





## Supplementary Material

### Pubmed Search

("Cellular Senescence"[mh] OR "cell ageing"[tiab] OR "cellular ageing"[tiab] OR "cell aging"[tiab] OR "Senescence-Associated Secretory Phenotype"[tiab] OR "Senescence Associated Secretory Phenotype"[tiab] OR ("cellular"[tiab] AND "senescence"[tiab]) OR "cellular senescence"[tiab] OR "cell senescence"[tiab] OR "CD28 Antigens"[mh] OR "CD28"[tiab] OR "CD57 Antigens"[mh] OR "CD57"[tiab] OR "KLRG1 protein, human" [Supplementary Concept] OR "KLRG1"[tiab] OR "Immunosenescence"[mh] OR "Immunosenescence"[tiab]) AND ("exercise"[mh] OR "exercise"[tiab] OR "physical training"[tiab] OR "physical activity"[tiab]) NOT (Review [Publication Type] OR "Review Literature as Topic" [mh])

### Web of Science Search

(TS= ("Cellular Senescence") OR TS= ("cell ageing") OR TS= ("cellular ageing") OR TS= ("cell aging") OR TS= ("Senescence-Associated Secretory Phenotype") OR TS= ("Senescence Associated Secretory Phenotype") OR TS= ("cellular senescence") OR TS= ("cell senescence") OR TS= ("cell senescence") OR TS= ("CD28 antigens") OR TS= ("CD57 Antigens") OR TS= ("KLRG1 protein, human") OR TS= ("Immunosenescence") OR AB= ("Cellular Senescence") OR AB= ("cell ageing") OR AB= ("cellular ageing") OR AB= ("cell aging") OR AB= ("Senescence-Associated Secretory Phenotype") OR AB= ("Senescence Associated Secretory Phenotype") OR AB= ("cellular senescence") OR AB= ("cell senescence") OR AB= ("cell senescence") OR AB= ("CD28 antigens") OR AB= ("CD57 Antigens") OR AB= ("KLRG1 protein, human") OR AB= ("Immunosenescence")) AND (TS= ("exercise") OR AB= ("exercise") OR AB= ("physical activity") OR AB= ("physical training") OR TI= ("exercise") OR TI= ("physical activity") OR TI= ("physical training")) AND LANGUAGE: (English) AND TYPE OF DOCUMENT: (Article)

### Embase Search

(('cell aging'/exp OR 'cell aging':ab,ti OR 'Senescence Associated Secretory Phenotype'/exp OR 'CD28 antigen'/exp OR 'CD57 antigen'/exp OR 'klrg1 protein'/exp OR 'immunosenescence'/exp OR cd28:ab,ti OR cd57:ab,ti OR klr1:ab,ti OR immunosenescence:ab,ti OR 'cellular aging':ab,ti OR 'Senescence-Associated Secretory Phenotype':ab,ti OR "Senescence Associated Secretory Phenotype":ab,ti OR ('cellular':ab,ti AND 'senescence':ab,ti) OR 'cellular senescence':ab,ti OR 'cell senescence':ab,ti)) AND ('kinesiotherapy'/exp OR 'exercise'/exp OR 'exercise':ab,ti OR 'physical activity':ab,ti OR 'physical training':ab,ti) AND ([article]/lim OR [article in press]/lim OR [data papers]/lim) AND 'human'/de

### Scopus Search

(TITLE-ABS-KEY("Cellular Senescence") OR TITLE-ABS-KEY("cell ageing") OR TITLE-ABS-KEY("cellular ageing") OR TITLE-ABS-KEY("cell aging") OR TITLE-ABS-KEY("Senescence-Associated Secretory Phenotype") OR TITLE-ABS-KEY("Senescence Associated Secretory Phenotype") OR TITLE-ABS-KEY("cellular senescence") OR TITLE-ABS-KEY("cell senescence") OR TITLE-ABS-KEY("CD28") OR TITLE-ABS-KEY("CD57") OR TITLE-ABS-KEY("KLRG1") OR TITLE-ABS-KEY("Immunosenescence")) AND (TITLE-ABS-KEY("exercise") OR TITLE-ABS-KEY("physical activity") OR TITLE-ABS-KEY("physical training")) AND ( LIMIT-TO ( DOCTYPE , "ar" )) AND ( LANGUAGE ( DOCTYPE , "ar" ))

### Cochrane Search

(("Cellular Senescence"):ti,ab,kw OR ("cell ageing"):ti,ab,kw OR ("cellular ageing"):ti,ab,kw OR ("cell aging"):ti,ab,kw OR ("Senescence-Associated Secretory Phenotype"):ti,ab,kw OR ("Senescence Associated Secretory Phenotype"):ti,ab,kw OR ("cell senescence"):ti,ab,kw OR ("CD28"):ti,ab,kw OR ("CD57"):ti,ab,kw

OR ("KLRG1"):ti,ab,kw OR ("Immunosenescence"):ti,ab,kw) AND (("exercise"):ti,ab,kw OR ("physical training"):ti,ab,kw OR ("physical activity"):ti,ab,kw)