

Parental smoking and the risk of middle ear disease in children

Jones, Laura L; Hassanien, Amal; Cook, Derek G; Britton, John; Leonardi-Bee, Jo

DOI:

[10.1001/archpediatrics.2011.158](https://doi.org/10.1001/archpediatrics.2011.158)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Jones, LL, Hassanien, A, Cook, DG, Britton, J & Leonardi-Bee, J 2012, 'Parental smoking and the risk of middle ear disease in children: a systematic review and meta-analysis', *Archives of Pediatrics and Adolescent Medicine*, vol. 166, no. 1, pp. 18-27. <https://doi.org/10.1001/archpediatrics.2011.158>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the author's accepted manuscript of the article published as above.

Eligibility for repository checked February 2015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **Parental Smoking and the Risk of Middle Ear Disease in Children**

2 ***A Systematic Review and Meta-analysis***

3

4 *Laura L. Jones, PhD; Amal Hassanien, MSc; Derek G. Cook, PhD; John Britton, FRCP; and Jo*

5 *Leonardi-Bee, PhD*

6

7 Author Affiliations: UK Centre for Tobacco Control Studies, Division of Epidemiology and Public

8 Health, University of Nottingham, Nottingham, UK (Dr Jones, Ms Hassanien, Professor Britton

9 and Dr Leonardi-Bee) and Division of Population Health Sciences and Education, St George's

10 University of London, London, UK (Professor Cook).

11

12 **Word Count:** (excluding title, abstract, references, figure legends, and tables): 2892

13

14

15

16 **Objectives:** A systematic review and meta-analysis of studies of the association between
17 passive smoking and middle ear disease (MED) in children.

18 **Data Sources:** MEDLINE, EMBASE and CAB Abstracts (to December 2010) and reference lists.

19 **Study Selection:** Sixty-one epidemiological studies of children assessing the effect of passive
20 smoke exposure on outcomes of MED were included. Articles were reviewed, data extracted
21 and synthesized by two researchers.

22 **Main Outcome Exposures:** Children's passive smoke exposure including: maternal smoking
23 during and after pregnancy, paternal and household.

24 **Main Outcome Measures:** Middle ear disease in children.

25 **Results:** Living with a smoker was associated with an increased risk of MED in children; by an
26 odds ratio (OR) of 1.62 (95% confidence interval (CI) 1.33 to 1.97) for maternal post-natal
27 smoking and by 1.37 (95% CI 1.25 to 1.50) for any household member smoking. Pre-natal
28 maternal smoking (OR 1.11, 95% CI 0.93 to 1.31) and paternal smoking (OR 1.24, 95% CI 0.98 to
29 1.57) were associated with a non-significant increase in the risk of MED in children. The
30 strongest effect was on the risk of surgery for MED, where maternal post-natal smoking
31 increased the risk by an OR of 1.86 (95% CI 1.31 to 2.63) and paternal smoking by 1.83 (95% CI
32 1.61 to 2.07).

33 **Conclusions:** Passive smoke exposure, particularly by the mother, significantly increases the risk
34 of MED in childhood; this risk is particularly strong for MED requiring surgery. We have shown
35 that 130,200 of child MED episodes per year are directly attributable to passive smoke
36 exposure.

37

38 Middle ear disease (MED) is a common illness among children that accounts for a large number
39 of physician visits, and which if untreated, can cause significant disability though hearing
40 impairment.¹ It is estimated that around 10% of children have three episodes of acute otitis
41 media (AOM) before their first birthday,² whilst middle ear effusion is the most common
42 reason for admission of young children to hospitals for surgery, putting a heavy financial
43 burden on health care services.³ Furthermore, adenoidectomy and particularly
44 adenotonsillectomy, which are surgical treatments for otitis media with effusion (OME), have
45 been associated with significant morbidity and mortality, including that arising from surgery.⁴
46 Middle ear effusion is associated with hearing loss in children, which may lead to delayed
47 linguistic and cognitive development.³ The prevalence of MED is higher among children with
48 learning impairment.⁵

49 In 1998, a systematic review by Strachan and Cook of papers published up to 1996
50 found a significant association between parental smoking and MED.⁶ However, the authors
51 concluded that few studies had compared the effect of smoking by the mother and father and
52 none had compared the effect of pre- and post-natal tobacco smoke exposure to MED. This
53 original review was commissioned for a UK government Scientific Committee on Tobacco and
54 Health (SCOTH),⁷ and was subsequently updated as part of the 2006 US Surgeon General's
55 report on the effects of involuntary exposure to tobacco smoke, which concluded that there
56 was sufficient evidence to infer a causal relationship between parental smoking and otitis
57 media in childhood.⁸ Since these early reviews of papers published up until 2001, the evidence
58 base on the association between parental smoking and MED in childhood has significantly
59 increased. To date however, these new studies have not been subject to meta-analysis. We

60 have therefore carried out a systematic review and meta-analysis of the epidemiological data to
61 provide contemporary estimates of the effects of smoking by parents and other household
62 members on the risk of middle ear disease in childhood. The work was carried out as part of a
63 more extensive review of the effects of passive smoking in children, for the Royal College of
64 Physicians.⁹

65

66 **METHODS**

67 **Systematic review methods**

68 Any analytical epidemiological study assessing the effect of passive smoke exposure (including
69 household smoking, paternal¹ smoking, maternal smoking after during and after pregnancy)
70 were included in the review. Outcomes of interest were MED, sub-divided into: middle ear
71 infections (including acute otitis media, otitis media with effusion, recurrent otitis media,
72 chronic otitis media); hearing impairment (including hearing loss, deafness, glue ear), and
73 surgery related to MED (including adenotonsillectomy, tonsillectomy, adenoidectomy, and
74 grommet insertion).

75 We searched MEDLINE, EMBASE and CAB Abstracts (from 1997 to December 2010),
76 using the keywords *Tobacco smoke, cigarette smoking, passive smoking, parental smoking,*
77 *maternal smoking, parental smoking, environmental tobacco smoking, second hand smoke,*
78 *children, infants, adolescents, pediatric, otitis media with effusion, deafness, adenoidectomy,*
79 *middle ear disease, adenotonsillectomy, acute otitis media, recurrent otitis media, middle ear*

¹ Please note that it was not possible in the current (or previous) study to identify studies that measured paternal smoking independently of maternal smoking i.e. father smokes but mother does not.

80 *effusion, glue ear, otitis, tympanum, tonsil, otitis interna*. Hand searching of reference lists was
81 also performed. No language restrictions were imposed during the searches, however, to be
82 consistent with the original review,⁶ we report only those studies published in English.

83 Titles and abstracts from the identified studies were reviewed independently by two of
84 three authors (AH & JLB or LLJ & JLB) to identify eligible studies. The full text of studies
85 potentially eligible for inclusion were sought and assessed independently by two of the three
86 authors. For included studies, two of the three authors independently extracted data using a
87 standard data extraction form and assessed methodological quality using the Cochrane
88 Collaboration Non-Randomized Studies Working Group recognized Newcastle-Ottawa Quality
89 Assessment Scale¹⁰ based on the following: selection of cases and controls, or cohort;
90 comparability of the cases and controls, or cohort, and assessment of exposure/outcome. A
91 score of six or more was chosen *a priori* to indicate higher methodological quality. In addition,
92 all studies included in the previous review⁶ were assessed for methodological quality using the
93 same methods. Disagreements were resolved through discussion.

94

95 **Statistical analysis**

96 Where possible, the data extracted were unadjusted odds ratios (OR), or in preference, OR
97 adjusted for potential confounding variables. Measures of uncertainty were also extracted
98 either in the form of standard errors or 95% confidence intervals (CI). Pooled estimates of
99 measures of association were estimated using random effect meta-analyses and presented as
100 pooled OR with 95% CI. Heterogeneity was assessed using recognized methods (I^2).¹¹ Random
101 effect meta-regression analyses were conducted to investigate the reasons for any

102 heterogeneity between epidemiological studies based on definition of MED (middle ear
103 infection, surgery and hearing impairment), methodological quality (higher versus lower), study
104 design (cohort, cross-sectional and case-control), ascertainment of passive smoke exposure
105 (biochemical vs. self-report) and by date of publication. Exposure was defined as household,
106 paternal and maternal; maternal was split into pre- and post-natal. Data were analyzed using
107 Review Manager, version 5.0.23 ((RevMan), Copenhagen, The Nordic Cochrane Centre, The
108 Cochrane Collaboration) and STATA MP/11.0 for Windows (StataCorp LP, 4905 Lakeway Drive,
109 College Station, TX 77845, USA). P values less than 0.05 were considered statistically significant.
110 The analysis was performed in accordance with the Meta-Analysis of Observational Studies in
111 Epidemiology (MOOSE) guidelines.¹²

112

113 **Population attributable fraction estimation**

114 We estimated the proportion of children in England who live in a household in which at least
115 one person smokes using data from the Health Survey for England,⁹ and used the formula $p(OR-$
116 $1)/[p(OR-1)+1]$, in which p is the proportion of the cohort exposed to passive smoking, and OR
117 the odds ratio for MED in children where a member of the household smokes, to estimate the
118 proportion of children whose MED is attributable to household smoking exposure. We then
119 used national MED prevalence⁹ data for England and Wales to estimate the number of disease
120 episodes generated as a result of household passive smoke exposure.

121

122 RESULTS

123 From 360 titles published since 1997 identified in the literature search, 55 abstracts were
124 deemed potentially eligible, and of these, 36 were included following the full-text review
125 (Figure 1). The reasons for exclusion were: not having a comparative group without the
126 outcome,¹³⁻¹⁵ not assessing passive smoke as an exposure,¹⁶ not assessing MED as an
127 outcome,^{17, 18} being published in a language other than English,¹⁹⁻²⁵ only reporting statistical
128 significance (p value) of the result without data,²⁶⁻²⁹ or only having passive smoking data as a
129 confounder in the analysis.³⁰ Combining the results from this updated search with the previous
130 review (25 studies) resulted in 61 epidemiological studies (Table 1, Figure 1).

131 Of the 61 studies included, 15³¹⁻⁴⁵ were a cross-sectional survey, 23⁴⁶⁻⁶⁸ were case-
132 control studies and 23⁶⁹⁻⁹¹ were cohort studies. Seventeen different disease outcomes were
133 reported within these studies: acute infection and serious otitis media,⁸³ acute otitis media,<sup>32,
134 35, 38, 68, 69, 72, 86, 87, 91</sup> chronic suppurative otitis media,⁵⁹ earache,⁸² glue ear,⁵¹ hearing loss,⁷⁰
135 middle ear disease,^{46, 71} otitis media,^{36, 39, 43, 48, 74, 81} otitis media with effusion,<sup>31, 34, 40, 42, 44, 45, 47,
136 50, 53, 66, 67, 75, 76, 78, 80, 84, 90</sup> otitis prone,⁶³ recurrent otitis media,^{49, 60, 64, 73, 77, 79, 88, 89} suppurative
137 otitis media,³³ surgery (adenoids/tonsils),⁴¹ surgery (otitis media with effusion),^{52, 57, 58, 61, 85}
138 surgery (otitis media),^{37, 55, 56} surgery (recurrent otitis media),⁶² and surgery (tonsils).^{54, 65}

139

140 Methodological Quality of Studies and Publication Bias

141 The methodological quality of the 61 studies included in the meta-analysis, as judged by the
142 Newcastle-Ottawa scale score, is presented in Table 1. The overall median score was 5.5 (range
143 2 to 8). Using the *a priori* threshold of six to indicate high methodological quality, we judged 34

144 of the studies to be of high quality; the remaining 27 were deemed to be of lower quality
145 primarily due to a combination of a lack of biochemical validation of passive smoke exposure,
146 lack of representativeness of the study sample, and/or lack of adjusted analyses. There was no
147 evidence of publication bias identified from funnel plots. The funnel plot for household
148 exposure and the risk of MED is presented in Figure 2.

149

150 **Effects of maternal post-natal smoking**

151 Meta-analysis of the 20 studies of post-natal maternal smoking showed a statistically significant
152 increase in the risk of MED in childhood by 1.62 (95% CI 1.33 to 1.97). High levels of
153 heterogeneity were present in this analysis ($I^2 = 93\%$). Pooled estimates for each of the
154 outcome categories showed that the increase in risk of MED was driven predominantly by an
155 increase in the risk of surgery for MED (OR 1.86, 95% CI 1.31 to 2.63; 5 studies; Figure 3) and to
156 a lesser extent hearing impairment (OR 1.74, 95% CI 1.08 to 2.81; 1 study) and middle ear
157 infection (OR 1.53, 95% CI 1.22 to 1.92; 14 studies). In a meta-regression based on method of
158 ascertainment of passive smoke exposure, studies that used self-reported data showed a higher
159 increase in disease risk (OR 1.70, 95% CI 1.29 to 2.25; 17 studies), than studies that used
160 biochemical validation (OR 1.29, 95% CI 0.86 to 1.94; 3 studies). In a sub-group analysis based
161 on study design, case-control studies showed a statistically significant increase in the risk of
162 MED in children (OR 2.09, 95% 1.19 to 3.66; 10 studies), unlike cohort (OR 1.19, 95% CI 0.94 to
163 1.49; 6 studies) and cross-sectional (OR 1.28, 95% CI 0.88 to 1.86; 4 studies) study designs
164 which were not statistically significantly associated with an increase in disease risk. Similar
165 pooled estimates were also shown for the meta-regression analysis based on methodological

166 quality and date of publication (see Table 2a). In a multiple meta-regression adjusting for study
167 design, publication date, ascertainment and methodological quality, none of the factors
168 independently predicted the odds ratio for maternal post-natal smoking.

169

170 **Effects of maternal pre-natal smoking**

171 All of the six studies of pre-natal maternal smoking were indentified from the updated search,
172 as they were published after 1996. Pre-natal maternal smoking was not associated with a
173 statistically significant increase in the risk of MED (OR 1.11, 95% CI 0.93 to 1.31; 6 studies);
174 however, high levels of heterogeneity were seen between the studies ($I^2 = 79\%$). Similarly, none
175 statistically significant pooled estimates were also seen for meta-regression analyses stratified
176 by study design, ascertainment of smoking status and methodological quality (see Table 2a).

177

178 **Effects of paternal smoking**

179 Exposure to paternal smoking was associated with a non-significant ($p=0.07$) increase in the
180 odds of middle ear disease in childhood by 1.24 (95% CI 0.98 to 1.57; 12 studies). Very high
181 levels of heterogeneity were seen in the analysis ($I^2 = 87\%$). Sub-group analysis based on the
182 definition of outcome showed that the increased risk of disease was due to a strong association
183 between paternal passive smoke exposure and the risk of surgery for MED (OR 1.83, 95% CI
184 1.61 to 2.07; 4 studies; Figure 4). The association between paternal smoking and middle ear
185 infection was not statistically significant (OR 1.06, 95% CI 0.91 to 1.24; 8 studies, $p=0.47$).
186 Similar pooled estimates were also seen for meta-regression analyses stratified by study design,
187 ascertainment of smoking status, date of publication and methodological quality (see Table 2b).

188 In a multiple meta-regression adjusting for study design, publication date, ascertainment and
189 methodological quality, none of the factors independently predicted the odd ratio for paternal
190 smoking.

191

192 **Effects of household smoking**

193 A pooled estimate derived from the 49 studies which defined exposure as household smoking
194 (the study by Jacoby et al.⁸¹ is shown in the Forest plot as two separate entries given the
195 differing estimates reported for the two samples: aboriginal vs. non-aboriginal) demonstrated a
196 statistically significant increase in the risk of middle ear disease by an OR of 1.37 (95% CI 1.25 to
197 1.50; 49 studies). High levels of heterogeneity were seen between the studies ($I^2 = 76\%$). Sub-
198 group analysis based on the definition of outcome showed that the increase in risk was mainly
199 attributable to a increase in risk of surgery for MED (OR 1.62, 95% CI 1.32 to 1.98; 11 studies;
200 Figure 5) and to a lesser extent middle ear infection (OR 1.32, 95% CI 1.20 to 1.45; 38 studies).
201 Meta-regression analysis based on study design showed varied pooled estimates, with case-
202 control studies showing the highest increase in disease risk (OR 1.55, 95% CI 1.35 to 1.77; 18
203 studies), followed by cross sectional studies (OR 1.33, 95% CI 1.10 to 1.60; 13 studies) and
204 cohort studies (OR 1.27, 95% CI 1.13 to 1.43; 18 studies). Similar pooled estimates were also
205 seen for analyses stratified by ascertainment of smoking status, date of publication and
206 methodological quality (see Table 2b). In a multiple meta-regression adjusting for study design,
207 publication date, ascertainment and methodological quality, none of the factors independently
208 predicted the odd ratio for household smoking.

209

210 **Population attributable fraction**

211 Health survey for England data indicate that in 2007, around 22% of children aged up to 15
212 years lived in a household in which someone smokes.{Royal College of Physicians, 2010 #561}
213 Using the odds ratio for household smoking (1.37) as the estimated relative risk of developing
214 MED, the proportion of children developing MED likely to be attributable to exposure to
215 smoking in the home is estimated at 7.5%. In 2008 there were about 1,735,710 episodes of
216 MED in children under the age of 16 years in the UK.⁹ A 7.5% attributable fraction translates
217 into approximately 130,200 new episodes of MED arising from exposure to smoking in the
218 home in the UK.

219

220 **DISCUSSION**

221 Middle ear disease is a significant cause of morbidities in children and has been shown to be
222 associated with parental passive smoke exposure.⁶ This relationship has been further explored
223 in the current systematic review and meta-analysis, providing novel findings which suggest that
224 maternal post-natal smoking, rather than maternal pre-natal or paternal smoking has the
225 strongest influence on disease risk. This may suggest that the effect is due to ambient smoke
226 pollution from the child's close proximity to the primary caregiver, not to development effects.
227 However, it important to consider that only six pre-natal studies met the inclusion criteria in the
228 current study and hence may be underpowered to detect an association. Therefore, further
229 well-conducted research studies are needed. We additionally found that smoking by any
230 household member was statistically significantly associated with an increased risk of disease in

231 children, which translates to an additional 130,200 episodes of MED per year in the UK, which
232 are directly attributable to passive smoking.

233 From meta-regression analysis exploring the different MED outcomes (middle ear
234 infection, surgery for middle ear infection, hearing impairment or hearing loss), we found that
235 the effect of passive smoke exposure was strongest for surgery for MED, with an increased risk
236 of 1.86 for maternal post-natal, 1.83 for paternal and 1.62 for household smoking. In addition,
237 maternal post-natal smoking was shown to increase the risk of hearing impairment by an odds
238 of 1.74 (95% CI 1.08 to 2.81), although this estimate is based on only one study of high
239 methodological quality.⁵¹

240 Our findings are likely to be representative estimates of the true effects of passive
241 smoking on the risk of MED in children since they are based on results of a comprehensive
242 search, including data identified through hand searching of reference lists and previous reviews.
243 However, there are limitations to this review. We elected to keep methods consistent with the
244 original strategy⁶ and only included studies written in English in the meta-analyses. Additionally,
245 we were inevitably limited in the range of confounding factors that could be adjusted for in our
246 analyses. Although the high quality studies generally adjusted for maternal age and
247 socioeconomic status; other potential confounders, such as smoking by other individuals in the
248 household, were not consistently adjusted for in the analyses of the individual effects of
249 paternal and maternal smoking. A further limitation of this meta-analysis was that high levels of
250 heterogeneity were observed for some comparisons, which suggests that there are unexplained
251 reasons for variation in the findings between studies over and above chance, which may result
252 in potentially misleading summary estimates. We investigated reasons for heterogeneity by

253 performing meta-regression analyses; however, these analyses revealed relatively consistent
254 findings. Generally, the pooled results did not differ appreciably between studies of different
255 methodological quality, publication date or study design.

256

257 **CONCLUSIONS**

258 This study confirms that household smoking, in particular, maternal post-natal smoking causes
259 a statistically significant increase in the risk of MED in childhood, and identifies that one of the
260 main consequences of children's exposure to passive smoke is the significant increase in risk of
261 having to have surgery for chronic MED. Surgical treatments for otitis media, such as grommet
262 insertion, have been shown to be questionable in their effectiveness, high risk and resource and
263 cost intensive.⁹² Therefore, primary prevention through the reduction of risk factors such as
264 exposure to passive smoke, are key to reducing the burden of MED in childhood. Although
265 evidence is emerging to suggest that the incidence of MED has been declining in recent years in
266 England,⁹³ perhaps as a reflection of a reduction in the number of parents who smoke, MED is
267 still a major public and child health concern, with a total of 1.74 million episodes estimated in
268 the UK each year.⁹ We have shown that 7.5% of these episodes (130,200) are directly
269 attributable to passive smoke exposure in the home, all of which are avoidable. The findings
270 from this study should encourage renewed efforts to promote smoke free environments for
271 children.

272

273 **Correspondence:** Dr Laura L. Jones, Division of Epidemiology and Public Health, University of
274 Nottingham, City Hospital Campus, Hucknall Road, Nottingham, NG5 1PB, UK
275 (laura.jones@nottingham.ac.uk)

276 **Author Contributions:** Dr Leonardi-Bee had full access to all the data in the study and takes
277 responsibility for the integrity of the data and the accuracy of the data analysis. *Study concept*
278 *and design:* Britton and Leonardi-Bee. *Acquisition of data:* Jones, Hassanien and Leonardi-Bee.
279 *Analysis and interpretation of data:* Jones and Leonardi-Bee. *Drafting of the manuscript:* Jones,
280 Hassanien and Leonardi-Bee. *Critical revision of the manuscript for important intellectual*
281 *content:* Jones, Hassanien, Cook, Britton and Leonardi-Bee. *Statistical analysis:* Leonardi-Bee.
282 *Administrative, technical, and material support:* Jones and Leonardi-Bee. *Study supervision:*
283 Jones and Leonardi-Bee.

284 **Financial Disclosure:** None reported.

285 **Funding/Support:** This work was supported by project grant C1512/A11160 from Cancer
286 Research UK, and by core funding to the UK Centre for Tobacco Control Studies
287 (www.ukctcs.org) from the British Heart Foundation, Cancer Research UK, Economic and Social
288 Research Council, Medical Research Council, and the Department of Health, under the auspices
289 of the UK Clinical Research Collaboration.
290

291 **REFERENCES**

- 292 1. World Health Organization. Protection from exposure to second hand smoke: policy
293 recommendations. 2007;
294 http://whqlibdoc.who.int/publications/2007/9789241563413_eng.pdf. Accessed
295 13/7/10.
- 296 2. McConnell TH. *The nature of disease: pathology for health professionals*: Lippincott
297 Williams and Wilkins; 2007.
- 298 3. US EPA. *Respiratory Health Effects of Passive Smoking (Also Known as Exposure to*
299 *Secondhand Smoke or Environmental Tobacco Smoke ETS)*: US Environmental Protection
300 Agency, Office of Research and Development, Office of Health and Environmental
301 Assessment, Washington, DC, EPA/600/6-90/006F; 1992.
- 302 4. Alpher C, Bluestone, CD, Casselbrant, ML, Dohar, JE, Mandell, EM. *Advanced therapy of*
303 *otitis media*. New York: BC Decker Inc; 2004.
- 304 5. Freeman BA, Parkins C. The prevalence of middle ear disease among learning impaired
305 children. Does a higher prevalence indicate an association? *Clin Pediatr. Apr*
306 *1979;18(4):205-212*.
- 307 6. Strachan DP, Cook DG. Health effects of passive smoking. 4. Parental smoking, middle
308 ear disease and adenotonsillectomy in children. *Thorax*. January 1, 1998 1998;53(1):50-
309 56.
- 310 7. Department of Health. Department of Health and Social Services Northern Ireland. The
311 Scottish Office Department of Health and Welsh Office. *Report of the Scientific*
312 *Committee on Tobacco and Health*. London: The Stationery Office; 1998.

- 313 8. U.S. Department of Health and Human Services. *The Health Consequences of Involuntary*
314 *Exposure to Tobacco Smoke: A Report of the Surgeon General.*: Centers for Disease
315 Control and Prevention, National Center for Chronic Disease Prevention and Health
316 Promotion, Office on Smoking and Health; 2006.
- 317 9. Royal College of Physicians. Passive smoking and children. A report by the Tobacco
318 Advisory Group. London: RCP; 2010.
- 319 10. Wells G, Shea B, O'Connell D, et al. Newcastle-Ottawa scale (NOS) for assessing the
320 quality of non randomised studies in meta-analysis. .
321 http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm. Accessed 23/10/2009.
- 322 11. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. Jun
323 15 2002;21(11):1539-1558.
- 324 12. Stroup DF, Thacker SB, Olson CM, Glass RM, Hutwagner L. Characteristics of meta-
325 analyses related to acceptance for publication in a medical journal. *J Clin Epidemiol*. Jul
326 2001;54(7):655-660.
- 327 13. Damoiseaux RA, Rovers MM, Van Balen FA, Hoes AW, de Melker RA. Long-term
328 prognosis of acute otitis media in infancy: determinants of recurrent acute otitis media
329 and persistent middle ear effusion. *Fam Pract*. Feb 2006;23(1):40-45.
- 330 14. Praveen CV, Terry RM. Does passive smoking affect the outcome of grommet insertion
331 in children? *J Laryngol Otol*. Jun 2005;119(6):448-454.
- 332 15. Tariq S, Memon, IA. Acute otitis media in children. *JCPSP*. 1999;9(12):507-510.

- 333 16. Homoe P. Otitis media in Greenland. Studies on historical, epidemiological,
334 microbiological, and immunological aspects. *Int J Circumpolar Health*. 2001;60 Suppl 2:1-
335 54.
- 336 17. Kolossa-Gehring M, Becker K, Conrad A, et al. German Environmental Survey for
337 Children (GerES IV)--first results. *Int J Hyg Environ Health*. Oct 2007;210(5):535-540.
- 338 18. Kukla L, Hrubá D, Tyrlik M. Influence of prenatal and postnatal exposure to passive
339 smoking on infants' health during the first six months of their life. *Cent Eur J Public*
340 *Health*. Sep 2004;12(3):157-160.
- 341 19. Badenska B, Czerwionka-Szaflarska, M. The analysis of effects of passive smoking
342 exposition in children in pre- and postnatal periods. *Prze Pediatr*. 2002;32(3):199-202.
- 343 20. Barbier C, Houdret, N, Vittrant, C, Deschildre, A, Turck, D. Evaluation of passive smoking
344 by measurement of urinary cotinine in a pediatric outpatient clinics of the North-Pas-de-
345 Calais region. *Arch Pediatr*. 2000;7(7):719-724.
- 346 21. Kukla L, Hrubá, D, Tyrlik, M. Smoking of Mothers after Delivery Plays Significant Role in
347 Higher Morbidity of Newborns and Sucklings. *Cesko-Slovenska Pediatrie*.
348 2004;59(5):225-228.
- 349 22. Pospiech L, Rak J, Jaworska M, Klemous J. [Epidemiology of secretory otitis media in
350 children examined at the Otolaryngologic Clinic in Wroclaw in 1996-1999]. *Wiad Lek*.
351 2002;55(5-6):296-300.
- 352 23. Veneziano A, Mayer, M, Greco, L. Passive smoke and morbidity in primary child care.
353 *Medico e Bambino*. 2000;19(5):300-302.

- 354 24. Vera G, Royo, GF, Pujol, PE, Sevillano, RF, Alvaro, OMJ. Acute otitis media during the first
355 year of life and its relationship with some risk factors. *An Esp Pediatr.* 1997;47(5):473-
356 477.
- 357 25. Wefring KW, Lie KK, Loeb M, Nordhagen R. [Nasal congestion and earache--upper
358 respiratory tract infections in 4-year-old children]. *Tidsskr Nor Laegeforen.* Apr 30
359 2001;121(11):1329-1332.
- 360 26. de O. Saes S, Goldberg, TBL, Montovani, JC. Secretion of middle ear in infants -
361 occurrence, recurrence and related factors. *J. Pediatr. (Rio J.)*. 2005;81(2):133-138.
- 362 27. Jackson JM, Mourino AP. Pacifier use and otitis media in infants twelve months of age or
363 younger. *Pediatr Dent.* Jul-Aug 1999;21(4):255-260.
- 364 28. Mukherjee D, Stephens, D. Otitis media in intellectually disabled children. *J Audiol Med.*
365 1997;6(1):10-23.
- 366 29. Kristjansson S, Skuladottir HE, Sturludottir M, Wennergren G. Increased prevalence of
367 otitis media following respiratory syncytial virus infection. *Acta Paediatr.*
368 2010;99(6):867-870.
- 369 30. Ladomenou F, Moschandreas J, Kafatos A, Tselentis Y, Galanakis E. Protective effect of
370 exclusive breastfeeding against infections during infancy: a prospective study. *Arch Dis*
371 *Child.* 2010;95(12):1004-1008.
- 372 31. Apostolopoulos K, Xenelis J, Tzagaroulakis A, Kandiloros D, Yiotakis J, Papafragou K. The
373 point prevalence of otitis media with effusion among school children in Greece. *Int J*
374 *Pediatr Otorhinolaryngol.* 10 1998;44(3):207-214.

- 375 32. Froom J, Culpepper L, Green LA, et al. A cross-national study of acute otitis media: risk
376 factors, severity, and treatment at initial visit. Report from the International Primary
377 Care Network (IPCN) and the Ambulatory Sentinel Practice Network (ASPN).[see
378 comment]. *J Am Board Fam Pract.* Nov-Dec 2001;14(6):406-417.
- 379 33. Gryczynska D, Kobos J, Zakrzewska A. Relationship between passive smoking, recurrent
380 respiratory tract infections and otitis media in children. *Int J Pediatr Otorhinolaryngol.*
381 05 1999;49(SUPPL. 1):275-278.
- 382 34. Hammaren-Malmi S, Tarkkanen J, Mattila PS. Analysis of risk factors for childhood
383 persistent middle ear effusion. *Acta Otolaryngo.* 2005;125(10):1051-1054.
- 384 35. Homoe P, Christensen RB, Bretlau P. Acute otitis media and sociomedical risk factors
385 among unselected children in Greenland. *Int J Pediatr Otorhinolaryngol.* 15
386 1999;49(1):37-52.
- 387 36. Lieu JEC, Feinstein AR. Effect of gestational and passive smoke exposure on ear
388 infections in children. *Arch Pediatr Adolesc Med.* 2002;156(2):147-154.
- 389 37. Lister SM, Jorm LR. Parental smoking and respiratory illnesses in Australian children
390 aged 0-4 years: ABS 1989-90 National Health Survey results. *Aust N Z J Public Health.*
391 Dec 1998;22(7):781-786.
- 392 38. Lubianca Neto JF, Burns AG, Lu L, Mombach R, Saffer M. Passive smoking and
393 nonrecurrent acute otitis media in children. *Otolaryngol Head Neck Surg.*
394 1999;121(6):805-808.
- 395 39. Rylander R, Megevand Y. Environmental risk factors for respiratory infections. *Arch*
396 *Environ Health.* 2000;55(5):300-303.

- 397 40. Safavi Naini A, Safavi Naini A, Vazirnezam M. Parental smoking and risk of otitis media
398 with effusion among children. *Tanaffos*. 2002;1(3):25-28.
- 399 41. Said G, Zalokar J, Lellouch J, Patois E. Parental smoking related to adenoidectomy and
400 tonsillectomy in children. *J Epidemiol Community Health*. 1978;32:97-101.
- 401 42. Saim A, Saim L, Saim S, Ruszymah BHI, Sani A. Prevalence of otitis media with effusion
402 amongst pre-school children in Malaysia. *Int J Pediatr Otorhinolaryngol*. 18
403 1997;41(1):21-28.
- 404 43. Shiva F, Nasiri M, Sadeghi B, Padyab M. Effects of passive smoking on common
405 respiratory symptoms in young children. *Acta Paediatr*. Dec 2003;92(12):1394-1397.
- 406 44. Strachan DP. Impedance tympanometry and the home environment in seven-year-old
407 children. *J Laryngol Otol*. 1990;104:4-8.
- 408 45. Xenellis J, Paschalidis J, Georgalas C, Davilis D, Tzagaroulakis A, Ferekidis E. Factors
409 influencing the presence of otitis media with effusion 16 months after initial diagnosis in
410 a cohort of school-age children in rural Greece: A prospective study. *Int J Pediatr*
411 *Otorhinolaryngol*. Dec 2005;69(12):1641-1647.
- 412 46. Adair-Bischoff CE, Sauve RS. Environmental tobacco smoke and middle ear disease in
413 preschool-age children. *Arch Pediatr Adolesc Med*. 1998;152(2):127-133.
- 414 47. Barr G. Passive smoking and otitis media with effusion. *BMJ*. 1992;304:382-383.
- 415 48. da Costa JL, Navarro A, Neves JB, Martin M. Household wood and charcoal smoke
416 increases risk of otitis media in childhood in Maputo. *Int J Epidemiol*. 2004;33(3):573-
417 578.

- 418 49. Daigler GE, Markello SJ, Cummings KM. The effect of indoor air pollutants on otitis
419 media and asthma in children. *Laryngoscope*. 1991;101:293-296.
- 420 50. Green MRE, Cooper MNK. Passive smoking and middle ear effusions in children of
421 British Servicemen in West Germany - a point prevalence survey of outpatient
422 attendance. *J R Army Med Corps*. 1991;137:31-33.
- 423 51. Haggard MP, Gannon MM, Birkin JA, et al. Selecting persistent glue ear for referral in
424 general practice: A risk factor approach. *Br J Gen Pract*. 2002;52(480):549-553.
- 425 52. Hinton AE. Surgery for otitis media with effusion in children and its relationship to
426 parental smoking. *J Laryngol Otol*. 1989;103:559-561.
- 427 53. Hinton AE, Buckley G. Parental smoking and middle ear effusions in children. *J Laryngol*
428 *Otol* 1988;102:992-996.
- 429 54. Hinton AE, Herdman RCD, Martin-Hirsch D, Saeed SR. Parental cigarette smoking and
430 tonsillectomy in children. *Clin Otolaryngol*. 1993;18:178-180.
- 431 55. Ilicali OC, Kclcj N, Deger K, Savaj I. Relationship of passive cigarette smoking to otitis
432 media. *Arch Otolaryngol Head Neck Surg*. Jul 1999;125(7):758-762.
- 433 56. Ilicali OC, Keles N, Deger K, Sagun OF, Guldiken Y. Evaluation of the effect of passive
434 smoking on otitis media in children by an objective method: Urinary cotinine analysis.
435 *Laryngoscope*. 2001;111(1):163-167.
- 436 57. Kitchens GG. Relationship of environmental tobacco smoke to otitis media in young
437 children. *Laryngoscope*. 1995;105(5 Part 2, Suppl 69):1-13.
- 438 58. Kraemer MJ, Richardson MA, Weiss NS, et al. Risk factors for persistent middle-ear
439 effusions. *JAMA*. 1983;249(8):1022-1025.

- 440 59. Lasisi AO, Olaniyan FA, Muibi SA, et al. Clinical and demographic risk factors associated
441 with chronic suppurative otitis media. *Int J Pediatr Otorhinolaryngol.* Oct
442 2007;71(10):1549-1554.
- 443 60. Pukander J, Luotonen J, Timonen M, Karma P. Risk factors affecting the occurrence of
444 acute otitis media among 2-3-year old urban children. *Acta Otolaryngology (Stockholm).*
445 1985;100:260-265.
- 446 61. Rowe-Jones JM, Brockbank MJ. Parental smoking and persistent otitis media with
447 effusion in children. *Int J Pediatr Otorhinolaryngol.* 1992;24:19-24.
- 448 62. Ståhlberg M-R, Ruuskanen O, Virolainen E. Risk factors for recurrent otitis media.
449 *Pediatr Infect Dis.* 1986;5(1):30-32.
- 450 63. Stenstrom C, Ingvarsson L. Otitis-prone children and controls: A study of possible
451 predisposing factors. 2. Physical findings, frequency of illness, allergy, day care and
452 parental smoking. *Acta Otolaryngol.* 1997;117(5):696-703.
- 453 64. Stenstrom R, Bernard PAM, Ben-Simhon H. Exposure to environmental tobacco smoke
454 as a risk factor for recurrent acute otitis media in children under the age of five years.
455 *Int J Pediatr Otorhinolaryngol.* 1993;27:127-136.
- 456 65. Willatt DJ. Children's sore throats related to parental smoking. *Clin Otolaryngol.*
457 1986;11:317-321.
- 458 66. Gultekin E, Develioglu ON, Yener M, Ozdemir I, Kulekci M. Prevalence and risk factors for
459 persistent otitis media with effusion in primary school children in Istanbul, Turkey. *Auris*
460 *Nasus Larynx.* 2010;37(2):145-149.

- 461 67. Sophia A, Isaac R, Rebekah G, Brahmadathan K, Rupa V. Risk factors for otitis media
462 among preschool, rural Indian children. *Int J Pediatr Otorhinolaryngo*. 2010;74(6):677-
463 683.
- 464 68. MacIntyre EA, Karr CJ, Koehoorn M, et al. Otitis media incidence and risk factors in a
465 population-based birth cohort. *Paediatr Child Health*. 2010;15(7):437-442.
- 466 69. Alho OP, Kilkku O, Oja H, Koivu M, Sorri M. Control of the temporal aspect when
467 considering risk factors for acute otitis media. *Arch Otolaryngol Head Neck Surg*.
468 1993;119(4):444-449.
- 469 70. Bener A, Eihakeem AAM, Abdulhadi K. Is there any association between consanguinity
470 and hearing loss. *Int J Pediatr Otorhinolaryngol*. Mar 2005;69(3):327-333.
- 471 71. Bennett KE, Haggard MP. Accumulation of factors influencing children's middle ear
472 disease: risk factor modelling on a large population cohort. *J Epidemiol Community*
473 *Health*. Dec 1998;52(12):786-793.
- 474 72. Bentdal YE, Karevold G, Nafstad P, Kvaerner KJ. Early acute otitis media: Predictor for
475 AOM and respiratory infections in schoolchildren? *Int J Pediatr Otorhinolaryngol*. Aug
476 2007;71(8):1251-1259.
- 477 73. Collet J-P, Larson CP, Boivin J-F, Suissa S, Pless IB. Parental smoking and risk of otitis
478 media in pre-school children. *Can J Public Health* 1995;86(4):269-273.
- 479 74. Daly KA, Pirie PL, Rhodes KL, Hunter LL, Davey CS. Early otitis media among Minnesota
480 American Indians: The little ears study. *Am J Public Health*. Feb 2007;97(2):317-322.
- 481 75. Engel J, Anteunis L, Volovics A, Hendriks J, Marres E. Risk factors of otitis media with
482 effusion during infancy. *Int J Pediatr Otorhinolaryngol*. 25 1999;48(3):239-249.

- 483 76. Etzel RA, Pattishall EN, Haley NJ, Fletcher RH, Henderson FW. Passive smoking and
484 middle ear effusion among children in day care. *Pediatrics*. 1992;90:228-232.
- 485 77. Ey JL, Holberg CJ, Aldous MB, et al. Passive smoke exposure and otitis media in the first
486 year of life. *Pediatrics*. 1995;95:670-677.
- 487 78. Gliddon ML, Sutton GJ. Prediction of 8-month MEE from neonatal risk factors and test
488 results in SCBU and full-term babies.[erratum appears in Br J Audiol 2001 Jun;35(3):219].
489 *Br J Audiol*. Feb 2001;35(1):77-85.
- 490 79. Hammaren-Malmi S, Saxen H, Tarkkanen J, Mattila PS. Passive smoking after
491 tympanostomy and risk of recurrent acute otitis media. *Int J Pediatr Otorhinolaryngol*.
492 Aug 2007;71(8):1305-1310.
- 493 80. Iversen M, Birch L, Lundqvist GR. Middle ear effusion in children and the indoor
494 environment: an epidemiological study. *Arch Environ Health*. 1985;40(2):74-79.
- 495 81. Jacoby PA, Coates HL, Arumugaswamy A, et al. The effect of passive smoking on the risk
496 of otitis media in Aboriginal and non-Aboriginal children in the Kalgoorlie-Boulder region
497 of Western Australia. *Med J Aust*. May 19 2008;188(10):599-603.
- 498 82. Lee DJ, Gaynor JJ, Trapido E. Secondhand smoke and earaches in adolescents: The
499 Florida youth cohort study. *Nicotine Tob Res*. Dec 2003;5(6):943-946.
- 500 83. Noakes P, Taylor A, Hale J, et al. The effects of maternal smoking on early mucosal
501 immunity and sensitization at 12 months of age. *Pediatr Allergy Immunol*. Mar
502 2007;18(2):118-127.

- 503 84. Paradise JL, Rockette HE, Colborn DK, et al. Otitis media in 2253 Pittsburgh-area infants:
504 Prevalence and risk factors during the first two years of life. *Pediatrics*. Mar
505 1997;99(3):318-333.
- 506 85. Rasmussen F. Protracted secretory otitis media. The impact of familial factors and day-
507 care center attendance. *Int J Pediatr Otorhinolaryngol*. 1993;26:29-37.
- 508 86. Salazar JC, Daly KA, Giebink GS, et al. Low cord blood pneumococcal immunoglobulin G
509 (IgG) antibodies predict early onset acute otitis media in infancy. *Am J Epidemiol*. 01
510 1997;145(11):1048-1056.
- 511 87. Stathis SL, O'Callaghan DM, Williams GM, Najman JM, Andersen MJ, Bor W. Maternal
512 cigarette smoking during pregnancy is an independent predictor for symptoms of
513 middle ear disease at five years' postdelivery. *Pediatrics*. Aug 1999;104(2):e16.
- 514 88. Tainio V-M, Savilahti E, Salmenperä L, Arjomaa P, Siimes MA, Perheentupa J. Risk factors
515 for infantile recurrent otitis media: atopy but not type of feeding. *Pediatr Res*.
516 1988;23(5):509-512.
- 517 89. Teele Dw, Klein JO, Rosner B, and the Greater Boston Otitis Media Study Group.
518 Epidemiology of otitis media during the first seven years of life in children in Greater
519 Boston: A prospective, cohort study. *J Infect Dis*. 1989;160(1):83-94.
- 520 90. Zielhuis GA, Heuvelmans-Heinen EW, Rach GH, Van Den Broek P. Environmental risk
521 factors for otitis media with effusion in preschool children. *Scand J Prim Health Care*.
522 1989;7:33-38.

- 523 91. Haberg SE, Bentdal YE, London SJ, Kvaerner KJ, Nystad W, Nafstad P. Prenatal and
524 postnatal parental smoking and acute otitis media in early childhood. *Acta Paediatr.*
525 2010;99(1):99-105.
- 526 92. Danhauer JL, Johnson CE, Rotan SN, Snelson TA, Stockwell JS. National survey of
527 pediatricians' opinions about and practices for acute otitis media and xylitol use. *J Am*
528 *Acad Audiol.* May 2010;21(5):329-346.
- 529 93. Fleming DM, Ross AM, Cross KW, Kendall H. The reducing incidence of respiratory tract
530 infection and its relation to antibiotic prescribing. *Br J Gen Pract.* Oct 2003;53(495):778-
531 783.
- 532
- 533
- 534

535 **FIGURE TITLES AND LEGENDS**

536 **Figure 1.** Flow diagram of included and excluded studies

537 **Figure 2.** Funnel plot for household second hand smoke exposure against middle ear disease.

538 (Plot shows the standard error of the odds ratio versus odds ratio for each study (random effects model). Vertical
539 dotted lines indicate pooled effect estimate; and dots, individual studies).

540 **Figure 3.** Relationship between passive smoke exposure by maternal smoking after birth and
541 the risk of middle ear disease using a meta-analysis of comparative epidemiologic studies (Data
542 are presented as odds ratios sub-grouped by the definition of middle ear disease outcome).

543 (Squares denote the odds ratio (OR) for a single study with horizontal lines denoting 95% confidence intervals. The
544 centre of the diamond denotes the pooled OR and the corners the 95% confidence intervals. An OR >1 indicates a
545 higher risk of the outcome in those exposed to passive smoke).

546 **Figure 4.** Relationship between paternal passive smoke exposure and the risk of middle ear
547 disease using a meta-analysis of comparative epidemiologic studies (Data are presented as odds
548 ratios sub-grouped by the definition of middle ear disease outcome).

549 (Squares denote the odds ratio (OR) for a single study with horizontal lines denoting 95% confidence intervals. The
550 centre of the diamond denotes the pooled OR and the corners the 95% confidence intervals. An OR >1 indicates a
551 higher risk of the outcome in those exposed to passive smoke).

552 **Figure 5.** Relationship between passive smoke exposure by any household member and the risk
553 of middle ear disease using a meta-analysis of comparative epidemiologic studies (Data are
554 presented as odds ratios sub-grouped by the definition of middle ear disease outcome).

555 (Squares denote the odds ratio (OR) for a single study with horizontal lines denoting 95% confidence intervals. The
556 centre of the diamond denotes the pooled OR and the corners the 95% confidence intervals. An OR >1 indicates a
557 higher risk of the outcome in those exposed to passive smoke).

TABLES

Table 1 Summary of studies included in the meta-analysis

| Study | Year of Publication* | Study Design | Study Location | Ascertainment | Exposure | Outcome | Methodological Quality[†] |
|------------------------------|-----------------------------|---------------------|-----------------------|-----------------------------|--|----------------------------|---|
| Adair-Bischoff ⁴⁶ | 1998 | Case control | Canada | Biochemical and self-report | Maternal post-natal, paternal, household | Middle ear disease | 8 |
| Alho ⁶⁹ | 1993 | Cohort | Finland | Biochemical | Household | Acute otitis media | 8 |
| Apostolopoulos ³¹ | 1998 | Cross sectional | Greece | Self-report | Household | Otitis media with effusion | 6 |
| Barr ⁴⁷ | 1992 | Case control | United Kingdom | Self-report | Maternal pre-natal, household | Otitis media with effusion | 5 |
| Bener ⁷⁰ | 2005 | Cohort | Qatar | Self-report | Maternal pre-natal | Hearing loss | 7 |
| Bennett ⁷¹ | 1998 | Cohort | United Kingdom | Self-report | Maternal post-natal | Middle ear disease | 6 |
| Bentdal ⁷² | 2007 | Cohort | Norway | Self-report | Household | Acute otitis media | 7 |
| Collet ⁷³ | 1995 | Cohort | Canada | Self-report | Household | Recurrent otitis media | 6 |
| da Costa ⁴⁸ | 2004 | Case control | Mozambique | Self-report | Household | Otitis media | 8 |
| Daigler ⁴⁹ | 1991 | Case control | United States | Self-report | Maternal post-natal, paternal | Recurrent otitis media | 8 |

Table 1 Continued

| Study | Year of Publication * | Study Design | Study Location | Ascertainment | Exposure | Outcome | Methodological Quality † |
|--------------------------|------------------------------|---------------------|--|----------------------|--|----------------------------|---------------------------------|
| Daly ⁷⁴ | 2007 | Cohort | United States | Self-report | Maternal post-natal, household | Otitis media | 6 |
| Engel ⁷⁵ | 1999 | Cohort | Netherlands | Self-report | Household | Otitis media with effusion | 7 |
| Etzel ⁷⁶ | 1992 | Cohort | United States | Biochemical | Household | Otitis media with effusion | 8 |
| Ey ⁷⁷ | 1995 | Cohort | United States | Biochemical | Maternal post-natal | Recurrent Otitis media | 8 |
| Froom ³² | 2001 | Cross sectional | United States, Canada, United Kingdom, Netherlands | Self-report | Household | Acute otitis media | 4 |
| Gliddon ⁷⁸ | 2001 | Cohort | United Kingdom | Self-report | Maternal post-natal, household | Otitis media with effusion | 3 |
| Green ⁵⁰ | 1991 | Case control | Germany | Self-report | Maternal post-natal, paternal | Otitis media with effusion | 5 |
| Gryczynska ³³ | 1999 | Cross sectional | Poland | Self-report | Household | Suppurative otitis media | 2 |
| Gultekin ⁶⁶ | 2010 | Case control | Turkey | Self-report | Maternal post-natal, paternal, household | Otitis media with effusion | 6 |

Table 1 continued

| Study | Year of Publication* | Study Design | Study Location | Ascertainment | Exposure | Outcome | Methodological Quality† |
|------------------------------|-----------------------------|---------------------|-----------------------|----------------------|--|--------------------------------------|--------------------------------|
| Haberg ⁹¹ | 2010 | Cohort | Norway | Self-report | Maternal pre-natal, maternal post-natal, household | Acute otitis media | 4 |
| Haggard ⁵¹ | 2002 | Case control | United Kingdom | Self-report | Maternal post-natal | glue ear | 6 |
| Hammaren-Malmi ³⁴ | 2005 | Cross sectional | Finland | Self-report | Maternal post-natal, paternal | Otitis media with effusion | 4 |
| Hammaren-Malmi ⁷⁹ | 2007 | Cohort | Finland | Self-report | Maternal post-natal, paternal | Recurrent otitis media | 8 |
| Hinton ⁵³ | 1988 | Case control | United Kingdom | Self-report | Household | Otitis media with effusion | 4 |
| Hinton ⁵² | 1989 | Case control | United Kingdom | Self-report | Household | Surgery (otitis media with effusion) | 2 |
| Hinton ⁵⁴ | 1993 | Case control | United Kingdom | Self-report | Maternal post-natal, paternal, household | Surgery (tonsils) | 6 |
| Homoe ³⁵ | 1999 | Cross sectional | Greenland | Self-report | Household | Acute otitis media | 4 |

Table 1 continued

| Study | Year of Publication* | Study Design | Study Location | Ascertainment | Exposure | Outcome | Methodological Quality† |
|------------------------|-----------------------------|---------------------|-----------------------|-----------------------------|--|--------------------------------------|--------------------------------|
| Ilicali ⁵⁵ | 1999 | Case control | Turkey | Self-report | Maternal post-natal, paternal, household | Surgery (otitis media) | 3 |
| Ilicali ⁵⁶ | 2001 | Case control | Turkey | Biochemical and self-report | Household | Surgery (otitis media) | 5 |
| Iversen ⁸⁰ | 1985 | Cohort | Denmark | Self-report | Household | Otitis media with effusion | 6 |
| Jacoby ⁸¹ | 2008 | Cohort | United States | Self-report | Household | Otitis media | 8 |
| Kitchens ⁵⁷ | 1995 | Case control | United States | Self-report | Maternal post-natal, paternal, household | Surgery (otitis media with effusion) | 5 |
| Kraemer ⁵⁸ | 1983 | Case control | United States | Self-report | Household | Surgery (otitis media with effusion) | 6 |
| Lasisi ⁵⁹ | 2007 | Case control | Nigeria | Self-report | Household | Chronic suppurative otitis media | 6 |
| Lee ⁸² | 2003 | Cohort | United States | Self-report | Household | Earache | 4 |

Table 1 continued

| Study | Year of Publication* | Study Design | Study Location | Ascertainment | Exposure | Outcome | Methodological Quality† |
|--------------------------|-----------------------------|---------------------|-----------------------|-----------------------------|--|--|--------------------------------|
| Lieu ³⁶ | 2002 | Cross sectional | United States | Biochemical and self-report | Maternal pre-natal, maternal post-natal, household | Otitis media | 7 |
| Lister ³⁷ | 1998 | Cross sectional | Australia | Self-report | Maternal post-natal | Surgery (otitis media) | 4 |
| Lubianca ³⁸ | 1999 | Cross sectional | Brazil | Self-report | Household | Acute otitis media | 4 |
| MacIntyre ⁶⁸ | 2010 | Case control | Canada | Self-report | Maternal pre-natal | Acute otitis media | 7 |
| Noakes ⁸³ | 2007 | Cohort | Australia | Biochemical | Maternal pre-natal | Acute infection and serious otitis media | 4 |
| Paradise ⁸⁴ | 1997 | Cohort | United States | Self-report | Household | Otitis media with effusion | 6 |
| Pukander ⁶⁰ | 1985 | Case control | Finland | Self-report | Household | Recurrent otitis media | 5 |
| Rasmussen ⁸⁵ | 1993 | Cohort | Sweden | Biochemical | Household | Surgery (otitis media with effusion) | 7 |
| Rowe-Jones ⁶¹ | 1992 | Case control | United Kingdom | Self-report | Household | Surgery (otitis media with effusion) | 3 |

Table 1 continued

| Study | Year of Publication* | Study Design | Study Location | Ascertainment | Exposure | Outcome | Methodological Quality† |
|----------------------------|-----------------------------|---------------------|-----------------------|----------------------|--|----------------------------------|--------------------------------|
| Rylander ³⁹ | 2000 | Cross sectional | Switzerland | Self-report | Household | Otitis media | 4 |
| Safavi Naini ⁴⁰ | 2002 | Cross sectional | Iran | Self-report | Household | Otitis media with effusion | 4 |
| Said ⁴¹ | 1978 | Cross sectional | France | Self-report | Maternal post-natal, paternal, household | Surgery (adenoids/tonsils) | 3 |
| Saim ⁴² | 1997 | Cross sectional | Malaysia | Self-report | Household | Otitis media with effusion | 4 |
| Salazar ⁸⁶ | 1997 | Cohort | United States | Self-report | Household | Acute otitis media | 7 |
| Shiva ⁴³ | 2003 | Cross sectional | Iran | Self-report | Household | Otitis media | 3 |
| Sophia ⁶⁷ | 2010 | Case control | India | Self-report | Household | Otitis media with effusion | 7 |
| Ståhlberg ⁶² | 1986 | Case control | Finland | Self-report | Household | Surgery (recurrent otitis media) | 4 |
| Stathis ⁸⁷ | 1999 | Cohort | Australia | Self-report | Maternal pre-natal, household | Acute otitis media | 7 |
| Stenstrom ⁶⁴ | 1993 | Case control | Canada | Self-report | Household | Recurrent otitis media | 6 |

Table 1 continued

| Study | Year of Publication* | Study Design | Study Location | Ascertainment | Exposure | Outcome | Methodological Quality† |
|-------------------------|-----------------------------|---------------------|-----------------------|----------------------|-------------------------------|----------------------------|--------------------------------|
| Stenstrom ⁶³ | 1997 | Case control | Sweden | Self-report | Maternal post-natal, paternal | Otitis prone | 5 |
| Strachan ⁴⁴ | 1990 | Cross sectional | United Kingdom | Self-report | Household | Otitis media with effusion | 6 |
| Tainio ⁸⁸ | 1988 | Cohort | Finland | Biochemical | Household | Recurrent otitis media | 8 |
| Teele ⁸⁹ | 1989 | Cohort | United States | Biochemical | Household | Recurrent otitis media | 7 |
| Willatt ⁶⁵ | 1986 | Case control | United Kingdom | Self-report | Household | Surgery (tonsils) | 6 |
| Xenellis ⁴⁵ | 2005 | Cross sectional | Greece | Self-report | Household | Otitis media with effusion | 2 |
| Zielhuis ⁹⁰ | 1989 | Cohort | Netherlands | Biochemical | Household | Otitis media with effusion | 6 |

* Studies published prior to 1996 from previous Strachan and Cook review⁶

† Methodological quality of the studies are based on the Newcastle-Ottawa Quality Assessment Scale¹⁰

Table 2a Summary of overall effect and meta-regression analysis of maternal pre- and post-natal passive smoke exposure on the risk of middle ear disease in childhood

| | | Maternal pre-natal | | | | | Maternal post-natal | | | | |
|------------------------|--------------------------------|--------------------|--------------|---------|----------------|------|---------------------|--------------|---------|----------------|------|
| | | OR | 95% CI | Studies | I ² | p‡ | OR | 95% CI | Studies | I ² | p‡ |
| Overall effect | | 1.11 | 0.93 to 1.31 | 6 | 79 | N/A | 1.62 | 1.33 to 1.97 | 20 | 93 | N/A |
| Outcome | Middle ear infection | 1.15 | 0.98 to 1.35 | 5 | 79 | 0.36 | 1.53 | 1.22 to 1.92 | 16 | 94 | 0.63 |
| | Surgery for middle ear disease | N/A | N/A | 0 | N/A | | 1.86 | 1.31 to 2.63 | 5 | 73 | |
| | Hearing loss | 0.61 | 0.35 to 1.08 | 1 | N/A | | N/A | N/A | 0 | N/A | |
| | Hearing impairment | N/A | N/A | 0 | N/A | | 1.74 | 1.08 to 2.81 | 1 | N/A | |
| Study Design | Cohort | 1.97 | 0.51 to 2.24 | 4 | 85 | 0.94 | 1.19 | 0.94 to 1.49 | 6 | 76 | 0.14 |
| | Cross sectional | 1.07 | 0.97 to 1.18 | 1 | N/A | | 1.28 | 0.88 to 1.86 | 4 | 94 | |
| | Case-control | 1.16 | 1.09 to 1.23 | 1 | N/A | | 2.09 | 1.19 to 3.66 | 10 | 92 | |
| Methodological Quality | High | 1.17 | 0.95 to 1.44 | 4 | 85 | 0.49 | 1.83 | 1.21 to 2.76 | 10 | 96 | 0.47 |
| | Low | 0.74 | 0.24 to 2.27 | 2 | 41 | | 1.47 | 1.12 to 1.94 | 10 | 88 | |
| Publication Date | Prior to 1996 | N/A | N/A | 0 | N/A | N/A | 1.48 | 1.22 to 1.80 | 7 | 31 | 0.58 |
| | Post 1996 | 1.11 | 0.93 to 1.31 | 6 | 79 | | 1.72 | 1.35 to 2.21 | 13 | 95 | |
| Ascertainment | Self-report | 1.17 | 0.87 to 1.57 | 4 | 85 | 0.56 | 1.70 | 1.29 to 2.25 | 17 | 94 | 0.56 |
| | Biochemical | 0.75 | 0.22 to 2.61 | 2 | 47 | | 1.29 | 0.86 to 1.94 | 3 | 82 | |

OR = odds ratio

95% CI = 95% confidence interval

I² = percentage of heterogeneity

N/A = not applicable

p‡ = p value from random effect meta-regression analysis

Table 2b Summary of overall effect and meta-regression analysis of household and paternal passive smoke exposure on the risk of middle ear disease in childhood

| | | Household | | | | | Paternal | | | | |
|------------------------|--------------------------------|-----------|--------------|---------|----------------|------|----------|--------------|---------|----------------|--------|
| | | OR | 95% CI | Studies | I ² | p‡ | OR | 95% CI | Studies | I ² | p‡ |
| Overall effect | | 1.37 | 1.25 to 1.50 | 49 | 76 | N/A | 1.24 | 0.98 to 1.57 | 12 | 87 | N/A |
| Outcome | Middle ear infection | 1.32 | 1.20 to 1.45 | 38 | 73 | 0.11 | 1.06 | 0.91 to 1.24 | 8 | 33 | <0.001 |
| | Surgery for middle ear disease | 1.62 | 1.32 to 1.98 | 11 | 41 | | 1.83 | 1.61 to 2.07 | 4 | 0 | |
| | Hearing loss | N/A | N/A | 0 | N/A | | N/A | N/A | 0 | N/A | |
| | Hearing impairment | N/A | N/A | 0 | N/A | | N/A | N/A | 0 | N/A | |
| Study Design | Cohort | 1.27 | 1.13 to 1.43 | 18 | 46 | 0.25 | 2.45 | 0.85 to 7.07 | 1 | 0.28 | |
| | Cross sectional | 1.33 | 1.10 to 1.60 | 13 | 90 | | 1.51 | 0.82 to 2.78 | 2 | | 66 |
| | Case-control | 1.55 | 1.35 to 1.77 | 18 | 25 | | 1.14 | 0.90 to 1.44 | 7 | | 39 |
| Methodological Quality | High | 1.30 | 1.19 to 1.43 | 28 | 67 | 0.40 | 1.19 | 0.94 to 1.50 | 5 | 8 | 0.82 |
| | Low | 1.41 | 1.18 to 1.68 | 21 | 71 | | 1.25 | 0.91 to 1.74 | 7 | 92 | |
| Publication Date | Prior to 1996 | 1.46 | 1.26 to 1.70 | 21 | 60 | 0.27 | 1.40 | 1.01 to 1.94 | 5 | 64 | 0.22 |
| | Post 1996 | 1.31 | 1.17 to 1.46 | 28 | 77 | | 1.11 | 0.92 to 1.35 | 7 | 49 | |
| Ascertainment | Self-report | 1.40 | 1.25 to 1.56 | 40 | 74 | 0.32 | 1.25 | 0.97 to 1.62 | 11 | 88 | 0.74 |
| | Biochemical | 1.22 | 1.03 to 1.45 | 9 | 66 | | 1.11 | 0.76 to 1.63 | 1 | N/A | |

OR = odds ratio

95% CI = 95% confidence interval

I² = percentage of heterogeneity

N/A = not applicable

p‡ = p value from random effect meta-regression analysis

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

