Psychiatric co-morbidities in patients with dilated cardiomyopathy

Rasoul, Debar; Potluri, Siri; Wong, Sam C.h.; Gorantla, Ramakrishna S.; Aziz, Amir; Chandran, Suresh; Uppal, Hardeep; Potluri, Rahul

DOI:

[10.1016/j.ijcard.2015.04.271](https://doi.org/10.1016/j.ijcard.2015.04.271)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Rasoul, D, Potluri, S, Wong, SCH, Gorantla, RS, Aziz, A, Chandran, S, Uppal, H & Potluri, R 2015, 'Psychiatric co-morbidities in patients with dilated cardiomyopathy', *International Journal of Cardiology*, vol. 191, pp. 71-73. <https://doi.org/10.1016/j.ijcard.2015.04.271>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published as Rasoul Debar, Potluri Siri, Wong Sam, Gorantla Ramakrishna S., Aziz Amir, Chandran Suresh, Uppal Hardeep, Potluri Rahul, Psychiatric comorbidities in patients with dilated cardiomyopathy, *International Journal of Cardiology* (2015), doi: 10.1016/j.ijcard.2015.04.271

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Psychiatric co-morbidities in patients with dilated cardiomyopathy

Debar Rasoul, Siri Potluri, Sam Wong, Ramakrishna S. Gorantla, Amir Aziz, Suresh Chandran, Hardeep Uppal, Rahul Potluri

PII: S0167-5273(15)01005-0
DOI: doi: [10.1016/j.ijcard.2015.04.271](https://doi.org/10.1016/j.ijcard.2015.04.271)
Reference: IJCA 20456

To appear in: *International Journal of Cardiology*

Received date: 27 April 2015
Accepted date: 30 April 2015

Please cite this article as: Rasoul Debar, Potluri Siri, Wong Sam, Gorantla Ramakrishna S., Aziz Amir, Chandran Suresh, Uppal Hardeep, Potluri Rahul, Psychiatric co-morbidities in patients with dilated cardiomyopathy, *International Journal of Cardiology* (2015), doi: [10.1016/j.ijcard.2015.04.271](https://doi.org/10.1016/j.ijcard.2015.04.271)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Psychiatric co-morbidities in patients with dilated cardiomyopathy

International Journal of Cardiology – Short Communication

Debar Rasoul ¹	MBChB
Siri Potluri ²	MBChB(in prog)
Sam Wong ¹	MBChB
Ramakrishna S Gorantla ³	MBBS
Amir Aziz ⁴	PhD
Suresh Chandran ⁵	FRCP
Hardeep Uppal ⁶	MRCPsych
Rahul Potluri ⁶	MRCP

¹Department of Medicine, Macclesfield District General Hospital, Macclesfield, UK²School of Medicine, Cardiff University, Cardiff, UK³Department of Internal Medicine, Bassett Medical Center, Columbia University of physicians and surgeons, Cooperstown, USA⁴Department of Cardiology, University of Birmingham, Birmingham, UK⁵Department of Acute Medicine, North Western Deanery, UK⁶ACALM Study Unit in collaboration with Aston Medical School, Aston University, Birmingham, UK**Key word(s):** – Dilated Cardiomyopathy, Psychiatric Comorbidities, Trends**Short title:** - Psychiatric comorbidities and DCM**Conflict of interest statement:** The authors report no relationships that could be construed as a conflict of interest.**Correspondence to:** Dr Rahul Potluri, Honorary Clinical Lecturer in Cardiology, Aston University, Aston Triangle, Birmingham, B4 7ET, UK. Email: rahulpotluri@outlook.com. Telephone: 00447877482778. Fax: 00447877482778

Dear Editor,

Various studies have investigated and reported on the association between physical and psychiatric conditions^{1,2}. Recently, research focussing on the link between cardiovascular disease and psychiatric conditions has been on the rise³. Carter et al³, found that one in four psychiatric patients also suffered from a cardiovascular co-morbidity. Whether these links are causative, or not, there is a need to quantify this data to provide the evidence on which better targeted, holistic care-pathways can be based.

It is well known that depression and anxiety are significant and commonly reported factors in heart failure⁴ and it has been shown that patients with non-ischaemic heart failure, such as hypertrophic cardiomyopathy (HCM)^{5,6} and dilated cardiomyopathy (DCM)⁷ experience emotional stress, anxiety and in some cases depression and other psychiatric co-morbidities. This is often attributed the fear of associated risk of sudden death and potential genetic risk to their own children^{5,8}. In addition, recent studies⁹ have shown an increasing trend in patients with psychiatric comorbidities alongside DCM, and have called for a lower threshold to both assess and detect these co-morbidities as they often go under-detected and under-diagnosed in hospital¹⁰.

In light of this and the increasing evidence of association between psychiatric conditions and cardiovascular disease, we investigated the prevalence of psychiatric co-morbidities and tendencies in patients with dilated cardiomyopathy (DCM) over a 14 year period. We do not believe any study has looked into DCM and the possible links to psychiatric conditions and tendencies over such a long period to date.

We compiled an entirely anonymous database of adult patients diagnosed with dilated cardiomyopathy across seven hospitals in the North of England, UK during the period 01/01/2000 to 31/03/2013. We analysed the data for prevalence of firstly psychiatric co-morbidities such as;

anxiety disorder, obsessive compulsive disorder, schizophrenia, schizoaffective disorder, bipolar disease, depression and dementia and secondly substance abuse and suicidal intent. We traced our patients with the ACALM (Algorithm for Co-morbidities, Associations, length of stay and Mortality) study protocol, which uses ICD-10 (International Classification of Disease, Version 10) and OPCS-4 (Office of population Censuses and Surveys Classification of Interventions and Procedures) procedure codes to correctly allocate patients for statistical analysis using SPSS Version 20.0. The prevalence data was assembled into five groups to aid analysis of trends over time. This methodology has been described and by our group and others previously^{1-3,9,11-23}.

Between the years 2000 and 2013, 929,552 patients were admitted, 710 of which (0.08%) were labelled with a diagnosis of dilated cardiomyopathy. The majority of the population were made up of male patients (69%) and the mean age was 54.9 years $\pm$ 13.9 years(S.D). The population consisted of mainly Caucasians (77.2%), with a South Asian population of 7.87%, an Afro-Caribbean population of 2.88% and an unknown ethnicity in 8.43%.

Of these 710 patients with dilated cardiomyopathy, 9% (n=63) had other psychiatric conditions. We mainly found that in our cohort 4.65% (n=33) of patients had depression, 0.85% (n=6) of patients were diagnosed with dementia and a further 1.13% (n=8) suffered from schizophrenia. In addition to this, in 11% (n=81) of cases in our cohort, we found that patient suffered from some form of substance misuse with 4.08% (n=29) abusing alcohol and a further 6.34% (n=45) whom smoke tobacco.

Steptoe et al⁷ used the Hospital Anxiety and Depression (HAD)²⁴ in 99 cases and found 52% of his cohort fulfilled the criteria for possible anxiety disorder and in comparison to medical in-patients found that anxiety scores were significantly higher making the conclusion that DCM patients report higher levels of anxiety than their counterpart population. This finding was much higher than what

we observed in our study in which 1.13% (n=8) of our patients had anxiety. It is noteworthy for comparison that our study observed patients diagnosed with an anxiety disorder by a healthcare professional as opposed to a patient questionnaire.

In addition, Steptoe et al⁷ also found that 22% of his DCM cohort fulfilled the criteria for clinical depression, similar to the rate of depression found in coronary artery disease (CAD)⁴. In our study, in comparison we found that 4.65% (n=33) of our patients were suffering from clinical depression.

It is well known that alcohol is a common cause of DCM. Various theories have been proposed to establish its causative effect and pathogenesis, but mainly it remains the thought that the toxic metabolites of ethanol disrupts cardiovascular metabolism and that there is increased cardiomyocyte loss secondary to oxidative damage²⁵.

In addition, stress, anxiety and depression has been shown to affect the cardiovascular system by increasing activity on the sympathetic nervous system and the hypothalamo-pituitary axis which both help to regulate cardio-respiratory function²⁶,

On the contrary, it is believed that the combination of 1) psychiatric treatments effecting the cardiovascular system, 2) patients with depression tend to be less compliant with prevention of disease and intervention and 3) patients with psychiatric conditions subscribe to a lifestyle which can have negative effect on cardiovascular wellbeing by ingesting alcohol and tobacco, well known risk factors for cardiovascular disease, are the reasons for the correlation between psychiatric comorbidities and DCM.

Ultimately, it remains debatable whether these psychiatric conditions and tendencies represent a causative or correlative link. We did however find a steady increase in the abuse tobacco and

alcohol, and the diagnosis of depression in relation to our DCM patients during our 14 year period. We found that 1 in 10 patients with DCM suffer from a psychiatric co-morbidity and that 1 in 10 patients with DCM are suffering from some form of substance abuse.

We believe, that a holistic approach to the care of patients with DCM should be used, and that this holistic approach would fall short if it did not include focussed history expanding into a patient's use of recreational drugs, suicidal intent and psychiatric wellbeing in form of a history into psychiatric co-morbidities.

Table 1: Psychiatric co-morbidities in patients with Dilated Cardiomyopathy

	Prevalence of co-morbidity during the time period (%)					
	2000- 2002	2003- 2005	2006- 2008	2009- 2011	2012- 2013	Total in 14 years
n=	118	166	239	159	28	
<u>Psychiatric Condition</u>						
Anxiety disorder	0.85%	1.81%	1.26%	0.63%	0%	1.13%
Phobic disorder	1.69%	0%	0.42%	0%	0%	0.42%
Obsessive compulsive disorder	0.85%	0%	0%	0%	0%	0.14%
Schizophrenia	2.54%	0.60%	0.42%	1.89%	0%	1.13%
Schizoaffective disorder	0.85%	0%	0%	0%	0%	0.14%
Bipolar disorder	0%	0.60%	0%	1.26%	0%	0.42%
Depression	2.54%	4.82%	4.18%	6.92%	3.57%	4.65%
Dementia	0%	0%	1.67%	1.26%	0%	0.85%
<u>Psychiatric Tendency</u>						
Tobacco use	0.85%	2.41%	7.11%	10.06%	25.00%	6.34%
Alcohol use	1.69%	4.82%	4.18%	4.40%	7.14%	4.08%
Cannabis use	0%	0%	0.42%	0%	0%	0.14%
Opioid use	0%	0.60%	0.84%	0%	0%	0.42%
Stimulant use	0%	0.60%	0%	0%	0%	0.14%
Cocaine use	0%	0%	0.42%	0%	0%	0.14%
Overdose	0%	0%	0%	0.63%	0%	0.14%
Suicide attempt	0%	0%	0%	0.63%	0%	0.14%

References

1. Ciputra, Regina Nathania, Yan Efrata Sembiring, Olivia Listiowati Prawoto, Niece Khouw, Mudassar Baig, Hardeep Uppal, Suresh Chandran, and Rahul Potluri. "Length of Stay in Hospital Is Longer in Ethnic Minority Patients after Coronary Artery Bypass Surgery." *International Journal of Cardiology* 172, no. 3 (April 1, 2014): e434–35. doi:10.1016/j.ijcard.2013.12.182.
2. Schoepf, Dieter, Hardeep Uppal, Rahul Potluri, and Reinhard Heun. "Physical Comorbidity and Its Relevance on Mortality in Schizophrenia: A Naturalistic 12-Year Follow-up in General Hospital Admissions." *European Archives of Psychiatry and Clinical Neuroscience* 264, no. 1 (February 2014): 3–28. doi:10.1007/s00406-013-0436-x.
3. Carter, Paul, Jake Mann, Jason Sangha, Hardeep Uppal, Ammar Natalwala, Amir Aziz, Reinhard Heun, and Rahul Potluri. "The Burden of Cardiovascular Disease amongst Psychiatric Patients." *International Journal of Cardiology* 169, no. 4 (November 15, 2013): e65–66. doi:10.1016/j.ijcard.2013.08.129.
4. Faris, R., H. Purcell, M. Y. Henein, and A. J. S. Coats. "Clinical Depression Is Common and Significantly Associated with Reduced Survival in Patients with Non-Ischaemic Heart Failure." *European Journal of Heart Failure* 4, no. 4 (August 2002): 541–51.
5. Morgan, John F., Ann C. O'Donoghue, William J. McKenna, and Martin M. Schmidt. "Psychiatric Disorders in Hypertrophic Cardiomyopathy." *General Hospital Psychiatry* 30, no. 1 (February 2008): 49–54. doi:10.1016/j.genhosppsych.2007.09.005.
6. Cox, S., A. C. O'Donoghue, W. J. McKenna, and A. Steptoe. "Health Related Quality of Life and Psychological Wellbeing in Patients with Hypertrophic Cardiomyopathy." *Heart (British Cardiac Society)* 78, no. 2 (August 1997): 182–87.

7. Steptoe, A., A. Mohabir, N. G. Mahon, and W. J. McKenna. "Health Related Quality of Life and Psychological Wellbeing in Patients with Dilated Cardiomyopathy." *Heart (British Cardiac Society)* 83, no. 6 (June 2000): 645–50.
8. Rosenfeld, Jonathan E. "Emotional and Psychiatric Issues in Hypertrophic Cardiomyopathy and Other Cardiac Patients." *Anadolu Kardiyoloji Dergisi: AKD = the Anatolian Journal of Cardiology* 6 Suppl 2 (December 2006): 5–8.
9. Teo, Roxanne, Nicholas D. Gollop, Mudassar Baig, Hardeep Uppal, Suresh Chandran, and Rahul Potluri. "The Burden and Trends of Psychiatric Co-Morbidities amongst Patients with Cardiomyopathy." *International Journal of Cardiology* 174, no. 2 (June 15, 2014): 398–99. doi:10.1016/j.ijcard.2014.04.062.
10. Joynt, Karen E., David J. Whellan, and Christopher M. O'connor. "Why Is Depression Bad for the Failing Heart? A Review of the Mechanistic Relationship between Depression and Heart Failure." *Journal of Cardiac Failure* 10, no. 3 (June 2004): 258–71.
11. Kimani KN, Potluri R, Natalwala A, Ghosh S, Heun R, Narendran P. Length of hospital stay is shorter in black and ethnic minority patients with diabetes. *Diabet Med.* 2012;29;830-1.
12. Khouw N, Wasim M, Aziz A, Uppal H, Chandran S, Potluri R. Length of hospital stay is shorter in South Asian patients with myocardial infarction. *Int J Cardiol.* 2014;171;e54-5.
13. Smith S, Gollop N, Uppal H, Chandran S, Potluri R. Length of hospital stay is shorter in South Asian patients with acute pulmonary embolism. *Heart Asia.* 2014;6;1-2.
14. Carter P, Rai G, Aziz A, Mann J, Chandran S, Uppal H, Potluri R. Trends of cardiovascular disease amongst psychiatric patients between 2001 and 2012 in Greater Manchester, UK. *Int J Cardiol.* 2014 May 15;173(3);573-4.

15. Lavu D, Uppal H, Katragadda A, Ahmed A, Potluri R. Effect of ethnicity on live birth rates after in vitro fertilisation or Intracytoplasmic sperm injection treatment: possible explanations and further observations. *BJOG*. 2014 Nov;121(12):1576-7.
16. Potluri R, Baig M, Mavi JS, Ali N, Aziz A, Uppal H, Chandran S, The role of angioplasty in patients with acute coronary syndrome and previous coronary artery bypass grafting. *Int J Cardiology*. 2014 Oct 20;176(3):760-3.
17. Thakker C, Gollop ND, Carter P, Lavu D, Uppal H, Chandran S, Potluri R. Prevalence and length of hospital stay in patients with aortic valve disease is lower amongst South Asians. *Int J Cardiol*. 2014 Nov 15;177(1):34-6.
18. Green B, Gollop ND, Baig M, Uppal H, Chandran S, Potluri R. Prolonged length of hospital stay in Far-East Asian and Afro-Caribbean patients with cardiomyopathy. *Int J Cardiol*. 2014 Sep;176(1):230-1.
19. Potluri R, Wasim M, Markandey B, Kapour A, Khouw N, Carter P, Uppal H, Chandran S. Length of hospital stay is shorter in South Asian patients with Ischaemic Stroke. *Int J Cardiol*. 2015 Mar 20;187:190-191.
20. Yssennagger L, Gollop ND, Gorantla RS, Nimmagadda M, Potluri S, Uppal H, Chandran S, Potluri R. Increasing burden of Psychiatric Comorbidities amongst patients with Ischaemic Heart Disease. *Int J Cardiol*. 2015 Mar 18;186:200-201.
21. Uppal H, Chandran S, Potluri R. Risk factors for mortality in Down syndrome. *J Intellect Disabil Res*. 2015 Apr 8. doi: 10.1111/jir.12196. [Epub ahead of print] PubMed PMID: 25851193.

22. Gorantla RS, Nimmagadda M, Potluri S, Uppal H, Chandran S, Potluri R. Ethnic variations in length of hospital stay in patients with atrial fibrillation. *Int J Cardiol.* 2015 Mar 30;187:542-544.
23. SKA Khan, O McConnell, J Khan, F Ziaei, R Bhayani, S Shakir, S Potluri, RS Gorantla, H Uppal, S Chandran, R Potluri. The impact of ethnicity on the prevalence and length of hospital stay in patients with mitral regurgitation. *Int J Cardiol.* 2015 (IN PRESS)
24. Zigmond, A. S. and Snaith, R. P. (1983), The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67: 361–370. doi: 10.1111/j.1600-0447.1983.tb09716.x
25. Walker, Robin K., Valerie M. Cousins, Nsini A. Umoh, Miara A. Jeffress, Delaram Taghipour, Mustafa Al-Rubaiee, and Georges E. Haddad. "The Good, the Bad, and the Ugly with Alcohol Use and Abuse on the Heart." *Alcoholism, Clinical and Experimental Research* 37, no. 8 (August 2013): 1253–60. doi:10.1111/acer.12109.
26. Williams, R. B., T. L. Haney, K. L. Lee, Y. H. Kong, J. A. Blumenthal, and R. E. Whalen. "Type A Behavior, Hostility, and Coronary Atherosclerosis." *Psychosomatic Medicine* 42, no. 6 (November 1980): 539–49.