Dermal bioaccessibility of flame retardants from indoor dust and the influence of topically applied cosmetics

Pawar, Gopal; Abdallah, Mohamed; V-de-Sáa, Eugenia; Harrad, Stuart

DOI:

[10.1038/jes.2015.84](https://doi.org/10.1038/jes.2015.84)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Pawar, G, Abdallah, M, V-de-Sáa, E & Harrad, S 2016, 'Dermal bioaccessibility of flame retardants from indoor dust and the influence of topically applied cosmetics', *Journal of Exposure Science & Environmental Epidemiology*, vol. 27, pp. 100-105. <https://doi.org/10.1038/jes.2015.84>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Final published version available at: <http://dx.doi.org/10.1038/jes.2015.84>

Checked January 2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Dermal Bioaccessibility of Flame Retardants from Indoor Dust and the Influence of Topically Applied Cosmetics

Gopal Pawar, BSc¹, Mohamed Abou-Elwafa Abdallah, PhD^{1,2}, Eugenia Villaverde de Saa,
BSc³, Stuart Harrad, PhD^{1*}

¹Division of Environmental Health and Risk Management, School of Geography, Earth
and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, UK.

²Department of Analytical Chemistry, Faculty of Pharmacy, Assiut University, 71526
Assiut, Egypt.

³Department of Analytical Chemistry, University of Santiago de Compostela , 15782 -
Santiago de Compostela, Spain

* Corresponding author

Email s.j.harrad@bham.ac.uk

Tel. +44 121 414 7298

Fax. +44 121 414 3078

1 Abstract

Despite extensive literature on their potential adverse health effects, there is a lack of information on human dermal exposure to organic flame retardant chemicals (FRs). This study applies an *in vitro* physiologically based extraction test to provide new insights into the dermal bioaccessibility of various FRs from indoor dust to synthetic sweat/sebum mixture (SSSM). The bioaccessible fractions of α -, β -, γ -hexabromocyclododecane (HBCD) and tetrabromobisphenol A (TBBPA) to 1:1 (sweat:sebum) mixture were 41%, 47%, 50% and 40%, respectively. For Tris-2-chloroethyl phosphate (TCEP), tris (1-chloro-2-propyl) phosphate (TCIPP) and tris-1,3-dichloropropyl phosphate (TDCIPP), bioaccessible fractions were 10%, 17% and 19%. Composition of the SSSM and compound-specific physicochemical properties were the major factors influencing the bioaccessibility of target FRs. Except for TBBPA, the presence of cosmetics (moisturizing cream, sunscreen lotion, body spray, and shower gel) had a significant effect ($P < 0.05$) on the bioaccessibility of the studied FRs. The presence of cosmetics decreased the bioaccessibility of HBCDs from indoor dust, while shower gel and sunscreen lotion enhanced the bioaccessibility of target PFRs. Our bioaccessibility data were applied to estimate the internal exposure of UK adults and toddlers to the target FRs via dermal contact with dust. Our worst-case scenario exposure estimates fell far below available health based limit values for TCEP, TCIPP and TDCIPP. However, future research may erode the margin of safety for these chemicals.

Keywords: dermal exposure; bioaccessibility; BFRs; PFRs; cosmetics; indoor dust.

43

44 2 Introduction

45 Organic flame retardants like polybrominated diphenyl ethers (PBDEs),
46 hexabromocyclododecane (HBCD), tetrabromobisphenol-A (TBBPA), novel brominated
47 flame retardants (NBFRs), and organophosphate flame retardants (PFRs) have found
48 widespread application in a plethora of consumer items ^{1, 2}. However, concerns exist
49 over possible adverse health impacts following numerous reports of exposure to BFRs
50 through inhalation, dermal contact and ingestion of both diet and settled dust ^{3, 4}. In a
51 recent review ⁵, we highlighted the potential importance of dermal uptake of FRs as an
52 exposure pathway. The lack of experimental information on human dermal uptake of
53 these chemicals from contact with organic films present on indoor surfaces, as well as
54 contact with dust particles and source materials may be attributed to ethical issues
55 associated with both *in vivo* and *in vitro* studies using human tissues. In addition,
56 uncertainties arise from interspecies variation and allometric scaling of dermatokinetic
57 data from animals to humans ⁵. These challenges further support the need for
58 alternative *in vitro* methods to study dermal availability of FRs in indoor dust to
59 humans.

60 Survey of existing literature reveals various modelling approaches for dermal risk
61 assessment including quantitative structure activity relationship (QSAR)-based
62 methods ^{6, 7} and pharmacokinetic (PK)-modelling methods ⁸. However, such approaches
63 have some limitations, for example QSAR-based approaches report uncertainties
64 associated with the relationship between K_m (the partition coefficient between the
65 exposure vehicle and *stratum corneum*) of the studied molecule and its K_{ow} , where the
66 extent to which K_{ow} is good predictor for K_m is questionable, especially when the
67 exposure vehicle is not water. Moreover, the thickness of the *stratum corneum* (SC)

varies between species and estimated values of the compound diffusivity through the skin based on extrapolation from other studies on different compounds can be misleading⁹.

On the other hand, PK modelling studies of FRs report uncertainties associated with the fraction of FR available for absorption following exposure via different pathways (i.e. ingestion, inhalation or dermal contact) in addition to the lack of reliable information on the elimination half-lives of different FRs from various tissues^{10, 11}. Moreover, the influence of physiological fluids (e.g. sweat, gastrointestinal fluid, etc) on the bioavailable fraction of FRs is often neglected.

Physiologically-based *in vitro* bioaccessibility tests have emerged as an alternative method to study the availability for dermal uptake of several xenobiotics including heavy metals¹²⁻¹⁶ and pesticides¹⁷. Such bioaccessibility tests have been incorporated in regulatory frameworks such as the European standard for the release of nickel in artificial sweat (BS EN 1811, 2011). Bioaccessibility may be defined as “*the fraction of the total dose of a specific chemical/contaminant present in a matrix that becomes liberated into the body fluids and hence, is available for absorption*”¹⁸. In other words, a combination of data on bioaccessibility and subsequent dermal uptake is required to determine the ability of a chemical (e.g. an FR) present in a matrix (e.g. dust), to be released from that matrix and be subsequently absorbed by an organ of the human body like the skin¹⁷. Bioaccessibility data from *in vitro* studies are conservative, because not all the mass of a given chemical released into the body fluid (i.e. the bioaccessible fraction) will likely be absorbed through the biological membrane (e.g. skin) to reach the systemic circulation (i.e. bioavailable)¹⁹. The outermost surface of the human skin, the *stratum corneum*, is covered with a skin surface film liquid (SSFL) mixture which consists of varying proportions of sweat and sebum^{20, 21}. Sweat is aqueous in nature

and secreted to regulate body temperature. It consists mainly of electrolytes, organic acids, amino acids, vitamins and other nitrogenous substances. Sebum is a clear, oily substance secreted by sebaceous glands and forms a 0.5 to >4.0 μm thick layer to protect the skin from drying out. It mainly consists of squalene, wax esters and triglycerides, as well as free fatty acids, with a small amount of cholesterol and cholesterol esters ²².

Cosmetics (e.g. sunscreen creams) may contain certain ingredients (e.g. surfactants) which can remain on the skin and become incorporated within the SSFL. This in turn, may alter the lipid domain of the skin, by interacting with the proteins in the barrier, or hydration, thereby increasing partitioning of chemicals to the SC ²³. Previous studies have shown certain sunscreen lotions to act as inadvertent penetration enhancers for potentially harmful chemicals ^{24, 25}. Therefore, it is important to investigate the effect of topically-applied cosmetics on the dermal bioaccessibility of FRs in indoor dust.

Against this background, we investigate *-for the first time-* the dermal bioaccessibility of selected organic FRs present in house dust including: TBBPA, α -, β - and γ -HBCD, Tris-2-chloroethyl phosphate (TCEP), tris (1-chloro-2-propyl) phosphate (TCIPP) and tris-1,3-dichloropropyl phosphate (TDCIPP). We quantify the bioaccessible fraction of these FRs from dust to varying physiologically-relevant mixtures of synthetic sweat and sebum, and examine the impact on bioaccessibility of various topically-applied cosmetic products.

3 Materials and Methods

3.1 Characterisation of the studied house dust

SRM 2585 (organics in house dust, particle size < 100 μm and total moisture content = 2.11 ± 0.06 %) was purchased from NIST (Gaithersburg, MD, USA). Aliquots (n=5,

~0.1 g each) of SRM2585 were analysed for target FRs using previously reported methods by our research group ^{26, 27}. Results compared well with the indicative and reported levels of target FRs in this SRM (Tables SI-1 to SI-5).

3.2 Preparation of synthetic sweat and sebum mixture

Physiologically-simulated artificial sweat and sebum mixture (SSSM) was prepared according to a previously reported method and US patent using over 25 different chemical components ^{22, 28} (see Table SI-6 for details). The pH was adjusted to that of normal human skin (5.3 ± 0.1) and preserved at 8 °C. Synthetic sweat and sebum were prepared separately, and then mixed in different physiologically-relevant proportions using Tween 80 to mimic the naturally secreted surface active agents in the SSSM ^{22, 28}.

3.3 Dermal bioaccessibility *in vitro* test protocol

Briefly, ~60 mg of NIST SRM2585 dust and (when tested) 6 mg of cosmetics (moisturising cream, sun screen lotion, shower gel and body spray were each examined separately) were accurately weighed and transferred into a clean dry test tube. In the absence of definitive data on the dust to sweat ratio on human skin (which is greatly influenced by variations of sweat secretion and dust loadings), we adopted a previously reported method ¹⁷ to mimic “wet skin conditions” using 1:100 w/v dust to sweat ratio (i.e. 6 mL of the SSSM were applied for each 60 mg of dust). The mixture was then gently agitated on a heated magnetic-stirrer plate maintained at physiological skin temperature (32 °C). After 1 hour, phase separation was achieved by centrifugation at 3000 rpm for 15 mins. The dust (solid residue) and SSSM (supernatant) samples were analysed separately.

3.4 Chemical Analysis

3.4.1 Determination of HBCDs and TBBPA

Dust/SSSM/cosmetic samples were spiked with 30 μL of ^{13}C -isotopically labelled α -HBCD, β -HBCD, γ -HBCD and TBBPA ($1\text{ ng }\mu\text{L}^{-1}$), prior to extraction with 3 mL of hexane: ethyl acetate (1:1 v/v) using a QuEChERS-based method. Sample tubes were vortexed on a multi-positional mixer for 5 mins, followed by ultra-sonication for 5 mins and centrifugation at 3000 rpm for 5 mins. The extraction cycle was repeated twice before the pooled supernatant was collected in a clean tube and evaporated to $\sim 1\text{ mL}$ under a stream of N_2 . The crude extract was washed with $\sim 2\text{ mL}$ of 95 % H_2SO_4 to remove lipids. The organic layer and washings were combined and evaporated to incipient dryness under N_2 . Target analytes were reconstituted in $150\text{ }\mu\text{L}$ of methanol containing $50\text{ pg }\mu\text{L}^{-1}$ of d_{18} - α -HBCD used as recovery determination standard (RDS) prior to LC-MS/MS analysis using previously reported methods ²⁹.

3.4.2 Determination of PFRs

Dust/SSSM/cosmetic samples were spiked with 30 μL of d_{15} -triphenyl phosphate (d_{15} -TPHP, $10\text{ ng }\mu\text{L}^{-1}$) used as internal (surrogate) standard prior to extraction with hexane: ethyl acetate (1:1 v/v, 3 mL) using the same procedure applied for HBCDs. The crude extract ($\sim 1\text{ mL}$) was cleaned up by loading onto a Florisil SPE cartridge (pre-conditioned with 6 mL of hexane). Fractionation was achieved by eluting with 8 mL of hexane (F1, discarded) followed by 10 mL of ethyl acetate (F2). F2 was evaporated to incipient dryness under N_2 . Target PFRs were reconstituted in $100\text{ }\mu\text{L}$ of isooctane containing ^{13}C -BDE-100 used as RDS prior to GC/MS analysis according to a previously reported method ³⁰.

3.5 Quality Assurance and Quality Control

All experiments were conducted in triplicate. Good IS recoveries were obtained for all samples (Table SI-7). One procedural blank was run every 6 samples. This consisted of anhydrous sodium sulfate (~0.1 g) exposed to the same experimental protocol as a dust sample. None of the target compounds were detected in procedural blanks. Identification and quantification of target analytes were performed according to the retention times and peak areas of the corresponding calibration standards injected before and after each sample batch. While the overall method performance for dust analysis was evaluated via replicate analysis (n=5) of SRM 2585, method performance for the analysis of SSSM/cosmetic samples was checked by a matrix spike exercise at 3 concentration levels. The results obtained (Table SI-8) indicated good accuracy and precision of the applied analytical method.

3.6 Assessment of dermal bioaccessibility

In this study, bioaccessibility is expressed as $f_{\text{bioaccessible}}$, calculated (equation 1) as the percentage of each target FR detected in the dust that was found in the supernatant at the end of each bioaccessibility experiments (all experiments were carried out in triplicate, hence average values were used) (tables SI-2 and SI-5):

$$f_{\text{bioaccessible}} (\%) = \frac{\text{Average mass of FR in supernatant}}{\text{Average mass of FR in dust}} \times 100 \dots (1)$$

3.7 Statistical Analysis and Data Processing

Statistical analysis of data was conducted using Microsoft Excel 2010™ and SPSS 22™ for Windows. Means of various datasets were estimated and compared using ANOVA and

Tukey's honestly significant difference *post hoc* test. *P* values less than 0.05 were considered significant.

4 Results and Discussion

4.1 Dermal bioaccessibility of FRs in indoor dust

The process of human dermal uptake of chemicals from house dust to the general circulation is limited by two main factors. These are the bioaccessibility and the penetration rate. In the human skin, the *stratum corneum* (outermost dead corneous layer) presents the major limiting factor for penetration of chemicals and passive diffusion is the main transport mechanism for organic chemicals. Therefore, the penetration rate across the *stratum corneum* is mainly controlled by compound-specific physico-chemical properties. However, for chemicals bound to particulate matter as in indoor dust, the chemical's release from particles into the body fluids on the skin surface can be more important^{17, 31, 32}. The hydrolipidic SSFL and other ingredients of topically applied cosmetics may enhance, or reduce the chemical release ($f_{\text{bioaccessible}}$) from particles adhered to the skin. Once the chemical passes through the corneous layer by passive diffusion, it follows the intracellular/intercellular routes of penetration in the epidermis and dermis layers and subsequently reaches the blood stream ($f_{\text{bioavailable}}$) (Figure 1). Our results show that none of the target FRs were 100% bioaccessible from indoor dust particles into any of the studied SSSM combinations (Table 1). This indicates that assumption of 100% absorption of intake via the dermal route could lead to a substantial overestimation of human exposure to FRs via indoor dust.

4.1.1 Dermal bioaccessibility of HBCDs and TBBPA

In general, $f_{\text{bioaccessible}}$ of HBCDs and TBBPA increased with increasing sebum content of the SSFL (Table 1). At 100% sweat, the $f_{\text{bioaccessible}}$ of γ -HBCD ($1.4 \pm 0.1\%$) was less than that of β -HBCD ($1.6 \pm 0.6\%$) and α -HBCD ($2.3 \pm 0.2\%$). However, the reverse trend was observed at 100% sebum, where the $f_{\text{bioaccessible}}$ was highest for γ -HBCD ($67.2 \pm 3.37\%$), followed by β -HBCD ($60.4 \pm 10.1\%$) and α -HBCD ($50.5 \pm 7.0\%$). This behavior is consistent with the lower water solubility of the γ -isomer ($2 \mu\text{g L}^{-1}$) compared to that of β -HBCD ($15 \mu\text{g L}^{-1}$) and α -HBCD ($49 \mu\text{g L}^{-1}$)¹⁹.

We recorded $f_{\text{bioaccessible}}$ values for TBBPA of $3.5 \pm 0.5 \%$ and $55.7 \pm 8.5 \%$ in 100% sweat and 100% sebum, respectively. Compared to HBCDs, the higher $f_{\text{bioaccessible}}$ value for TBBPA in 100% sweat is likely attributable to the higher water solubility of TBBPA ($1.26 \times 10^3 \mu\text{g L}^{-1}$).

Compared to the aqueous-based sweat, the substantially higher bioaccessibility of the studied BFRs in sebum can be attributed to the enhanced solubility of these lipophilic chemicals in the oily sebum.

4.1.2 Dermal bioaccessibility of PFRs

In general, PFRs were more bioaccessible in sebum than sweat. In 100 % sweat, $f_{\text{bioaccessible}}$ values for the studied PFRs were $16.0 \pm 1.2\%$ (TCEP), $12.4 \pm 4.4\%$ (TCIPP) and $11.9 \pm 3.6\%$ (TDCIPP); while in 100% sebum, the corresponding values were $22.3 \pm 2.3\%$ (TCEP), $26.9 \pm 6.4\%$ (TCIPP), and $28.1 \pm 0.6 \%$ (TDCIPP). This concurs with the physicochemical properties of our target PFRs (Table SI-9). In particular, the water solubility of TCEP, TCIPP and TDCIPP was reported as 7×10^3 , 1.6×10^3 and 1.5 mg L^{-1} , respectively³³. Compared to the studied BFRs, PFRs show higher bioaccessibility in sweat and lower bioaccessibility in sebum (Table 1), which can be attributed to the differences in $\log K_{\text{ow}}$ and water solubility among these two classes of FRs (Table SI-9).

Overall, at the most realistic SSFL composition (1:1 sweat:sebum) studied here, BFRs showed higher dermal bioaccessibility than PFRs, which may be attributed to increased partitioning of the more lipophilic BFRs from dust to the oily sebum.

4.1.3 Effect of cosmetics on the dermal bioaccessibility of FRs in indoor dust

To investigate the influence of commonly applied cosmetics on the dermal bioaccessibility of FRs in indoor dust, we determined $f_{\text{bioaccessible}}$ values of target FRs from reference dust into 1:1 sweat:sebum mixture, in the presence of (separately) moisturizing cream, sunscreen lotion, body spray, and shower gel. Results for each target compound were compared to a control group comprising reference dust exposed only to 1:1 sweat: sebum mixture without any surfactant or cosmetics. Except for TBBPA, statistically significant differences ($P < 0.05$; ANOVA) were observed between $f_{\text{bioaccessible}}$ values of target FRs in the presence of various cosmetics compared to the control group (Figure 2). Interestingly, the presence of cosmetics seems to decrease the bioaccessibility of HBCDs from indoor dust (Figure 2). This is in agreement with the reported slight decrease in dermal bioaccessibility of PCBs from house dust in the presence of skin cream ¹⁷, which was attributed to possible retention of the lipophilic chemicals by skin cream lipids. Our results also show that while shower gel and sunscreen lotion enhanced the bioaccessibility of target PFRs, body spray significantly decreased the $f_{\text{bioaccessible}}$ value of TDCIPP from indoor dust (Figure 2).

To summarise, our results agree with previous reports that cosmetics contain various ingredients that can alter the composition of the SSFL and affect the availability of dust-bound FRs for dermal uptake. However, it is also evident that the nature and magnitude of this effect is substance-specific and highly dependent on the composition of the cosmetic preparation. The effect of surfactants - that are common ingredients of most

cosmetics - on the dermal absorption of various chemicals has been previously highlighted ^{24, 25}. In addition, we hypothesize that the lipid content, ionic strength and skin contact period of these cosmetics can also influence the bioaccessibility of FRs from indoor dust. Detailed studies are required to test this hypothesis and fully investigate the factors affecting the bioaccessibility of FRs and ultimately their dermal uptake in the presence of various cosmetic preparations.

4.1.4 Comparison of digestive and dermal bioaccessibility

Despite the vast differences between the digestive and dermal body fluids in terms of both composition and function, it is instructive to compare our results to previously reported bioaccessibilities of target FRs via the oral route. This can shed some light on the relative importance of dermal uptake versus ingestion as pathways of human exposure to FRs in indoor dust.

Abdallah et al. ¹⁹ reported on the gut bioaccessibility of HBCDs and TBBPA from indoor dust using a colon enhanced-physiologically based extraction test (CE-PBET). On average, $f_{\text{bioaccessible}}$ values of 92%, 80%, 72% and 94% were reported for α -, β -, γ -HBCDs and TBBPA, respectively. These are almost twice the dermal $f_{\text{bioaccessible}}$ values for the same BFRs in our study (Table 1). The gut bioaccessibility of PFRs following ingestion of indoor dust was also studied using a modified version of the CE-PBET mentioned above ³⁴. Mean $f_{\text{bioaccessible}}$ values for TCEP, TCIPP and TDCIPP from 17 house dust samples were 80%, 82% and 85%, respectively, which are substantially higher than the corresponding dermal $f_{\text{bioaccessible}}$ values for the same PFRs (Table 1).

The substantially higher gut bioaccessibility of FRs may be attributed to several factors. These include the strong acidic medium in the stomach (pH = 1), the bile salts and digestive enzymes in the small intestine, the presence of carbohydrates to simulate the

fed status, coupled with the long contaminant residence time in the models used (~13 - 21.5 hours)^{19, 34} compared to the 1 h dermal exposure period used in this study. More research is required to fully understand the influence of prolonged dermal exposure times on the bioaccessibility of FRs from indoor dust and examine the kinetics of the release of various FRs from indoor dust to the sweat/sebum mixture.

5 Assessment of human dermal exposure to FRs in indoor dust

The results of dermal bioaccessibility experiments obtained in this study (Table 1) were used to gain some insight on the internal dose of the target FRs arising from dermal exposure to contaminated indoor dust. Results revealed $f_{\text{bioaccessible}}$ values for the studied FRs in indoor dust were significantly influenced by the presence of various cosmetic preparations. However, incorporation of our data into risk assessment models is hampered by the current lack of reliable information on the exact amount of cosmetics remaining on the skin after application and on the skin residence time of such formulations. Therefore, exposure assessment estimations were performed without such data.

Human dermal exposure to our target FRs was estimated using the general equation:

$$DED = C \times BSA \times DAS \times FA \times IEFBW \times 1000 \dots 2$$

Where **DED** = Daily exposure dose (ng/kg bw/day), **C** = FR concentration in dust (ng/g), **BSA** = Body surface area exposed (cm²), **DAS** = Dust adhered to skin (mg/cm²), **FA** = fraction absorbed by the skin (unitless), **IEF** = indoor exposure fraction (hours spent over a day in an indoor environment) (unitless), **BW** = Body weight (kg).

We estimated the dermal exposure of 2 age groups (adults and toddlers) using three exposure scenarios. We used data previously reported by our research group on the minimum, median and maximum concentrations (Table SI-10) of target FRs in indoor dust from several UK microenvironments^{26, 35} to estimate low, average and high exposure, respectively. The parameter F_A in equation 2 was replaced by the experimental values of $f_{\text{bioaccesssible}}$ obtained in this study for each target FR at the most physiologically abundant sweat:sebum mixture (1:1) (Table 1). Values for other parameters in equation 2 were obtained from the USEPA exposure factors handbook³⁶ and summarized in Table 2.

Our dermal exposure estimates (Table 3) highlight the potential importance of the dermal route as a pathway of human exposure to FRs in indoor dust. The average scenario estimate of dermal exposure of UK adults and toddlers to the target BFRs ranged from (99-110%) and (44-59%) respectively, of their estimated exposure via dust ingestion²⁶ (Figure 3). For PFRs, the estimated average dermal exposure corresponded to (26-42%) and (28-45%) of previously reported exposure via dust ingestion³⁵. However, it should be noted that our dermal exposure estimates assume a fixed body area undergoing constant exposure to FRs in indoor dust for a constant period daily at a fixed absorbed fraction derived from 1 h dermal contact time with indoor dust. Such rigid assumptions are likely unrealistic and introduce uncertainty to our estimates of dermal exposure. A further significant caveat is that our estimates account only for bioaccessibility – i.e. the efficiency of release of FRs from dust into sweat/sebum. While this is important, reliable data are not yet available on the subsequent dermal transfer of the studied FRs from sweat/sebum across the epidermis to the systemic circulation. Such transfer will very likely be <100%, and thus the true influence of dermal exposure to dust will likely be appreciably lower than the values

shown in Table 3. While noting this caveat, we also note that our estimates of exposure via dust ingestion assume 100% efficiency of transfer from dust into gut fluids and thence across the gastro-intestinal tract.

In a risk assessment context, an extensive survey of the available literature revealed a No Significant Risk Level (NSRL) of 5.4 µg/day for TDCIPP listed as a carcinogen under the State of California safe drinking water and toxic enforcement act of 1986, PROPOSITION 65³⁷. No other health based limit values (HBLVs) of legislative standing for our target FRs were found in the literature. However, based on a chronic no observed adverse effect level (NOAEL) divided by an uncertainty factor of 1,000, HBLVs of 22,000 and 80,000 ng/kg bw/day were derived for TCEP and TCIPP respectively³⁸. Our worst-case scenario exposure estimates for dermal exposure of adults and toddlers fall far below these HBLV values even under our high-end dermal exposure scenario. However, as noted by Ali et al.³⁸, the HBLV values cited here were based on relatively old toxicological studies and it is possible that future research may erode the margin of safety.

In conclusion, notwithstanding the various caveats noted above, the results of this *in vitro* bioaccessibility study provide some important first insights into human dermal exposure to various FRs present in indoor dust. The composition (i.e. sweat:sebum ratio) of skin fluids, as well as the presence/absence of commonly used skin cosmetics is demonstrated to exert a substantial influence on the efficiency with which our target FRs are released from dust and rendered available for dermal uptake.

6 Acknowledgements

The research leading to these results has received funding from the European Union Seventh Framework Programme FP7/2007-2013 under grant agreement no. 316665

(A-TEAM) & grant agreement no. 327232 (ADAPT). E. Villaverde de Saa also acknowledges funding from the Spanish Ministry of Science and Innovation (FPI grant BES-2011-047887).

7 Supplementary information

Supplementary information is available at the Journal of Exposure Science and Environmental Epidemiology website.

8 References

1. Ghosh R, Hageman KJ, Björklund E Selective pressurized liquid extraction of three classes of halogenated contaminants in fish. *Journal of Chromatography A* 2011; 1218: 7242-7247.
2. van der Veen I, de Boer J Phosphorus flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* 2012; 88: 1119-1153.
3. Ali N, Harrad S, Goosey E, Neels H, Covaci A "Novel" brominated flame retardants in Belgian and UK indoor dust: implications for human exposure. *Chemosphere* 2011; 83: 1360-1365.
4. van Leeuwen SP, de Boer J Brominated flame retardants in fish and shellfish - levels and contribution of fish consumption to dietary exposure of Dutch citizens to HBCD. *Mol Nutr Food Res* 2008; 52: 194-203.
5. Abdallah MA, Pawar G, Harrad S Evaluation of in vitro vs. in vivo methods for assessment of dermal absorption of organic flame retardants: a review. *Environ Int* 2015; 74: 13-22.
6. Fitzpatrick D, Corish J, Hayes B Modelling skin permeability in risk assessment--the future. *Chemosphere* 2004; 55: 1309-1314.
7. Chen L, Han L, Lian G Recent advances in predicting skin permeability of hydrophilic solutes. *Advanced Drug Delivery Reviews* 2013; 65: 295-305.
8. Anissimov YG, Jepps OG, Dancik Y, Roberts MS Mathematical and pharmacokinetic modelling of epidermal and dermal transport processes. *Adv Drug Deliv Rev* 2013; 65: 169-190.

413 9. Van de Sandt JJ, Dellarco M, Van Hemmen JJ From dermal exposure to internal dose. J Expo
414 Sci Environ Epidemiol 2007; 17 Suppl 1: S38-47.
415

416 10. Lorber M Exposure of Americans to polybrominated diphenyl ethers. J Expo Sci Env Epid
417 2008; 18: 2-19.
418

419 11. Abdallah MA-E, Harrad S Tetrabromobisphenol-A, hexabromocyclododecane and its
420 degradation products in UK human milk: Relationship to external exposure. Environ Int 2011;
421 37: 443-448.
422

423 12. Stefaniak AB, Duling MG, Geer L, Virji MA Dissolution of the metal sensitizers Ni, Be, Cr in
424 artificial sweat to improve estimates of dermal bioaccessibility. Environmental Science:
425 Processes & Impacts 2014; 16: 341-351.
426

427 13. Hedberg Y, Midander K, Wallinder IO Particles, sweat, and tears: a comparative study on
428 bioaccessibility of ferrochromium alloy and stainless steel particles, the pure metals and their
429 metal oxides, in simulated skin and eye contact. Integr Environ Assess Manag 2010; 6: 456-468.
430

431 14. Kulthong K, Srisung S, Boonpavanitchakul K, Kangwansupamonkon W, Maniratanachote R
432 Determination of silver nanoparticle release from antibacterial fabrics into artificial sweat. Part
433 Fibre Toxicol 2010; 7: 8.
434

435 15. Duling M, Stefaniak A, Lawrence R, Chipera S, Abbas Virji M Release of beryllium from
436 mineral ores in artificial lung and skin surface fluids. Environ Geochem Health 2012; 34: 313-
437 322.
438

439 16. Hillwalker WE, Anderson KA Bioaccessibility of metals in alloys: evaluation of three
440 surrogate biofluids. *Environ Pollut* 2014; 185: 52-58.
441

442 17. Ertl H, Butte W Bioaccessibility of pesticides and polychlorinated biphenyls from house dust:
443 in-vitro methods and human exposure assessment. *J Expo Sci Env Epid* 2012; 22: 574-583.
444

445 18. Ruby MV, Davis A, Schoof R, Eberle S, Sellstone CM Estimation of lead and arsenic
446 bioavailability using a physiologically based extraction test. *Environ Sci Technol* 1996; 30: 422-
447 430.
448

449 19. Abdallah MA, Tilston E, Harrad S, Collins C In vitro assessment of the bioaccessibility of
450 brominated flame retardants in indoor dust using a colon extended model of the human
451 gastrointestinal tract. *J Environ Monit* 2012; 14: 3276-3283.
452

453 20. Buckley WR, Lewis CE The "ruster" in industry. *J Occup Med* 1960; 2: 23-31.
454

455 21. Nicolaides N Skin lipids: their biochemical uniqueness. *Science* 1974; 186: 19-26.
456

457 22. Stefaniak AB, Harvey CJ. Artificial skin surface film liquids. In: Google Patents, 2008.
458

459 23. Lane ME Skin penetration enhancers. *Int J Pharm* 2013; 447: 12-21.
460

461 24. Pont AR, Charron AR, Brand RM Active ingredients in sunscreens act as topical penetration
462 enhancers for the herbicide 2,4-dichlorophenoxyacetic acid. *Toxicol Appl Pharmacol* 2004; 195:
463 348-354.
464

25. Walters KA, Brain KR, Howes D, James VJ, Kraus AL, Teetsel NM *et al* Percutaneous penetration of octyl salicylate from representative sunscreen formulations through human skin in vitro. Food Chem Toxicol 1997; 35: 1219-1225.
26. Abdallah MA, Harrad S, Covaci A Hexabromocyclododecanes and tetrabromobisphenol-A in indoor air and dust in Birmingham, U.K: implications for human exposure. Environ Sci Technol 2008; 42: 6855-6861.
27. Brommer S, Harrad S, Van den Eede N, Covaci A Concentrations of organophosphate esters and brominated flame retardants in German indoor dust samples. J Environ Monitor 2012; 14: 2482-2487.
28. Stefaniak AB, Harvey CJ Dissolution of materials in artificial skin surface film liquids. Toxicol in Vitro 2006; 20: 1265-1283.
29. Abdallah MA, Uchea C, Chipman JK, Harrad S Enantioselective Biotransformation of Hexabromocyclododecane by in Vitro Rat and Trout Hepatic Sub-Cellular Fractions. Environ Sci Technol 2014; 48: 2732-2740.
30. Abdallah MA, Covaci A Organophosphate flame retardants in indoor dust from Egypt: implications for human exposure. Environ Sci Technol 2014; 48: 4782-4789.
31. Qiao GL, Brooks JD, Riviere JE Pentachlorophenol dermal absorption and disposition from soil in swine: Effects of occlusion and skin microorganism inhibition. Toxicol Appl Pharm 1997; 147: 234-246.

32. Williams RL, Reifenrath WG, Krieger RI Artificial sweat enhances dermal transfer of chlorpyrifos from treated nylon carpet fibers. *J Environ Sci Heal B* 2005; 40: 535-543.
33. van der Veen I, de Boer J Phosphorus flame retardants: Properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* 2012; 88: 1119-1153.
34. Fang M, Stapleton HM Evaluating the Bioaccessibility of Flame Retardants in House Dust Using an In Vitro Tenax Bead-Assisted Sorptive Physiologically Based Method. *Environ Sci Technol* 2014; 48: 13323-13330.
35. Brommer S Characterising human exposure to organophosphate ester flame retardants. PhD thesis University of Birmingham ethesesbhamacuk/5292/5/Brommer14PhDpdf 2014.
36. USEPA Exposure factors handbook. www.epa.gov/ncea/efh/pdfs/efh-complete.pdf 2011.
37. OEHHA Office of Environmental Health Hazard Assessment, State of California, Environmental Protection Agency, Safe drinking water and toxic enforcement act of 1986 safe drinking water and toxic enforcement act of 1986, PROPOSITION 65. http://oehha.ca.gov/prop65/prop65_list/files/P65single082515.pdf 2015.
38. Ali N, Dirtu AC, Eede NV, Goosey E, Harrad S, Neels H *et al* Occurrence of alternative flame retardants in indoor dust from New Zealand: Indoor sources and human exposure assessment. *Chemosphere* 2012; 88: 1276-1282.

9 Tables

Table 1: Effect of the composition of synthetic sweat and sebum mixture (SSSM) on the bioaccessibility ($f_{\text{bioaccessible}}$) of target FRs from indoor dust

	$f_{\text{bioaccessible}}$ (%) for different SSSM compositions						
Compound	100% Sweat	99:1 sweat:sebum	95:5 sweat:sebum	9:1 sweat:sebum	8:2 sweat:sebum	1:1 sweat:sebum	100% Sebum
α -HBCD	2.3 ± 0.2	2.4 ± 0.2	12.0 ± 6.0	20.0 ± 2.8	36.1 ± 2.7	40.9 ± 2.9	50.5 ± 7.0
β -HBCD	1.6 ± 0.6	3.6 ± 0.7	10.1 ± 1.3	14.5 ± 5.7	29.7 ± 0.6	46.9 ± 3.4	60.4 ± 10.1
γ -HBCD	1.4 ± 0.1	4.1 ± 1.8	11.4 ± 2.2	19.0 ± 5.4	23.2 ± 6.5	49.6 ± 5.8	67.2 ± 3.37
Σ -HBCD	1.8 ± 0.2	3.3 ± 0.89	11.47 ± 4.1	18.7 ± 4.0	30.0 ± 4.2	45.2 ± 4.1	58.5 ± 5.7
TBBPA	3.5 ± 0.5	4.8 ± 1.9	9.6 ± 4.2	25.2 ± 7.1	32.4 ± 5.4	39.5 ± 4.3	55.7 ± 8.5
TCEP	16.0 ± 1.22	15.8 ± 0.8	14.8 ± 0.9	12.2 ± 1.0	11.2 ± 1.4	10.4 ± 1.8	22.3 ± 2.3
TCIPP	12.4 ± 4.4	15.4 ± 2.8	20.6 ± 3.2	8.4 ± 2.2	11.9 ± 1.9	17.4 ± 2.7	26.9 ± 6.4
TDCIPP	11.9 ± 3.6	12.0 ± 0.5	13.0 ± 0.4	10.5 ± 0.4	12.4 ± 0.3	18.6 ± 0.8	28.1 ± 0.6

*Experiments were performed in triplicate, results are presented as average ± SD

519 **Table 2: Parameters used in dermal exposure assessment of target FRs in indoor dust ³⁶.**

Parameter	Adult	Toddler
Age	>18 years	2-3 years
Body weight	70 Kg	15 Kg
Body surface area	1.94 m ²	0.6 m ²
Skin surface exposed	4615 cm ² (head, forearms, hands and feet)	2564 cm ² (head, extremities including hands and feet)
Dust adhered to skin	0.01 mg/cm ²	0.04 mg/cm ²
Indoor exposure fraction ²⁶		
House	63.8%	86.1%
Office	22.3%	-
Car	4.1%	4.1%

520

521

522

523

524 **Table 3: Assessment of human dermal exposure (ng/kg bw/day) to FRs present in indoor dust upon contact with a skin surface**
525 **film composed of 1:1 sweat:sebum**

FR/ Scenario	UK Adult			UK Toddler		
	Low	Average	High	Low	Average	High
α-HBCD	0.1	1.0	14.1	0.1	1.1	16.9
β-HBCD	<0.1	0.3	6.7	<0.1	0.4	7.9
γ-HBCD	0.2	3.0	25.8	0.2	3.3	29.7
TBBPA	<0.1	0.1	0.7	<0.1	0.1	0.9
TCEP	<0.1	0.1	3.7	<0.1	0.5	17.4
TCIPP	0.3	0.5	6.4	3.9	4.7	46.8
TDCIPP	<0.1	0.1	2.2	<0.1	0.9	19.2

526

527

528

529

530

531

532

533

534 **10 Figures**

535 **Figure 1:** Schematic illustration depicting the structure of the skin and the absorption process for FRs in indoor dust in the presence of
536 sweat/sebum mixture and topically applied cosmetics.

537

538

539

540

541

542 **Figure 2: Effect of applied cosmetics on the bioaccessibility ($f_{\text{bioaccessible}}$ %) of target FRs from indoor dust.**

543

544 * Denotes a statistically significant difference ($P < 0.05$) from the control group.

545

546

547 **Figure 3: Comparison for (a) UK adults and (b) toddlers of exposure (ng/kg bw. day) to**
 548 **FRs in indoor dust via dermal contact (this study, average exposure scenario) and dust**
 549 **ingestion** ^{26, 35}.