On the origin of AMS "cooking Organic Aerosol" at a rural site

Dall'Osto, M.; Paglione, M.; Decesari, S.; Facchini, M. C.; O'Dowd, C.; Plass-Duellmer, Christian; Harrison, Roy M.

DOI:

[10.1021/acs.est.5b02922](https://doi.org/10.1021/acs.est.5b02922)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Dall'Osto, M, Paglione, M, Decesari, S, Facchini, MC, O'Dowd, C, Plass-Duellmer, C & Harrison, RM 2015, 'On the origin of AMS "cooking Organic Aerosol" at a rural site', *Environmental Science & Technology*, vol. 49, no. 24, pp. 13964-13972. <https://doi.org/10.1021/acs.est.5b02922>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Copyright © 2015 American Chemical Society

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

ON THE ORIGIN OF AMS “COOKING ORGANIC AEROSOL” AT A RURAL SITE

M. Dall’Osto^{1,2,3*}, M. Paglione⁴, S. Decesari⁴,
C. Facchini⁴, C. O’Dowd³, Christian Plass-Duelli⁵ and Roy M.
Harrison^{2,6}

¹Institut de Ciències del Mar, CSIC
Pg Marítim de la Barceloneta 37-49 08003 Barcelona, Spain

²National Centre for Atmospheric Science
School of Geography, Earth & Environmental Sciences
University of Birmingham
Edgbaston, Birmingham, United Kingdom

³School of Physics, National University of Ireland
Galway, Ireland

⁴National Research Council (CNR)
Institute of Atmospheric Sciences and Climate (ISAC)
Via Gobetti 101, 40129 Bologna, Italy

⁵Meteorologische Observatorium Hohenpeißenberg
Albin Schwaiger Weg 10
82383 Hohenpeißenberg
Germany

⁶Department of Environmental Sciences /
Center of Excellence in Environmental Studies
King Abdulaziz University, PO Box 80203
Jeddah, 21589, Saudi Arabia

* Corresponding Author: Manuel Dall’Osto; Email: [dallost@icm.csic.es](mailto:dallosto@icm.csic.es)

Abstract

A number of field observations employing aerosol mass spectrometers (AMS) have demonstrated that organic matter rich in monocarboxylic acids and aliphatic carbonyls originating from cooking activities (the COA factor) contributes significantly to ambient Organic Matter (OM) in urban environments. Little is known about the contribution and nature of COA in rural localities. We studied the correlation of COA with chemical tracers at a rural site in the Po Valley, Italy. Our statistical approach, based on Positive Matrix Factorization (PMF) shows that the COA factor was clearly linked to local emissions of chloride and methane sulphonic acid (MSA), chemical tracers not associated with cooking emissions, or with combustion sources. While the association with Cl is not understood at this stage, the emission of reduced sulphur compounds, aliphatic carbonyls and monocarboxylic acids is consistent with several agricultural practices (e.g., manure storage) and waste disposal systems (e.g., landfills) which characterize the sub-urban and rural areas of the Po Valley and of other many populated environments. It is concluded that the nature and origins of the AMS COA factor measured at a rural site are complex and include far more than the emissions from food cooking.

KEY WORDS: Aerosol mass spectrometer; cooking organic aerosol; COA; ATOFMS

69 1. INTRODUCTION

70
71 Urban air quality provides one of the main drivers for the study of atmospheric science,
72 especially due to growing urbanization. Although considerable progress in improving air
73 quality has been made, it is imperative to better understand the source attribution of
74 particles and health effects in the urban atmosphere¹. Among the constituents, organic
75 aerosol (OA) accounts for a large fraction of urban particulate matter². Primary OA (POA)
76 is directly emitted from fossil fuel combustion, biomass burning, and other sources, but the
77 atmospheric evolution of POA after emission remains poorly characterized³⁻⁵. Whilst road
78 traffic is often a major contributor to aerosol in urban areas next to major roads, OA in
79 urban background areas is not easily associated with a specific source. Few studies have
80 sought to characterise the sources of OA in rural areas, remote from major sources.

81
82 In recent years Mass Spectrometry of Atmospheric Aerosol (MSAA) has become one of
83 the fastest growing area of aerosol research⁶. Such techniques have greatly enhanced our
84 capacity of observing the atmospheric processes responsible for the formation and
85 evolution of airborne particles. In this regard, the Aerodyne Aerosol Mass Spectrometer
86 (AMS) is able not only to measure the OA concentration and size distribution but also to
87 provide its mass spectrum⁷. Analysis of AMS data collected at a large number of sites has
88 revealed that the oxidised OA component (OOA) tends to dominate everywhere, including
89 some heavily urbanized regions^{2,8}.

90
91 Many AMS studies have focused on the application of factor analysis to the organic mass
92 fraction in an attempt to deconvolve it into descriptive sub-components, namely a
93 hydrocarbon-like organic aerosol (HOA) factor, an oxygenated organic aerosol (OOA)
94 factor, and a semi-volatile factor (SV-OOA)^{4,9}. Such assignments are based primarily
95 upon the main mass spectral components and the diurnal profile. In some cases,

96 supporting evidence from chamber experiments is available. Recently, a previously
97 reported AMS spectrum^{10,11} has been firmly associated with primary emissions from
98 cooking activities. Such emissions have been claimed to account for up to half of the total
99 primary urban OA at sites in Europe¹²⁻¹⁴, Asia¹⁵ and America¹⁶. However, there is a
100 possibility for a factor to include a combination of factors associated with different
101 sources^{11,17}, and some recent studies report a COA factor lacking the diurnal profile
102 expected from cooking activities, although possible explanations have been given^{13,18}.
103 Other studies have also expressed the view that it is most reasonable to characterize the
104 COA component as "cooking influenced" but not purely from cooking sources¹⁹. In a
105 comparison of source apportionment by AMS-PMF with a Chemical Mass Balance model,
106 the AMS estimate of COA for a site in London exceeded the CMB-derived concentration of
107 cooking aerosol by a factor of ca 1.6²⁰.

108

109 Taken together, evidence for source attribution of the AMS COA factor is currently
110 incomplete and most importantly not well supported by other measurement techniques.
111 This lack of knowledge needs to be filled, and the goal of this paper is to use results of a
112 source apportionment study using six state-of-the-art spectrometric techniques deployed
113 at a rural site in the Po Valley (Italy) in 2009 to investigate the nature of the AMS COA
114 factor. Previous work²¹ provides a comprehensive picture of the nature of organic and
115 inorganic aerosols and aerosol precursors at a European rural site with an unprecedented
116 level of detail. Figure S1 shows an example of the high time resolution particle mass
117 spectrometry instruments deployed. In this study Positive Matrix Factorization (PMF) has
118 been applied to all the high time resolution data in order to better elucidate aerosol
119 sources not clearly identified when analyzing results from individual aerosol techniques on
120 their own. Particular attention is given to the AMS COA factor, but a complete overview of
121 the aerosol sources apportioned is also presented and discussed.

2. EXPERIMENTAL

2.1 Study Site

The Po Valley is located in Northern Italy between two mountain ranges, the Alps in the north and west and the Apennines in the south. The Po Valley has 20 million inhabitants spread over an area of 48,000 km² (Fig S2). In the present study we used the EMEP Station of San Pietro Capofiume (SPC, 44°23'N 11°22'E, 11 m a.s.l.), a rural background site (distance from major pollution sources: 10 – 50 km²²). The measurements were conducted from 26 June to 15 July 2009.

2.2 Aerosol Mass Spectrometry Techniques

Four on-line aerosol spectrometers were used: the TSI Aerosol Time-Of-Flight Mass Spectrometer (ATOFMS), the Aerodyne High-Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-TOF-AMS) and Soot Particle Aerosol Mass Spectrometer (SP-AMS), and finally a Thermal Desorption Aerosol Gas Chromatograph AMS (TAG-AMS). Twelve-hour resolution proton nuclear magnetic resonance (H-NMR) spectra were also obtained by off-line analysis. Finally, information on gas-phase precursors of secondary aerosols was acquired using Chemical Ionization Mass Spectrometry (CIMS). A set of other measurements are described in detail elsewhere²¹. Overall, the state-of-the-art instruments used for this analysis measure different species with different accuracy and precision, as discussed in great detail in other publications²³⁻²⁵.

2.3 Factor Analysis

Factor analysis involves a wide set of multivariate statistical techniques that have been extensively used in atmospheric science. Its aim in this field is to apportion aerosol sources on the basis of the internal correlations of observational data collected at a

155 measurement point, called the “receptor site”. Receptor modeling by factor analysis does
156 not need any detailed a prior knowledge of source profiles and it is therefore very useful
157 for determination of aerosol fractions of secondary origin. “Positive Matrix Factorization”
158 (PMF)^{26,27} has been used extensively for source apportionment of ambient particulate
159 matter (PM), and in particular is so far the most widespread tool for AMS spectral data
160 analysis²⁸. In this study the EPA open-source software EPA-PMF v3.0 was used for PMF
161 analysis. PMF was applied on the hourly data obtained by the analytical techniques
162 described above and also by the previous PMF analysis on spectroscopic data reported
163 elsewhere²¹.

164

165 The criteria adopted in this study for input data treatment, including the uncertainty matrix,
166 and for the evaluation of the PMF solutions are discussed extensively in the
167 supplementary material.

168

169 **3. RESULTS**

170 PMF analysis was performed for one to ten factors, and it was found that the solutions with
171 factor numbers greater than five provided no new meaningful information for both
172 datasets. Rotational constraint was explored with Fpeak. The solutions ultimately selected
173 was based on an Fpeak value of zero. Additional factors resulted in a splitting of the
174 existing factors (mainly splitting of secondary organic and inorganic components (Figures
175 S5 and S6; Table S2 and S3). Table S4 shows there is no clear improvement with the six
176 factor solutions from either of the two datasets. A full list of variables is found in Table S1.
177 The figures in brackets after the PMF input variable represent the percent contribution of
178 that atmospheric variable within the whole five factor solution (% of species). A description
179 of the strong agreement between the solutions from the two datasets is presented at the

180 end of this section. More details on the PMF diagnostics are presented in supplementary
181 material.

182

183

184 **3.1 Dataset 1 : Results from the Full Campaign ("ALL")**

185

186 Five factors were found very well describing the long time series dataset (16 days). This
187 dataset included the first week of the study when the anticyclonic conditions favoured
188 transport of aerosol at the regional scale and the recirculation of pollutants in the Po
189 basin²¹. Figure 1 and table 1 show all the PMF factor profiles of this solution, whereas the
190 temporal trends and the diurnal profiles can be seen in Figures 2 and S9a, respectively.

191 The individual factors are as follows:

- 192 • Oxidised Organic Aerosol from photooxidation processes "Photox OOA" (30% of the
193 total, Figure 1a). It represents oxidised organic aerosols (accounting for a substantial
194 percentage of AMS_LV-OOA MO 48% and AMS_LV-OOA LO 49%), AMS_sulphate
195 (48%), Ozone (83%) and CIMS_MA (84%). This is associated with photochemically
196 aged secondary aerosols, and the temporal trends (Figure 2, S9a) show this aerosol
197 component occurring mainly during daytime and during the first week of the study.
- 198 • Nitrate "NIT-Regional" (18%, Figure 1b). This factor is strongly associated with
199 AMS_Nitrate (79%) and ATOFMS_NIT-Regional (32%). There is an important
200 component of AMS_Cl (38%) and AMS_SV-OOA (27%) associated with it. This factor
201 is associated with nitrate containing aerosol of regional origin²⁹, and related to high
202 RH (Figure 2), with time trends modulated by the diurnal variations of temperature and
203 relative humidity which regulate the concentration of semivolatile compounds in the
204 aerosol (ammonium nitrate and chloride, SV-OOA).
- 205 • Sulphate "SUL-Regional" (23%, Figure 1c). This aerosol source is mainly composed of
206 sulphate (ATOFMS_SUL-Reg 90%), ATOFMS_EC-Reg (72%), and AMS_OOA-MO

(32%). This aerosol source accounted for 35% of explained variation during the first week of the study (stagnant air), but only 3% during the second week (Figure S9a). Part of this component is thought to be the aerosol core from the evaporating regional nitrate during daytime^{21,29}. This factor seems to be linked to the transport of sulphate and BC during the days of anticyclonic conditions.

- "NO" (12%, Figure 1d). This represents the smallest aerosol source contribution as expected given the rural location of San Pietro Capofiume. The factor is well described primarily by NO (90%), followed by minor contributions from AMS_HOA (23%) and NO₂ (25%). BC is only partially described by this factor (6%). Although the diurnal profile (Figure 2) shows a main morning spike during the urban rush hour (7am), processes other than traffic and certainly including photolysis of NO₂ and of HONO most probably contributed to NO at the site. Other studies^{21,30,31} have already demonstrated the presence of a traffic-influenced component at SPC station with contributions ranging between 14-24% of total OA. This non-negligible contribution and its diurnal trend could be explained both by the proximity of main transportation routes and by the accumulation of the primary emissions overnight because of the reduced atmospheric mixing and dispersion. However the NO apportioned in this study could be also formed from photochemical activity (photolysis of NO₂ in the first hours after the sunrise) and so we prefer to label this factor simply as "NO", the most represented species, without assigning a specific source.

- "Cooking" Organic Aerosol "COA" (17%, Figure 1e). This factor accounts for a high proportion of AMS COA (77%) and ATOFMS OC-SUL-NIT (77%). Other interesting variables describing this factor includes CIMS MSA (41% of the total), AMS Cl (35%) and NO₂ (31%). The temporal trends show a major occurrence during the second week (24%) rather than the first regional pollution week (14%). The diurnal profile shows a minor spike during evening time (7-9pm), followed by sustained

concentrations overnight. There is a lack of the midday peak seen associated with COA factors in many studies^{13,18}.

The analysis of the correlation of the five factor concentrations with local wind direction (Figure S10a) shows that the first three factors are associated with sustained winds from either east (OOA), west (NIT-regional) or south-west (SUL-regional) directions, while the COA factor is most concentrated in calm conditions. Therefore, the COA is more influenced by local sources compared to the first three factors which are mainly transported to the site. The NO factor shows an intermediate behaviour, with a component from the west (possibly NO_x transport) and another local (possibly soil emissions³²).

3.2 Dataset 2: Three Measurement Days Including SP-AMS Data (“HIGH” Resolution)

The factors of the dataset_2 (HIGH) PMF solution can be seen in Table 1 and Figure 3. In the last days of the campaign, which were covered by the measurements described in this section, the weather conditions were unfavourable for pollutant recirculation and the aerosol concentrations were fairly low in day-time when the planetary boundary layer was fully developed. Conversely, the concentrations at night-time and in the early morning continued to be sustained by local sources in the Po Valley²¹. The analysis of this campaign period is therefore particularly useful for the source apportionment of the aerosol components originating from emissions in the Po Valley, such as cooking aerosols, because the interference from the background particles is smaller with respect to the first part of the campaign²¹. The diurnal profiles can be seen in Figure 2 and the temporal trends in Figure S9b. Three of the five factors presented previously (Section 3.1, ALL) were also found in this second solution, including:

- 259 • Oxygenated Organic Aerosol "Photox OOA" (38%, Figure 3a). This described
 260 Secondary Organic aged aerosols accounting for a high percentage of AMS LV-OOA
 261 MO (81%), SP-AMS "SV-OOA Day" (89%), ozone (83%) and sulphate from both HR-
 262 AMS (68%) and SP-AMS (67%) instruments.
- 263 • Nitrate regional "NIT-Regional" (17%). A strong nitrate signal from all the on-line
 264 spectrometers including HR-AMS (72%), SP-AMS (45%) and ATOFMS (NIT-Reg
 265 71%) can be seen (Figures S9b and 2). Additionally, an hydrocarbon like contribution
 266 of HOA (21%) and BC (20%) is notable.
- 267 • "NO" (17%). This shows a strong NO signature (93%) followed by HOA (22%) and BC
 268 (11%) (Figure 3c). The diurnal profile seen in Figures 2 and S9b shows a morning
 269 peak likely to be associated with traffic and/or photolysis.

270

271 Unlike the "ALL" solution, in this 3-day dataset_2 (AMS, ATOFMS, SP, CIMS, gas, 33
 272 variables, last 3 days of the field study, dataset_2, HIGH) a factor related to sulphate of
 273 regional origin was not identified. This is very likely due to the fact that during the second
 274 week of the field study there was little contribution of regional pollution, so the solution was
 275 not able to extract the SUL-Reg factor, mainly seen during the first week (Figure S9a). By
 276 contrast, during nighttime the relatively higher concentration of nitrate allowed identification
 277 of the NIT-Reg. factor. During daytime the SUL-Reg., if present, is likely to be incorporated
 278 in the regional OOA factor.

279

280 However, the second solution with more variables (33 in total) was able to provide a better
 281 description of the COA factor, because the statistical correlations of AMS COA
 282 concentrations with the other variables are much more clear in the HIGH dataset than in
 283 the ALL one (Table S6). The factor analysis of the HIGH set of variables actually provided
 284 two different aerosol sources associated with the AMS "cooking aerosols" and specifically:

- "Cooking" organic aerosol "COA-MSA-Cl" (seen in Figure 3d). This factor presents a strong AMS COA signal (54%), with CIMS_MSA (36%) and a strong inorganic AMS Cl signature (70%). A strong semi-volatile component is also present (AMS SV-OOA and SP-AMS "SV-OOA Night", both 58%). Additionally, the factor ATOFMS OC-SUL-NIT is also associated with this factor (32%). In summary, this COA factor is associated with MSA, chloride and semi-volatile OA components.
- "Cooking" organic aerosol "COA-MSA-HOA" (seen in Figure 3e). This factor shares the part of the AMS_COA signal (46%) not identified by the previous factor (54%) and a strong association with CIMS MSA (37%). However, this factor is more associated with AMS HOA (38%) SP-AMS HOA (42%) and contains almost no semi-volatile component. Interestingly, this factor is associated with nitrate of local origin (ATOFMS NIT-Local, 42%). In support of this, it accounts for the highest percentage of NO₂ (33%) of all factors. In summary, this second "COA" factor is associated with anthropogenic HOA and somehow related to nitrate locally formed, organic nitrate and nitrogenous gaseous²⁹. This fingerprint could be characteristic of urban sources near to the site.

3.3 Comparison Among the two PMF Solutions and External Correlations

Support

The two PMF solution (ALL and HIGH) provided three common factors (OOA, NIT-Reg and Traffic). When correlation of the overlapping temporal trends of the last three days are examined, Table S4 shows a very good agreement between the two PMF solutions: OOA (34-38%, $R^2=0.95$), Nitrate regional (14-17%, $R^2=0.94$) and NO (14-17%, $R^2=0.85$). Sulphate Regional cannot be temporally compared, although the apportionment

contributions is similar for both solutions (0-3%). The "Cooking" factor solution for the dataset_ALL is found to contribute less (25%) than the sum of the two "cooking" aerosol factors of the dataset_HIGH (38%). However, it is important to note that the temporal correlation between "COA" from dataset_ALL and the sum of COAs from dataset_HIGH ("COA-MSA-CI" and "COA-MSA-HOA" together) is very high ($R^2=0.85$).

The factors presented are strongly supported by correlations with external measurements presented elsewhere²¹. Table S5 shows that factor Photox. OOA correlates very well with NMR_PMF_F4 (Aged humic-like substances, $R^2=0.90$), whereas factor SUL-Reg. correlates well with NMR_PMF_F3 (organic sulphate and less aged humic-like substances, $R^2=0.57$). Furthermore, whilst COA-MSA-CI correlates only with NMR_PMF_F1 (aliphatic amines and unspeciated aliphatic compounds, $R^2=0.77$), factor COA-MSA-HOA correlates only with NMR_PMF_F2 (aliphatic alkanolic acids and oxo-acids, $R^2=0.95$). These are robust external correlations which strongly support our PMF solutions. Correlations reported in Table S4 are much stronger ($R^2=0.7-1.0$) than those previously reported ($R^2=0.3-0.6$)²¹ because the chemical profiles of the PMF solutions herein presented are less dependent upon specific markers, but rather show a combination of markers that better describe an organic aerosol source.

4. DISCUSSION

4.1 Major Constituents

The analysis of the correlations between the "regional" aerosol components (i.e., the components expected to exhibit an extended source footprint) suggests that the OOA atmospheric evolution over north Italy is completely distinct from that of nitrate of regional origin. Factor "Photox. OOA" was found mainly associated with oxidised organic aerosols (48-88%) and sulphate (48-68%). This is in line with previous studies³³ who reported aged

340 OOA detected above the Po Valley column to be secondary in nature and highly oxidized
341 in the regional aerosol, with much higher amounts of sulphate and organics relative to
342 nitrate. By contrast, NIT-Reg. is found to account for the majority of nitrate aerosol
343 detected (AMS_Nit 72-79%), of regional nature (ATOFMS NIT-Reg 32-71%). The organic
344 component in this factor is found to be of primary origin: AMS_HOA (21-45%). The
345 association of NIT-Reg. with HOA may suggest a larger size mode of nitrate during the
346 night acting as a coagulation sink for fresher finer local aerosols. Alternatively, the HOA
347 may in part be subjected to transport across the Po Valley basin together with nitrate. This
348 factor is associated with a specific mass spectrum and a characteristic temporal dynamic
349 already reported in London²⁹, Wales³⁴ and Barcelona³⁵. The LRT-NIT particle type is
350 volatile, going into the gas phase during the day and leaving a non-volatile internally
351 mixed core mostly composed of sulphate, elemental and organic carbon. The SUL-Reg is
352 mainly composed of EC-Sulphate (90%), BC (42%) and sulphate (27%). During this study
353 we find local, less aged nitrate of finer size mode of local origin (ATOFMS_NIT-local)
354 associated with COA-MSA-HOA.

355
356
357 HOA and BC are usually related to fresh traffic emissions in many AMS studies. This work
358 shows that traffic makes only a small contribution to BC (11%) and a modest one to HOA
359 (23%). By contrast, the majority of BC is related to the core of the regional nitrate, being
360 internally mixed with sulphate and BC (42% of total BC). As a result, it is important to
361 stress that the HOA and BC is mainly seen during nighttime and mainly associated with
362 nitrate.

363
364
365
366

367 **4.2 The "Cooking" Aerosol Factor(s)**

368
369

370 The mass spectrum of the AMS COA factor obtained from San Pietro Capofiume was
371 converted in m/z unit mass resolution and compared with other factors in the literature. All
372 factors were compared at unit mass resolution. The COA factor did not correlate ($R^2 < 0.1$)
373 with any of the HOA, SV-OOA or LV-OOA in the literature. A good correlation (R^2) was
374 found with AMS COA reported in seven previous studies: 0.84 (laboratory studies¹¹); 0.74
375 and 0.83 (London¹²); 0.75 (Manchester¹²); 0.80 (Paris^{14,36}); 0.75 (Zurich¹⁰); 0.85 (Cork¹⁸).
376 Therefore, the same name "COA" was kept for this study.

377

378 The composition of the aerosol and of the air masses associated with the AMS COA
379 factors are key indicators of its origins. A first, unexpected finding is that the PMF analyses
380 from both datasets (ALL and HIGH) show the association of AMS COA with secondary
381 organic aerosol components (CIMS MSA and ATOFMS OC-SUL-NIT) rather than with
382 markers of primary combustion (or thermal) processes. More specifically, the analysis of
383 the HIGH dataset decomposes the "COA" factor into two, with one (COA-MSA-HOA)
384 retaining a partial fingerprint for primary sources (HOA, BC, NO_x) together with that of
385 secondary components (nitrate, MSA) while the other factor (COA-MSA-CI) shows very
386 small correlations with the primary tracers. Fig. S11 shows the main differences among the
387 two COA PMF factors. The low correlation of COA with tracers of urban pollution in this
388 last factor was unexpected because cooking aerosols should be transported in urban
389 plumes to the countryside along with the products of traffic emissions (BC, NO_x, HOA). It is
390 important to stress that very little is known on the transformation of COA in the
391 atmosphere³⁷. Preliminary findings of meat cooking aerosol ageing show an increase of
392 the AMS O:C ratio from 0.1 up to 0.3³⁸. By looking at the prevalent wind directions
393 associated with the two COA factors in the HIGH datasets, the COA-MSA-HOA shows a
394 transport component from the west, while the COA-MSA-CI has a more local footprint.
395 Therefore, it is very unlikely that the COA-MSA-CI originated from cooking activities in

396 urban areas. The local sources to be indicated could be cooking in rural houses and small
397 towns in the vicinity of SPC, but this does not explain the low correlation with the traffic
398 markers (as people who live there still use cars). Most importantly, the source profile
399 emerging from COA-MSA-Cl is inconsistent with the known composition of cooking
400 aerosol, in which particles are mainly organic in nature with inorganic ions being present
401 only in trace amounts³⁷, while here the aerosol mass AMS Cl/ AMS COA ratio is 1.63.
402 These findings indicate that either the published emission composition for cooking
403 emissions does not apply to this environment, or cooking is not a major source for the
404 organic aerosol included in the COA-MSA-Cl factor.

405

406 Explaining the correlation between COA and aerosol chloride is challenging. Aerosol
407 chloride in this study is mainly non-sea salt chloride, which may originate from various
408 anthropogenic sources including industry, combustion and incinerators^(39,45). There are no
409 industrial plants around SPC. Incinerators emit hydrochloric acid which comes from the
410 combustion of plastic material^(39,45). However, the single particle mass spectra associated
411 with this factor (ATOFMS_OC-NIT-SUL) does not have metals previously identified with
412 Cl-containing aerosols emitted by waste incinerators³⁹. Also the SP-AMS did not observe
413 any mass fragments from metals such as lead which typically characterize the emissions
414 of incinerators (Dr. J. Allan, personal communication).

415

416 Finally, the source profile of COA-MSA-Cl indicates that the same source is responsible for
417 the emission of reduced sulfur species (precursors of gas-phase MSA measured by CIMS)
418 which is inconsistent with the hypothesis of a high-temperature combustion process such
419 as in an incinerator. This is because in thermal oxidizers such as industrial incinerators
420 chemically reduced pollutants are generally destroyed via combustion forming more
421 oxidised species such as CO₂, SO₂ and H₂O. On the contrary, the production of reduced

sulfur species points to thermal processes occurring at low temperature (including some cooking practices) or to emissions at ambient temperature⁴⁰. Dimethylsulfide emissions from marine biota are believed to be largely responsible for global MSA production⁴¹; however, in this study gas-phase MSA was not associated with marine air masses. On the contrary the MSA concentrations reach a maximum at nighttime when the circulation is mainly from the inland to the sea²¹. A recent review on cooking aerosol tracers actually does not mention sulphur compounds as cooking tracers³⁷. However, it is worth mentioning that roasting of coffee beans may be a source of reduced sulphur compounds⁴² and also meat cooking can produce several thioethers⁴³. However, cooking is only one of the possible anthropogenic sources of sulfur compounds. Urban sources of DMS include aerobic composting of food wastes^{40,44-47}. More generally, volatile organic sulphur compounds can readily form from the aerobic and anaerobic degradation of organic matter, including solid wastes, wastewater, manure and livestock excreta and feed⁴⁴⁻⁴⁸. If we quantify such sources using co-emitted compounds such as ammonia, then emissions in the Po Valley are dominated by the agricultural practices and animal husbandry activities. A prevalent source of MSA precursors in rural areas, with emissions from manure management, livestock and agricultural land is consistent with the appearance of the COA-MSA-Cl factor as a local source at SPC unrelated to urban sources. The origin of organic particulate matter associated with the aerobic/anaerobic organic matter decomposition in agricultural land is poorly documented. Broadly, air pollutants emitted from the agricultural sector are mainly methane (CH₄) and other VOCs, nitrous oxide (N₂O) and ammonia (NH₃). Agriculture is also a main source of PM, both primary and secondary in origin originating from livestock production, application of fertilizers and pesticides, land preparation, harvesting and field burning of agricultural waste⁴⁶⁻⁴⁸. It is known that some husbandry activities (poultry) are strong point sources of PM₁₀ and PM_{2.5}, but this aerosol source has never been related to the AMS COA prior to

448 this study. By contrast, the emissions of VOCs from livestock and manure management
449 have been characterized in some detail and often associated with the production of
450 reduced sulfur species⁵⁰⁻⁵¹. Such VOCs comprise low-molecular weight organic acids, but
451 also C₆-C₁₀ aliphatic aldehydes and monocarboxylic acids. These chemical compounds
452 are analogous to the organic compounds believed to contribute to the AMS COA but
453 exhibit too low a molecular weight to exist in the particulate phase. Therefore, the
454 production of COA from these VOCs is possible only via a gas-phase oxidation step. C_{>8}
455 n-alkanals in particular can form SOA under low NO_x conditions⁵², while unsaturated
456 alkanals⁵³ are good SOA precursors also in NO_x-rich environments⁴⁹. According to this
457 hypothesis, the COA in the rural Po Valley may be contributed from secondary organic
458 compounds in the same manner that MSA is produced by the oxidation of reduced sulfur
459 species. It is worthy of note that MSA and COA concentrations peak at nighttime, when
460 OH concentrations are near zero while the nitrate radical NO₃ becomes the major
461 oxidant⁵²⁻⁵⁴. Unlike OH, the concentrations of NO₃ depend on the availability of NO_x, which
462 can explain why the COA-MSA factors show a component of westerly transport, i.e., from
463 the more polluted sector of the valley. It can also explain the correlation of COA with urban
464 tracers (including NO_x) in the factor COA-MSA-HOA. The correlation of COA with aerosol
465 chloride cannot be fully explained by this hypothesis. Possible point sources for
466 hydrochloric acid, reduced sulfur species and VOCs from organic matter degradation are
467 landfills, and there are actually two at ten-twenty km west of SPC.

468

469 In conclusion, the origin of factor COA-MSA-Cl (21% of total aerosol, 54% of AMS COA)
470 can be explained by hypothesizing emissions from agricultural/husbandry activities with a
471 potential additional contribution from waste disposal at urban sites. A second AMS COA
472 source (COA-MSA-HOA, 17% of the total aerosol, 46% of the remaining AMS COA) was

also linked with secondary components (nitrate, MSA), although retaining a partial fingerprint for primary traffic sources (HOA, BC, NO_x).

Quantification of such sources is challenging and calls for more research. However, the results of this study suggests that the current estimates of OA sources from organic matter degradation processes in agricultural and waste systems in populated areas may be substantially underestimated. The COA factor as determined by the AMS includes far more sources and processes than solely primary emissions from cooking activities.

Associated content

Supporting Information

A description and details of the field campaign discussed in the manuscript; PMF analysis, temporal trends of the PMF solutions, external correlation between this PMF study and previous ones are discussed. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Author information

Corresponding author

Manuel Dall'Osto: Phone: +34 644 053 801, Fax: +34 93 2309555, Email: dallosto@icm.csic.es

Notes

The authors declare no competing financial interest.

Acknowledgements. This work was funded by European integrated project on aerosol cloud climate and air quality interactions (no. 036833-2, EUCAARI). Manuel Dall'Osto and Roy M. Harrison thank the UK National Centre for Atmospheric Science for financial support. We acknowledge funding from the EU FP7-ENV-2013 programme "impact of

500 Biogenic vs. Anthropogenic emissions on Clouds and Climate: towards a Holistic
501 UnderStanding” (BACCHUS), project no. 603445. Programme Ramon y Cajal (RYC-2012-
502 11922) also provided financial support. Dr. James Allan (NCAS, University of Manchester,
503 England) is particularly acknowledge for very fruitful discussion which much improved the
504 manuscript.

505

506

507 **References**

508

509

510 (1) Pope, C. A. and Dockery, D. W.: Health effects of fine particulate air pollution: Lines
511 that connect, *J. Air Waste Manage.*, **2006**, 56, 709–742.

512

513 (2) Zhang, Q., Jimenez, J.L., Canagaratna, M.R., Allan, J.D., Coe, H., Ulbrich, I.,
514 Alfarra, M.R., Takami, A., Middlebrook, A.M., Sun, Y.L., et al. Ubiquity and dominance of
515 oxygenated species in organic aerosols in anthropogenically influenced northern
516 hemisphere mid-latitudes. *Geophys. Res. Lett.*, **2007**, 34 L13801.

517

518 (3) Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M.,
519 Grieshop, A. P., Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking organic aerosols:
520 Semivolatile emissions and photochemical aging, *Science*, **2007**, 315, 1259–1262.

521

522 (4) Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll,
523 J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., et al.: Evolution of organic aerosols in
524 the atmosphere, *Science*, **2009**, 326, 1525–1529.

525

526 (5) Dall'Osto, M., Thorpe, A., Beddows, D. C. S., Harrison, R. M., Barlow, J. F., Dunbar, T.,
527 Williams, P. I., and Coe, H.: Remarkable dynamics of nanoparticles in the urban
528 atmosphere, *Atmos. Chem. Phys.*, **2011**, 11, 6623–6637, doi:10.5194/acp-11-6623-2011

529

530 (6) Laskin A, Laskin J, Nizkorodov SA. Mass spectrometric approaches for chemical
531 characterization of atmospheric aerosols: critical review of the most recent advances.
532 *Environ Chem* **2012**;9:163–89.

533

534 (7) Jayne, J. T., Leard, D. C., Zhang, X., Davidovits, P., Smith, K. A., Kolb, C. E., and
535 Worsnop, D. R.: Development of an aerosol mass spectrometer for size and composition
536 analysis of submicron particles, *Aerosol Sci. Technol.*, **2000**, 33, 49–70.

537

538 (8) Donahue, N.M., Robinson, A.L. and Pandis, S.N. Atmospheric Organic Particulate
539 Matter: From Smoke to Secondary Organic Aerosol. *Atmos. Environ.*, **2009**, 43: 94–106.

540

541 (9) Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.:
542 Interpretation of organic components from Positive Matrix Factorization of aerosol mass
543 spectrometric data, *Atmos. Chem. Phys.*, **2009**, 9, 2891–2918, doi:10.5194/acp-9-2891-
544 2009.

545

- (10) Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., and Prevot, A. S. H.: Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, *Atmos. Chem. Phys.*, **2007**, 7, 1503–1522, doi:10.5194/acp-7-1503-2007.
- (11) Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of primary organic aerosol emissions from meat cooking, trash burning, and motor vehicles with High-Resolution Aerosol Mass Spectrometry and comparison with ambient and chamber observations, *Environ. Sci. Technol.*, **2009**, 43, 2443–2449, doi:10.1021/es8011518.
- (12) Allan, J. D., Williams, P. I., Morgan, W. T., Martin, C. L., Flynn, M. J., Lee, J., Nemitz, E., Phillips, G. J., Gallagher, M. W., and Coe, H.: Contributions from transport, solid fuel burning and cooking to primary organic aerosols in two UK cities, *Atmos. Chem. Phys.*, **2010**, 10, 647–668, doi:10.5194/acp-10-647-2010.
- (13) Mohr, C., DeCarlo, P. F., Heringa, M. F., Chirico, R., Slowik, J.G., Richter, R., Reche, C., Alastuey, A., Querol, X., Seco, et al.: Identification and quantification of organic aerosol from cooking and other sources in Barcelona using aerosol mass spectrometer data, *Atmos. Chem. Phys.*, **2012**, 12, 1649–1665, doi:10.5194/acp-12-1649-2012.
- (14) Crippa, M., DeCarlo, P. F., Slowik, J. G., Mohr, C., Heringa, M.F., Chirico, R., Poulain, L., Freutel, F., Sciare, J., Cozic, J., et al.: Wintertime aerosol chemical composition and source apportionment of the organic fraction in the metropolitan area of Paris, *Atmos. Chem. Phys.*, 2013, 13, 961–981, doi:10.5194/acp-13-961-2013.
- (15) Huang, X.-F., He, L.-Y., Hu, M., Canagaratna, M. R., Sun, Y., Zhang, Q., Zhu, T., Xue, L., Zeng, L.-W., Liu, X.-G., Zhang, Y.-H., et al.: Highly time-resolved chemical characterization of atmospheric submicron particles during 2008 Beijing Olympic Games using an Aerodyne High-Resolution Aerosol Mass Spectrometer, *Atmos. Chem. Phys.*, **2010**, 10, 8933–8945, doi:10.5194/acp-10-8933-2010.
- (16) Sun, Y. L., Zhang, Q., Schwab, J. J., Yang, T., Ng, N. L., and Demerjian, K. L.: Factor analysis of combined organic and inorganic aerosol mass spectra from high resolution aerosol mass spectrometer measurements, *Atmos. Chem. Phys.*, **2012**, 12, 8537–8551, doi:10.5194/acp-12-8537-2012.
- (17) He, L. Y., Huang, X. F., Xue, L., Hu, M., Lin, Y., Zheng, J., Zhang, R., and Zhang, Y. H. L.: Submicron aerosol analysis and organic source apportionment in an urban atmosphere in Pearl River Delta of China using high-resolution aerosol mass spectrometry, *J. Geophys. Res.*, **2011**, 116, D12304, doi:10.1029/2010JD014566.
- (18) Dall'Osto M, Ovadnevaite J, Ceburnis D, Martin D, Healy RM, O'Connor IP, et al. Characterization of urban aerosol in Cork city (Ireland) using aerosol mass spectrometry. *Atmos Chem Phys*, **2013**;13:4997–5015. <http://dx.doi.org/10.5194/acp-13-4997-2013>.
- (19) Hayes, P. L., A. M. Ortega, M. J. Cubison, K. D. Froyd, Y. Zhao, S. S. Cliff, W. W. Hu, D. W. Toohey, J. H. Flynn, B. L. Lefer, et al., Organic aerosol composition and sources in Pasadena, California, during the 2010 CalNex campaign, *J. Geophys. Res.*, **2013**, doi:10.1002/jgrd.50530.

- (20) Yin, J., Cumberland, S. A., Harrison, R. M., Allan, J., Young, D. E., Williams, P. I., and Coe, H.: Receptor modelling of fine particles in southern England using CMB including comparison with AMS-PMF factors, *Atmos. Chem. Phys.*, **2015**, 15, 2139-2158, doi:10.5194/acp-15-2139-2015.
- (21) Decesari, S., Allan, J., Plass-Duelmer, C., Williams, B. J., Paglione, M., Facchini, M. C., O'Dowd, C., Harrison, R. M., Gietl, J. K., Coe, H., et al.: Measurements of the aerosol chemical composition and mixing state in the Po Valley using multiple spectroscopic techniques, *Atmos. Chem. Phys.*, **2014**, 14, 12109-12132, doi:10.5194/acp-14-12109-2014.
- (22) Larssen, S., R. Sluyter, C. Helmis, Criteria for EUROAIRNET, the EEA Air Quality Monitoring and Information Network, **1999**.
- (23) Berresheim, H., Elste, T., Plass-Dülmer, C., Eisele, F. L., and Tanner, D. J.: Chemical ionization mass spectrometer for long-term measurements of atmospheric OH and H₂SO₄, *Int. J. Mass Spectrom.*, 202, 91–109, 2000.
- (24) Song, X. H., Hopke, P. K., Fergenson, D. P., and Prather, K. A.: Classification of single particles analyzed by ATOFMS using an artificial neural network, ART-2A, *Anal. Chem.*, 71, 860–865, 1999.
- (25) Onasch, T. B., Trimborn, A., Fortner, E. C., Jayne, J. T., Kok, G. L., Williams, L. R., Davidovits, P., and Worsnop, D. R.: Soot particle aerosol mass spectrometer: development, validation, and initial application, *Aerosol Sci. Tech.*, 46, 804–817, 2012.
- (26) Paatero, P. and Tapper, U.: Positive matrix factorization: A nonnegative factor model with optimal utilization of error estimates of data values, *Environmetrics*, **1994**, 5, 111–126
- (27) Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, **1994**, 5, 111–126
- (28) Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometrics and Intelligent Laboratory Systems*, **1997**, 37, 23–35.
- (29) Dall'Osto, M., Harrison, R. M., Coe, H., Williams, P. I., and Allan, J. D.: Real time chemical characterization of local and regional nitrate aerosols, *Atmos. Chem. Phys.*, **2009**, 9, 3709-3720.
- (30) Gilardoni, S., Massoli, P., Giulianelli, L., Rinaldi, M., Paglione, M., Pollini, F., Lanconelli, C., Poluzzi, V., Carbone, S., Hillamo, R., et al.: Fog scavenging of organic and inorganic aerosol in the Po Valley, *Atmos. Chem. Phys.*, 14, 6967-6981, doi:10.5194/acp-14-6967-2014, 2014.
- (31) Saarikoski, S., Carbone, S., Decesari, S., Giulianelli, L., Angelini, F., Canagaratna, M., Ng, N. L., Trimborn, A., Facchini, M. C., Fuzzi, et al.: Chemical characterization of springtime submicrometer aerosol in Po Valley, Italy, *Atmos. Chem. Phys.*, 12, 8401–8421, doi:10.5194/acp-12-8401-2012, 2012.

- (32) Yienger, J. J., and H. Levy II. 1994. Empirical model of global soil biogenic NO_x emissions. *Journal of Geophysical Research* 100:11 447–11 464.
- (33) Crosier, J. Allan, J.D., Coe, H., Bower, K. N., Formenti P. and Williams P. I., Chemical composition of summertime aerosol in the Po Valley (Italy), northern Adriatic and Black Sea, *Q. J. R. Meteorol. Soc.*, **2007**, 133: (S1) 61–75
- (34) Dall'Osto, M., M. J. Booth, W. Smith, R. Fisher and R. M. Harrison. A study of the size distributions and the chemical characterization of airborne particles in the vicinity of a large integrated steelworks. *Aerosol Science And Technology*, **2008**, 42(12): 981-991.
- (35) Dall'Osto, M., Querol, X., Alastuey, A., Minguillon, M. C., Alier, M., Amato, F., Brines, M., Cusak, M., Grimalt, J. O., Karanasiou, et al.: Presenting SAPUSS: Solving Aerosol Problem by using synergistic strategies at Barcelona, Spain, *Atmos. Chem. Phys.*, **2013**, 12, 18741-18815,
- (36) Crippa, M., El Haddad, I., Slowik, J. G., DeCarlo, P. F., Mohr, C., Heringa, M., Chirico, R., Marchand, N., Sciare, J., Baltensperger, U., et al.: Identification of marine and continental aerosol sources in Paris using high resolution aerosol mass spectrometry, *J. Geophys. Res.*, **2013**, 118, 1950–1963, doi:1910.1002/jgrd.50151.
- (37) Abdullahi X. and Harrison Roy. M. Emissions and indoor concentrations of particulate matter and its specific chemical components from cooking: A review., *Atmospheric Environment*, **2013**, 71, 260-294
- (38) Kaltsonoudis, C., Kostenidou, E., Louvaris, E., Psichoudaki, M., Tsiligiannis, E., Florou, K. , and S.N. Pandis, Characterization of fresh and aged organic aerosol emissions from meat charbroiling. EAC 2015, Milan 6-11 September 2015
- (39) Moffet RC, Desyaterik Y, Hopkins RJ, Tivanski AV, Gilles MK, Wang Y, et al. Characterization of aerosols containing Zn, Pb, and Cl from an industrial region of Mexico City. *Environ Sci Technol* **2008**;42:7091–7. <http://dx.doi.org/10.1021/es7030483>
- (40) Zhang H., Schuchardt F., Li G., Yang Y., Yang Q. Emission of volatile sulfur compounds during composting of municipal solid waste (MSW). *Waste Management*, **2013**, 33, 957–963.
- (41) Rinaldi, M., S. Decesari, E. Finessi, L. Giulianelli, C. Carbone, S. Fuzzi, C. D. O'Dowd, D. Ceburnis, and M. C. Facchini. Primary and secondary organic marine aerosol and oceanic biological activity: Recent results and new perspectives for future studies, *Adv. Meteorol.*, **2010**, 310682, doi:10.1155/2010/310682.
- (42) Kabir and Kim. An investigation on hazardous and odorous pollutant emission during cooking activities. *Journal of Hazardous Materials*, **2013**, 188 (2011) 443–454
- (43) Byrne, D. V., Bredie, W. L. P., Mottram, D. S., and Martens, M. Sensory and Chemical Investigations on the Effect of Oven Cooking on Warmed-Over Flavour Development in Chicken Meat. *Meat Sci.*, **2001**, 61:127–139.
- (44) Wu, T., Wang, X.M., Li, D.J., Yi, Z.G., Emission of volatile organic sulfur compounds (VOSCs) during aerobic decomposition of food wastes. *Atmos. Environ.*, **2010**, 44, 5065–5071.

- (45) Komilis, D.P., Ham, R.K., Park, J.K., Emission of volatile organic compounds during composting of municipal solid wastes. *Water Res.*, **2004**, 38, 1707–1714.
- (46) Trabue, S., K. Scoggin, F. Mitloehner, H. Li, R. Burns, H. Xin, Field sampling method for quantifying volatile sulphur compounds from animal feeding operations, *Atmos. Environ.*, **2008** 42, 3332–3341.
- (47) Song, S.-K., Z.-H. Shon, K.-H. Kim, Photochemical oxidation and dispersion of gaseous sulfur compounds from natural and anthropogenic sources around a coastal location, *Atmos. Environ.*, **2009**, 43, 3015–3023.
- (48) Kim, K.-H., R. Pal, J.-W. Ahn, Y.-H. Kim, Food decay and offensive odorants: A comparative analysis among three types of food, *Waste Management*, **2009**, 29 1265–1273.
- (49) Chan, A. W. H., Chan, M. N., Surratt, J. D., Chhabra, P. S., Loza, C. L., Crounse, J. D., Yee, L. D., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Role of aldehyde chemistry and NO_x concentrations in secondary organic aerosol formation, *Atmos. Chem. Phys.*, **2010**, 10, 7169–7188, doi:10.5194/acp-10-7169-2010.
- (50) Alanis, P., M. Sorenson, M. Beene, C. Krauter, B. Shamp, A. S. Hasson, Measurement of non-enteric emission fluxes of volatile fatty acids from a California dairy by solid phase micro-extraction with gas chromatography/mass spectrometry, *Atmos. Environ.*, **2008**, 42, 6417–6424.
- (51) Howard et al., Direct Measurements of the Ozone Formation Potential from Livestock and Poultry Waste Emissions, *Environ. Sci. Technol.* **2010**, 44, 2292–2298.
- (52) Aimanant, S., and P. J. Ziemann, Chemical Mechanisms of Aging of Aerosol Formed from the Reaction of n-Pentadecane with OH Radicals in the Presence of NO_x, *Aerosol Science and Technology*, **2013**, 47:979–990
- (53) Kim, K.-H., and S.-Y. Park, A comparative analysis of malodor samples between direct (olfactometry) and indirect (instrumental) methods, *Atmos. Environ.*, **2008**, 42 5061–5070.

746 **TABLE LEGEND**

747
748 **Table 1.** Percentages of factors over the same time interval (9-12/07/2009) for the
749 dataset_1 (all field study) and dataset_2 (high, only second part).
750

751
752
753 **FIGURE LEGENDS**

754
755 **Figure 1.** PMF factors of the dataset 1 (ALL field study).
756

757 **Figure 2.** Diurnal profiles of dataset_1 (ALL field study) and dataset_2 (second part,
758 HIGH). Figure 3 (1-2) shows again the COA factors coming from dataset_1
759 (COA) and dataset_2 (COA-MSA-CI and COA-MSA-HOA).
760

761 **Figure 3.** PMF factors of the dataset_2 for the second part of the field study (HIGH).
762
763
764
765
766
767
768
769
770
771

772 **Table 1.** Percentages of factors over the same time interval (9-12/07/2009) for the
773 dataset_1 (all field study) and dataset_2 (high, only second part)
774
775
776
777

Time period	dataset type	OOA-Reg.	NIT-Reg.	SUL-Reg.	NO	COA-MSA-CI (from ALL)	COA-MSA-CI (from HIGH)	COA-MSA-HOA (from HIGH)
Whole study	ALL	30	18	23	12	17		
Only 9-12/07/2009	ALL	38	17	3	17	25		
Only 9-12/07/2009	HIGH	34	14	0	14		21	17

778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813

814
815

(a)

(b)

821
822

(c)

(d)

(e)

Figure 1. PMF factors of the dataset 1 (ALL field study)

Figure 2. Diurnal profiles of dataset_1 (ALL field study) and dataset_2 (second part, HIGH). Figure (1-2) shows again the COA factors coming from dataset_1 (COA) and dataset_2 (COA-MSA-CI and COA-MSA-HOA)

Photox. OOA

(a)

NIT-Reg.

(b)

(c)

(d)

(e)

Figure 3. PMF factors of the dataset_2 for the second part of the field study (HIGH)