Association between peripheral plasma markers and left atrial anatomy in patients with atrial fibrillation

Nedios, Sotirios; Seewöster, Timm; Lip, Gregory H.y.; Husser, Daniela; Hindricks, Gerhard; Bollmann, Andreas; Kornej, Jelena

DOI:

[10.1016/j.ijcard.2015.11.022](https://doi.org/10.1016/j.ijcard.2015.11.022)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Nedios, S, Seewöster, T, Lip, GHY, Husser, D, Hindricks, G, Bollmann, A & Kornej, J 2016, 'Association between peripheral plasma markers and left atrial anatomy in patients with atrial fibrillation', *International Journal of Cardiology*, vol. 203, pp. 621-623. <https://doi.org/10.1016/j.ijcard.2015.11.022>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Eligibility for repository: checked 10/02/16

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Association between peripheral plasma markers and left atrial anatomy in patients with atrial fibrillation

Sotirios Nedios, Timm Seewöster, Gregory Lip, Daniela Husser, Gerhard Hindricks, Andreas Bollmann, Jelena Kornej

PII: S0167-5273(15)30832-9
DOI: doi: [10.1016/j.ijcard.2015.11.022](https://doi.org/10.1016/j.ijcard.2015.11.022)
Reference: IJCA 21554

To appear in: *International Journal of Cardiology*

Received date: 31 October 2015
Accepted date: 4 November 2015

Please cite this article as: Nedios Sotirios, Seewöster Timm, Lip Gregory, Husser Daniela, Hindricks Gerhard, Bollmann Andreas, Kornej Jelena, Association between peripheral plasma markers and left atrial anatomy in patients with atrial fibrillation, *International Journal of Cardiology* (2015), doi: [10.1016/j.ijcard.2015.11.022](https://doi.org/10.1016/j.ijcard.2015.11.022)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Association between peripheral plasma markers and
left atrial anatomy in patients with atrial fibrillation**

Sotirios Nedios,¹ MD, FHRS, Timm Seewöster,¹ MD, Gregory Lip,² MD, Daniela Husser,¹ MD,
Gerhard Hindricks,¹ MD, Andreas Bollmann,¹ MD, PhD, Jelena Kornej,^{1,2} MD

¹ University of Leipzig, Heart Center, Department of Electrophysiology, Leipzig, Germany

² University of Birmingham, Centre for Cardiovascular Sciences, City Hospital, Birmingham, UK

Words (total): 903

Address for correspondence:

Sotirios Nedios, MD
Heart Center, University of Leipzig
Department of Electrophysiology
Strümpellstr. 39, 04289 Leipzig, Germany
Phone: +49 341 865 1431
Fax: +49 341 865 1460
Email: snedios@gmail.com

Conflict of interest: None.

Disclosure: Dr. Kornej was supported by the German Cardiac Society St. Jude Medical Stipend

Keywords: atrial fibrillation; left atrial anatomy; inflammation; hsIL-6

Left atrial (LA) size has been associated with cardiovascular outcomes and the success of different therapy strategies in patients with atrial fibrillation (AF) [1, 2]. The assessment of LA is therefore recommended in the clinical routine in all AF patients [3]. There are several imaging modalities to assess the LA, though echocardiography is widely available and thus the most frequently used. However, the echocardiographic LA diameter (LA-D) does not reliably reflect the true size of LA anatomy, as pathological LA is often enlarged asymmetrically during AF progression [4-6].

Computer tomography (CT) is a modality that has been increasingly used to obtain three-dimensional (3D) images prior to AF catheter ablation. This provided new insights on the LA shape and volume, which have been proven to be better predictors of AF recurrences after LA ablation in comparison to the commonly used anterior-posterior LA diameter [4-6]. Recently we demonstrated that LA dilatation is more pronounced on the coronal plane, as represented by the transversal LA diameter (LA-TV) [7]. Furthermore, LA-TV was associated with AF recurrences and remained stronger predictor for rhythm outcomes compared with the echocardiographic LA diameter. We also demonstrated strong association between pro-inflammatory plasma markers and AF recurrences [8, 9]. However, whether LA size parameters are associated with peripheral plasma markers of inflammation is unknown.

Consequently, the present study (approved by the ethics committee) recruited 51 consecutive patients presenting for their first AF catheter ablation at Heart Center Leipzig. All patients gave informed consent according to institutional guidelines and the Declaration of Helsinki. Echocardiography and cardiac-CT with a multidetector 64-row helical system (Brilliance 64, Philips, Best, Netherlands) were performed (2 ± 1 days) before the procedure. CT data were

reviewed using 3D reconstruction (EnSite Verismo, SJM, MN) and LA volume (LAV) was determined after exclusion of the atrial appendage (LAA) and the pulmonary veins (PV). LA was then centered on all three cutting planes and the superior–inferior (SI), transversal (TV) and anterior–posterior (AP) diameters were measured. Measurements were performed offline by an experienced observer and were repeated 4 weeks later by the same investigator and a second blinded reviewer. High sensitive interleukin 6 (hsIL-6) was analyzed from pre-procedural blood samples using a commercially available assay.

Catheter ablation was performed as previously described [10], with circumferential ablation of the ipsilateral pulmonary veins, verified with a multipolar circular catheter. In patients with persistent AF, additional linear lesions were added at the mitral isthmus and the posterior LA wall to create a “box” lesion. Follow-up was performed with repeated 7-day-Holter ECG recordings at 6, 12, 24 and 36 months. Recurrence was defined as any documented atrial tachycardia or fibrillation episodes of ≥ 30 s (after a 3 month blanking period).

Statistical analyses were performed with SPSS 17 (SPSS Inc., Chicago, USA). Parameters with a p-value < 0.1 in the univariable analysis (UV), were introduced in multivariable analyses (MV) in order to identify with hsIL-6 levels independently associated parameters. A two-tailed p value < 0.05 was considered significant.

The clinical characteristics of the study population are presented in **Table 1**. The intra- and inter-observer correlation coefficients were ≥ 0.88 . We found a significant correlation between peripheral hsIL-6 and LA-TV ($r^2=0.34$, $p=0.017$) but not with LAV ($p=N.S.$, **Figure 1**). On univariable analysis, advanced age, higher BMI and lower eGFR as well as as well as with left atrial dimensions – e.g. LA-AP, LA-TV and LAD (but not LAV) – were significantly associated

with hsIL-6 levels. On multivariable analysis, the levels of hsIL-6 remained associated with age ($r^2=0.28$, $p=0.029$), BMI ($r^2=0.33$, $p=0.011$), renal dysfunction ($eGFR<60$ ml/min/1.73m², $r^2=0.29$, $p=0.019$), and LA-TV ($r^2=0.38$, $p=0.012$) but not with LAD or LA-AP.

Both electrical-anatomical remodeling and inflammation are known factors in AF pathogenesis. The present study demonstrates a significant association between different left atrial measurements and peripheral pro-inflammatory markers and adds to our understanding about the remodeling processes in patients with AF. The role of inflammation in AF pathogenesis is a well-known factor. Pro-inflammatory reactions may modulate the treatment success and, as previously demonstrated by our group, positively correlate with AF recurrences after radiofrequency catheter ablation [8, 9]. Recently, we also showed that anatomical LA changes carry a predictive value that is better represented with LA-TV or LAV than with the commonly used LAD [7]. The present data extend these studies and reveal the strong correlation between the plasma marker hsIL-6 and the anatomical atrial remodeling, as represented by the transversal (LA-TV) and not just the global (LAV) dilatation. These findings link pro-inflammatory changes with the left atrial remodeling and emphasize the close relationship between these important AF co-variables.

The small size of this study cohort and the limitations (loss) of follow-up prohibited further analysis of the impact of LA remodeling and hsIL-6 on ablation outcomes. The use of CT for the measurement of LA-TV prohibits its use in repeated routine follow-up studies, but since newer techniques, such as magnetic resonance imaging or 3D echocardiography, are becoming more widely available, soon this information could be assessed with no radiation.

These findings should be seen as hypothesis generating and point out the potential importance and scientific prominence of inflammation state in patients with AF. Certainly, larger clinical studies are needed to examine the practical implications of hsIL-6 or LA measurements for better patient selection and treatment strategy in early or advanced AF stages. Since a pro-inflammatory state is associated with anatomical LA remodeling, it remains to be examined whether AF patients might profit from anti-inflammatory treatment during or after invasive treatment.

References

- [1] Berruezo A, Tamborero D, Mont L, et al. Pre-procedural predictors of atrial fibrillation recurrence after circumferential pulmonary vein ablation. *Eur Heart J*. 2007;28:836-41.
- [2] Abecasis J, Dourado R, Ferreira A, et al. Left atrial volume calculated by multi-detector computed tomography may predict successful pulmonary vein isolation in catheter ablation of atrial fibrillation. *Europace*. 2009;11:1289-94.
- [3] Camm AJ, Lip GY, De Caterina R, et al. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation. Developed with the special contribution of the European Heart Rhythm Association. *Eur Heart J*. 2012;33:2719-47.
- [4] Nedios S, Tang M, Roser M, et al. Characteristic changes of volume and three-dimensional structure of the left atrium in different forms of atrial fibrillation: predictive value after ablative treatment. *J Interv Card Electrophysiol*. 2011;32:87-94.
- [5] Nedios S, Koutalas E, Kosiuk J, et al. Impact of asymmetrical dilatation of the left atrium on the long-term success after catheter ablation of atrial fibrillation. *Int J Cardiol*. 2015;184:315-7.
- [6] Bisbal F, Guiu E, Calvo N, et al. Left atrial sphericity: a new method to assess atrial remodeling. Impact on the outcome of atrial fibrillation ablation. *J Cardiovasc Electrophysiol*. 2013;24:752-9.
- [7] Nedios S, Kosiuk J, Koutalas E, et al. Comparison of left atrial dimensions in CT and echocardiography as predictors of long-term success after catheter ablation of atrial fibrillation. *J Interv Card Electrophysiol*. 2015;43:237-44.

- [8] Kornej J, Dinov B, Blann AD, et al. Effects of radiofrequency catheter ablation of atrial fibrillation on soluble P-selectin, von Willebrand factor and IL-6 in the peripheral and cardiac circulation. *PLoS One*. 2014;9:e111760.
- [9] Kornej J, Reinhardt C, Kosiuk J, et al. Response of high-sensitive C-reactive protein to catheter ablation of atrial fibrillation and its relation with rhythm outcome. *PLoS One*. 2012;7:e44165.
- [10] Esato M, Hindricks G, Sommer P, et al. Color-coded three-dimensional entrainment mapping for analysis and treatment of atrial macroreentrant tachycardia. *Heart Rhythm*. 2009;6:349-58.

Table 1. Baseline characteristics of the study population

Variables	Study population
	n=51
Age, years	62 ± 10
Females	37 %
Persistent AF	59 %
BMI	30 ± 4.6
eGFR, ml/min/1.73 m ²	98 ± 30
Hypertension	88 %
Diabetes mellitus	18 %
CHA ₂ DS ₂ -VASs score	2 (2 – 3)
LA diameter, mm (Echo)	44 ± 6.5
EF, %	57 ± 11
LA-TV, mm (CT)	77 ± 11
LA-SI, mm (CT)	64 ± 8
LA-AP, mm (CT)	51 ± 12
LA-V, ml (CT)	141 ± 50
hsIL-6, ng/ml	1.6 (0.9 – 2.9)

Abbreviations: AF – atrial fibrillation, BMI - body mass index, eGFR - estimated glomerular filtration rate, TE - thromboembolic events, IQR - interquartile range, EF - ejection fraction, LA - left atrial, hsIL-6 - high sensitive interleukin 6 (hsIL-6)

Figure 1. Left atrial (LA) parameters and high sensitive interleukin 6 (hsIL-6) of 2 patients with the same LA volume. Anatomical changes on the coronal plane (bigger transversal diameter, LA-TV) are associated with higher hsIL-6 values (B, D).