Intensive Exercise Does Not Preferentially Mobilize Skin-Homing T Cells and NK Cells

Turner, James E; Wadley, Alex J; Aldred, Sarah; Fisher, James P; Bosch, Jos A; Campbell, John P

DOI:

[10.1249/MSS.0000000000000914](https://doi.org/10.1249/MSS.0000000000000914)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Turner, JE, Wadley, AJ, Aldred, S, Fisher, JP, Bosch, JA & Campbell, JP 2016, 'Intensive Exercise Does Not Preferentially Mobilize Skin-Homing T Cells and NK Cells', *Medicine and Science in Sports and Exercise*.
<https://doi.org/10.1249/MSS.0000000000000914>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is a non-final version of an article published in final form in *Medicine and Science in Sports and Exercise*.

Checked March 2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Medicine & Science IN Sports & Exercise

The Official Journal of the American College of Sports Medicine

www.acsm-msse.org

... Published ahead of Print

Intensive Exercise Does Not Preferentially Mobilize Skin-Homing T Cells and NK Cells

James E. Turner¹, Alex J. Wadley^{2,3}, Sarah Aldred³, James P. Fisher³,
Jos A. Bosch^{4,5}, and John P. Campbell⁶

¹Department for Health, University of Bath, Bath, United Kingdom; ²Institute of Science and the Environment, University of Worcester, Worcester, United Kingdom; ³School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, United Kingdom; ⁴Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands; ⁵Mannheim Institute of Public Health, Social and Preventive Medicine (MIPH), University of Heidelberg, Heidelberg, Germany; ⁶Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, United Kingdom

Accepted for Publication: 3 January 2016

Medicine & Science in Sports & Exercise, **Published ahead of Print** contains articles in unedited manuscript form that have been peer reviewed and accepted for publication. This manuscript will undergo copyediting, page composition, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered that could affect the content.

Intensive Exercise Does Not Preferentially Mobilize Skin-Homing T Cells and NK Cells

James E. Turner¹, Alex J. Wadley^{2,3}, Sarah Aldred³, James P. Fisher³,

Jos A. Bosch^{4,5}, and John P. Campbell⁶

¹Department for Health, University of Bath, Bath, United Kingdom; ²Institute of Science and the Environment, University of Worcester, Worcester, United Kingdom; ³School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, United Kingdom; ⁴Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands; ⁵Mannheim Institute of Public Health, Social and Preventive Medicine (MIPH), University of Heidelberg, Heidelberg, Germany; ⁶Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, United Kingdom

Corresponding author details:

John P. Campbell

Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, UK. B15
2TT.

Telephone: +44 7530595912

Email: j.p.campbell@bham.ac.uk

CONFLICTS OF INTEREST: None declared. The results of this present study do not constitute endorsement by ACSM. The flow cytometric analyses for this study were funded by the University of Birmingham Clinical Immunology Service.

Running head: CLA+ lymphocytes and exercise

ACCEPTED

Abstract

Purpose: This study investigated whether natural killer (NK) cells and CD8⁺ T cells expressing cutaneous lymphocyte antigen (CLA) – a homing molecule for endothelial cell leukocyte adhesion molecule 1 (ELAM-1), which enables transmigration to the skin – are selectively mobilised in response to acute exercise. **Methods:** Nine healthy males (mean $\pm$ SD age: 22.1 $\pm$ 3.4 years) completed two exercise sessions: high-intensity continuous cycling ('continuous exercise' at 80% $\dot{V}O_{2\text{ MAX}}$ for 20 min) and low-volume high-intensity interval exercise ('HIIE' at 90% $\dot{V}O_{2\text{ MAX}}$ 10 $\times$ 1 min repetitions with 1 min recovery intervals). Blood was collected before, immediately- and 30 min post-exercise for cryo-preservation of peripheral blood mononuclear cells. CLA⁺ and CLA[−] cells were quantified within NK subpopulations (CD56^{bright} 'regulatory' and CD56^{dim} 'cytotoxic' cells) as well as the following CD8⁺ T cell subpopulations: naive ('NA'; CD45RA+CCR7+), central memory ('CM'; CD45RA−CCR7+), effector-memory ('EM'; CD45RA−CCR7−) and CD45RA-expressing effector-memory cells ('EMRA'; CD45RA+CCR7−). **Results:** CLA⁺ NK cells and CD8⁺ memory T cells increased in response to both exercise bouts, but, overall, their numerical contribution to the exercise lymphocytosis was inferior to CLA[−] cells, which increased to a much greater extent during exercise. Tellingly, the most exercise-responsive cells – effector memory CD8⁺ cells and CD56^{dim} cells – were CLA[−]. **Conclusions:** A small subset of CLA⁺ lymphocytes are mobilised into blood during acute intensive exercise, but CLA⁺ cells are not major contributors to exercise lymphocytosis, thus providing preliminary evidence that the skin is not a major origin, or homing-destination, of exercise-sensitive lymphocytes.

Key words: T cells, inflammation, skin, homing

INTRODUCTION

Memory CD8⁺ T cells and natural killer (NK) cells are the largest contributors to the acute and transient lymphocytosis that occurs in peripheral blood during intensive exercise (6, 20, 36). Largely driven by their relative high density cell surface expression of β 2-adrenergic receptors that induce detachment from the vascular endothelium upon stimulation by catecholamines, the mobilisation of NK cell and CD8⁺ T cell subsets is considered to be an evolved mechanism that facilitates effector cell recruitment to sites of potential or ongoing injury (10). Built on findings from animal models and using an analogy of immune cells as ‘soldiers’, it is proposed that naïve and central memory CD8⁺ T cells – which are slightly increased during exercise – traffic from the boulevards (i.e., ‘bloodstream’) back to the ‘barracks’ (i.e., the spleen, lymph nodes). On the other hand, tissue-specific memory CD8⁺ T cells – which mobilise to a larger extent during exercise – home to specific peripheral tissues (e.g., lungs, gut) to conduct immune-surveillance against potential pathogenic challenge. Finally, it is hypothesised that effector CD8⁺ T cells – which are mobilised to the greatest extent during exercise – are redeployed to ‘battlefield’ sites of wound healing (e.g. in the skin) following acute exercise (9, 10). However, whether these highly cytotoxic effector NK and CD8⁺ T cells, which are selectively increased in an acute and transient manner by acute exercise, have the propensity to migrate to peripheral cutaneous sites remains unknown.

In support of this model, it has been demonstrated in murine models that exercise redeploys large numbers of T cells to the Peyer’s patches, lungs and bone marrow, reinforcing the idea that T cells are mobilised to sites of potential antigen encounter (i.e., lungs, gut), as well as to serve other important functions such as the provision of additional stimuli for

haematopoiesis in the bone marrow (20). However, the aforementioned study was unable to measure homing to cutaneous sites or areas of ongoing inflammation. It has been shown that T cells and NK cells do show increased infiltration into sites of experimental inflammation (i.e., subcutaneous implantation of a surgical sponge treated with the lymphocyte-specific chemokine lymphotactin) in mice 24 to 48 hours after acute psychological stress (40). However, it remains unclear whether this increased cell infiltration is driven by stress-induced effector cell redistribution, or other mechanisms occurring in the days after the acute stressor.

An approach that is commonly used to investigate lymphocyte homing propensity in humans is the assessment of cell surface adhesion molecule expression on cells; this approach can be used to reveal the probable trafficking patterns of cells mobilised into the bloodstream during exercise. For example, studies have shown that there is a selective influx of CD8⁺ memory cells into the bloodstream that exhibit lower levels of lymphoid homing markers such as CD62L and CCR7 (6), thus providing evidence that exercise mobilises CD8⁺ T cells with a homing capacity for peripheral tissues. Further research has shown that these cells mobilised by exercise express adhesion molecules such as CD11a (15, 22), CD11b (17, 18), VLA-4 (very late antigen-4) and LPAM-1 (lymphocyte Peyer's patch adhesion molecule-1) (15), which enable migration to peripheral sites including the bone marrow (23), Peyer's patches (43) and lungs (38). However, the aforementioned adhesion molecules cannot be used exclusively for identifying skin-homing potential and it remains uncertain whether exercise-responsive CD8⁺ T cells and NK cells have a skin-homing phenotype.

Cutaneous lymphocyte antigen (CLA) expression on lymphocytes can be used to determine whether lymphocytes preferentially mobilised during intensive exercise exhibit a homing phenotype for cutaneous sites. CLA is a specialised form of P-selectin glycoprotein ligand-1 (PSGL-1; CD162), a surface glycoprotein expressed constitutively on all human peripheral-blood T cells. After post-translational modification, PSGL-1 bears a Sialyl-Lewis^X (sLe^X) moiety, termed CLA, which avidly binds CD62E (E-selectin), an adhesion molecule which initiates the transmigration cascade to the skin, and which is also upregulated during cutaneous inflammation (4, 27, 28, 32). Thus, most T cells in both normal and diseased cutaneous sites are CLA⁺ (7, 29, 30). In support, leukocyte infiltration to inflammatory sites can be largely inhibited by a CLA modifier (11). Taken together, analyses of CLA⁺ cell mobilisation can be used as proxy marker to reveal whether cells mobilised by exercise have the phenotypic capacity to migrate to ‘battlefield’ sites in the skin (10).

The primary objective of this study was to investigate whether CD8⁺ T cells preferentially mobilised during intensive exercise stress exhibit a homing phenotype for cutaneous sites. To fulfil this objective, we investigated the number of CLA-positive cells among CD8⁺ T cell subsets, to establish the contribution of CLA-positive cells to the stepwise CD8⁺ lymphocytosis pattern previously observed in response to exercise (6). We also extended the analyses of CLA⁺ and CLA[−] cells to NK cell populations – the largest responders to exercise stress (6). The second objective of this study was to compare the magnitude of CD8⁺ T cell and NK cell mobilisation in response to continuous high-intensity exercise (continuous exercise) and high-intensity intermittent exercise (HIIE). HIIE is often referred to as High Intensity Interval Training (HIIT) when repeated frequently over several weeks or months. This form of exercise

has received considerable attention as a short-duration and low-volume means of achieving similar health benefits to continuous exercise. Given the increasing health benefits that have been observed with HIIE, and although we have established in a prior report that CD8⁺ T cell and NK cell subset mobilisation is intensity dependent (6), the effects of HIIE on changes to these cell subsets remains unknown.

METHODS

Participants

Ten healthy males were recruited to take part in this study as previously described (41, 42). Peripheral blood mononuclear cells (PBMCs) were available from nine of the ten participants, and were isolated from blood samples collected before, immediately after, and 30 min after two different forms of exercise, described below. All nine participants (age: 22.1 ± 3.4 years; height: 180.5 ± 6.1 cm; weight: 78.1 ± 11.0 kg; body mass index: 24.0 ± 3.1 kg.m²; $\dot{V}O_{2\text{ MAX}}$: 43.8 ± 4.1 ml.kg.min⁻¹) included in this study were non-smokers, and refrained from taking vitamin supplements and anti-inflammatory medication for fourteen days, and did not exercise and consume alcohol or caffeine for two days prior to experimental trials. All participants provided written informed consent and the study was approved by the Science, Technology, Engineering and Mathematics Ethical Review Committee at University of Birmingham (reference: ERN_12-0830).

Preliminary measurements

Height and weight were assessed using standard methods and cardiorespiratory fitness ($\dot{V}O_{2\text{ MAX}}$) was measured during an incremental exercise test on an electromagnetically braked

cycle ergometer (*Lode Excalibur Sport*, Groningen, Netherlands). Following a three-minute warm up at 30 watts, workload was increased by 30 watts every minute, until volitional exhaustion. A facemask was fitted throughout preliminary exercise tests in each main trial, and expired air measured breath-by-breath averaged every 20 seconds for oxygen uptake and carbon dioxide production (*Oxycon Pro*, Jaeger, Wuerzburg, Germany). Heart rate (HR) was monitored every minute using a Polar Vantage heart rate monitor (Polar Vantage, Kempele, Finland); HR_{peak} represents the maximum heart rate achieved during each trial. The following criteria were used to indicate that $\dot{V}O_{2\ MAX}$ had been reached: a fall in cadence below 60 rpm, a respiratory exchange ratio ($\dot{V}CO_2 / \dot{V}O_2$) >1.10-1.15, a plateau in oxygen consumption and a heart rate >220 beats min^{-1} minus age.

Experimental trials

Experimental trials were undertaken at least seven days after preliminary measurements, in the morning, and following an overnight fast. Each trial was separated by at least three days in a randomised design. Prior to each trial, but after the baseline blood sample, participants undertook a warm up (5 minutes) at a workload eliciting 40% $\dot{V}O_{2\ MAX}$. Exercise trials were either vigorous steady state cycling at 80% $\dot{V}O_{2\ MAX}$ for 20 minutes ('continuous exercise') or high intensity interval exercise ('HIIE') trial. HIIE consisted of ten 1 minute cycling phases at a workload to elicit 90% $\dot{V}O_{2\ MAX}$, with 1 minute of low intensity cycling at 40% $\dot{V}O_{2\ MAX}$ between each phase. Workload was expressed as watts, and relative to body mass (i.e., watts/kg). Values presented are the average workload over the entire exercise protocol (i.e., in the vigorous trial: the duration of cycling at 80% $\dot{V}O_{2\ MAX}$; and in the HIIE protocol: each 1-minute sprint phase of cycling at 90% $\dot{V}O_{2\ MAX}$). Values were obtained directly from the electromagnetically braked

cycle ergometer (Lode Excalibur Sport, Groningen, Netherlands). Energy expenditure (Kcal) was estimated by indirect calorimetry from calculations of fat and carbohydrate oxidation (g/min) and data were averaged over the 1-minute phases of the protocols and summed to provide total energy expenditure from the entire duration of each trial, expressed relative to body mass (Kcal/kg). Ratings of perceived exertion (RPE) were recorded using the Borg Scale every 1 minute during continuous exercise, and after each 90% $\dot{V}O_{2\text{ MAX}}$ interval during HIIE; the highest RPE score was selected as the final RPE result for each exercise trial (RPE_{peak}).

Blood sampling

An intravenous cannula (*Becton & Dickson, Oxford, UK*) was inserted into an antecubital vein and blood samples were drawn into potassium ethylene diaminetetraacetic acid (EDTA) vacutainer tubes (*Becton & Dickson, Oxford, UK*). The cannula was kept patent with saline (0.9% NaCl). The leukocyte differential was assessed using an automated haematology analyser (*Coulter Analyser, Beckman-Coulter, High Wycombe, UK*).

Blood Cell Isolation

Approximately 15 ml of blood from each time point (pre-exercise, post-exercise and 30 min post-exercise) was diluted 1:1 with Roswell Park Memorial Institute Media (RPMI), and then layered on top of Ficoll paque PLUS (GE Healthcare) (2 blood : 1 Ficoll), before centrifuging at $500 \times g$ for 30 minutes at 21°C. PBMCs were aspirated and washed three times in RPMI by centrifuging at $400 \times g$ for 5 minutes. The cell pellet was re-suspended in 1-ml of freezing mixture (70% RPMI, 20% fetal calf serum (FCS) and 10% dimethyl sulfoxide (DMSO))

and frozen at -1°C / min using a freezing container (Nalgene “Mr Frosty” Thermoscientific). Cells were stored at -80°C and analysed within six months.

Flow cytometry

Samples were thawed rapidly at 37°C and washed twice in PBS containing 2% FCS and 2mM EDTA by centrifuging at $400 \times g$ for 5 minutes. Cells were counted using a haemocytometer and approximately 300,000 PBMCs were added to tubes for incubation with fluorescently conjugated antibodies to identify specific lymphocyte populations using eight-colour flow cytometry (FACS-CANTO, Becton–Dickinson, San Jose, USA). The following monoclonal antibodies (mAbs) were used: anti-CD45RA-FITC clone # HI100, anti-CD197 (CCR7)-PE clone # 150503, anti-CD56-PE-Cy7 clone # B159, anti-CD8-APC clone # RPA-T8, anti-CD3-APC-Cy7 clone # SK7 (BD Pharmingen, San Diego, USA), anti-CLA-V450 clone # HECA-452 (BioLegend, San Diego, USA) and anti-CD16-V500 clone # 3G8 (BD Horizon, San Diego, USA). In addition, 7-aminoactinomycin D (7-AAD; PerCP channel; BD Pharmingen, San Diego, USA) was used to exclude necrotic and apoptotic cells. For validation purposes, fluorescence-minus-one (FMO) tubes were used on separate samples from three healthy donors to establish negative and positive gating strategies for CLA expression.

Flow cytometry data were analysed using FlowJo version 7.6.5 (FlowJo LLC, Oregon, USA). Briefly, lymphocytes were gated on the forward *versus* sideways scatter, and 7AAD– cells were divided into CD3+ or CD3– cells. Cytotoxic T cells were identified as being CD3+CD8+ and further differentiated into naïve (‘NA’; CCR7+CD45RA+), central memory (‘CM’; CCR7+CD45RA–), effector memory (‘EM’; CCR7–CD45RA–) or CD45RA+ effector

memory cells ('EMRA'; CCR7-CD45RA+). CD3- cells were further differentiated into CD56^{dim} 'cytotoxic' NK cells or CD56^{high} 'regulatory' NK cells, as proposed by Cooper and colleagues (2001), using CD16 and CD56 dotplot gating (8). Within each subpopulation identified above, cells were subdivided based on their expression of CLA (CLA+ or CLA-) in SSC (side scatter) versus V450-CLA mode. Absolute cell counts for each subpopulation were computed by multiplying the percentage of cells within the CD3+ or CD3- gates by the total lymphocyte count (obtained via the Coulter principle). This value and subsequent absolute subpopulation counts were multiplied by percentage values for gated daughter subpopulations.

Statistical analyses

All results are presented as mean $\pm$ standard deviation unless otherwise stated. Statistical calculations were performed on IBM SPSS for Windows Version 21. Kolmogorov-Smirnov tests confirmed that all data were normally distributed at all time points and for all variables. Repeated Measures Analysis of Variance (ANOVA) tests were used to contrast changes over time (baseline, exercise and 30 min post-exercise), and between the two exercise conditions (HIIE and continuous exercise) for each cell type. Within-trial main effects of time for each cell type were calculated in separate individual ANOVAs. Post-hoc pairwise comparisons were made with Bonferroni adjustments for multiple comparisons. Between trial differences at a given time point were assessed by paired *t*-tests. Statistical significance was accepted at the $p < .05$ level.

RESULTS

All participants completed both the continuous exercise and HIIE tasks. Total energy expenditure was significantly higher after continuous exercise compared to HIIE (HIIE: $2.69 \pm$

0.31 Kcal/kg; continuous exercise: 3.54 ± 0.48 Kcal/kg; $F_{(1,17)} = 20.0$, $p < .05$). Similarly, RPE_{peak} was significantly higher after continuous exercise compared to HIIE (HIIE: 16 ± 2 (Borg scale); continuous exercise: 18 ± 1 (Borg scale); $F_{(1,17)} = 4.6$, $p < .05$). Workload was higher after HIIE compared to continuous exercise (HIIE: 2.74 ± 0.54 Watts/kg; continuous exercise: 2.18 ± 0.33 Watts/kg; $F_{(1,17)} = 7.1$, $p < .05$). No differences were found between trials for HR_{peak} (HIIE: 178 ± 14 beats/min; continuous exercise: 187 ± 7 beats/min; $p > .05$).

Effects of HIIE and continuous exercise on the number of total leukocytes and leukocyte subpopulations

Total leukocyte counts increased ($p < .05$) from baseline after both the HIIE (5.31 ± 1.15 [$\times 10^9/L$] to 8.92 ± 2.77) and continuous exercise (5.57 ± 0.94 to 10.73 ± 2.14) conditions. Similarly, total lymphocyte counts also increased ($p < .05$) from baseline after both the HIIE (1.83 ± 0.47 to 3.64 ± 1.46) and continuous exercise (1.91 ± 0.41 to 4.83 ± 1.33) conditions, with a larger mean change observed during continuous exercise, compared to HIIE, but these differences were not significant ($p > .05$). 30 min after each exercise condition, lymphocyte numbers between trials were similar (HIIE: 1.67 ± 0.37 ; continuous exercise: 1.84 ± 0.35) and had returned to pre-exercise levels ($p > .05$, compared to baseline).

Effects of HIIE and continuous exercise on the number of total CD3+ T cells and CD3+CD8+ T cells

Table 1 displays total CD3+ T cells and CD3+CD8+ T cell numbers during both exercise tasks. Total CD3+ T cells increased significantly during both conditions ($p < .01$), with larger mean increases during continuous exercise, compared to HIIE, but these between trial

differences were not significant ($p > .05$). 30 min after each condition, CD3+ T cell numbers between trials were similar and had returned to pre-exercise levels. CD3+CD8+ T cells increased significantly during both conditions ($p < .05$), with larger increases during continuous exercise compared to HIIE; again, these differences between trials were not significant ($p > .05$). 30 min after each condition, CD3+CD8+ numbers between trials were similar and had returned to pre-exercise levels. Replicating previous findings (6, 39), CD3+CD8+ T cell subsets were mobilised in a stepwise manner, with CD3+CD8+ EMRA T cells mobilising more than EM, CM and NA cells. Within these subsets, there were no significant differences in the magnitude of mobilisation between continuous exercise or HIIE trials.

Effects of HIIE and continuous exercise on the numbers of CLA+ and CLA- CD3+ T cells and CD3+CD8+ T cells

Approximately 14% of total CD3+ T cells were CLA+ at baseline, and despite an increase in the number of CLA+ CD3+ T cells during both exercise conditions ($p < .05$), the proportion of CLA+ CD3+ T cells decreased slightly during exercise due to a larger influx of CLA- CD3+ T cells ($p = .05$) (see Table 1). At 30 minutes post-exercise the number and proportion of CLA+ CD3+ T cells returned to pre-exercise values.

Approximately 9% of CD3+CD8+ T cells were CLA+ at baseline, and, in a similar manner to total CD3+ T cells, the number of CLA+ CD3+CD8+ T cells increased during both exercise conditions ($p < .05$), before returning to baseline levels at 30 minutes post-exercise (see Table 1). However, there was no change ($p > .05$) in the proportion of CLA+ CD3+CD8+ T cells

during exercise. No differences were observed in the magnitude of mobilisation between CLA⁺ and CLA⁻ cells in any of these broad T cell subsets between HIIE and continuous exercise.

Effects of HIIE and continuous exercise on the numbers of CLA⁺ and CLA⁻ CD3⁺CD8⁺ T cell subsets

In accordance with prior literature (7, 31), very few of the CD3⁺CD8⁺ NA cell population expressed CLA (median = 1.35%) and the few CLA⁺ naïve cells did not change during exercise (data not shown). The largest proportion of CLA⁺ CD3⁺CD8⁺ T cells were of the CM (~20%) and EM (~23%) phenotypes, with only a small fraction of EMRA cells expressing CLA (<7%). As illustrated in Figure 1, the percentage change in the number of CLA⁺ cells increases in a stepwise mobilisation pattern (i.e., EMRA > EM > CM) in response to both exercise conditions, and this pattern is also evident in CLA⁻ cells. However, as illustrated in Figure 2, the proportion of CLA⁺ cells per memory CD3⁺CD8⁺ T cell subset actually shows a marginal and significant ($p < 0.05$) decline during exercise, before returning to baseline levels at 30 min post-exercise. Thus, CLA⁺ cells do not appear to be major contributors to the large influx of effector cells observed in the peripheral bloodstream during exercise; this is illustrated by a greater mobilisation of CLA⁻, compared to CLA⁺, EMRA CD3⁺CD8⁺ T cells in Figure 1.

Effects of HIIE and continuous exercise on the numbers of NK cells, and CLA⁺ and CLA⁻ cells within the NK cell subsets

Table 2 displays changes in the numbers of CD3⁻CD56⁺ NK cells in response to continuous exercise and HIIE. As expected, NK cells were extremely sensitive to exercise stress, exhibiting significant differences between the two exercise modalities ($p < .05$), with HIIE

resulting in a ~400% increase and continuous exercise resulting in a ~600% change during exercise. With regards to the major CD56⁺ NK cell subsets, CD56^{dim} ‘cytotoxic’ cells were the most sensitive to exercise stress, with HIIE resulting in a 550% increase and continuous exercise resulting in a 725% increase ($p < .05$). As expected, CD56^{bright} ‘regulatory’ NK cells were less sensitive to exercise stress and no differences were found between exercise conditions ($p > .05$); HIIE resulted in a 100% increase and continuous exercise resulted in a 200% increase.

At baseline, approximately ~22% of CD3⁺CD56⁺ cells were CLA⁺, with a higher proportion of CD56^{bright} ‘regulatory’ NK cells positive for CLA (~67%) than CD56^{dim} ‘cytotoxic’ NK cells (~17.5%). During exercise, the total number of CLA⁺ NK cells increased (HIIE: ~230%; continuous exercise: ~350%), and, as expected, this was driven by a larger increase in CLA⁺ CD56^{dim} cells compared to the CLA⁺ CD56^{bright} cells. As a consequence of a greater influx of CLA⁺ NK cells (HIIE: ~500%; continuous exercise: ~650% compared to baseline), primarily comprised of CLA⁺CD56^{dim} cells during exercise (Figure 3), the proportion of CLA⁺ cells in the total NK cell pool was actually reduced during exercise (Figure 4). Thus, CLA⁺ NK cells do not appear to be major contributors to the large influx of effector cells observed in the peripheral bloodstream during exercise.

DISCUSSION

This study assessed the mobilisation propensity of CLA⁺ CD8⁺ T cells into the peripheral bloodstream in response to intensive exercise. A preferential mobilisation of this cell phenotype would fulfil a component of the stress redistribution theory (10), which hypothesises that effector memory cells preferentially mobilised by exercise have a phenotype that enables

post-exercise migration to cutaneous sites of wound healing or active inflammation (e.g., injured skin). We examined the number of CLA⁺ and CLA⁻ cells because CLA avidly binds CD62E; CD62E is thought to be the primary initiator of routine CD8⁺ T cell transmigration to the skin, and is also central to T cell tethering to cytokine-activated endothelium at inflamed cutaneous sites (14). We found that exercise resulted in an increase in the number of memory CLA⁺ CD3⁺CD8⁺ T cells. However, the numerical contribution of CLA⁺ memory cells to exercise-induced lymphocytosis was inferior to CLA⁻ cells, and, as a consequence, the proportion of CLA⁺ cells amongst the total memory CD8⁺ T cell pool showed a decline during exercise. Furthermore, the most exercise-sensitive subset (EMRA CD8⁺ T cells) were largely CLA⁻, suggesting that the most exercise-responsive T cell subset mobilised by exercise do not have a phenotype that would enable rapid transmigration to sites of active cutaneous inflammation.

Until now, the effect of exercise on the mobilisation of CLA⁺ cells has not yet been investigated. A study investigating the effects of acute psychological stress on circulating CLA⁺ CD8⁺ cells found a marginal decline in the number of CLA⁺ CD8⁺ cells during the stressor (2). These aforementioned results differed to the findings of our study, as we observed an approximate doubling in the number of CLA⁺ CD3⁺CD8⁺ cells during exercise. It was proposed by Atanackovic et al. (2006) that CLA⁺ CD8⁺ T cells had already initiated transmigration to sites of inflammation during acute stress. However, evidence exists to suggest that elevated levels of epinephrine selectively decreases adhesion of CD8⁺ T cells (12) and NK cells (3) to endothelial cells, and may thus be a contributing mechanism for the maintenance of CLA⁺ cells in the peripheral bloodstream, as observed in our exercise trials. We extended our CLA⁺ analyses to four distinct CD8⁺ T cell subsets conventionally identified using the cell-surface

markers CCR7 and CD45RA (17). In agreement with prior literature showing that the majority of CLA⁺ CD3⁺ cells are CD45RO⁺ (7), we found that few naïve (CCR7⁺CD45RA⁺) CD8⁺ T cells expressed CLA (31). This did not change in response to exercise. In further agreement with Clark et al. (2006), we found a majority of CLA⁺ cells were of the central memory (CCR7⁺CD45RA⁻) phenotype and fewer were effector memory (CCR7⁻CD45RA⁻) CD8⁺ T cells; these CLA⁺ effector memory cells showed the largest increase during exercise, though the numerical increase was small.

On one hand, the findings of this study are supportive of the stress redistribution model in that exercise evoked an increase in the total number of peripheral blood CLA⁺ CD8⁺ T cells, which are phenotypically consistent with skin-homing T cells. On the other hand, the absolute number of CLA⁺ cells mobilised was small, and the relative magnitude of mobilisation was much less than CLA⁻ CD8⁺ cells. Furthermore, few EMRA CD8⁺ T cells – conventionally the most exercise-sensitive CD8⁺ subset – expressed CLA, and did not increase in number as much as CLA⁻ EMRA CD8⁺ T cells. Thus, the CD8⁺ T cells with the greatest propensity for exercise-induced mobilisation and extravasation are CLA⁻ and do not exhibit a phenotype characteristic for rapid homing to cutaneous sites or sites of endothelial inflammation. Given that the large majority of CLA⁺ cells reside in the skin (7), it is possible to conclude that the skin is not a major contributor/source of cells to exercise-induced lymphocytosis. The findings of this study are not surprising: CLA⁺ cells are not as susceptible to immunosenescence (25), and are thus strikingly different to the most exercise-sensitive cells, which have shorter telomeres, high expression of CD57 (35, 36); and exhibit exaggerated exercise-induced mobilisation in participants seropositive for cytomegalovirus (39). In addition, it has been demonstrated in

rodents that exercise redeploys large numbers of T cells to the lungs (20). Lung-derived T cells are all CLA⁻, but positive for other homing molecules including CCR5 and CXCR3 (5). Similarly, exercise also redeploys T cells to the Peyer's patches and bone marrow (19, 20) and this is likely governed by other site-specific homing molecules like LPAM-1 (37) or VLA-4 (23, 26) rather than CLA. This allocation of certain cells to defined parts of the body might represent a homeostatic immune-surveillance response (10), or, instead, it has been hypothesised that senescent T cells are mobilised into the blood to facilitate their subsequent apoptosis in peripheral tissues (21, 34), which may contribute to progenitor cell mobilisation after exercise (24). With regards to cutaneous surveillance against tumours and stressed tissue cells, it is unlikely this is tasked by $\alpha\beta$ CD8⁺ T cells alone (13), and is likely supported by $\gamma\delta$ T cells and NK cells, which are highly responsive to acute stress (1, 6), and have the migratory capacity to enter cutaneous sites (13). We found that CLA⁺ CD56⁺ NK cells were substantially mobilised by exercise, but, much like CD8⁺ T cells, were outnumbered by the mobilisation of CLA⁻ NK cells.

This is the first study to investigate and compare the effects of HIIE (also referred to as HIIT when repeated over several weeks or months) and continuous exercise on T cell and NK cell subset mobilisation responses. HIIE typically involves shorter and more intense bouts of exercise than more traditional forms (e.g., 30 minutes of moderate intensity running or cycling) and is thought to be more attractive and better tolerated by participants. HIIE has received considerable recent attention as an effective means of achieving certain physiological adaptations (e.g., improved insulin sensitivity and cardiorespiratory fitness, but probably not weight loss) in

healthy and diseased populations that are similar or superior to traditional endurance-based exercise (16).

Although we found a trend whereby a greater mobilisation of lymphocytes was observed during continuous exercise compared to HIIE, these differences were not significant for the majority of lymphocyte subpopulations investigated. We found that CD56⁺ NK cells – conventionally the most exercise responsive cells – were mobilised to a greater extent during continuous exercise compared to HIIE, a response that was driven by larger increases to CD56^{dim} NK cells. It may be that similar intensity-dependent effects would have been found for CD8⁺ T cells, and other subsets, if a larger group of participants had been tested.

A limitation of this study was that post-exercise blood samples were collected 30 minutes after exercise cessation, rather than after 60 minutes, when NK and CD8⁺ T cell lymphocytopenia is more pronounced (6, 39). This was a consequence of practical and logistical constraints imposed by the broader study that was being undertaken (41). At 30 min post-exercise, we found that all cell phenotypes were present in peripheral blood in similar numbers to pre-exercise levels. It would be of interest to investigate the contribution of CLA⁺ T cells to lymphocytopenia at later time points. An additional practical limitation was the cryopreservation of PBMCs, which may disproportionately affect the viability of some cell phenotypes – however this is unlikely to affect the within-subject model used in our study. A further consideration for future studies surrounds the ongoing debate over the optimal phenotyping of skin-homing CD8⁺ T cells. In addition to CLA, CCR4 and CCR8 may be useful in differentiating between CD8⁺ T cells involved in normal cutaneous immune-surveillance to those involved in acute or chronic

inflammation (7, 33). Finally, our investigations were limited to healthy individuals with no apparent cutaneous inflammation. Investigation of exercise-induced lymphocyte skin-homing in experimental models of infection or in clinical populations including psoriasis is warranted, and more invasive techniques such as biopsy may provide further insights into exercise-induced homing to cutaneous or inflammatory sites.

CONCLUSION

A greater number of CLA⁻ T cells and NK cells were mobilised into peripheral blood than CLA⁺ counterpart T cells and NK cells during exercise. Furthermore, the majority of EMRA T cells and CD56^{dim} cells – i.e., conventionally the most exercise-responsive cells – did not express CLA. Together, these findings demonstrate that CLA⁺ cells are not major contributors to exercise lymphocytosis, thus providing preliminary evidence that the skin is not a major origin, or homing-destination, of exercise-sensitive lymphocytes. We conclude that the most exercise-sensitive lymphocytes likely migrate from, and to, non-cutaneous sites post-exercise.

ACKNOWLEDGEMENTS: The results of this present study do not constitute endorsement by ACSM. The flow cytometric analyses for this study were funded by the University of Birmingham Clinical Immunology Service.

CONFLICTS OF INTEREST: None declared

ACCEPTED

REFERENCES

1. Anane LH, Edwards KM, Burns VE et al. Mobilization of gammadelta T lymphocytes in response to psychological stress, exercise, and beta-agonist infusion. *Brain, behavior, and immunity*. 2009;23(6):823-9.
2. Atanackovic D, Schnee B, Schuch G et al. Acute psychological stress alerts the adaptive immune response: stress-induced mobilization of effector T cells. *Journal of neuroimmunology*. 2006;176(1-2):141-52.
3. Benschop RJ, Nijkamp FP, Ballieux RE, Heijnen CJ. The effects of beta-adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium. *British journal of pharmacology*. 1994;113(4):1311-6.
4. Berg EL, Yoshino T, Rott LS et al. The cutaneous lymphocyte antigen is a skin lymphocyte homing receptor for the vascular lectin endothelial cell-leukocyte adhesion molecule 1. *The Journal of experimental medicine*. 1991;174(6):1461-6.
5. Campbell JJ, Brightling CE, Symon FA et al. Expression of chemokine receptors by lung T cells from normal and asthmatic subjects. *J Immunol*. 2001;166(4):2842-8.
6. Campbell JP, Riddell NE, Burns VE et al. Acute exercise mobilises CD8+ T lymphocytes exhibiting an effector-memory phenotype. *Brain, behavior, and immunity*. 2009;23(6):767-75.
7. Clark RA, Chong B, Mirchandani N et al. The vast majority of CLA+ T cells are resident in normal skin. *J Immunol*. 2006;176(7):4431-9.
8. Cooper MA, Fehniger TA, Caligiuri MA. The biology of human natural killer-cell subsets. *Trends in immunology*. 2001;22(11):633-40.

9. Dhabhar FS. Psychological stress and immunoprotection versus immunopathology in the skin. *Clinics in dermatology*. 2013;31(1):18-30.
10. Dhabhar FS, Malarkey WB, Neri E, McEwen BS. Stress-induced redistribution of immune cells--from barracks to boulevards to battlefields: a tale of three hormones--Curt Richter Award winner. *Psychoneuroendocrinology*. 2012;37(9):1345-68.
11. Dimitroff CJ, Kupper TS, Sackstein R. Prevention of leukocyte migration to inflamed skin with a novel fluorosugar modifier of cutaneous lymphocyte-associated antigen. *The Journal of clinical investigation*. 2003;112(7):1008-18.
12. Dimitrov S, Lange T, Born J. Selective mobilization of cytotoxic leukocytes by epinephrine. *J Immunol*. 2010;184(1):503-11.
13. Ebert LM, Meuter S, Moser B. Homing and function of human skin gammadelta T cells and NK cells: relevance for tumor surveillance. *J Immunol*. 2006;176(7):4331-6.
14. Fuhlbrigge RC, Kieffer JD, Armerding D, Kupper TS. Cutaneous lymphocyte antigen is a specialized form of PSGL-1 expressed on skin-homing T cells. *Nature*. 1997;389(6654):978-81.
15. Gannon GA, Rhind SG, Shek PN, Shephard RJ. Differential cell adhesion molecule expression and lymphocyte mobilisation during prolonged aerobic exercise. *European journal of applied physiology*. 2001;84(4):272-82.
16. Gibala MJ, Little JP, Macdonald MJ, Hawley JA. Physiological adaptations to low-volume, high-intensity interval training in health and disease. *The Journal of physiology*. 2012;590(Pt 5):1077-84.

17. Hamann D, Baars PA, Rep MH et al. Phenotypic and functional separation of memory and effector human CD8⁺ T cells. *The Journal of experimental medicine*. 1997;186(9):1407-18.
18. Hoshino T, Yamada A, Honda J et al. Tissue-specific distribution and age-dependent increase of human CD11b⁺ T cells. *J Immunol*. 1993;151(4):2237-46.
19. Kruger K, Lechtermann A, Fobker M, Volker K, Mooren FC. Exercise-induced redistribution of T lymphocytes is regulated by adrenergic mechanisms. *Brain, behavior, and immunity*. 2008;22(3):324-38.
20. Kruger K, Mooren FC. T cell homing and exercise. *Exercise immunology review*. 2007;13:37-54.
21. Kruger K, Mooren FC. Exercise-induced leukocyte apoptosis. *Exercise immunology review*. 2014;20:117-34.
22. Kurokawa Y, Shinkai S, Torii J, Hino S, Shek PN. Exercise-induced changes in the expression of surface adhesion molecules on circulating granulocytes and lymphocytes subpopulations. *European journal of applied physiology and occupational physiology*. 1995;71(2-3):245-52.
23. Mazo IB, Honczarenko M, Leung H et al. Bone marrow is a major reservoir and site of recruitment for central memory CD8⁺ T cells. *Immunity*. 2005;22(2):259-70.
24. Mooren FC, Kruger K. Apoptotic lymphocytes induce progenitor cell mobilization after exercise. *J Appl Physiol (1985)*. 2015:jap 00287 2015.
25. Neuber K, Schmidt S, Mensch A. Telomere length measurement and determination of immunosenescence-related markers (CD28, CD45RO, CD45RA, interferon-gamma and

- interleukin-4) in skin-homing T cells expressing the cutaneous lymphocyte antigen: indication of a non-ageing T-cell subset. *Immunology*. 2003;109(1):24-31.
26. Papayannopoulou T, Priestley GV, Nakamoto B, Zafiropoulos V, Scott LM. Molecular pathways in bone marrow homing: dominant role of alpha(4)beta(1) over beta(2)-integrins and selectins. *Blood*. 2001;98(8):2403-11.
27. Picker LJ, Kishimoto TK, Smith CW, Warnock RA, Butcher EC. ELAM-1 is an adhesion molecule for skin-homing T cells. *Nature*. 1991;349(6312):796-9.
28. Picker LJ, Michie SA, Rott LS, Butcher EC. A unique phenotype of skin-associated lymphocytes in humans. Preferential expression of the HECA-452 epitope by benign and malignant T cells at cutaneous sites. *The American journal of pathology*. 1990;136(5):1053-68.
29. Picker LJ, Treer JR, Ferguson-Darnell B, Collins PA, Bergstresser PR, Terstappen LW. Control of lymphocyte recirculation in man. II. Differential regulation of the cutaneous lymphocyte-associated antigen, a tissue-selective homing receptor for skin-homing T cells. *J Immunol*. 1993;150(3):1122-36.
30. Robert C, Kupper TS. Inflammatory skin diseases, T cells, and immune surveillance. *The New England journal of medicine*. 1999;341(24):1817-28.
31. Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature*. 1999;401(6754):708-12.
32. Santamaria Babi LF, Moser R, Perez Soler MT, Picker LJ, Blaser K, Hauser C. Migration of skin-homing T cells across cytokine-activated human endothelial cell layers involves interaction of the cutaneous lymphocyte-associated antigen (CLA), the very late antigen-

- 4 (VLA-4), and the lymphocyte function-associated antigen-1 (LFA-1). *J Immunol.* 1995;154(4):1543-50.
33. Schaerli P, Ebert LM, Moser B. Comment on "The vast majority of CLA+ T cells are resident in normal skin". *J Immunol.* 2006;177(3):1375-6; author reply 6-7.
34. Simpson RJ. Aging, persistent viral infections, and immunosenescence: can exercise "make space"? *Exercise and sport sciences reviews.* 2011;39(1):23-33.
35. Simpson RJ, Cosgrove C, Chee MM et al. Senescent phenotypes and telomere lengths of peripheral blood T-cells mobilized by acute exercise in humans. *Exercise immunology review.* 2010;16:40-55.
36. Simpson RJ, Cosgrove C, Ingram LA et al. Senescent T-lymphocytes are mobilised into the peripheral blood compartment in young and older humans after exhaustive exercise. *Brain, behavior, and immunity.* 2008;22(4):544-51.
37. Strauch UG, Lifka A, Gossler U, Kilshaw PJ, Clements J, Holzmann B. Distinct binding specificities of integrins alpha 4 beta 7 (LPAM-1), alpha 4 beta 1 (VLA-4), and alpha IEL beta 7. *International immunology.* 1994;6(2):263-75.
38. Thatte J, Dabak V, Williams MB, Braciale TJ, Ley K. LFA-1 is required for retention of effector CD8 T cells in mouse lungs. *Blood.* 2003;101(12):4916-22.
39. Turner JE, Aldred S, Witard OC, Drayson MT, Moss PM, Bosch JA. Latent cytomegalovirus infection amplifies CD8 T-lymphocyte mobilisation and egress in response to exercise. *Brain, behavior, and immunity.* 2010;24(8):1362-70.
40. Viswanathan K, Dhabhar FS. Stress-induced enhancement of leukocyte trafficking into sites of surgery or immune activation. *Proceedings of the National Academy of Sciences of the United States of America.* 2005;102(16):5808-13.

41. Wadley AJ, Chen YW, Bennett SJ et al. Monitoring changes in Thioredoxin and over-oxidised Peroxiredoxin in response to exercise in humans. *Free radical research*. 2015;49(3):290-8.
42. Wadley AJ, Chen YW, Lip GY, Fisher JP, Aldred S. Low volume-high intensity interval exercise elicits antioxidant and anti-inflammatory effects in humans. *Journal of sports sciences*. 2015;27:1-9.
43. Williams MB, Butcher EC. Homing of naive and memory T lymphocyte subsets to Peyer's patches, lymph nodes, and spleen. *J Immunol*. 1997;159(4):1746-52.

FIGURE CAPTIONS

Figure 1. Mean $\pm$ SEM changes in the proportion of CD8⁺ T cell subsets (CM = central memory; EM = effector memory; EMRA = CD45RA⁺ effector memory) that were either positive or negative for cell surface CLA, at baseline and immediately after Continuous exercise. It was found that CLA⁺ CD45RA⁺ effector memory CD8⁺ T cells were mobilised to a greater magnitude than CLA⁻ CD45RA⁺ effector memory CD8⁺ T cells; similar patterns were found for HIIE (data not shown).

* Indicates significant difference between CLA⁺ and CLA⁻ ($p < .05$).

Figure 2. Mean $\pm$ SEM changes in the proportion of CD8⁺ T cell subsets (NA = naïve; CM = central memory; EM = effector memory; EMRA = CD45RA⁺ effector memory) that were positive for CLA, from pre- to immediately post-continuous exercise. A reduction in the proportion of CLA⁺ CM, EM and EMRA cells was observed immediately after continuous exercise compared to baseline; similar patterns were observed during HIIE (data not shown).

* Indicates significant differences in the proportion of CLA⁺ cells between pre- and post-continuous exercise ($p < .05$).

Figure 3. Mean $\pm$ SEM changes in the proportion of NK cell subsets that were either CLA⁺ or CLA⁻, at baseline and immediately after continuous exercise. A trend was observed whereby CLA⁻ CD56^{dim} cells were mobilised to a greater extent than CLA⁺ CD56^{dim} NK cells ($p = .058$); similar patterns were found for HIIE (data not shown).

Figure 4. Mean $\pm$ SEM changes in the proportion of NK cell subsets that were CLA⁺, from pre- to immediately post-continuous exercise. A reduction in the proportion of CLA⁺ CD56^{bright} and

CD56^{dim} NK cells was observed from pre- to post-exercise; similar patterns were observed during HIIE (data not shown).

* Indicates significant differences in the proportion of CLA+ cells between pre- and post-continuous exercise ($p < .05$).

ACCEPTED

Figure 1

Figure 2

Figure 3

Figure 4

Table 1. Changes in the numbers of T-lymphocytes and T-lymphocyte subpopulations in response to HIIE and continuous exercise (mean $\pm$ SD).

Cells per μ L	Condition	Baseline	Exercise	+30min	Main effects of time	Time $\times$ task interaction
CD3+	HIIE	1494.4 $\pm$ 406.2	2508.8 $\pm$ 1076.5*	1330.2 $\pm$ 293.2	$F_{(2,16)} = 16.18; p = .002$	$F_{(2,32)} = 2.06; p = \text{NS}$
	Continuous Ex	1454.1 $\pm$ 321.9	3049.6 $\pm$ 794.4***	1416.3 $\pm$ 363.1	$F_{(2,16)} = 42.97; p < .001$	
CD3+ CLA+	HIIE	189.2 $\pm$ 89.7	316.2 $\pm$ 237.0	186.8 $\pm$ 95.7	$F_{(2,16)} = 6.80; p = .030$	$F_{(2,32)} = .37; p = \text{NS}$
	Continuous Ex	183.0 $\pm$ 91.4	350.6 $\pm$ 166.6**	196.7 $\pm$ 123.2	$F_{(2,16)} = 26.06; p < .001$	
CD3+ CLA-	HIIE	1305.2 $\pm$ 376.2	2192.6 $\pm$ 894.2**	1143.4 $\pm$ 260.2	$F_{(2,16)} = 17.72; p = .002$	$F_{(2,32)} = 2.33; p = \text{NS}$
	Continuous Ex	1271.0 $\pm$ 293.3	2699.0 $\pm$ 276.0***	1219.6 $\pm$ 291.7	$F_{(2,16)} = 41.40; p < .001$	
CD3+ CD8+	HIIE	509.5 $\pm$ 181.9	1000.9 $\pm$ 654.7	463.1 $\pm$ 174.7	$F_{(2,16)} = 9.46; p = .014$	$F_{(2,32)} = .89; p = \text{NS}$
	Continuous Ex	491.2 $\pm$ 162.3	1185.0 $\pm$ 398.8***	490.4 $\pm$ 195.4	$F_{(2,16)} = 48.99; p < .001$	
CD3+ CD8+ CLA+	HIIE	41.3 $\pm$ 21.6	77.9 $\pm$ 65.6	43.2 $\pm$ 29.3	$F_{(2,16)} = 6.22; p = .035$	$F_{(2,32)} = .28; p = \text{NS}$
	Continuous Ex	40.7 $\pm$ 26.2	86.9 $\pm$ 53.8**	43.8 $\pm$ 32.7	$F_{(2,16)} = 23.95; p = .001$	
CD3+ CD8+ CLA-	HIIE	468.2 $\pm$ 163.4	923.0 $\pm$ 591.5	420.0 $\pm$ 148.7	$F_{(2,16)} = 9.76; p = .013$	$F_{(2,32)} = .95; p = \text{NS}$
	Continuous Ex	450.4 $\pm$ 141.2	1098.1 $\pm$ 355.0***	446.5 $\pm$ 166.9	$F_{(2,16)} = 48.96; p < .001$	
Naïve CD8+	HIIE	239.9 $\pm$ 93.1	341.6 $\pm$ 144.1*	192.8 $\pm$ 62.2	$F_{(2,16)} = 14.14; p < .001$	$F_{(2,32)} = 1.51; p = \text{NS}$
	Continuous Ex	215.9 $\pm$ 76.1	377.9 $\pm$ 69.0***	198.5 $\pm$ 56.4	$F_{(2,16)} = 55.18; p < .001$	
Naïve CD8+ CLA+	HIIE	3.4 $\pm$ 1.4	5.2 $\pm$ 2.6	3.0 $\pm$ 1.4	$F_{(2,16)} = 10.20; p = .001$	$F_{(2,32)} = .97; p = \text{NS}$
	Continuous Ex	3.5 $\pm$ 1.8	6.3 $\pm$ 3.4*	3.5 $\pm$ 2.2	$F_{(2,16)} = 17.67; p = .002$	
Naïve CD8+ CLA-	HIIE	236.5 $\pm$ 92.2	336.4 $\pm$ 142.3*	189.8 $\pm$ 62.0	$F_{(2,16)} = 14.13; p < .001$	$F_{(2,32)} = 1.49; p = \text{NS}$
	Continuous Ex	212.4 $\pm$ 75.8	370.5 $\pm$ 70.3***	195.0 $\pm$ 55.4	$F_{(2,16)} = 54.09; p < .001$	
CM	HIIE	118.8 $\pm$ 37.2	221.8 $\pm$ 97.1*	124.7 $\pm$ 55.9	$F_{(2,16)} = 21.50; p = .001$	$F_{(2,32)} = 3.85; p = \text{NS}$
	Continuous Ex	121.9 $\pm$ 64.2	278.6 $\pm$ 96.2***	132.3 $\pm$ 85.4	$F_{(2,16)} = 101.54; p < .001$	
CM CD8+ CLA+	HIIE	20.2 $\pm$ 8.3	35.6 $\pm$ 25.0	21.7 $\pm$ 12.7	$F_{(2,16)} = 7.01; p = .026$	$F_{(2,32)} = .29; p = \text{NS}$
	Continuous Ex	20.0 $\pm$ 11.8	38.9 $\pm$ 20.7**	21.3 $\pm$ 14.7	$F_{(2,16)} = 25.16; p < .001$	
CM CD8+ CLA-	HIIE	98.6 $\pm$ 30.4	186.2 $\pm$ 72.9**	103.1 $\pm$ 44.2	$F_{(2,16)} = 27.81; p < .001$	$F_{(2,32)} = 5.69; p = .017$
	Continuous Ex	101.9 $\pm$ 53.5	239.7 $\pm$ 77.4***	110.6 $\pm$ 71.8	$F_{(2,16)} = 123.76; p < .001$	
EM	HIIE	64.4 $\pm$ 29.9	144.9 $\pm$ 70.8**	69.5 $\pm$ 37.2	$F_{(2,16)} = 26.74; p = .001$	$F_{(2,32)} = 3.30; p = \text{NS}$
	Continuous Ex	66.6 $\pm$ 39.3	184.8 $\pm$ 71.0***	75.6 $\pm$ 52.0	$F_{(2,16)} = 80.73; p < .001$	
EM CD8+ CLA+	HIIE	12.3 $\pm$ 10.5	24.7 $\pm$ 27.5	13.7 $\pm$ 13.4	$F_{(2,16)} = 4.73; p = \text{NS}$	$F_{(2,32)} = .13; p = \text{NS}$
	Continuous Ex	12.0 $\pm$ 10.8	26.9 $\pm$ 24.5*	13.6 $\pm$ 14.0	$F_{(2,16)} = 11.22; p = .010$	
EM CD8+ CLA-	HIIE	52.1 $\pm$ 20.5	120.2 $\pm$ 44.7***	55.8 $\pm$ 24.6	$F_{(2,16)} = 40.39; p < .001$	$F_{(2,32)} = 5.08; p = .026$
	Continuous Ex	54.6 $\pm$ 29.7	157.8 $\pm$ 51.3***	62.2 $\pm$ 40.4	$F_{(2,16)} = 90.05; p < .001$	
EMRA	HIIE	88.3 $\pm$ 87.0	292.6 $\pm$ 390.3	76.1 $\pm$ 69.7	$F_{(2,16)} = 4.07; p = \text{NS}$	$F_{(2,32)} = .13; p = \text{NS}$
	Continuous Ex	86.6 $\pm$ 57.7	344.6 $\pm$ 296.6*	83.8 $\pm$ 52.1	$F_{(2,16)} = 9.16; p = .016$	
EMRA CD8+ CLA+	HIIE	5.4 $\pm$ 4.0	12.4 $\pm$ 12.6	4.9 $\pm$ 3.7	$F_{(2,16)} = 5.35; p = .047$	$F_{(2,32)} = .29; p = \text{NS}$
	Continuous Ex	5.4 $\pm$ 4.4	14.6 $\pm$ 10.9**	5.4 $\pm$ 4.6	$F_{(2,16)} = 17.13; p = .003$	
EMRA CD8+ CLA-	HIIE	80.9 $\pm$ 84.0	280.2 $\pm$ 378.7	71.2 $\pm$ 66.8	$F_{(2,16)} = 4.02; p = \text{NS}$	$F_{(2,32)} = .12; p = \text{NS}$
	Continuous Ex	81.4 $\pm$ 54.4	329.9 $\pm$ 291.0*	78.4 $\pm$ 48.6	$F_{(2,16)} = 8.75; p = .018$	

* $p < .05$ in comparison to baseline

** $p < .01$ in comparison to baseline

*** $p < .001$ in comparison to baseline

NS $p > .05$

Table 2. Changes in NK cell numbers in response to HIIE and continuous exercise (mean $\pm$ SD).

Cells per μ L	Condition	Baseline	Exercise	+30min	Main effects of time	Time $\times$ task interaction
CD56+	HIIE	112.4 $\pm$ 44.8	591.9 $\pm$ 361.0**	115.2 $\pm$ 59.5	$F_{(2,16)} = 18.27; p = .003$	$F_{(2,32)} = 4.47; p = .05$
	Continuous Ex	164.3 $\pm$ 144.2	1130.6 $\pm$ 707.5**	157.6 $\pm$ 159.9	$F_{(2,16)} = 22.64; p = .001$	
CD56+CLA+	HIIE	28.2 $\pm$ 10.4	92.7 $\pm$ 47.8**	29.2 $\pm$ 12.4	$F_{(2,16)} = 23.96; p = .001$	$F_{(2,32)} = 3.37; p = \text{NS}$
	Continuous Ex	31.7 $\pm$ 9.5	142.1 $\pm$ 69.2**	31.9 $\pm$ 10.9	$F_{(2,16)} = 26.10; p = .001$	
CD56+CLA-	HIIE	84.1 $\pm$ 44.6	499.3 $\pm$ 352.6*	86.0 $\pm$ 56.5	$F_{(2,16)} = 14.13; p = .005$	$F_{(2,32)} = 3.75; p = \text{NS}$
	Continuous Ex	132.6 $\pm$ 136.6	988.4 $\pm$ 698.6**	125.7 $\pm$ 153.2	$F_{(2,16)} = 18.14; p = .003$	
CD56 ^{bright}	HIIE	11.0 $\pm$ 7.4	23.9 $\pm$ 19.4*	15.2 $\pm$ 13.0	$F_{(2,16)} = 9.98; p = .011$	$F_{(2,32)} = 2.17; p = \text{NS}$
	Continuous Ex	12.8 $\pm$ 7.0	36.0 $\pm$ 25.7*	15.6 $\pm$ 7.8	$F_{(2,16)} = 11.05; p = .009$	
CD56 ^{bright} CLA+	HIIE	7.6 $\pm$ 5.5	15.0 $\pm$ 12.7*	10.2 $\pm$ 9.1	$F_{(2,16)} = 9.05; p = .015$	$F_{(2,32)} = 1.32; p = \text{NS}$
	Continuous Ex	8.4 $\pm$ 4.7	20.3 $\pm$ 15.8*	10.2 $\pm$ 5.7	$F_{(2,16)} = 8.88; p = .015$	
CD56 ^{bright} CLA-	HIIE	3.4 $\pm$ 2.0	8.9 $\pm$ 7.2*	5.0 $\pm$ 4.0	$F_{(2,16)} = 10.14; p = .011$	$F_{(2,32)} = 1.79; p = \text{NS}$
	Continuous Ex	4.4 $\pm$ 2.7	15.8 $\pm$ 14.4	5.4 $\pm$ 2.5	$F_{(2,16)} = 6.35; p = .035$	
CD56 ^{dim}	HIIE	101.4 $\pm$ 44.5	568.0 $\pm$ 343.6**	99.9 $\pm$ 55.6	$F_{(2,16)} = 18.41; p = .003$	$F_{(2,32)} = 4.49; p = .05$
	Continuous Ex	151.5 $\pm$ 47.2	1094.5 $\pm$ 229.9**	141.9 $\pm$ 53.1	$F_{(2,16)} = 22.72; p = .001$	
CD56 ^{dim} CLA+	HIIE	20.6 $\pm$ 10.0	77.7 $\pm$ 43.8**	19.0 $\pm$ 7.6	$F_{(2,16)} = 22.68; p = .001$	$F_{(2,32)} = 3.21; p = \text{NS}$
	Continuous Ex	23.3 $\pm$ 7.8	121.8 $\pm$ 60.8**	21.7 $\pm$ 8.9	$F_{(2,16)} = 25.51; p = .001$	
CD56 ^{dim} CLA-	HIIE	80.7 $\pm$ 44.0	490.3 $\pm$ 345.8*	81.0 $\pm$ 55.0	$F_{(2,16)} = 14.16; p = .005$	$F_{(2,32)} = 3.75; p = \text{NS}$
	Continuous Ex	128.2 $\pm$ 135.3	972.7 $\pm$ 689.1**	120.3 $\pm$ 152.3	$F_{(2,16)} = 18.19; p = .003$	

* $p < .05$ in comparison to baseline** $p < .01$ in comparison to baselineNS $p > .05$