Hydroclimatic influences on peatland CO₂ exchange following upland forest harvesting on the boreal plains

Plach, J.M.; Petrone, R. M.; Waddington, J.M.; Kettridge, Nicholas; Devito, K. J.

DOI:
[10.1002/eco.1750](https://doi.org/10.1002/eco.1750)

License:
Other (please specify with Rights Statement)

Document Version
Peer reviewed version

Citation for published version (Harvard):
Plach, JM, Petrone, RM, Waddington, JM, Kettridge, N & Devito, KJ 2016, 'Hydroclimatic influences on peatland CO₂ exchange following upland forest harvesting on the boreal plains', *Ecohydrology*.
<https://doi.org/10.1002/eco.1750>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Hydroclimatic Influences on Peatland CO₂ Exchange Following Upland Forest Harvesting on the Boreal Plains, which has been published in final form at 10.1002/eco.1750. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

**Hydroclimatic Influences on Peatland CO₂ Exchange
Following Upland Forest Harvesting on the Boreal Plains**

Journal:	<i>Ecohydrology</i>
Manuscript ID	ECO-15-0169.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Plach, Janina; University of Waterloo, Faculty of Environment Petrone, Richard; University of Waterloo, Department of Geography and Environmental Management Waddington, James; McMaster University, School of Geography, Earth and Environmental Sciences Kettridge, Nick; University of Birmingham, School of Geography, Earth and Environmental Sciences Devito, Kevin; University of Alberta, Department of Biological Sciences
Keywords:	forest harvesting, peatland, CO ₂ , NEE, microclimate , soil moisture, boreal forest

SCHOLARONE™
Manuscripts

Hydroclimatic Influences on Peatland CO₂ Exchange Following Upland Forest Harvesting on the Boreal Plains

J.M. Plach^{1*}, R.M. Petrone¹, J.M. Waddington², N. Kettridge³ and K.J. Devito⁴

¹ Department of Geography & Environmental Management, University of Waterloo, 200 University Ave West, Waterloo, Ontario, Canada, N2L 3G1.

² School of Geography & Earth Sciences, McMaster University, Hamilton, Ontario, Canada

³ School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK

⁴Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

**corresponding author phone: (905) 630-6979; e-mail: janinaplach@gmail.com; postal address: Faculty of Environment, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada*

1 Abstract

2 A comparative study of forest clear-cut logging effects on daily growing season (May to
3 October) net ecosystem CO₂ exchange (*NEE*) of adjacent peatlands was conducted in two
4 neighbouring forest upland-peatland complexes over four-years (2005 to 2008) on the Boreal
5 Plains (BP) of Alberta, Canada. Higher vapour pressure deficit at the harvested-upland (H-U)
6 peatland, reflecting increased turbulent mixing after adjacent upland forest removal (2007 and
7 2008), resulted in increased peatland evapotranspiration rates that contributed to a seasonal
8 decline in soil moisture (*VMC*) influencing *NEE*. Overall, a significant change in mid-season
9 *NEE* occurred at the H-U peatland one-year post-harvesting, greater than *NEE* changes at the
10 neighboring intact-upland peatland. However, two years post-harvesting, mid-season *NEE*
11 returned to within range of pre-harvesting variability (-0.54 to 1.34 g CO₂-C m⁻² d⁻¹). Results of
12 this study demonstrate that BP peatland *NEE* is largely regulated by site-specific water
13 availability, which in turn, may be influenced in the short-term by shifting microclimate and soil
14 moisture patterns due to clear-cut logging. As such, predicting long-term carbon storage function
15 of BP peatlands will require careful consideration of changing hydroclimatic conditions due to
16 rapid expansion of BP deforestation, given that these ecosystems already exist in a state of
17 hydrologic risk in this moisture deficit eco-region.

18
19 Key Words: *NEE, CO₂, peatland, forest harvesting, boreal forest, soil moisture, microclimate*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Introduction**

2 Peatlands and wetlands cover up to 50% of the land surface on the Boreal Plains (BP) and store a
3 significant portion of carbon in Canada (Timoney, 2003; Kuhry *et al.*, 1993). The BP of western
4 Canada is experiencing extensive deforestation by timber harvesting (clear-cut) as well as road
5 and corridor construction from industrial oil and gas expansion (Timoney, 2003). Forests
6 regulate the microclimatic and hydrologic conditions (incoming solar radiation, wind velocity,
7 turbulence, temperature and moisture of the air and soils) of edge and adjacent ecosystems (Chen
8 *et al.*, 1993; Flesch and Wilson, 1999; Petrone *et al.*, 2007; Markfort *et al.*, 2014). Rapid
9 harvesting of upland forests may threaten the hydroclimatic stability of the already
10 hydrologically tenuous adjacent peatland ecosystems on the BP (Solondz *et al.*, 2008; Brown *et*
11 *al.*, 2010). While BP forest disturbance research has focused primarily on harvested areas and
12 the associated water and carbon dynamics of the disturbed forest soils (e.g. Amiro *et al.*, 2006;
13 Petrone *et al.*, 2015; Whitson *et al.*, 2005; Carrera-Hernández *et al.*, 2011), the potential impacts
14 on the terrestrial-atmosphere exchange of water and carbon dioxide (CO₂) in adjacent peatlands
15 remains unknown. Given the importance of peatlands for BP carbon storage and water supply
16 (Ferone and Devito, 2004; Smerdon *et al.*, 2005; Solondz *et al.*, 2008;), understanding how these
17 peatlands respond to upland clear-cut logging is fundamental to improving the design and
18 implementation of landscape management and forestry practices across this eco-region (Johnson
19 and Miyanishi, 2008).

20 Previous studies examining carbon exchange in peatlands show that soil temperature and
21 moisture conditions are well coupled to carbon losses (plant respiration and soil decomposition)
22 and carbon uptake (plant productivity) at both the plot-scale and ecosystem-level (Solondz *et al.*,
23 2008; Bubier *et al.*, 1998, 2003; McNeil and Waddington. 2003; Petrone *et al.*, 2011). The net

balance between CO₂ uptake and release (Net Ecosystem Exchange, *NEE*) is generally highest (i.e. increased net CO₂ release) under the most favourable conditions for microbial decomposition (i.e. warm, low moisture oxic peat) (Solondz *et al.*, 2008; Bubier *et al.*, 1998; Silvola *et al.*, 1996). Although the impact of land use changes on peatland water cycling and *NEE* are widely investigated in peatlands on the Boreal shield of eastern Canada (e.g. Waddington and Price 2000; Tuittila *et al.*, 1999), limited work on the response of peatland hydrology and/or trace gas exchange to anthropogenic disturbances exists on the BP of western Canada (Strack *et al.*, 2014). Heterogeneous glacial deposits along with the sub-humid climate of the BP, whereby precipitation roughly equaling potential evapotranspiration, results in water table positions and soil moisture gradients that are not under topographic control (Devito *et al.*, 2005). As such, it is unknown if shifts in peatland moisture conditions, soil temperature and carbon dynamics in response to disturbance events observed in the runoff-dominated shield can be extrapolated to peatlands in the complex hydrology of the sub-humid climate of the BP.

Forest cutblocks experience higher wind speeds, short-wave radiation, air temperatures and lower atmospheric moisture relative to within forest canopy stands (Chen *et al.*, 1993). Abrupt transitions between flat surfaces (e.g. cutblocks) and forest canopies or between vegetation types within a landscape can dynamically alter the atmospheric boundary layer and turbulent flow patterns across the transition zones (Markfort *et al.*, 2014; Yang *et al.*, 2006; Flesch and Wilson, 1999). As such, clear-cut logging has the potential to alter the microclimate conditions in adjacent peatlands to influence evapotranspiration (*ET*) and ecosystem water loss (Petrone *et al.*, 2007; Helgason and Pomeroy, 2005; Helgason and Pomeroy, 2012; Wharton *et al.*, 2010; Monteith, 1965). Given that subtle changes in *ET* can result in soil moisture deficits in this sub-humid climate (Devito *et al.*, 2005), shifts in *ET* are likely to be important to the CO₂ sink or

1
2
3 1 source status of BP peatlands. Due to the multitude of compounding hydrological interactions
4
5 2 and feedbacks in northern peatlands (Waddington *et al.*, 2015), applying an integrated
6
7 3 monitoring approach of hydrology, microclimate and CO₂ exchange is essential to evaluate the
8
9 4 natural baseline ecohydrological conditions of BP peatlands (e.g. Solondz *et al.*, 2008) and to
10
11 5 compare the potential impact of adjacent forest disturbance on the complex hydroclimatic and
12
13 6 biogeochemical factors governing peatland *NEE*.
14
15
16
17
18 7 Long-term CO₂ exchange is coupled to atmospheric processes (Lafleur *et al.*, 1997). Increased
19
20 8 summer warming in recent decades observed in western Canada (Gullet and Skinner, 1992)
21
22 9 along with projections of rising global temperatures and greater drought frequency in Boreal
23
24 10 regions suggests a future reduction in CO₂ sequestration by Boreal peatlands (Intergovernmental
25
26 11 Panel on Climate Change (IPCC) 2014; Gorham, 1991). As such, understanding the short-term
27
28 12 *NEE* response of BP peatlands to forest disturbances in the context of climate variability is
29
30 13 essential to facilitate effective predictions of the long-term fate of these large carbon stores. Due
31
32 14 to the large area of the BP covered by peatlands, establishing the relationships between clear-cut
33
34 15 logging, peatland hydroclimate and *NEE* at the peatland-scale could provide the means to
35
36 16 simplify and extrapolate the carbon functioning of these ecosystems to the landscape-scale and
37
38 17 generalize peatland responses to disturbance by monitoring clear-cut areas across the eco-region.
39
40 18 As such, the objectives of this study were to examine: (1) *NEE* of BP peatlands during the snow-
41
42 19 free period; (2) the relative impact of upland clear-cut logging on adjacent peatland *NEE*,
43
44 20 including the hydrologic and microclimatological controls on this exchange; and (3) estimate the
45
46 21 sustainability of BP peatlands' functionality as a CO₂ source or sink in context of periodic land
47
48 22 use disturbances and climate change.
49
50
51
52
53
54
55
56
57
58
59
60

23 **Methods**

1 Site Description

2 Two adjacent forested-peatlands, one with an intact adjacent forested upland (I-U) and one with
3 a harvested upland (H-U), were examined in this study located in the Utikuma Region Study
4 Area (URSA) near Utikuma Lake, north-central Alberta, Canada (56° 20'N, 115° 30' W; Figure
5 1). The two complexes are located on a disintegration ice moraine landform (Paulen *et al.*, 2004),
6 within the Central Mixedwood Natural sub-region of the Boreal Forest in Alberta (Natural
7 Regions Committee 2006) or Mid-Boreal Uplands Ecoregion of the Boreal Plains Ecozone
8 Alberta, Canada (Ecological Stratification Working Group 1996). The climate in this region is
9 characterized by short, warm summers and long, cold winters with a 30-year average annual
10 temperature, precipitation and potential evapotranspiration (PET) for the region as 1.7 °C, 485
11 mm and 515 mm, respectively (Environment Canada, 2007). The mean annual temperature and
12 precipitation at the study site for 2005, 2006, 2007 and 2008 were 2.5, 2.5, 1.5 and 0.9 °C and
13 491, 432, 530 and 504 mm, respectively making the study period slightly warmer and drier
14 during pre-harvest and slightly cooler and wetter post-harvest than the 30-year normal. The
15 prevailing wind direction across the study sites was from the South during the four-year study
16 period (Supporting Information S1).

17 The two peatlands surround shallow ponds (< 1 m depth) and are adjacent to hillslopes with
18 aspen-dominated uplands (up to 7 m above the pond surface) with a canopy height of
19 approximately 17 m to 21 m on average (Brown *et al.*, 2013) and canopy coverage averaging
20 68% (Chasmer *et al.*, 2010). The peatlands and shallow ponds are located in a recharge zone, and
21 water tables typically grade away from the peatlands into the hillslope (Ferone and Devito, 2004;
22 Redding and Devito, 2008). Vegetation in the peatlands are comparable, composed of a shrub
23 layer comprising mostly *Ledum groenlandicum*, *Vaccinium vitisidaea* and *Chamaedaphne*

1 *calyculata* as well as groundcover dominated by bryophyte and lichen species characteristic of
2 poor fen communities, mainly *Sphagnum* species and feather mosses (Solondz *et al.*, 2008;
3 Petrone *et al.*, 2011). A similar open canopy of black spruce (canopy coverage averaging 36 to
4 60% (Chasmer *et al.*, 2010)), approximately 2 m in height, occurs within these peatlands. Peat
5 physical characteristics (e.g. bulk density, hydraulic conductivity) do not differ between the
6 peatlands (Petrone *et al.*, 2008). Minimal peat subsidence occurs at these sites and does not
7 readily respond to water level changes in the peat or adjacent ponds (Petrone *et al.*, 2008).

8 *CO₂ Field Measurements*

9 Chamber CO₂ data was collected at ten sites within the harvested-upland (H-U) peatland and six
10 in the intact-upland (I-U) peatland over the four snow-free seasons (Figure 1). At each site 20 cm
11 (diameter) polyvinylchloride (PVC) collars were placed in adjacent lawns (classified as
12 topographically high mounds) and depressions (low lying areas) to capture the range of
13 microtopography in the peatlands and associated differences in CO₂ exchange (Petrone *et al.*,
14 2011). CO₂ exchange was measured using a dynamic closed chamber system with an Infrared
15 Gas Analyzer (IRGA) (EGM-4, PP Systems, Maryland, USA) (Solondz *et al.*, 2008). Removable
16 clear lexan chambers were fitted to the permanently installed collars, with coolant tubes and fans
17 operating to mimic ambient air temperatures and gradients (Solondz *et al.*, 2008; Welles *et al.*,
18 2001). For each sample, concentrations of CO₂ ppm were measured at 30 second intervals for 2.5
19 minutes during 0900-1600 h approximately ten times per month each season. Sampling times at
20 each location were randomly selected during each sampling day to ensure measurements were
21 taken over a wide range of light and temperature regimes that may occur throughout the day.
22 Chambers were covered with an opaque neoprene shroud when measuring the gross respiration
23 (R_{tot} = autotrophic and heterotrophic). Gross ecosystem productivity (*GEP*) was calculated as the

1 difference between NEE and R_{tot} ,

$$2 \quad GEP = NEE - R_{tot} \quad (1)$$

3 Negative values indicate a net CO_2 uptake by the peatland, and positive values indicate a net CO_2
4 release by respiration into the atmosphere.

5 *Environmental parameters*

6 Air (T_a) temperature and relative humidity (RH) (PP Systems, Maryland, USA), soil (T_{soil})
7 temperatures (Omega Engineering, Inc., Connecticut, USA) and photosynthetically active
8 radiation (PAR) (Quantum Sensor; LiCor Inc., Nebraska, USA) were recorded at the same
9 temporal and spatial scales as the CO_2 fluxes. T_a , RH and PAR were measured both inside and
10 outside of the chamber at approximately 0.5 m above the surface during each 2.5-min chamber
11 measurement. Soil (T_{soil}) temperatures were measured at 2, 5, and 10 cm depths and averaged for
12 the values at the three depths. VMC was measured beside each collar using time domain
13 reflectometry (TDR) (Hydrosense Probe, Campbell Scientific, Inc, Utah, USA) to give a bulk
14 soil moisture value over the top 10 cm of the soil profile. The TDR was calibrated in the lab by
15 drying representative undisturbed peat samples to different moisture contents (Solondz *et al.*,
16 2008). The field point measurements were applied to the chamber CO_2 measurements to
17 determine the modeling of GEP and R_{tot} . Water Table depth (WT), measured in meters below the
18 surface, was recorded weekly using PVC pipe wells (5 cm O.D.) in the peatland and upland at
19 each I-U and H-U site (Figure 1).

20 Meteorological towers (MET) located in each peatland (Figure 1) continuously measured
21 environmental parameters during the snow-free period (Day of Year (DOY)): 120 to 280) of

1 each year. Average VMC in the upper 30 cm of the peat was recorded using water content
2 reflectrometry (CS616, Campbell Scientific Inc, Utah, USA) placed vertically in both a lawn and
3 depression. Net radiation (Q^*) was measured at 1.5 m above the peat surface using a net
4 radiometer (NRLite, Kipp and Zonen, The Netherlands). The T_a and RH (Vaisala, Finland) were
5 measured at the same height. RH was not available at the H-U peatland mid season (DOY 201 to
6 243) in 2007 and therefore; vapour pressure deficit (VPD) was gap filled according to a
7 regression (using all available VPD data from MET and manual chamber measurements during
8 the post-harvesting period) as a function of T_{air} ,

9
$$VPD = 0.4748e(0.075T_{air}), r^2 = 0.87 \quad (2)$$

10 T_{soil} was recorded at 2, 5 and 10 cm using thermocouples (Omega copper-constantin, Campbell
11 Scientific Inc, Logan, Utah, USA) in a lawn and depression. Ground heat flux (Q_G) was
12 measured according to the calorimetric method (Halliwell and Rouse, 1987; Petrone and Rouse,
13 2000; Petrone *et al.*, 2007) using the soil temperature profile and heat capacity calculations for
14 each soil layer (2 to 5 cm and 5 to 10 cm) accounting for changes in moisture content and state
15 (Sutherland *et al.*, 2014). Published values for heat capacities of peat soils under a range of
16 moisture conditions were used in the calculation of ground heat flux (Q_g) (Brown *et al.*, 2010;
17 Oke, 1987; Halliwell and Rouse, 1987; Petrone and Rouse, 2000; Petrone *et al.*, 2007).
18 Horizontal wind speed (u) measurements were collected using cup anemometers (014A, Met
19 One, Oregon, USA) at 1.4 m at both the H-U and I-U sites (Figure 1).

20 Downstream from a surface discontinuity, such as that from a clear-cut forest to peatland, will
21 create horizontal differences in roughness lengths (Helgason and Pomeroy, 2005). Such
22 differences in momentum sinks will cause large horizontal wind variances in the peatland, which

means that mean average wind speeds may not increase but will become more variable in intensity (Helgason and Pomeroy, 2012; Wharton *et al.*, 2010). Thus, horizontal turbulence intensity (I_u) for the peatlands was calculated according to (Turnispeed *et al.*, 2003),

$$I_u = \frac{\sigma_u}{U} \quad (3)$$

where σ_u is the standard deviation of the mean daily horizontal wind speed (m s^{-1}) and U is the mean of the mean daily horizontal wind speed (m s^{-1}) for the peatlands for each year of the study period. Wharton *et al.*, (2010) suggest that this is a more reliable means of assessing changes in turbulent regimes as a result of changing surface condition than more traditional approaches based largely on the friction velocity (u^*), which under these conditions may suggest weak turbulent conditions while actual horizontal turbulent fluxes may be large.

Evapotranspiration

Surface conductance and aerodynamic measurements from the peatland *MET* and wind velocity stations (see section above) were utilized to estimate *ET* by applying a standardized reference Penman-Monteith equation (Chasmer *et al.*, 2011; Temesgen *et al.*, 2005; ASCE-EWRI, 2005),

$$ET = \frac{0.408\Delta(Q^* - Q_G) + \gamma \frac{900}{T_a + 273} u (e_s - e_a)}{\Delta + \gamma(1 + 0.34u)} \quad (4)$$

where *ET* is the daily evapotranspiration rate (mm d^{-1}), Q^* is the daily net radiation ($\text{MJ m}^{-2} \text{d}^{-1}$), Q_G soil heat flux ($\text{MJ m}^{-2} \text{d}^{-1}$), T_a is the mean daily air temperature ($^{\circ}\text{C}$), u is the mean daily wind speed (m s^{-1}), e_s is the mean saturation vapor pressure (kPa), e_a is the actual vapor pressure (kPa),

1
2
3
4 1 γ psychrometric constant (kPa °C⁻¹), 900 and 0.34 are constants for reference type and
5
6 2 calculation time (mm d⁻¹) (Chasmer *et al.*, 2011; Fournier *et al.*, 2007; Temesgen *et al.*, 2005;
7
8 3 Banaszuk and Kamocki, 2008). Evaluating surface resistance (r_s) from individual chambers
9
10 4 based on a Mann-Whitney Rank Sum Test showed no significant difference in r_s between the
11
12 5 sites before or after harvesting ($U = 34584$, $p < 0.05$). Further, this *ET* model approach was
13
14 6 validated by comparing the seasonal average evaporation rates calculated in this study with
15
16 7 previously published *ET* values, from combined methods of eddy covariance and Priestley–
17
18 8 Taylor model, at the H-U peatland in 2005 and 2006 (Brown *et al.*, 2010; Petrone *et al.*, 2007).

22
23
24 9 *Peatland CO₂ modelling*

25
26
27 10 At each peatland, growing season (DOY 120 to 280) CO₂ exchange was estimated using the field
28
29 11 point flux measurements of *GEP* and R_{tot} (i.e. combined lawns and depressions). The relationship
30
31 12 between *GEP* and *PAR* was fitted empirically using a rectangular hyperbola regression (Whiting,
32
33 13 1994; Waddington and Roulet, 1996),

34
35
36
37 14
$$GEP = \frac{PAR \cdot Q \cdot GP_{max}}{(PAR \cdot Q + GP_{max})} \quad (5)$$

38
39
40
41 15 where *PAR* is measured $\mu\text{mol m}^2 \text{s}^{-1}$, *Q* is the quantum efficiency that describes the initial slope
42
43 16 of the *GEP* versus *PAR* hyperbola, GP_{max} is the theoretical maximum rate of *GEP*, representing
44
45 17 the asymptote of the hyperbola. Ecosystem R_{tot} was modeled using a linear regression with
46
47 18 average T_{soil} (5 cm depth) according to,

48
49
50
51 19
$$R_{tot} = aT_{soil} + b \quad (6)$$

52
53
54
55 20 where *a* and *b* are parameters fitted by least squares regression. Peatland respiration is strongly

correlated with 5 cm soil temperatures at these sites (Solondz *et al.*, 2008; Petrone *et al.*, 2011) and frequently observed to correlate with soil temperature in other Boreal peatlands (e.g. Bubier *et al.*, 1998). In this study, T_{soil} (5 cm depth) also showed the best overall correlation with R_{tot} of the measured environmental parameters across the four-year study and thus equation 6 was applied to each year for consistency in the model. The daily CO_2 exchange was estimated over the 160-day growing season by applying equations 5 and 6 to daily average PAR and T_{soil} measurements collected from the *MET* stations at each peatland (see Figure 1) in 2005 through 2008. Although variability in VMC within peatlands may influence T_{soil} , there was a general agreement between T_{soil} and VMC measured at the *MET* stations (i.e. T_{soil} used for R_{tot} modeling) with T_{soil} and VMC measurements at the chamber sites (Figure S2a and S2b). As such, *MET* station T_{soil} and VMC were used when analyzing temperature and moisture conditions between years and study sites. With the aim of highlighting differences between peatlands and the response to disturbance (rather than quantifying the exact carbon budgets), modeled GEP and R_{tot} parameters described the field point flux measurements fairly well for most sampling years (Table 1) and were comparable to scatter in GEP and R_{tot} models previously reported (e.g. Bubier *et al.*, 1998; Lafleur, 1999; Petrone *et al.*, 2011; Strack *et al.*, 2014). Residuals from the regressions showed no systematic bias thus, the NEE models did not over- or under- estimate the effect of the harvesting treatment. Uncertainty estimates for NEE were assessed by assigning regression standard errors for the different models used each year.

Statistical analyses

Microclimatological and carbon exchange rate (NEE , GEP and R_{tot}) differences between the two peatlands (i.e. I-U versus H-U) were analyzed using Kruskal-Wallis One-Way analysis of variance (ANOVA) on Ranks and *post hoc* Tukey Test (TT), and within each peatland using

1
2
3 1 Friedman Repeated Measures on ANOVA on Ranks and TT over the four-year study between
4
5 2 day of year (DOY) 201 to 243 (i.e. mid-season) due to missing microclimatological data outside
6
7 3 this period in some years. Microclimatological parameters with unequal sample sizes within this
8
9 4 time period were analyzed using Kruskal-Wallis One-Way ANOVA and *post hoc* test Dunn's
10
11 5 Method (DM). Variations in *NEE*, *GEP* and R_{tot} (all available data for each year) were plotted
12
13 6 against each environmental parameter to isolate relationships.
14
15
16
17

18 **RESULTS**
19

20 *Climate and environmental variables*
21
22
23

24
25 9 Minimum variability in the *WT* of the H-U peatland occurred between years, whereby median
26
27 10 *WT* ranged from only 11 to 16 cm depth below surface (DBS) between the pre- and post-
28
29 11 harvesting period (Figure 2). Despite relatively small *WT* fluctuations, large inter-annual
30
31 12 variability in peatland volumetric moisture content (*VMC*) was measured. During pre-harvesting
32
33 13 (2005 and 2006), median peatland *VMC* was high (0.69 and $0.52 \text{ m}^3 \text{ m}^{-3}$, respectively) and
34
35 14 responsive to precipitation events (i.e. increase *VMC*; Figure 2). However, post-harvesting (2007
36
37 15 and 2008) peatland *VMC* showed a consistent seasonal decline, despite greater and more evenly
38
39 16 distributed precipitation. Median *VMC* in 2007 and 2008 was 0.43 and $0.47 \text{ m}^3 \text{ m}^{-3}$, respectively
40
41 17 and did not respond as strongly to precipitation events compared to the pre-harvesting years. In
42
43 18 contrast to the H-U peatland, *WT* variability of the I-U peatland was much greater (Figure 2),
44
45 19 where median *WT* ranged from 38 to 18 cm DBS across the four years. Although *WT*
46
47 20 fluctuations were relatively large, variability in peatland *VMC* was minimal (0.24 to $0.26 \text{ m}^3 \text{ m}^{-3}$
48
49 21 ³). Overall, *VMC* at the I-U peatland was significantly different from the H-U peatland [$H =$
50
51
52
53
54
55
56
57
58
59
60

130.806, d.f. = 6, $p < 0.001$], with consistently lower [*post hoc* test (DM), $p < 0.05$] median *VMC* in both the pre- and post-harvest periods.

T_{soil} at the H-U peatland was most similar to *T_a* during the late growing season of the post-harvesting years only (2007 and 2008) when soil moisture declined to its lowest values (i.e. *VMC* $< 0.45 \text{ m}^3 \text{ m}^{-3}$; Figure 3). In contrast, at the drier I-U peatland, a relatively stronger relationship between *T_{soil}* and *T_a* was observed across years (i.e. *T_{soil}* approximately equal to *T_a*). *T_a* was similar between sites, with differences in median *T_a* varying $< 3^\circ\text{C}$ during each year and showing minimal changes in each peatland between years (i.e. $< 6^\circ\text{C}$). Similarly, minimal differences in median photosynthetic active radiation (*PAR*) were observed between peatlands (i.e. $< 2 \text{ W m}^{-2}$) and within each peatlands (i.e. $< 40 \text{ W m}^{-2}$) across the four-year study period (Figure S3).

Significant differences in vapour pressure deficit (*VPD*) occurred at the H-U peatland [$H = 94.520$, d.f. = 3, $p < 0.001$] across years (Figure 4a). Pre-harvesting (2005 and 2006) median *VPD* was 0.37 and 0.82 kPa, respectively, and did not differ significantly [$H = 151.730$, d.f. = 7, $p < 0.001$, *post hoc* test (DM), $p < 0.05$] from *VPD* at the I-U peatland. However, post-harvesting (2007 and 2008), median *VPD* increased [*post hoc* test (DM), $p < 0.05$] to 1.53 and 1.62 kPa, respectively, and was significantly different [*post hoc* test (DM), $p < 0.05$] from the I-U peatland. The largest seasonal fluctuations in *VPD* and the highest recorded *VPD* (i.e. $> 2 \text{ kPa}$) were measured at the H-U peatland in the post-harvesting years. In contrast, median *VPD* at the I-U peatland remained low (i.e. $\leq 0.79 \text{ kPa}$) during each year of the study. Wind speed at the H-U site significantly differed across years [$\chi^2(3) = 39.921$, $p < 0.001$] (Figure 4b); however, *post hoc* tests (TT) showed that wind speed (2007) did not significantly differ from pre-harvesting (2005). Median wind speed at the H-U site varied from 0.8 to 1.4 m s^{-1} across years, and showed similar variability to wind speed at the I-U site (median 1.1 to 1.6 m s^{-1}). Despite minimal changes in

1 wind speed, turbulence intensity (I_u) at the H-U site increased post-harvesting from 0.26 and 0.38
 2 in 2005 and 2006 to 0.60 and 0.52 in 2007 and 2008, respectively (Figure 4b). In contrast, I_u
 3 remained low at the I-U site across years, ranging from 0.24 to 0.34. Significant differences in
 4 ET [$H = 79.355$, d.f. = 3, $p < 0.001$] occurred at the H-U peatland across the four-years (Figure
 5 4c). Similar to peatland VDP, *post hoc* tests (DM) showed ET was significantly different ($p <$
 6 0.05) between pre- and post-harvesting years. Although ET also significantly increased [$H =$
 7 15.892, d.f. = 3, $p < 0.001$, *post hoc* tests (DM), $p < 0.05$] at the I-U peatland in 2008 compared
 8 to 2005, overall larger increases in ET occurred at the H-U peatland between the pre- and post-
 9 harvesting years, whereby median ET increased from 1.4 and 2.0 mm d⁻¹ in 2005 and 2006 to 3.4
 10 and 4.3 mm d⁻¹ in 2007 and 2008, respectively.

11 *Seasonal variation in peatland carbon exchange*

12 During the pre-harvesting period (2005 and 2006), inter-annual variability of NEE between
 13 peatlands and within each peatland was low (Figure 5). NEE rates at the H-U peatland were
 14 consistently lower (i.e. greater carbon uptake) and significantly different [$H = 285.517$, d.f. = 7,
 15 *post hoc* test (TT), $p < 0.05$] from NEE at the I-U peatland. The H-U peatland fluctuated between
 16 a slight carbon source and carbon sink (median 0.52 and -0.03 g CO₂-C m⁻² d⁻¹ in 2005 and 2006,
 17 respectively) while the I-U peatland functioned as a consistent slight carbon source (median 1.30
 18 and 1.65 g CO₂-C m⁻² d⁻¹ in 2005 and 2006, respectively) (Figure 5). Pre-harvesting R_{tot} was
 19 lower at the H-U peatland (median 5.28 and 7.96 g CO₂ m⁻² d⁻¹ in 2005 and 2006, respectively)
 20 and significantly different [$H = 282.365$, d.f. = 7, *post hoc* test (TT), $p < 0.05$] from the I-U
 21 peatland (median 10.12 and 12.10 g CO₂ m⁻² d⁻¹ in 2005 and 2006, respectively; Figure 6a). In
 22 contrast, GEP was slightly higher at the H-U peatland in 2005 and 2006 (median 3.72 and 8.18 g
 23 CO₂ m⁻² d⁻¹, respectively) from the I-U peatland (median 2.03 and 6.27 g CO₂ m⁻² d⁻¹,

respectively) but did not significantly differ [$H = 282.520$, d.f. = 7, *post hoc* test (TT), $p < 0.05$] between peatlands (Figure 6b).

One year post-harvesting (2007), both peatlands functioned as a net carbon sink from the atmosphere at the beginning of the season and showed a steady decline in carbon uptake (i.e. higher *NEE*) towards mid-season (Figure 5). *NEE* rates were lower (i.e. greater carbon uptake) and significantly different [H-U peatland, $\chi^2(3) = 91.660$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$; I-U peatland, $\chi^2(3) = 99.865$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] from *NEE* in the pre-harvesting years at both peatlands. The H-U peatland functioned as a consistent net carbon sink from the atmosphere (median $-1.34 \text{ g CO}_2\text{-C m}^{-2} \text{ d}^{-1}$) and significantly differed [$H = 285.517$, d.f. = 7, $p < 0.001$, *post hoc* test (TT) $p < 0.05$] from *NEE* at the I-U peatland, which was a consistent net carbon source (median $0.62 \text{ g CO}_2\text{-C m}^{-2} \text{ d}^{-1}$). R_{tot} at the H-U peatland was lower (median $5.83 \text{ g CO}_2\text{-C m}^{-2} \text{ d}^{-1}$) and significantly different [$\chi^2(3) = 118.619$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] from 2006, but did not significantly differ [*post hoc* test (TT), $p < 0.05$] from 2005 (Figure 6a). No significant change [$\chi^2(3) = 70.898$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] in R_{tot} (median $10.89 \text{ g CO}_2\text{-C m}^{-2} \text{ d}^{-1}$) occurred at the I-U peatland relative to 2005 or 2006. Consistent with trends of the pre-harvesting period, R_{tot} at the H-U peatland was lower and significantly different [$H = 282.365$, d.f. = 7, *post hoc* test (TT), $p < 0.05$] from the I-U peatland. *GEP* increased within both peatlands and significantly differed [H-U peatland, $\chi^2(3) = 100.033$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$; I-U peatland, $\chi^2(3) = 104.526$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] from 2005 and 2006 (Figure 6b). Similar to pre-harvesting, *GEP* at the H-U peatland remained slightly higher (median $10.63 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) but was not significantly different [$H = 282.520$, d.f. = 7, *post hoc* test (TT) $p < 0.05$] from the I-U peatland in 2007 (median $9.01 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$).

Two years post-harvesting (2008), both peatlands showed a steady decline in net carbon uptake from the beginning to mid-season (Figure 5). *NEE* at the H-U peatland still remained lower (median $0.18 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) and significantly different [H-U peatland, $\chi^2(3) = 91.660$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] from *NEE* in 2005, however, was no longer significantly different [*post hoc* test (TT) $p < 0.05$] from 2006. In contrast, *NEE* at the I-U peatland was higher (median $2.67 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) and significantly different [$\chi^2(3) = 99.865$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] from 2005 and 2006. *NEE* at the I-U peatland was significantly different [$H = 285.517$, d.f. = 7, *post hoc* test (TT), $p < 0.05$] from the H-U peatland in 2008, showing the greatest net carbon release from either site observed over the four-year study period. R_{tot} in both peatlands significantly differed [H-U peatland, $\chi^2(3) = 118.619$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$; I-U peatland, $\chi^2(3) = 70.898$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] from the pre-harvesting years (2005 and 2006; Figure 6a). Although the highest R_{tot} was observed in both peatlands during 2008, R_{tot} at the H-U peatland ($11.92 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) still remained lower and significantly different [$H = 282.365$, d.f. = 7, *post hoc* test (TT), $p < 0.05$] from the I-U peatland ($15.98 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$). *GEP* in both peatlands increased and were significantly different [H-U peatland, $\chi^2(3) = 100.033$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$; I-U peatland, $\chi^2(3) = 104.526$, $p < 0.001$, *post hoc* test (TT), $p < 0.05$] from 2005 and 2006 (Figure 6b). *GEP* was significantly higher [$H = 282.520$, d.f. = 7, *post hoc* test (TT), $p < 0.05$] at the H-U peatland relative to the I-U peatland (median *GEP* = 11.69 and 6.73 $\text{g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$, respectively; Figure 6b).

Despite large degree of uncertainty in the peatland *NEE* models, the scatter is typical of CO_2 flux models reported in other studies (e.g. Bubier *et al.*, 1998; Lafleur, 1999; Petrone *et al.*, 2011; Strack *et al.*, 2014). Mean standard errors for the *NEE* models ranged from ± 0.73 to $1.6 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$. Although the uncertainty for the models (i.e. maximum and minimum estimates)

overlapped during the undisturbed period, the overall discrepancy in *NEE* between peatlands remained large after harvesting; therefore the interpretation of shifts in CO_2 exchange at the H-U peatland relative to the I-U peatland due to disturbance remains valid.

Hydroclimatic influence on peatland carbon exchange

NEE at the H-U peatland correlated well with changes in *VMC* during the post-harvesting period only (Figure 7a). Greater net carbon uptake (i.e. lower *NEE*) with increasing peatland soil moisture occurred in 2007 and 2008. In contrast, poor relationships between peatland *NEE* and *VMC* were observed during pre-harvesting (2005 or 2006). Minimal variability in *VMC* at the I-U peatland within growing seasons and across the study period resulted in no relationship between peatland *NEE* and *VMC* at the I-U site and as such, was not shown in Figure 7.

The distinct relationship between *NEE* and *VMC* at the H-U peatland in 2007 and 2008 reflected strong moisture linkages with *GEP* and R_{tot} (Figure 7b and 7c). For example, *GEP* in the H-U peatland increased with higher *VMC* (i.e. greater productivity with soil moisture) in 2006 to 2008 (Figure 7b). The strongest relationships between *GEP* and *VMC* occurred during the post-harvesting years (2007 and 2008). No correlation between *GEP* and *VMC* was observed in 2005. Similar to *GEP*, R_{tot} in the H-U peatland also correlated with *VMC*; however, the relationship only occurred during the post-harvesting years (2007 and 2008) and was seasonally dependent (Figure 7c). For example, during the early growing season of 2007 and 2008, R_{tot} showed a strong quadratic relationship with *VMC* peaking at 7.4 and 4.3 $\text{g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$, respectively, when the peatland soil was generally wet (Figure 2) and cool (Figure 3). By the end of the growing seasons, maximum R_{tot} peaked higher at 12.5 and 6.8 $\text{g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ in 2007 and 2008, respectively when the peatland soil was generally dry and relatively warmer.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

Natural variability in BP peatland carbon dynamics

Throughout the study period, the H-U peatland functioned mainly as a net carbon sink while the I-U peatland was frequently a small net source. *NEE* rates in both peatlands ranged from -1.51 to 2.12 g CO₂-C m⁻² d⁻¹ during the undisturbed pre-harvest period (2005 and 2006) and were comparable to previous reports of natural *NEE* rates in Boreal peatlands (e.g. Bellisario *et al.*, 1998; Shurpali *et al.*, 1995; Humphreys *et al.*, 2006).

Differences in *NEE* between the H-U and I-U peatlands during the undisturbed period indicates the potential for large natural variability in carbon cycling of peatlands on the Boreal Plains (BP), even peatlands in relatively close proximity to each other as observed in this study (i.e. < 1 Km distance). Higher inter-annual *R*_{tot} and frequently larger seasonal variability in *R*_{tot} at the I-U peatland relative to the H-U peatland likely reflected the consistently drier (i.e. lower water table and soil moisture) and warmer soils at that site. Greater CO₂ release has been observed under lower water table positions in both laboratory settings (Moore and Dalva, 1993; Van de Reit *et al.*, 2013) and in situ studies (Silvola *et al.*, 1996; Kim and Verma, 1992; Gažovič *et al.*, 2013; Helfter *et al.*, 2015). Further, changes in peat temperature can alter decomposition rates and CO₂ emissions (Waddington *et al.*, 2001) whereby even minor soil temperature increases in high quality soils can provide optimal conditions for decomposition, thus increasing respiration (Solondz *et al.*, 2008).

Soil moisture is also an important factor influencing peatland *Sphagnum* productivity (McNeil and Waddington. 2003), as was reflected by the slightly higher *GEP* at the wetter H-U peatland that contributed to the overall lower *NEE* (i.e. greater net carbon uptake) during the undistributed

1 period. However, despite *WT* and soil *VMC* being low and disconnected at the I-U peatland,
2 maintenance of a sufficient water supply by dew or precipitation as well as moisture retention by
3 the moss (Strack and Price, 2009) may have moderated the impact of low soil water availability
4 on moss carbon uptake. This will maintain a relatively productive moss cover at the I-U peatland
5 (i.e. median *GEP* at the I-U peatland was 54% to 76% of median *GEP* at H-U peatland).
6 Together, differences in *NEE* linked to natural variations in site hydrology suggest the potential
7 for spatially variable seasonal and inter-annual carbon exchange of peatlands on the BP,
8 consistent with previous studies reporting soil moisture controls on CO₂ balances in peatlands
9 (Lafleur *et al.*, 2001; Strack *et al.*, 2009; Trudeau *et al.*, 2014). Therefore, assessing potential
10 impacts of land use disturbances on *NEE* of BP peatlands requires careful consideration of site-
11 specific natural variability in water availability influencing respiration and productivity across
12 this eco-region.

13 *Impact of forest harvesting on peatland hydroclimatic and carbon exchange*

14 Shifts in peatland microclimate at the H-U site post-harvest may have reflected the loss of a
15 protective sheltering effect created by the adjacent upland forest (Chen *et al.*, 1993). Post-
16 harvesting, higher vapour pressure deficit (*VPD*) measured in the newly exposed H-U peatland
17 may have resulted from decreased stability of the surface boundary layer and increased mixing
18 with the upper atmosphere due to an increase in fetch within the newly formed adjacent cutblock.
19 Dynamic turbulent flow patterns occur across landscape transitions (e.g. Markfort *et al.*, 2014;
20 Yang *et al.*, 2006; Flesch and Wilson, 1999). Although significant increases in wind speed were
21 not observed in the H-U peatland after harvesting, wind speeds are expected to be more variable
22 and turbulence increase with an increased momentum sink over the peatland relative to the clear-
23 cut forested area (Helgason and Pomeroy, 2005; Helgason and Pomeroy, 2012; Turnispeed *et al.*,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2003), likely contributing towards higher peatland *VPD* and *ET* (Petrone *et al.*, 2007). Even slight shifts in *ET* can alter the water balance of a peatland in this water deficit eco-region, given that *ET* is the dominant component of peatland water budgets on the BP (Devito *et al.*, 2005; Ferone and Devito, 2004; Smerdon *et al.*, 2005). As such, the greater evaporative water loss in the H-U peatland post-harvesting may have contributed to the seasonal decline of peatland *VMC* at that site, despite a relatively consistent supply of precipitation and overall wetter conditions in 2007 and 2008.

Post-harvesting, *VMC* became a limiting control on *NEE* at the H-U peatland, as indicated by the strong relationship with peatland *NEE* occurring in 2007 and 2008 only. Moisture conditions are frequently well correlated with both carbon losses (plant respiration and soil decomposition) and carbon fixation (plant production) (Davidson *et al.*, 2000; Bubier *et al.*, 2003). Although the overall range of soil *VMC* was similar in each year of the four-year study period (~0.40 to 0.70 m³ m⁻³), the timing of moisture loss (i.e. consistent seasonal decline) appeared critical to influencing seasonal patterns of respiration and productivity during the post-harvesting period. For example, low *VMC* near the end of the 2007 and 2008 growing seasons corresponded with the warmest soil temperatures and thus, contributed to higher *R_{tot}* (i.e. greater carbon release to the atmosphere) relative to the wetter cool soils at the beginning of the season. In contrast, frequent re-wetting by precipitation events during the pre-harvesting period resulted in no seasonal trend between *VMC*, *T_{soil}* and *R_{tot}* in 2005 and 2006.

Strong correlations between increasing *GEP* and increasing *VMC* were observed at the H-U peatland from 2006 to 2008. The lack of a relationship between *GEP* and *VMC* in 2005 likely reflected the predominantly high moisture conditions that year. The strongest relationship between *GEP* and *VMC*, as well as higher *GEP* for a given *VMC* (i.e. more efficient carbon

uptake), occurred during the post-harvesting years, likely reflecting the coinciding timing of high soil *VMC* and peak *PAR* during that period. For example, the highest *GEP* occurred during the beginning and mid-season, corresponding to the highest *VMC* and near maximum *PAR*, thus contributed toward the large net carbon accumulation. In contrast, lower *GEP* during pre-harvesting may reflect that the highest *VMC* occurred later in the growing season when *PAR* was generally lower. Consistent with our results, Griffis *et al.*, (2000) found that large carbon accumulation in a subarctic fen was the result of wetter conditions and early snowmelt during the warm spring period, even when drier conditions persisted for the majority of the growing season.

Implications for land use management on the Boreal Plains

BP peatlands exist in a moisture deficit region, and are in a state of hydrologic risk. Results of this study indicate the importance of soil moisture influencing peatland productivity and respiration. Thus, any land-use disturbances impacting peatland water availability, such as changes to peatland microclimate observed in this study, are likely to have a direct influence on the CO_2 exchange of BP peatlands. Large inter-annual variability in CO_2 exchange is common in northern peatlands (e.g. Aurela *et al.*, 2009), including shifts between a net CO_2 sink and source due to natural variations in hydrological and microclimatic conditions (Joiner *et al.*, 1999; Shurpali *et al.*, 1995; Lund *et al.*, 2012). In this study, shifts in mid-season *NEE* one-year post-harvesting were greater than the natural *NEE* variability of the pre-harvesting period at both peatlands. However, a larger relative change in *NEE* (i.e. greater net carbon uptake) occurred at the H-U site after forest removal. This suggests clear-cut logging may modify adjacent peatland microclimates and soil moisture conditions to influence *GEP* and R_{tot} in the short-term. However, given that mid-season *NEE* returned to within the range of the pre-harvest period two years post-harvesting suggests the relatively small-scale clear cutblock in this study may not sufficiently

1
2
3 1 alter *NEE* outside that of natural variation due to climate and/or site hydrology and/or
4
5 2 microclimatic conditions. As such, the long-term carbon exchange function of BP peatlands may
6
7 3 largely be influenced by changes in water availability resulting from drier conditions expected by
8
9 4 future climate warming (IPCC, 2014). However, careful consideration of larger-scale logging
10
11 5 due to rapid expansion of deforestation across this region (Timoney, 2003) may compound
12
13 6 anticipated drought conditions induced by climate change. In particular, consideration of forest
14
15 7 cutblock size (e.g. Flesch and Wilson, 1999) as well as orientation of cutblocks relative to the
16
17 8 dominant wind direction may enhance alterations to adjacent peatland hydroclimatic conditions
18
19 9 and CO₂ exchange dynamics and thus, impact stability of BP peatland water supply and carbon
20
21 10 stores.
22
23
24
25
26
27

28 11 **Conclusion**
29
30

31 12 Peatland growing season ecosystem CO₂ exchange data suggest the potential for large variability
32
33 13 in carbon cycling of undisturbed peatlands on the BP linked to natural hydrologic differences
34
35 14 between sites (i.e. higher R_{tot} and *NEE* in the naturally drier peatland). The changes to peatland
36
37 15 moisture soil conditions, linked to alterations in microclimate (i.e. increased turbulent mixing,
38
39 16 vapour pressure deficit and evapotranspiration) by adjacent upland clear-cut logging, shifted
40
41 17 peatland respiration and productivity patterns and thus, demonstrate that utilizing an integrated
42
43 18 hydrometeorological approach is fundamental to ecosystem monitoring as well as designing
44
45 19 landscape management and forestry strategies for the protection of BP peatland ecosystem
46
47 20 function. The results of short-term alterations to peatland hydroclimatic and carbon exchange
48
49 21 dynamics by clear-cut logging indicates that in addition to climate change, the sustainability of
50
51 22 BP peatlands ecosystem function may also depend on periodic forest disturbances expected with
52
53 23 the rapid expansion of deforestation, while the particular peatland response is likely to be site-
54
55
56
57
58
59
60

specific due to natural variability in terrestrial-atmosphere exchange of water and CO₂ within these hydrologically tenuous ecosystems in this moisture deficit region.

Acknowledgements

The authors wish to acknowledge R.P. Van Haarlem, S.M. Brown and S. Solondz for their technical assistance in the field. Funding was provided by the Sustainable Forest Management Network, Natural Science and Engineering Research Council, Alberta Pacific Forest Industries, TOLKO Forest Products, Ducks Unlimited Canada, and Syncrude Canada Ltd.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

Amiro, B. D., Barr, A. G., Black, T. A., Iwashita, H., Kljun, N., McCaughey, J. H., Morgenstern, K., Murayma, S., Nesic, Z., Orchansky, A.L., and Saigusa, N. 2006. Carbon, energy and water fluxes at mature and disturbed forest sites, Saskatchewan, Canada. *Agricultural and forest meteorology* **136**(3): 237-251.

ASCE-EWRI. 2005. "The ASCE Standardized Reference Evapotranspiration Equation. Report of the Task Committee on Standardization of Reference Evapotranspiration". Environmental and Water Resources Institute of the American Society of Civil Engineers, p. 4.

Aurela, M., Lohila, A., Tuovinen, J. P., Hatakka, J., Riutta, T., and Laurila, T. 2009. Carbon dioxide exchange on a northern boreal fen. *Boreal Environment Research* **14**(4): 699-710.

Banaszuk, P. and Kamocki, A. 2008. Effects of climatic fluctuations and landuse changes on the hydrology of temperate fluviogenous mire. *Ecological Engineering* **32**(2): 133-146.

Bellisario, L. M., Morre, T. R., and Bubier, J. L. 1998. Net ecosystem CO₂ exchange in a boreal peatland, northern Manitoba. *Ecoscience* :534-541.

Brown, S. M., R. M. Petrone, L. Chasmer, C. Mendoza, M. S. Lazerjan, S. M. Landhäusser, U. Silins, J. Leach, and K. J. Devito. 2013. Atmospheric and soil moisture controls on evapotranspiration from above and within a Western Boreal Plain aspen forest. *Hydrological Processes* **28**(15) 4449-4462.

Brown, S. M., Petrone, R. M., Mendoza, C., and Devito, K. J. 2010. Surface vegetation controls on evapotranspiration from a sub-humid Western Boreal Plain wetland. *Hydrological Processes*

- 1 **24**(8): 1072-1085.
- 2 Bubier, J. L., Crill, P. M., Moore, T. R., Savage, K., and Varner, R. K. 1998. Seasonal patterns
3 and controls on net ecosystem CO₂ exchange in a boreal peatland complex. *Global*
4 *Biogeochemical Cycles* **12**(4): 703-714.
- 5 Bubier, J. L., Bhatia, G., Moore, T. R., Roulet, N. T., and Lafleur, P. M. 2003. Spatial and
6 temporal variability in growing-season net ecosystem carbon dioxide exchange at a large
7 peatland in Ontario, Canada. *Ecosystems*, **6**(4): 353-367.
- 8 Carrera-Hernández, J. J., Mendoza, C. A., Devito, K. J., Petrone, R. M., and Smerdon, B. D.
9 2011. Effects of aspen harvesting on groundwater recharge and water table dynamics in a
10 subhumid climate. *Water Resources Research* **47**(5).
- 11 Chasmer, L., Petrone, R., Brown, S., Hopkinson, C., Mendoza, C., Diiwu, J., Quinton, W., and
12 Devito, K. 2010. Sensitivity of modelled actual evapotranspiration to canopy characteristics
13 within the Western Boreal Plain, Alberta. Remote Sensing and Hydrology 2010 (Proceedings of
14 a symposium held at Jackson Hole, 1 Wyoming, USA, September 2010) (IAHS Publ., 352:337-
15 340, 2011).
- 16 Chen, J., Franklin, J. F., and Spies, T. A. 1993. Contrasting microclimates among clearcut, edge,
17 and interior of old-growth Douglas-fir forest. *Agricultural and forest meteorology* **63**(3): 219-
18 237.
- 19 Davidson, E. A., Verchot, L. V., Cattanio, J. H. 2000. Effects of soil water content on soil
20 respiration in forest and cattle pastures of eastern Amazonia. *Biogeochemistry* **48**: 53–79.

- 1
2
3 1 Devito, K. J., Creed, I. F., and Fraser, C. J. D. 2005. Controls on runoff from partially harvested
4
5 2 aspen- forested headwater catchment, boreal plain, Canada. *Hydrological Processes* **19**(1): 3–25.
6
7
8
9 3 Environment Canada. 2007. Canadian Climate Normals, Meteorological service of Canada.
10
11
12 4 Ecological Stratification Working Group. 1996. A National Ecological Framework for Canada.
13
14 5 Agriculture and Agri-Food Canada, Research Branch, Centre for Land and Biological Resource
15
16 6 Research and Environment Canada, State of the Environment Directorate, Ecozone Analysis
17
18 7 Branch, Ottawa/Hull. 125 pp. Map at 1: 7,500,000.
19
20
21
22
23 8 Ferone, J., and Devito, K. J. 2004. Shallow groundwater–surface water interactions in pond–
24
25 9 peatland complexes along a Boreal Plains topographic gradient. *Journal of Hydrology* **292**: 75-
26
27 10 95.
28
29
30
31 11 Flesch, T. K., and Wilson, J. D. 1999. Wind and remnant tree sway in forest cutblocks.: I.
32
33 12 Measured winds in experimental cutblocks. *Agricultural and forest meteorology* **93**(4): 229-242.
34
35
36
37 13 Fournier, V., Larocque, M. and Pellerin, S. 2007. Water budget of the Covey Hill Peatland.
38
39 14 Ottawa 2007, The Diamond Jubilee, Breaking Ground in the Nation's Capital.
40
41
42
43 15 Gažovič, M., Forbrich, I., Jager, D. F., Kutzbach, L., Wille, C., and Wilmking, M. 2013.
44
45 16 Hydrology-driven ecosystem respiration determines the carbon balance of a boreal peatland.
46
47 17 *Science of the Total Environment* **463**: 675-682.
48
49
50
51 18 Gorham E. 1991. Northern peatlands: role in the carbon cycle and probable responses to climatic
52
53 19 warming. *Ecological Applications* **1**(2): 182 – 195.
54
55
56
57 20 Griffis, T. J., Rouse, W. R. and Waddington, J. M. 2000. Interannual variability of net
58
59
60

- ecosystem CO₂ exchange at a subarctic fen. *Global Biogeochemical Cycles* **14**(4): 1109–1121.
- Gullet D. W., Skinner, W. R., 1992. The state of Canada's climate: temperature change in Canada, 1895–1991. Environment Canada, Ottawa, Ontario. State of the Environment Report, #92-2.
- Halliwel, D. H., and Rouse, W. R. 1987. Soil heat flux in permafrost: characteristics and accuracy of measurement. *Journal of Climatology* **7**: 571–584.
- Helfter, C., Campbell, C., Dinsmore, K. J., Drewer, J., Coyle, M., Anderson, M., Skiba, U., Nemitz, E., Billett, M. F., and Sutton, M. A. 2015. Drivers of long-term variability in CO₂ net ecosystem exchange in a temperate peatland. *Biogeosciences* **12**: 1799–1811, doi:10.5194/bg-12-1799-2015.
- Helgason, W.D., and Pomeroy, J.W. 2005. Uncertainties in estimating turbulent fluxes to melting snow in a Mountain Clearing. *62nd Eastern Snow Conference*, Waterloo, Ontario, Canada.
- Helgason, W.D., and Pomeroy, J.W. 2012. Characteristics of the near-surface boundary layer within a mountain valley during winter. *Journal of Applied Meteorology and Climatology* **51**: 583–597.
- Humphreys, E. R., Lafleur, P. M., Flanagan, L. B., Hedstrom, N., Syed, K. H., Glenn, A. J., and Granger, R. 2006. Summer carbon dioxide and water vapor fluxes across a range of northern peatlands. *Journal of Geophysical Research: Biogeosciences* (2005–2012), 111(G4).
- IPCC (Inter Governmental Panel on Climate Change) 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the

- 1 Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A.
2 Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
- 3 Johnson, E. A. and Miyanishi, K. 2008. Creating new landscapes and ecosystems: The Alberta
4 oil sands. *Annals of the New York Academy of Sciences*. 1134, 120-145. (All Projects –
5 Regional/Global Synthesis).
- 6 Joiner, D. W., Lafleur, P. M., McCaughey, J. H., and Bartlett, P. A. 1999. Interannual variability
7 in carbon dioxide exchanges at a boreal wetland in the BOREAS northern study area. *Journal of*
8 *Geophysical Research: Atmospheres* (1984–2012), **104**(D22), 27663-27672.
- 9 Kim, J., and Verma, S. B. 1992. Soil surface CO₂ flux in a Minnesota peatland. *Biogeochemistry*
10 **18**(1): 37-51.
- 11 Kuhry, P., Nicholson, B. J., Gignac, L. D., Vitt, D. H., and Bayley, S. E. 1993. Development of
12 Sphagnum dominated peatlands in boreal continental Canada. *Canadian Journal of Botany*:
13 **71**(1), 10–22.
- 14 Lafleur, P. M., Roulet, N. T., and Admiral, S. W. 2001. Annual cycle of CO₂ exchange at a bog
15 peatland. *Journal of Geophysical Research: Atmospheres* (1984–2012), **106**(D3): 3071-3081.
- 16 Lafleur PM. 1999. Growing season energy and CO₂ exchange at a subarctic boreal woodland.
17 *Journal of Geophysical Research* **104**(8): 9571 – 9580.
- 18 Lafleur, P. M., McCaughey, J. H., Joiner, D. W., Bartlett, P. A., and Jelinski, D. E. 1997.
19 Seasonal trends in energy, water, and carbon dioxide fluxes at a northern boreal wetland. *Journal*
20 *of Geophysical Research-Atmospheres* **102**(D24): 29009–29020.

- 1 Lund, M., Christensen, T. R., Lindroth, A., and Schubert, P. 2012. Effects of drought conditions
2 on the carbon dioxide dynamics in a temperate peatland. *Environmental Research Letters* 7(4):
3 045704.
- 4 Markfort, C. D., Porté-Agel, F., and Stefan, H. G. 2014. Canopy-wake dynamics and wind
5 sheltering effects on Earth surface fluxes. *Environmental Fluid Mechanics* 14(3): 663-697.
- 6 McNeil, P., and Waddington, J. M. 2003. Moisture controls on Sphagnum growth and CO₂
7 exchange on a cutover bog. *Journal of Applied Ecology* 40(2): 354-367.
- 8 Monteith, J. L. 1965. Evaporation and environment. In Symp. Soc. Exp. Biol. Vol. 19, No. 205-
9 23, p. 4.
- 10 Moore, T. R., and Dalva, M. 1993. The influence of temperature and water table position on
11 carbon dioxide and methane emissions from laboratory columns of peatland soils. *Journal of Soil*
12 *Science* 44(4): 651-664.
- 13 Natural Regions Committee. 2006. Natural Regions and Subregions of Alberta. Compiled by
14 D.J. Downing and W.W. Pettapiece. Government of Alberta. Pub. No. T/852.
- 15 Oke, T.R. 1987. Boundary Layer Climates, 2nd edn. University Press: Cambridge.
- 16 Paulen, R. C., Pawlowicz, J. G., and Fenton, M. M. 2004. Surficial geology of the Utikuma Area
17 (NTS 83O/NW), Map 312, 1: 100,000. Alberta Geologic Survey, Alberta Energy and Utilities
18 Board, Edmonton, AB, Canada
- 19 Petrone, R. M., Chasmer, L., Hopkinson, C., Silins, U., Landhäusser, S. M., Kljun, N., and
20 Devito, K. J. 2015. Effects of harvesting and drought on CO₂ and H₂O fluxes in an aspen-

1 dominated western boreal plain forest: early chronosequence recovery. *Canadian Journal of*
2
3 1 Forest Research **45**(1): 87-100.
4
5 2
6
7
8
9 3 Petrone, R. M., Solondz, D. S., Macrae, M. L., Gignac, D., and Devito, K. J. 2011.
10
11 4 Microtopographical and canopy cover controls on moss carbon dioxide exchange in a western
12
13 5 Boreal Plain peatland. *Ecohydrology* **4**(1): 115-129.
14
15
16
17 6 Petrone, R.M., Devito, K.J., Silins, U., Mendoza, C., Brown, S.C., Kaufman, S.C. and Price, J.S.,
18
19 7 2008. Transient peat properties in two pond-peatland complexes in the sub-humid Western
20
21 8 Boreal Plain, Canada. *Mires Peat*, **3**(5), pp.1-13.
22
23
24
25 9 Petrone, R. M., Silins, U., and Devito, K.J., 2007. Dynamics of evapotranspiration from a
26
27 10 riparian pond complex in the Western Boreal Forest, Alberta. *Hydrological Processes* **21**: 1391–
28
29 11 1401.
30
31
32
33 12 Petrone, R. M., and Rouse, W. R., 2000. Synoptic controls on the surface energy and water
34
35 13 budgets in subarctic regions of Canada. *International Journal of Climatology* **20**: 1149–1165.
36
37
38
39 14 Redding, T. E. and Devito, K. J. 2008. Lateral flow thresholds for aspen forested hillslopes on
40
41 15 the Western Boreal Plain, Alberta, Canada. *Hydrological Processes* **22**: 4287-4300.
42
43
44
45 16 Shurpali, N. J., Verma, S. B., Kim, J., and Arkebauer, T. J. 1995. Carbon dioxide exchange in a
46
47 17 peatland ecosystem. *Journal of Geophysical Research: Atmospheres* (1984–2012), **100**(D7):
48
49 18 14319-14326.
50
51
52
53
54
55
56
57
58
59
60

- 1 Silvola, J., Alm, J., Ahlholm, U., Nykanen, H., and Martikainen, P. J. 1996. CO₂ fluxes from
2 peat in boreal mires under varying temperature and moisture conditions. *Journal of Ecology*
3 :219-228.
- 4 Smerdon, B., Devito, K., and Mendoza, C. 2005. Interaction of groundwater and shallow lakes
5 on outwash sediments in the sub-humid Boreal Plains of Canada. *Journal of Hydrology* **314**:
6 246-62.
- 7 Solondz, D. S., Petrone, R. M., and Devito, K. J. 2008. Forest floor carbon dioxide fluxes within
8 an upland□peatland complex in the Western Boreal Plain, Canada. *Ecohydrology* **1**(4): 361-376.
- 9 Strack, M., Keith, A. M., and Xu, B. 2014. Growing season carbon dioxide and methane
10 exchange at a restored peatland on the Western Boreal Plain. *Ecological Engineering* **64**: 231-
11 239.
- 12 Strack, M., Waddington, J. M., Lucchese, M. C., and Cagampan, J. P. 2009. Moisture controls on
13 CO₂ exchange in a Sphagnum□dominated peatland: results from an extreme drought field
14 experiment. *Ecohydrology* **2**(4): 454-461.
- 15 Strack, M., and Price, J. S. 2009. Moisture controls on carbon dioxide dynamics of peat□
16 Sphagnum monoliths. *Ecohydrology* **2**(1): 34-41.
- 17 Sutherland, G., Chasmer, L. E., Petrone, R. M., Kljun, N., and Devito, K. J. 2014. Evaluating the
18 use of spatially varying versus bulk average 3D vegetation structural inputs to modelled
19 evapotranspiration within heterogeneous land cover types. *Ecohydrology*, **7**(6), 1545-1559.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Temesgen, B., Eching, S., Davidoff, B. Z. and Frame, K. 2005. Comparison of some reference
2 evapotranspiration equations for California. *J. Irrig. Drain. Engng* **131**(1): 73–84.

3 Timoney, K. P. 2003. The changing disturbance regime of the boreal forest of the Canadian
4 Prairie Provinces. *The Forestry Chronicle* **79**(3): 502-516.

5 Trudeau, N. C., Garneau, M., and Pelletier, L. 2014. Interannual variability in the CO₂ balance of
6 a boreal patterned fen, James Bay, Canada. *Biogeochemistry* **118**(1-3): 371-387.

7 Tuittila, E. S., Komulainen, V. M., Vasander, H., and Laine, J. 1999. Restored cut-away peatland
8 as a sink for atmospheric CO₂. *Oecologia* **120**(4): 563-574.

9 Turnispeed, A.A., Anderson, D.E., Blanken, P.D., Baugh, W.M., and Monson, R.K. 2003.
10 Airflows and turbulent flux measurements in mountainous terrain Part 1. Canopy and local
11 effects. *Agricultural and Forest Meteorology* **119**: 1-21.

12 Van de Riet, B. P., Hefting, M. M., and Verhoeven, J. T. A. 2013. Rewetting drained peat
13 meadows: risks and benefits in terms of nutrient release and greenhouse gas exchange. *Water,*
14 *Air, & Soil Pollution* **224**(4): 1-12.

15 Waddington, J. M., Morris, P. J., Kettridge, N., Granath, G., Thompson, D. K., and Moore, P. A.
16 2015. Hydrological feedbacks in northern peatlands. *Ecohydrology* **8**(1): 113-127.

17 Waddington, J. M., Rotenberg, P. A., and Warren, F. J. 2001. Peat CO₂ production in a natural
18 and cutover peatland: implications for restoration. *Biogeochemistry* **54**(2): 115-130.

19 Waddington, J. M., and J. S. Price. 2000. Effect of peatland drainage, harvesting, and restoration
20 on atmospheric water and carbon exchange. *Physical Geography* **21**(5): 433-451.

- 1 Waddington, J. M., and Roulet, N. T. 1996. Atmosphere-wetland carbon exchanges: Scale
2 dependency of CO₂ and CH₄ exchange on the developmental topography of a peatland. *Global*
3 *Biogeochemical Cycles* **10**(2): 233–245.
- 4 Welles, J. M., Demetriades-Shah, T. H., and McDermitt, D. K. 2001. Considerations for
5 measuring ground CO₂ effluxes with chambers. *Chemical Geology* **177**: 3–13.
- 6 Wharton, S., Paw, K.T., Schroeder, M., Bible, K., and Falk, M. 2010. Turbulence considerations
7 for comparing ecosystem exchange over old-growth and clear-cut stands with limited fetch and
8 complex canopy flows. *29th Conference on Agricultural and Forest Meteorology*, Keystone,
9 Colorado, USA, August 1- 6, 2010.
- 10 Whiting, G. J. 1994. Seasonal CO₂ exchange in communities of the Hudson Bay lowlands.
11 *Journal of Geophysical Research* **99**(D1): 1519-1528.
- 12 Whitson, I. R., Chanasyk, D. S., and Prepas, E. E. 2005. Effect of forest harvest on soil
13 temperature and water storage and movement patterns on Boreal Plain hillslopes. *Journal of*
14 *Environmental Engineering and Science* **4**(6): 429-439.
- 15 Yang, B., Raupach, M. R., Shaw, R. H., and Morse, A. P. 2006. Large-eddy simulation of
16 turbulent flow across a forest edge. Part I: flow statistics. *Boundary-Layer Meteorology* **120**(3):
17 377-412.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Table 1: Parameters for *NEE* models at the I-U and H-U Peatland

<i>GEP</i> parameters				<i>R</i> _{tot} parameters		
Site	GP _{max} (g CO ₂ m ⁻² d ⁻¹)	<i>Q</i>	<i>R</i> ²	<i>a</i>	<i>b</i>	<i>R</i> ²
I-U Peatland						
2005	4.36	0.15	0.22	0.47	4.59	0.25
2006	13.19	0.45	0.54	0.32	7.87	0.10
2007	24.79	0.37	0.79	0.95	-0.74	0.64
2008	33.93	0.10	0.55	1.08	1.33	0.74
H-U Peatland						
2005	20.55	0.06	0.34	0.57	0.01	0.55
2006	18.46	0.48	0.49	0.60	1.05	0.52
2007	24.52	0.66	0.71	0.38	1.22	0.59
2008	31.92	0.39	0.54	0.86	1.28	0.83

2

Figure 1. Harvested-upland (H-U) and intact-upland (I-U) study complexes, located on the Boreal Plains Ecozone region of Northern Alberta. Shown are the ponds, surrounding peatlands and the adjacent uplands with locations of the meteorological stations (*MET*) and groundwater wells. Circles represent site locations of the point field measurements.

Figure 2. Water table (*WT*) position (dotted lines) from the peatland groundwater wells (see Figure 1) and soil moisture content (*VMC*) of the peatland (solid lines) measured at the meteorological stations (see Figure 1) and total daily precipitation (vertical bars) at the harvested-upland (H-U) and intact-upland (I-U) complexes in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada.

Figure 3. Average daily air temperature (T_a) and soil temperature (T_{soil}) (averaged 2, 5 and 10 cm depths) measured at the meteorological stations (see Figure 1) at the harvested-upland (H-U) and intact-upland (I-U) complexes in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada.

Figure 4. Mid-season (DOY 201 to 243) average daily (a) vapor pressure deficit (*VPD*), (c) wind speed and horizontal turbulence intensity (I_u), and (d) evapotranspiration (*ET*) measured at the meteorological stations (see Figure 1) at the harvested-upland (H-U) and intact-upland (I-U) complexes in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada. N/A indicates data was not available.

1
2
3
4 1
5
6
7 2 Figure 5. Daily total net ecosystem CO₂ exchange (*NEE*) (error bars 95% confidence interval) in
8
9 3 the harvested-upland (H-U) peatland and intact-upland (I-U) peatland in 2005, 2006, 2007 and
10
11 4 2008, Utikuma Region Study Area, Alberta, Canada. Dotted lines represent the minimum and
12
13 5 maximum uncertainty estimates for the *NEE* models.
14
15
16
17 6

18
19
20
21 7 Figure 6. Daily average (a) total respiration (*R*_{tot}) and (b) gross ecosystem productivity (*GEP*) in
22
23 8 the harvested-upland (H-U) peatland and intact-upland (I-U) peatland in 2005, 2006, 2007 and
24
25 9 2008, Utikuma Region Study Area, Alberta, Canada. For comparative purposed, seasonal *R*_{tot}
26
27 10 were grouped into four different time periods (e.g. Solondz et al., 2008): early green (EG: DOY
28
29 11 120 to 160), green (G: DOY 161 to 218), late green (LG: DOY 219 to 250) and senescence (S:
30
31 12 DOY 251 to 278). N/A indicates data was not available.
32
33
34
35
36 13

37
38
39 14 Figure 7. Variations in (a) total net ecosystem exchange (*NEE*) with soil moisture content
40
41 15 (*VMC*), (b) variation in gross ecosystem production (*GEP*) with *VMC*, and (c) variation in total
42
43 16 respiration (*R*_{tot}) with *VMC* in the harvested-upland (H-U) peatland in 2005, 2006, 2007 and
44
45 17 2008, Utikuma Region Study Area, Alberta, Canada. Daily *NEE*, *GEP* and *R*_{tot} rates
46
47 18 corresponding to daily *VMC* values were binned to improve data clarity, using an average value
48
49 19 for each sample within 0.01 m³ m⁻³ interval. Seasonal *R*_{tot} were grouped into two different time
50
51 20 periods: early season (DOY 140 to 179) and late season (DOY 180 to 278).
52
53
54
55
56
57
58
59
60

Figure 1

Figure 1. Harvested-upland (H-U) and intact-upland (I-U) study complexes, located on the Boreal Plains Ecozone region of Northern Alberta. Shown are the ponds, surrounding peatlands and the adjacent uplands with locations of the meteorological stations (MET) and groundwater wells. Circles represent site locations of the point field measurements.

190x142mm (300 x 300 DPI)

Figure 2

Figure 2. Water table (WT) position (dotted lines) from the peatland groundwater wells (see Figure 1) and soil moisture content (VMC) of the peatland (solid lines) measured at the meteorological stations (see Figure 1) and total daily precipitation (vertical bars) at the harvested-upland (H-U) and intact-upland (I-U) complexes in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada.
190x142mm (300 x 300 DPI)

Figure 3

Figure 3. Average daily air temperature (T_a) and soil temperature (T_{soil}) (averaged 2, 5 and 10 cm depths) measured at the meteorological stations (see Figure 1) at the harvested-upland (H-U) and intact-upland (I-U) complexes in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada.
190x142mm (300 x 300 DPI)

Figure 4

Figure 4. Mid-season (DOY 201 to 243) average daily (a) vapor pressure deficit (VPD), (c) wind speed and horizontal turbulence intensity (I_u), and (d) evapotranspiration (ET) measured at the meteorological stations (see Figure 1) at the harvested-upland (H-U) and intact-upland (I-U) complexes in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada. N/A indicates data was not available.
190x142mm (300 x 300 DPI)

Figure 5

Figure 5. Daily total net ecosystem CO₂ exchange (NEE) (error bars 95% confidence interval) in the harvested-upland (H-U) peatland and intact-upland (I-U) peatland in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada. Dotted lines represent the minimum and maximum uncertainty estimates for the NEE models.
190x254mm (300 x 300 DPI)

Figure 6

Figure 6. Daily average (a) total respiration (R_{tot}) and (b) gross ecosystem productivity (GEP) in the harvested-upland (H-U) peatland and intact-upland (I-U) peatland in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada. For comparative purposes, seasonal R_{tot} were grouped into four different time periods (e.g. Solondz et al., 2008): early green (EG: DOY 120 to 160), green (G: DOY 161 to 218), late green (LG: DOY 219 to 250) and senescence (S: DOY 251 to 278). N/A indicates data was not available.

190x142mm (300 x 300 DPI)

Figure 7

Figure 7. Variations in (a) total net ecosystem exchange (NEE) with soil moisture content (VMC), (b) variation in gross ecosystem production (GEP) with VMC, and (c) variation in total respiration (R_{tot}) with VMC in the harvested-upland (H-U) peatland in 2005, 2006, 2007 and 2008, Utikuma Region Study Area, Alberta, Canada. Daily NEE, GEP and R_{tot} rates corresponding to daily VMC values were binned to improve data clarity, using an average value for each sample within $0.01 \text{ m}^3 \text{m}^{-3}$ interval. Seasonal R_{tot} were grouped into two different time periods: early season (DOY 140 to 179) and late season (DOY 180 to 278). 190x142mm (300 x 300 DPI)