A two-stage, two-organism process for biohydrogen from glucose

Redwood, MD; Macaskie, Lynne

DOI:

[10.1016/j.ijhydene.2006.06.018](https://doi.org/10.1016/j.ijhydene.2006.06.018)

Citation for published version (Harvard):

Redwood, MD & Macaskie, L 2006, 'A two-stage, two-organism process for biohydrogen from glucose', *International Journal of Hydrogen Energy*, vol. 31, pp. 1514-1521. <https://doi.org/10.1016/j.ijhydene.2006.06.018>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

A two-stage, two-organism process for biohydrogen from glucose

Mark D. Redwood, Lynne E. Macaskie*

School of Biosciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

* Corresponding author. Tel.: +44 1214145889; fax: +44 1214145925.

E-mail address: L.E.Macaskie@bham.ac.uk (L.E. Macaskie).

Abstract

H₂ can potentially be produced in a two-stage biological process: the fermentation of glucose by *Escherichia coli* HD701 and the photofermentation of the residual medium by *Rhodobacter sphaeroides* O.U. 001. In a typical batch fermentation, *E. coli* consumed glucose and produced H₂, organic end-products and biomass. Organic end-products and residual glucose were removed during subsequent photofermentation by *R. sphaeroides*, with associated growth and neutralization of pH. However, photoproduction of H₂ did not occur during photofermentation of the residual liquor per se due to the presence of fixed nitrogen compounds. Nevertheless, this two-stage approach could be applied to dispose of sugar-containing industrial wastes, H₂ being used for on-site power generation.

Keywords: Biohydrogen; *Rhodobacter sphaeroides*; *Escherichia coli*; Organic acids; Cross-feeding

1. Introduction

Biohydrogen is anticipated to play an important role in the future hydrogen economy, as it can be produced from readily available renewable substrates. Sugars are promising substrates for biological H₂ production, being readily and renewably available and potentially giving a high yield of H₂.

The stoichiometric yield of 12 mol H₂ per mol hexose represents the ultimate target for biohydrogen and it was suggested previously that a yield of 8 mol/mol is sufficient for economic application [3], although the economic calculation does not take into account the escalating cost of waste disposal via landfilling. No single organism is capable of performing the conversion with this efficiency. In fact the theoretical maximum yield for fermentation is 4 mol/mol as illustrated in Eq. (2) [2]. It is predicted that yields can be significantly improved by integrating fermentation and photoheterotrophy in a two stage, two-organism system.

Eqs. (2) and (3) describe an ideal situation in which all carbon substrate is processed along the appropriate pathways and none is diverted to the formation of biomass or alternative metabolites. Practical fermentation yields tend to be around 2 mol H₂ per mol hexose while yields approaching 4 mol/mol have been achieved using thermophilic fermentation [5]. Reported photoheterotrophic yields exceed 80% [4].

Fig. 1. (Overall scheme) In reactor 1, *E. coli* HD701 (hydrogen overproducing strain) ferments glucose to produce H₂. Organic end-products (e.g. lactate, acetate, and ethanol) are cross-fed to reactor 2, where they are utilised by *R. sphaeroides* in photoproduction of H₂. CO₂ and biomass are formed as by-products.

Fermentation at moderate temperatures (mesophilic fermentation) is a prevalent approach, permitting biohydrogen production at a high rate and on a large scale using substrates found in organic wastes (e.g. from the food industry). This method has been extensively studied using *Clostridia* or extant microbial consortia from the feedstock [6–10]. Although formation of H₂ during mixed-acid fermentation by *Escherichia coli* is well documented historically [11], efforts to develop the H₂-producing capacity of *E. coli* have only recently achieved prominence. Unlike many other fermentative bacteria the Enterobacteria produce hydrogen by the action of the formate hydrogen lyase (FHL) system which cleaves formate to H₂ and CO₂ under pH stress [12]:

E. coli represents a convenient vector for metabolic engineering and significantly increased rates of H₂ production have been achieved through derepression of the FHL system [13–15]. The H₂ yield from mesophilic fermentation is thermodynamically limited to 4 mol H₂ per mol hexose [2], while experimental yields tend to be in the region of 2 mol/mol [16–18].

The accumulation of fermentation end-products (e.g. ethanol and organic acids) and fall in pH can exert sufficient ‘stress’ to halt mesophilic H₂ production in an excess of substrate. Furthermore, the presence of organic acids in the residual medium presents a disposal issue. Utilization of fermentation end-products (for further H₂ production) in a second stage would increase the economic potential of an H₂-producing process by improvement of the H₂ yield and reduction of the organic content of the final waste.

Anoxygenic photosynthetic bacteria (e.g. *Rhodobacter* spp.) can assimilate organic acids and produce H₂ under photoheterotrophic conditions (light, anaerobiosis, high C/N ratio). Experimental yields for H₂ production

from organic acids by *Rhodobacter* spp. can approach 100% [4,19]. H₂ is produced as a by-product of the nitrogenase system, which functions in the fixation of N₂ as ammonia, to provide a growth advantage under nitrogen limitation. Due to the high metabolic cost of maintaining active nitrogenase, its synthesis and activity are very tightly regulated with respect to the nitrogen status of the cell. The reversible inhibition of nitrogenase activity ('nitrogenase switch-off') occurs in response to micromolar concentrations of ammonium ion. However, nitrogenase is not regulated in response to the availability of its substrate (N₂), and H₂ production is normally performed under an argon atmosphere in order to eliminate product inhibition. In the absence of N₂, ammonia is not formed but H₂ production continues, as shown in Eqs. (5) and (6) [20,21]:

In the presence of N₂:

In the absence of N₂ (e.g. under argon):

It may be practical to integrate mesophilic fermentation and photosynthesis in order to produce hydrogen from sugars with high efficiency. The use of co-cultures was reviewed in [19]. This approach is limited by the difficulty of maintaining optimum conditions for both modes of metabolism in a single reactor. It is considered preferable to operate a two-stage system: a separate fermenter and photobioreactor with cross-feeding of fermentation end-products (Fig. 1).

Kataoka et al. [16] predicted that a two-stage system for co-operative hydrogen production by *Clostridium butyricum* and photosynthetic bacteria could achieve an overall yield of 5.6 mol H₂ per mol glucose. Kim et al. [17] performed a two-stage reaction with *C. butyricum* and *R. sphaeroides* and achieved an overall conversion of 1.64 mol/mol. The photosynthetic step operated at approximately 7% substrate conversion efficiency and contributed only 22% of the H₂ yield. Yokoi et al. [18] reported an overall yield of 6.6 mol/mol using a twostage system in which the supernatant from a *C. butyricum*/*Enterobacter enterogenes* co-culture was fed to *R. sphaeroides* M-19.

The present study represents the first instance of a two-stage system combining *E. coli* and *R. sphaeroides* in H₂ production and remediation of the primary fermentation effluent. The chosen *E. coli* strain is the metabolically engineered, H₂-overproducing HD701 [13] and the chosen *R. sphaeroides* strain is O.U.001 which has been studied extensively as a H₂-producer [22–27].

2. Experimental

2.1. Micro-organisms and culture conditions

The H₂-overproducing strain *E. coli* HD701 was kindly provided by Professor A. Böck (Lehrstuhl für Mikrobiologie, Munich, Germany) and cultured aerobically on nutrient broth (Oxoid) (30 °C, 200 rpm).

Rhodobacter sphaeroides O.U.001 (DSMZ 5864) was held in stock at –80 °C (in 15% glycerol v/v), revived on nutrient agar (30 °C) and cultured anaerobically under fluorescent illumination (39.5 uE/m²/s measured with a PAR light meter SKP200, Skye Instruments Ltd.) at 30 °C using the SyA medium described by Hoekema et al. [28].

2.2. Determination of biomass concentration

The optical density (*E. coli* OD600, *R. sphaeroides* OD660) was measured using an UltraspecIII variable wavelength spectrophotometer (1 cm light path). OD values were used to calculate the biomass concentration (g dry biomass/l) with reference to a dried standard, prepared in triplicate by recording optical densities from dense cultures after various dilutions in deionised water. Cultures were washed twice by centrifugation and resuspension (2400 × g, 20 min, 4 °C, 50 ml deionised H₂O) before drying at 60 °C and weighing to constant mass.

2.3. *E. coli* glucose fermentation

E. coli glucose fermentation was performed in a 6 l vessel (Electrolab, UK). The initial reactor contents were 2.5 l *E. coli* culture in nutrient broth (Oxoid), 2.5 l phosphate-buffered saline (1.43 g Na₂HPO₄, 0.2gKH₂PO₄, 1.0 g NaCl, 0.2 g KCl per l) and 0.55 l 1M glucose. The initial culture pH was 6.80. Anaerobiosis was established by sparging with argon (1 h), after which off-gas was collected over a solution of 1M NaOH. The temperature was maintained at 30.0 °C and the culture was stirred continuously (300 rpm).

2.4. Preparation of liquor

Fermentation proceeded for 24 h, after which the residual medium was treated by centrifugation (4435 × g, 20°C, 10 min), and ultrafiltration of the supernatant (Acrodisc 32 mm; 0.2 µm supor membrane). The liquor was diluted 1 in 2 with sterile deionised water before inoculating with *R. sphaeroides*.

2.5. Photofermentation of *R. sphaeroides* on liquor

R. sphaeroides was harvested from the late exponential phase of growth (OD660 = 1.04) by centrifugation (4435 × g, 20°C, 10 min). A cell concentrate containing 4 g dry biomass/l was prepared by resuspending the pellet in a sterile solution of 50 mg/l MgSO₄ · 7H₂O and 25 mg/l CaCl₂ with micronutrients as described by Hoekema et al. [28]. This suspension (5 ml) was inoculated

into 195 ml pre-degassed liquor to give an initial biomass concentration of 0.2 g/l. The culture was sealed using butyl rubber stoppers with aluminium crimp seals and sparged with argon (20 min) to remove N₂ and to establish anaerobiosis. Light was provided by two fluorescent 10W lamps (Eterna). The intensity of photosynthetically active radiation (measured as 2.1) was 10 $\mu\text{E}/\text{m}^2/\text{s}$ (47W/m²). Reactors were stirred magnetically. The temperature was maintained at 30 °C using a water bath.

2.6. Measurement of hydrogen production

Exit gas from *R. sphaeroides* reactors was directed through a bubbler, which contained a solution of 200 g/l NaCl (pH 2) and then into a graduated glass tube containing 1M NaOH. The bubbler functioned to maintain anaerobiosis and the NaOH served to remove CO₂ from the exit gas so its displacement from the tube would indicate the volume of H₂ produced.

2.7. Chemical analyses

5ml samples were withdrawn at intervals from 200 ml *R. sphaeroides* cultures. OD₆₆₀ and pH were measured immediately before cells were separated by centrifugation (13000 $\times$ g, 20°C, 4 min). Supernatants were stored at -20 °C before analysis. Organic acids were measured by anion HPLC ([Table 1](#)); glucose was assayed using the colorimetric dinitrosalicylic acid assay [29]; ammonia was determined using the colorimetric Nessler assay [30]; protein was measured using the colorimetric bicinchoninic acid assay (Sigma procedure TPR0562) and ethanol was determined colorimetrically by monitoring the enzymatic reduction of NAD using alcohol dehydrogenase (Sigma A-6338, assay concentration: 2.64 U/ml) after pre-removal of aldehyde by aldehyde dehydrogenase (Sigma A-7011, assay concentration: 16.18 U/l).

Table 1. Conditions for anion chromatography

Columns	IonPac AS11-HC Analytical (4mm) IonPac AG11-HC Analytical (4mm)				
Eluent	A Deionised water (18.2 MΩ-cm) B 5 mM NaOH C 100 mM NaOH D 250 mM NaOH				
Gradient (%)	Time (min)	A	B	C	D
	-2	60	40	0	0
	0 (inj.)	60	40	0	0
	1	88	12	0	0
	10	88	12	0	0
	20	80	0	20	0
	23	20	0	0	80
	24	20	0	0	80
	24.01	60	40	0	0
	27	60	40	0	0
Flow rate	15 ml/min				
Inj. volume	15 µl				
Detection	Suppressed conductivity, ASRS Autosuppression recycle mode				

3. Results and discussion

3.1. First stage fermentation by *E. coli* and second stage photofermentation

The initial fermentation using *E. coli* (see materials and methods) was described in detail previously [31]. In this 24 h batch fermentation, H₂ was produced at a rate of 52.0 ml/l/h and at a molar yield of 0.376. Using an improved methodology (pH controlled at 5.5, fed-batch, washed cells pre-grown in the presence of 5 g/l sodium formate) the H₂ yield was improved to 2.4 mol/mol (M.D. Redwood, D.W. Penfold and L.E. Macaskie unpublished). Liquor from the 24 h batch fermentation was used to test the remediation-potential of *R. sphaeroides* via photofermentation. The cell-free liquor was inoculated with *R. sphaeroides* and monitored for growth (Fig. 2A), neutralization of pH (Fig. 2B), removal of organic components (Fig. 3) and H₂ production.

Fig. 2. Growth of *R. sphaeroides* on *E. coli* spent fermentation liquor and neutralization of pH: (A) Growth of *R. sphaeroides* on liquor from *E. coli* fermentation (Table 2) adjusted to different pH values prior to inoculation and (B) pH profile during growth at different initial pH values.

Fig. 3. Removal of organic compounds from fermentation liquor by *R. sphaeroides*. Data represent means from at least three replicate experiments. Bars represent standard error of the mean.

3.2. Cultivation of *R. sphaeroides* on *E. coli* fermentation liquor

Fermentation by *E. coli* resulted in acidification of the medium from pH 7.0–4.5 due to the formation of organic acids (Table 2). As shown in Fig. 2 spent liquor at pH 4.5 did not permit growth of *R. sphaeroides*. However, when the pH was adjusted to 5.5, growth and further neutralization of pH occurred rapidly (growth rate: 0.20 g/l/day). The highest growth rate (0.41 g/l/day) was observed after the pH was adjusted to 6.5. These observations were in agreement with previous reports [25].

While the required pre-adjustment of pH was only 1 pH unit, this would introduce an additional cost in largescale cultivation of photoheterotrophic bacteria. Further work would be required to achieve acid-tolerant remediation of *E. coli* fermentation liquor.

3.3. Removal of organic components from *E. coli* fermentation liquor

The composition of typical *E. coli* spent fermentation liquor is shown in Table 2. The low turbidity, high organic content and high nitrogenous content make this a suitable substrate for cultivation of *R. sphaeroides*. As shown in Fig. 3 the primary carbon sources for growth were glucose, acetate and lactate, which were removed at average rates of 0.46, 0.30, and 0.17 mmol/day/reactor, respectively. Although succinate is usually a preferred carbon source for growth of purple non-sulphur bacteria [32], it was not consumed in these experiments, possibly due to its low initial concentration. The consumption of ethanol was secondary to the other substrates, commencing only after ~100 h. By combining the removal rates for glucose, acetate and lactate a total removal rate of 3.87 mmol C/day/reactor was calculated.

These results show that the two-stage system described here could be applied for remediation of industrial

Table 2: Properties of fermentation liquor.

Fermentation properties		General properties	
Glucose	40 mM	pH	4.5
Ethanol	20 mM	Turbidity	Low
Total organic acids	38 mM	Ammonia	4.4 mM
Acetate	20 mM	Protein	2.30 g/l
Lactate	15 mM	Chloride	46 mM
Succinate	3 mM	Phosphate	5.2 mM

^a Liquor was diluted 1 in 2 before inoculating with *R. sphaeroides*.

sugary wastes. Suitable organic feedstocks for *E. coli* can be obtained at low cost, also helping to forestall landfill as a disposal issue [13,31]. Out-flow from the photobioreactor was recycled back into the primary *E. coli* fermenter showing that metabolic products of *R. sphaeroides* are not inhibitory to further H₂ production by *E. coli* since a second burst of H₂ evolution was observed (M.D. Redwood, D.W. Penfold and L.E. Macaskie, unpublished).

3.4. Potential for H₂ production from E. coli fermentation liquor

Theoretically, the organic acids present in fermentation liquor could be converted to H₂ to yield 0.19 mol (4.83 l) H₂ per litre fermentation waste (excluding that from ethanol and glucose). Although *R. sphaeroides* assimilated organic acids from the fermentation liquor (Fig. 3), H₂ formation was not observed. This was attributed to the presence of fixed nitrogen sources (4.4mM ammonium ion, 2.3 g/l protein: Table 2). A positive control was performed using a synthetic medium lacking nitrogenous components, but containing glucose, ethanol and organic acids at concentrations identical to those in the *E. coli* spent liquor. H₂ was produced at a rate of 124 ml H₂/l culture/day (10.5 ml/h/g dry biomass).

Repressive concentrations of ammonium ion were present in the liquor (together with significant protein: Table 2) and the rate of removal (38 $\mu\text{mol NH}_4^+$ /l/day) was insufficient to overcome repression of nitrogenase activity. Nitrogenase fixes N₂ gas as ammonia to overcome nitrogen-limitation, forming H₂ as a by-product. This metabolically demanding process is repressed in the presence of fixed nitrogen (see earlier). The repressibility of nitrogenase is a common bottleneck in photobiological H₂ production, which may be overcome by the use of various strategies, for example:

- Denitrification of the fermentation liquor could be achieved by alkalisation followed by boiling. This method was used to prepare food processing wastewater and sewage sludge for use as fermentation feed [17]. Alternatively, classical deionisation techniques could potentially remove NH₄⁺ with some selectivity, but either approach would add significant process costs.
- A two-part photobioreactor permitting ammonia consumption for growth in the first reactor followed by H₂ production in a nitrogen-limited second reactor [33]. Wild-type bacterial strains could be employed in this approach, but process costs would be increased as compared to those for a single reactor arrangement.
- The use of nitrogenase-derepressed mutants which exhibit nitrogenase expression in the presence of

fixed nitrogen [34,35]. Although industry is often reluctant to accept engineered strains, this approach could offer the efficient photoproduction of H₂ from a nitrogenous feed in a single stage.

- A membrane separation system could be used to separate the organic acids. Preliminary tests using this approach have suggested that this permits H₂-evolution in the *R. sphaeroides* reactor receiving organic acids from the *E. coli* primary fermentation (M.D. Redwood and L.E. Macaskie, unpublished). Elucidation of mass balances and consideration of process economics are in progress.

In conclusion, there is significant potential for H₂ production from fermentation liquor. It should be noted that the *R. sphaeroides* photofermentation was not performed under optimum conditions at this stage. This was suggested by a relatively low rate of H₂ production observed for the positive control. This may be attributed to the use of fluorescent light, whereas tungsten/halogen lamps are preferable due to the emission of wavelengths in the range of 800 and 850 nm, corresponding to the absorption maxima of bacteriochlorophylls [4]. The use of an improved light source (tungsten; 250 $\mu\text{m}^2/\text{s}$) could permit higher rates of substrate removal than those reported here, but the high nitrogen content would still prevent photoproduction of H₂.

4. Conclusions

R. sphaeroides was able to grow on fermentation liquor (after pH adjustment), simultaneously removing organic components and neutralizing the pH. Nitrogenase activity (H₂ production) was inhibited by the high nitrogen content of the liquor but was shown in nitrogen-free medium. This data highlights that inhibition of nitrogenase is the foremost challenge stalling the development of a two-stage system for biohydrogen, but numerous strategies have the potential to overcome this. Future work will aim to achieve photoproduction of H₂ from fermentation liquor by the use of membraneseparation technology.

Acknowledgements

This work was supported by a BBSRC studentship (to MDR: no.10703) and by a BBSRC/Royal Society Industry Fellowship to LEM. We thank Prof A. Livingston (Imperial College London) for useful discussions on membrane separations, Dr P. Yong for help with analysis and Dr D.W. Penfold for provision of the *E. coli* fermentation liquor.

References

- [1] Macler BA, Pelroy RA, Bassham JA. Hydrogen formation in nearly stoichiometric amounts from glucose by a *Rhodospseudomonas* mutant. J Bacteriol 1979;138(2):446–52.
- [2] Thauer R. Limitation of microbial H₂-formation via

- fermentation. In: Schlegel HG, Barnea J, editors. Microbial energy conversion. Oxford: Pergamon; 1977. p. 201–4.
- [3] Keasling JD, Benemann JR, Pramanik J, Carrier TA, Jones KL, Van Dien SJ. A toolkit for metabolic engineering of bacteria: application to hydrogen production. In: Zaborski OR, Benemann JR, Matsunaga T, Miyake J, San Pietro A, editors. Biohydrogen. New York: Plenum Press; 1997. p. 87.
- [4] Rocha JS, Barbosa MJ, Wijffels RH. Hydrogen production by photoheterotrophic bacteria: culture media. In: Miyake J, Matsunaga T, San Pietro A, editors. Biohydrogen II: an approach to environmentally acceptable technology. Oxford: Pergamon; 2001. p. 3–32.
- [5] van Niel EWJ, Budde MAW, de Haas GG, van der Wal FJ, Claassen PAM, Stams AJM. Distinctive properties of high hydrogen producing extreme thermophiles, *Caldicellulosiruptor saccharolyticus* and *Thermotoga elfii*. Int J Hydrogen Energy 2002;27(11–12):1390–8.
- [6] Hussy I, Hawkes FR, Dinsdale R, Hawkes DL. Continuous fermentative hydrogen production from sucrose and sugarbeet. Int J Hydrogen Energy 2005;30:471–83.
- [7] Nandi R, Sengupta S. Microbial production of hydrogen: an overview. Crit Rev Microbiol 1998;24(1):61–84.
- [8] Collet C, Adler N, Schwitzguebel J-P, Peringer P. Hydrogen production by *Clostridium thermolactum* during continuous fermentation of lactose. Int J Hydrogen Energy 2004;29: 1479–85.
- [9] Yokoi H, Tokushige T, Hirose J, Hayashi S, Takasaki Y. H₂ production from starch by a mixed culture of *Clostridium butyricum* and *Enterobacter aerogenes*. Biotechnol Lett 1998;20(2):143–7.
- [10] Kim S-H, Han S-K, Shin H-S. Feasibility of biohydrogen production by anaerobic co-digestion of food waste and sewage sludge. Int J Hydrogen Energy 2004;29:1607–16.
- [11] Blackwood AC, Neish AC, Ledingham GA. Dissimilation of glucose at controlled pH values by pigmented and nonpigmented strains of *Escherichia coli*. J Bacteriol 1956;72: 497–9.
- [12] Sauter M, Bohm R, Bock A. Mutational analysis of the operon (*hyc*) determining hydrogenase-3 formation in *Escherichia coli*. Mol Microbiol 1992;6(11):1523–32.
- [13] Penfold DW, Forster CF, Macaskie LE. Increased hydrogen production by *Escherichia coli* strain HD701 in comparison with the wild-type parent strain MC4100. Enz Microb Tech 2003;33(2–3):185–9.
- [14] Sode K, Yamamoto S, Tomiyama M. Metabolic engineering approaches for the improvement of bacterial hydrogen production based on *Escherichia coli* mixed acid fermentation. In: Miyake J, Matsunaga T, San Pietro A, editors. Biohydrogen II: an approach to environmentally acceptable technology. Oxford: Pergamon; 1997. p. 195–204.
- [15] Yoshida A, Nishimura T, Kawaguchi H, Inui M, Yukuwa H. Enhanced hydrogen production from formic acid by formate hydrogen lyase-overexpressing *Escherichia coli* strains. Appl Environ Microbiol 2005;71(11):6762–8.
- [16] Kataoka N, Miya A, Kiriya K. Studies on hydrogen

- production by continuous culture system of hydrogen-producing anaerobic bacteria. *Water Sci Tech* 1997;36(6–7):41–7.
- [17] Kim M-S, Lee TJ, Yoon YS, Lee IG, Moon KW. Hydrogen production from food processing wastewater and sewage sludge by anaerobic dark fermentation combined with photofermentation. In: Miyake J, Matsunaga T, San Pietro A, editors. *Biohydrogen II: an approach to environmentally acceptable technology*. Oxford: Pergamon; 2001. p. 263–72.
- [18] Yokoi H, Saitsu A, Uchida H, Hirose J, Hayashi S, Takasaki Y. Microbial hydrogen production from sweet potato starch residue. *J Biosci Bioeng* 2001;91(1):58–63.
- [19] Sasikala K, Ramana CV, Rao PR, Kovacs KL. Anoxygenic phototrophic bacteria: Physiology and advances in hydrogen production technology. *Adv Appl Microbiol* 1993;38:211–95.
- [20] Akkerman I, Janssen M, Rocha J, Wijffels RH. Photobiological hydrogen production: photochemical efficiency and bioreactor design. *Int J Hydrogen Energy* 2002;27(11–12):1195–208.
- [21] Koku H, Eroglu I, Gunduz U, Yucel M, Turker L. Aspects of the metabolism of hydrogen production by *Rhodobacter sphaeroides*. *Int J Hydrogen Energy* 2002;27:1315–29.
- [22] Eroglu I, Aslan K, Gunduz U, Yucel M, Turker L. Substrate consumption rates for hydrogen production by *Rhodobacter sphaeroides* in a column photobioreactor. *J Biotechnol* 1999;70:103–13.
- [23] Hallenbeck PC. Integration of hydrogen evolving systems with cellular metabolism: the molecular biology and biochemistry and electron transport factors and associated reductases. In: Miyake J, Matsunaga T, San Pietro A, editors. *Biohydrogen II: an approach to environmentally acceptable technology*. Oxford: Pergamon; 1997. p. 171–81.
- [24] Koku H, Eroglu I, Gunduz U, Yucel M, Turker L. Kinetics of biological hydrogen production by the photosynthetic bacterium *Rhodobacter sphaeroides* O.U. 001. *Int J Hydrogen Energy* 2003;28:381–8.
- [25] Sasikala K, Ramana CV, Rao PR. Environmental regulation for optimal biomass yield and photoproduction of hydrogen by *Rhodobacter sphaeroides* O.U.001. *Int J Hydrogen Energy* 1991;16(9):597–601.
- [26] Turkarslan S, Yigit OD, Aslan K, Eroglu I, Gunduz U. Photobiological hydrogen production by *R. sphaeroides* O.U.001 by utilisation of waste water from milk industry. In: Zaborski OR, Benemann JR, Matsunaga T, Miyake J, San Pietro A, editors. *Biohydrogen*. New York: Plenum Press; 1997. p. 143.
- [27] Yetis M, Gunduz U, Eroglu I, Yucel M, Turker L. Photoproduction of hydrogen from sugar refinery wastewater by *Rhodobacter sphaeroides* O.U. 001. *Int J Hydrogen Energy* 2000;25:1035–41.
- [28] Hoekema S, Bijmans M, Janssen M, Tramper J, Wijffels RH. A pneumatically agitated flat-panel photobioreactor with gas re-circulation: anaerobic photoheterotrophic cultivation of a purple non-sulfur bacterium. *Int J Hydrogen Energy* 2002;27(11–12):1331–8.
- [29] Chaplin MF. Monosaccharides. In: Chaplin MF, Kennedy JF, editors. *Carbohydrate analysis: a practical approach*. Oxford:

IRL Press; 1986. p. 3.

[30] Mendam J, Denney RC, Barnes JD, Thomas M. Vogel's quantitative chemical analysis. 6th ed., Englewood Cliffs, NJ: Prentice Hall; 2000.

[31] Penfold DW. A dual anaerobic system for bioremediation of metal and organic waste, chapter VI.5. PhD thesis, University of Birmingham, UK, 2005.

[32] Biebl H, Pfennig N. Isolation of members of the family *Rhodospirillaceae*. In: Starr MP, Stolp H, Truper HG, Balows A, Schlegel HG, editors. The prokaryotes. New York: Springer; 1981. p. 267–73.

[33] Fascetti I, Todini O. *Rhodobacter sphaeroides* RV cultivation and hydrogen production in a one- and two-stage chemostat. Appl Microbiol Biotechnol 1995;44:300–5.

[34] Zinchenko VV, Kopteva AV, Belavina NV, Mitronova TN, Frolova VD, Shestakov SV. The study of *Rhodobacter sphaeroides* mutants of different type with derepressed nitrogenase. Genetika 1991;27(6):991–9.

[35] Zinchenko VV, Babykin M, Glaser V, Mekhedov S, Shestakov SV. Mutation in *ntnC* gene leading to the derepression of nitrogenase synthesis in *Rhodobacter sphaeroides*. FEMS Microbiol Lett 1997;147:57–61.