Social Anxiety in Cornelia de Lange Syndrome

Richards, Caroline; Moss, Joanna; O'Farrell, L; Kaur, G; Oliver, Christopher

DOI:

[10.1007/s10803-009-0730-7](https://doi.org/10.1007/s10803-009-0730-7)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Richards, C, Moss, J, O'Farrell, L, Kaur, G & Oliver, C 2009, 'Social Anxiety in Cornelia de Lange Syndrome', *Journal of Autism and Developmental Disorders*, vol. 39, no. 8, pp. 1155-1162. <https://doi.org/10.1007/s10803-009-0730-7>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10803-009-0730-7>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Social Anxiety in Cornelia de Lange Syndrome

Caroline Richards, Jo Moss, Laura O' Farrell,
Gurmeash Kaur and Chris Oliver

*Cerebra Centre for Neurodevelopmental Disorders,
School of Psychology,
University of Birmingham*

Please use this reference when citing this work:

Richards, C., Moss, J., O'Farrell, L., Kaur, G., & Oliver, C. (2009). Social Anxiety in Cornelia de Lange Syndrome. *Journal of Autism and Developmental Disorders*, 39, 1155-1162.

Abstract

In this study we assessed the behavioral presentation of social anxiety in Cornelia de Lange syndrome (CdLS) using a contrast group of Cri du Chat syndrome (CdCS). Behaviors indicative of social anxiety were recorded in twelve children with CdLS (mean age = 11.00; SD = 5.15) and twelve children with CdCS (8.20; SD = 2.86) during social interaction. Lag sequential analysis revealed that participants with CdLS were significantly more likely to evidence behavior indicative of anxiety in close temporal proximity to the point at which they maintained eye contact or spoke. Individuals with CdLS demonstrate a heightened probability of anxiety related behavior during social interaction but only at the point at which social demand is high.

Introduction

Cornelia de Lange syndrome (CdLS) is a rare genetic syndrome with an estimated prevalence of 1:50,000 births (Beck, 1976; Beck & Fenger, 1985) and is caused by a deletions on chromosomes 5, 10 and X (Gillis *et al.*, 2004; Krantz *et al.*, 2004; Tonkin, Wang, Lisgo, Bamshad & Strachan, 2004; Musio *et al.*, 2006; Deardorff *et al.*, 2007). The physical phenotype includes growth retardation, upper limb abnormalities, hypertrichosis and facial dysmorphism (Jackson, Kline, Barr & Koch, 1993). CdLS is associated with intellectual disability ranging from mild to profound, health problems including vision and hearing impairments and gastro-oesophageal reflux (Berney, Ireland & Burn, 1999; Jackson *et al.*, 1993; Collis *et al.*, 2008; Hall, Arron, Sloneem and Oliver, 2008; Oliver, Arron, Hall & Sloneem, 2008). Behavioral problems associated with CdLS include impulsivity, repetitive and self injurious behaviors (Berney *et al.*, 1999; Hyman, Oliver and Hall, 2002, Moss *et al.*, In press).

Recent research indicates a heightened prevalence of autism spectrum disorder (ASD), in CdLS with estimates ranging from 55 to 66% (Basile, Villa, Selicorni & Molteni, 2007; Bhyuian *et al.*, 2006; Moss *et al.*, 2008; Oliver, *et al.*, 2008; Moss *et al.*, In press). Fine-grained investigation has indicated that the presentation of the triad of impairments in CdLS may not be typical of that observed in idiopathic ASD. Specifically, social impairment in CdLS may be characterised by selective mutism, extreme shyness and social anxiety (Goodban, 1993; Collis, Oliver & Moss, 2006; Moss *et al.*, 2008). This presentation of social impairment appears similar to the social anxiety and shyness that is reported in Fragile X syndrome (Dykens & Volkmar, 1997). Arron *et al.* (2006) also described a high prevalence of socially avoidant behaviors such as ‘wriggling out of physical contact’ and ‘attempting to move away

during an interaction' in fourteen out of sixteen individuals with CdLS. Collectively these studies indicate that social anxiety and social avoidance may be evident in individuals with CdLS. Additionally social anxiety has been found to be prevalent in individuals with ASD (Simonoff *et al.* 2008). However, unlike in CdLS, selective mutism is a less common expression of social anxiety, and social anxiety itself is not a core feature or diagnostic symptom of ASD.

Social anxiety is difficult to assess in individuals with developmental disorders as introspection and self report may be compromised (Kim, Szatmari, Bryson, Streiner & Wilson, 2000; Gillott, Furniss & Walter, 2001). Consequently, observation of behavioral markers of social anxiety has been employed in empirical studies. Hall, DeBernadis & Reiss (2006) manipulated social and performance demands in order to examine social escape behaviours in 114 children with Fragile X syndrome. All children were exposed to four video taped conditions of varying social demand ranging from silent reading to an interview condition. In the interview condition, participants had to converse with an experimenter and particular emphasis was placed on the participant maintaining eye contact to increase social interaction. Social escape behaviours such as fidgeting and low levels of eye contact were coded, and were found to be most prevalent in the interview condition. The presence of these behaviours was also associated with increased levels of salivary cortisol suggesting that an interaction between biological and environmental factors may contribute to social escape behaviours in Fragile X syndrome.

Lesniak-Karpiak, Mazzocco and Ross (2003) used a similar methodology to assess social withdrawal and avoidance of social interaction in Turner syndrome and Fragile

X syndrome, compared to social interaction in typically developing individuals. They constructed eight behavioural markers of social anxiety which included eye contact avoidance, fidgeting and wringing of hands, and measured the occurrence of these in role play interactions. The role play interactions included conditions of conversing with familiar adults and unfamiliar adults. Again, individuals with Fragile X syndrome, engaged in more hand movements during social interaction than the group with Turner syndrome, or the typically developing control group. This methodology is promising and is used in this study to assess social anxiety in children with CdLS.

Contrasts between syndrome groups are useful for determining the specificity of behavioral phenotypes (Hodapp & Dykens, 2001). In this study a matched comparison group of children with Cri du Chat syndrome (CdCS) was employed. CdCS is caused by a deletion on the short arm of chromosome 5 (5p15; Goodhart *et al.*, 1994; Overhauser *et al.*, 1994) and occurs in approximately 1 in 50,000 births. Individuals with CdCS have similar levels of intellectual disability, expressive and receptive communication to those identified in CdLS (Cornish & Munir, 1988; Cornish, Bramble, Munir & Pigram, 1999). In contrast to individuals with CdLS, nonverbal communication skills and social interaction skills are reported to be a relative strength in CdCS (Cornish & Pigram, 1996; Cornish, Munir & Bramble, 1998; Sarimski, 2002) although, as is the case in CdLS, speech is compromised. These shared general and specific characteristics, makes CdCS an appropriate contrast group for this study. In this study we use fine grained observations of social interaction to investigate the behavioral indicators of social anxiety in children with CdLS and CdCS during periods of high social demand.

Method

Recruitment and Participants

Participants were recruited from a study of nerve function in CdLS and CdCS (Oliver *et al.*, 2007)¹. The twelve most able participants with CdLS from this group were selected based on Vineland Adaptive Behavior Scales composite standard scores (VABS; Sparrow, Balla & Cicchetti, 1994). Twelve participants with CdCS were selected based on the matching criteria of age (+/- 2 years), gender and VABS scores. A measure of cognitive functioning derived from scores on the British Picture Vocabulary Scale (BPVS; Dunn, Dunn, Whetton & Burley, 1997) was available for some participants. However, these data were missing for three participants with CdLS and one participant with CdCS. In addition, two further participants with CdLS and four further participants with CdCS scored at floor level on the BPVS. Consequently, meaningful data was only available for fourteen of the total twenty four participants. Therefore, as no other data were available on cognitive functioning, it was determined that VABS data would be more useful to match the groups. Table 1 shows participants characteristics. The CdLS group did not differ from the CdCS group on any variables.

(Insert table 1 about here)

Measures

Structured social interactions. Participant behaviour was coded from video recordings made during an experimental analogue situation (similar to those described by Carr and Durand, 1985). As previous literature had indicated that social anxiety behaviours were most prevalent during periods of conversation with researchers,

participants were observed during high attention conditions of the analogue situation (Hall *et al.*, 2006; Lesniak-Karpiak *et al.*, 2003). Two, ten minute high attention conditions were assessed, during which the researcher maintained high levels of verbal attention toward the participant, whilst sustaining physical proximity. No task demands were placed on the participant during this condition. This condition was deemed to present the participants with high levels of social expectation and social demand since this condition involved the examiner repeatedly directing conversation towards the participant providing frequent opportunities for the participant to engage in social interaction. Participant behaviors during these interactions were operationally defined and coded from video recorded observations. In order to confirm that individuals in each participant group were given a comparable opportunity to engage in social interaction during this condition, the percentage duration of examiner communication during the 10 minute high attention conditions was compared between participant groups. No significant differences in examiner communication were identified (CdLS mean=71.10, SD=9.05; CdCS mean=72.08, SD=9.85). Participant behaviors during these interactions were operationally defined and coded from video recorded observations.

Behavior Definitions and Video Coding

Definitions of behavior indicative of social anxiety were constructed from reviewing studies that used similar methods (Hall *et al.*, 2006; Lesniak-Karpiak *et al.*, 2003). Behavioral codes focused on verbal and eye contact, and hand movements, these had been identified as indicators of anxiety in previous studies, with levels of the behavior differing significantly between those experiencing social anxiety, and those not (Hall *et al.*, 2006; Lesniak-Karpiak *et al.*, 2003). According to Hall *et al.* (2006), the increased

salivary cortisol levels associated with fidgeting in individuals with Fragile X syndrome give a physiological indication that this type of non-functional, non-rhythmic motor movement behavior is related to anxiety rather than any other underlying variable. Consequently the behavior code for hand movement in this study identified only hand movement behaviour that was non-functional and did not include stereotypic or repetitive rhythmic behaviors.

Table 2 shows the behaviors and operational definitions employed. Behaviors were either coded as duration variables (eye contact, participant and examiner communication), where the behavior onset and offset was recorded, or as event variables, where only the behavior occurrence was recorded (moving hands). Using these behavioral codes, each session was coded in real time using Obswin software (Martin, Oliver & Hall, 2000).

(Insert table 2 about here)

In order to analyse the relationship between social demand and social anxiety behaviors, behaviors were analysed as criterion or target variables. A criterion variable indicates a behavior which may increase the experience of social demand or pressure. Therefore, participant communication is a criterion variable. When a participant is engaging in verbal communication, social demand to communicate effectively is high. Likewise, participant eye contact is also a criterion variable, as engaging in eye contact increases social demand to participate in a verbally communicative manner. However, eye contact is also analysed as a target variable as it is identified by Hall *et al.*, (2006) to indicate anxiety in social encounters. Target variables are behaviors which are

predicted to change when social demand is high, indicating anxiety. Consequently, in addition to eye contact, hand movement behaviour is also a target behavior.

Interrater Reliability

20% of the video footage, an equal amount from both participant groups, was independently coded by a second observer. Kappa coefficients for 3s time windows were calculated for all coded behaviors (see Table 2). The mean Kappa score for behavioral codes was .88 (range .80 – .96) which indicates a good level of interrater reliability.

Data Analysis

Lag sequential analyses were used to investigate temporal relationships between points of high social demand and social anxiety behaviors. The analysis considers the unconditional probability of the participant engaging in the target behavior (e.g. moving hands) against the conditional probability of the participant engaging in the target behavior *given* the presence of a criterion behavior (e.g. moving hands *given* eye contact). Lags were examined for six, 3s intervals prior to and after the presence of the target behavior. Lag zero indicates the display of the criterion behavior. The lag analyses were *restricted* to the next occurrence of the criterion behavior. Consequently any association with the target behavior indicative of social anxiety can be isolated to that display of the criterion behavior. A significant degree of difference between the unconditional and conditional probabilities was deemed evident by a z score of greater than 4.5 ($p < .00001$) to avoid type 1 errors (see Moss *et al.*, 2005).

Results

Analysis of behaviors indicative of social anxiety

Analyses were conducted to test for differences in the frequency and duration of social anxiety behaviors between the CdLS and CdCS groups. No significant differences were found between the groups for any behaviors (See Table 3 for results).

(Insert table 3 about here)

The effects of social interaction upon behaviors indicative of social anxiety

Lag sequential analyses investigated the temporal relationship between eye contact and engaging in verbal communication with the examiner (points of high social demand) and participant hand movements. The results are shown in Figure 1.

(Insert figure 1 about here)

The upper panels of Figure 1 display the unconditional probability of each group engaging in eye contact and the conditional probability of each group engaging in eye contact *given* that they are engaging in verbal communication. The graphs show that both groups (CdLS and CdCS) engage in eye contact immediately prior to speaking to the researcher, and continue to engage in eye contact whilst speaking, only ceasing the eye contact after they have finished speaking. However, for the participants with CdLS, significant increases in eye contact occur closer to communicating and cease earlier after communicating. This suggests that the participants with CdLS engage in eye contact as temporally close to communicating as possible, rather than before or after. This analysis also identifies this point as characterised by high social demand i.e. there is both eye contact and the participant speaks

The central panels of Figure 1 display the unconditional probability of each group moving their hands, plotted alongside the conditional probability of each group moving their hands *given* that they are engaging in eye contact. The graphs indicate that the participants with CdLS move their hands immediately prior to and after engaging in eye contact significantly more frequently than they would usually move their hands. They continue to move their hands whilst engaging in eye contact although the probability decreases slightly, suggesting that the most anxiety provoking time is just prior to and after the social engagement. This contrasts with the graph for the participants with CdCS, who do not show a pronounced relationship between hand moving and engaging in eye contact.

Finally, the lower panels of Figure 1 display the unconditional probability of each group moving their hands, plotted against the conditional probability of each group moving their hands *given* that they are engaging in verbal communication. The participants with CdLS display a consistent pattern of increasing hand movements prior to engaging in verbal communication with the examiner, this peaks during communication and decreases after the social interaction has finished. The participants with CdCS show a less pronounced pattern and display no difference between the unconditional and conditional probabilities.

Discussion

In this study we examined behavioral indicators of social anxiety in a group of participants with CdLS and a matched contrast group of participants with CdCS. The use of the contrast group with similar characteristics and, specifically, restricted speech

enabled the specificity of these behavioral indicators to be evaluated in CdLS. We employed observational assessments of behavioral indicators of social anxiety similar to those used by Hall *et al* (2006) and Lesniak-Karpiak *et al* (2003) in order to investigate the prevalence of these behaviors and their association with high social demand. No significant differences were identified between behavioral indicators of social anxiety displayed by individuals with CdLS compared to those with CdCS in terms of overall levels. However, more detailed investigation of the pattern of behavioral responses before and after points of high social demand revealed that individuals with CdLS were significantly more likely to display behaviors indicative of social anxiety, such as hand movements, in close temporal proximity to periods of social interaction. The findings also demonstrated that the use of eye contact in CdLS was far more fleeting during these points of high social demand in comparison to individuals with CdCS. Given that communicative ability is thought to be comparable between the two participant groups (Cornish & Munir, 1988; Cornish *et al.*, 1999) it is assumed that any differences in communication between the groups are not attributable simply to communication impairment, but rather reflect group differences in response to social demand.

The results of this study add to a growing body of literature describing the presence of social anxiety in CdLS and demonstrate a more refined methodology for evaluating behavioral indices of social anxiety. Previous research had demonstrated that during periods of social interaction, individuals with CdLS engaged in socially avoidant behaviors in order to terminate social interaction (Arron *et al.*, 2006). The present study has added to this research by further delineating social behavior in CdLS with two characteristic findings. Firstly, that there is no quantitative difference in the amount of behaviors indicative of social anxiety displayed by children with CdLS

compared to children with CdCS. This is in marked contrast to the profile of social anxiety in other genetic syndromes such as Fragile X (Lesniak-Karpiak *et al.*, 2003) where the prevalence of social anxiety behaviors are reported to be heightened in comparison to those with Turner syndrome and typically developing individuals. The comparable nature of social interaction in CdLS and CdCS is further highlighted by the initial lag analyses which demonstrate that the pattern or sequence of social interaction behaviors in CdLS appears to be very similar to that of individuals with CdCS. Both groups of children engage in eye contact prior to and during verbal communication with a researcher. However, this analysis also revealed that although the participants with CdLS can and do engage in eye contact whilst communicating, they only do so in close temporal proximity to their verbal communication. This may indicate that social interaction combining eye contact and verbal communication is anxiety provoking for individuals with CdLS, and therefore they avoid engaging in it until the last possible moment and as briefly as possible.

This initial difference between the two experimental groups becomes more apparent when the behavior of ‘moving hands’ was analysed. Previous literature in people with Fragile X indicated that this behavior (or behaviors similar to this; ‘fidgeting’, ‘wringing hands’) was indicative of social anxiety (Lesniak-Karpiak *et al.*, 2003; Hall *et al.*, 2006). The lag analyses revealed that in the CdLS group, hand movements indicative of social anxiety were more likely to occur when social demand was heightened, specifically when the individual was engaging in eye contact or verbal communication with the examiner. This pattern of behaviour was not observed in the CdCS group. Whilst the two groups displayed a comparable frequency and duration of this behavioral indicator of anxiety, the participants with CdLS displayed hand movement differentially during social interaction. This may suggest that individuals

with CdLS experience heightened levels of anxiety during periods of high social demand. The finding that children with CdLS display social anxiety through small, non verbal behaviours should not be surprising, given that due to deficits in expressive communication, individuals with CdLS tend to rely on nonverbal communicative acts to express themselves (Sarimski, 1997).

This study was limited by two factors. Firstly the sample size of the groups was relatively small, and the participants were young. Previous studies have indicated that social anxiety in CdLS may be more prevalent amongst young adults with CdLS (Collis *et al.*, 2006). It may be the case that the subtle indicators of social anxiety identified in the current study sample of younger individuals may become stronger with age and this should be investigated. Secondly, the presence of hand movements as indicative of anxiety has been assumed, in part based on the clustering of these behaviors around periods of high social demand. However, previous studies have used physiological assessment to validate the presence of hand moving behaviors as indicative of social anxiety (Hall *et al.*, 2006). This type of validation was not possible in this study but should be considered in future studies of social anxiety in CdLS.

References

- Arron, K., Oliver, C., Hall, S., Sloneem, J., Forman, D. & McClintock, K. (2006). Effects of social context on social interaction and self-injurious behavior in Cornelia de Lange syndrome. *American Journal on Mental Retardation*, 111, 184-192.
- Basile, E., Villa, L., Selicorni, A. & Molyeni, M (2007). The behavioural phenotype of Cornelia de Lange Syndrome: A study of 56 individuals. *Journal of Intellectual Disability Research*, 51, 671-681.
- Beck, B. (1976). Epidemiology of Cornelia de Lange's Syndrome. *Acta Paediatirca Scandinavia*, 65, 631-638.
- Beck, B. & Fenger, K. (1985). Mortality, pathological findings and causes of death in the de Lange Syndrome. *Acta Paediatrica Scandinavia*, 74, 765-769.
- Berney, T. P., Ireland, M. & Burn, J. (1999) Behavioural phenotype of Cornelia de Lange syndrome. *Archives of Disease of Childhood*, 81, 333-336.
- Bhyuian, Z.A., Klein, M., Hammond, P., van Haeringen, A., Mannens, M.A.M., Van Berckelaer-Onnes, I. & Hennekam, R.C.M. (2006). Genotype-phenotype correlations of 39 patients with CdLS: The Dutch experience. *Journal of Medical Genetics*, 46, 568-575.
- Carr, E. G. & Durand, V. M. (1985). Reducing behavior problems through functional communication training. *Journal of Applied Behavior Analysis*, 18, 111-126.
- Collis, L., Oliver, C. & Moss, J. (2006). Low mood and social anxiety in Cornelia de Lange Syndrome. *Journal of Intellectual Disability Research*, 50, 792.

- Collis, L., Moss, J., Jutley, J., Cornish, K. & Oliver, C. (2008). Facial expression in Cornelia de Lange and Cri du Chat syndromes. *Journal of Intellectual Disability Research*. **52**, 207-215.
- Cornish, K., Bramble, D., Munir, F. & Pigram, J. (1999). Cognitive functioning in children with typical cri du chat (5p-) syndrome. *Developmental Medicine & Child Neurology*, *41*, 263-266.
- Cornish, K. & Munir, F. (1998). Receptive and expressive language skills in children with Cri-du Chat syndrome. *Journal of Communication Disorders*, *31*, 73-81.
- Cornish, K., Munir, F., & Bramble, D. (1998). Adaptive and Maladaptive Behaviour in Children with Cri-du- Chat syndrome. *Journal of Applied Research in Intellectual Disabilities*, *11*, 239-246.
- Cornish, K. & Pigram, J. (1996). Developmental and behavioral characteristics of Cri du Chat syndrome. *Archives of Diseases in Childhood*, *75*, 448-450.
- Deardorff, M. A., Kaur, M., Yaeger, D., Rampuria, A., Korolev, S., Pie, J., Gil-Rodríguez, C., Arnedo, M., Loeys, B., Klime, A, D., Wilson, M., Lillquist, K., Siu, V., Ramos, F, J., Musio, A., Jackson, L, S., Dorsett, D. & Krantz, I, D. (2007). Mutations in cohesion complex members SMC3 and SMC1A cause a mild variant of Cornelia de Lange Syndrome with predominant mental retardation. *The American Journal of Human Genetics*, *80*, 475-494.
- Dunn, L.M., Dunn, L.M., Whetton, C., Burley, J. (1997). *British Picture Vocabulary Scales* (2nd Edition). Windsor: Nfer Nelson.

- Dykens, E. M. & Volkmar, F. R. (1997). Medical conditions associated with autism. In D.J.Cohen & F. R. Volkmar (Eds.), *Handbook of Autism and Pervasive Developmental Disorders* (2 ed., pp. 388-407). New York: John Wiley & Sons Inc.
- Gillis, L.A., McCallum, J., Kaur, M., DeScipio, C., Yaeger, D., Mariani, A., Kline, A.D., Li, H., Devoto, M., Jackson, L.G., Krantz, I.D. (2004). NIPBL mutational analysis in 120 individuals with Cornelia de Lange syndrome and evaluation of genotype-phenotype correlations. *American Journal of Human Genetics*, 75, 610-623.
- Gillott, A., Furniss, F. & Walter, A. (2001). Anxiety in high-functioning children with autism. *Autism*, 5, 277-286.
- Goodban, M. T. (1993). Survey of speech and language skills with prognostic indicators in 116 patients with CdLS. *American Journal of Medical Genetics*, 47, 1059-1063.
- Goodhart, S.A., Simmons, A.D., Grady, D., Rojas, K., Moysis, R.K., Lovett, M., & Overhauser, J. (1994). A yeast artificial chromosome contig of the critical region for Cri du Chat syndrome. *Genomics*, 24, 63-68.
- Hall, S., Arron, K., Sloneem, J. and Oliver, C. (2008). Health and sleep problems in Cornelia de Lange Syndrome: A case control study. *Journal of Intellectual Disability Research*, 52, 458-468.
- Hall, S., DeBernardis, M. & Reiss, A. (2006). Social escape behaviors in children with

- Fragile X syndrome. *Journal of Autism and Developmental Disorders*, 36, 935-947.
- Hodapp, R.M. & Dykens, E.M. (2001). Strengthening behavioral research on genetic mental retardation syndromes. *American Journal on Mental Retardation*, 106, 4-15.
- Hyman, P., Oliver, C., & Hall, S. (2002). Self-Injurious Behavior, Self-Restraint, and Compulsive Behaviors in CdLS. *American Journal on Mental Retardation*, 107, 146-154.
- Jackson, L., Kline, A. D., Barr, M. A. & Kock, S. (1993). de Lange Syndrome: A clinical review of 310 individuals. *American Journal of Medical Genetics*, 47, 940-946.
- Kim, J. A., Szatmari, P., Bryson, S. E., Streiner, D. L. & Wilson, F. J. (2000). The prevalence of anxiety and mood problems among children with autism and Asperger syndrome. *Autism*, 4, 117-132.
- Krantz, I. D., McCallum, J., DeScipio, C., Kaur, M., Gillis, L. A., Yaeger, D. *et al.* (2004). Cornelia de Lange syndrome is caused by mutations in NIBPL, the human homolog of *Drosophila melanogaster* Nipped-B. *Nature Genetics*, 6, 631-635.
- Lesniak-Karpiak, K., Mazzocco, M. M. M. & Ross, J. L. (2003). Behavioral assessment of social anxiety in females with Turner or Fragile X syndrome. *Journal of Autism and Developmental Disorders*, 33, 55-67.
- Martin N., Oliver C. & Hall S. (2000) *Obswin*, Version 3.0. University of Birmingham, Birmingham.

- Moss, J., Oliver, C., Arron, K., Burbidge, C. and Berg, K. (In press). The prevalence and phenomenology of repetitive behavior in genetic syndromes. *Journal of Autism and Developmental Disorders*. (10.1007/s10803-008-0655-6)
- Moss, J., Oliver, C., Hall, S., Arron, K., Sloneem, J. & Petty, J. (2005). The association between environmental events and self-injurious behaviour in Cornelia de Lange syndrome. *Journal of Intellectual Disability Research*, 49, 269-277.
- Moss, J., Oliver, C., Jephcott, L., Berg, K., Kaur, G. & Cornish, K. (2008). Prevalence of autistic spectrum phenomenology in Cornelia de Lange and Cri du Chat syndromes. *American Journal on Mental Retardation*, 113, 278-291.
- Musio, A., Selicorni, A., Focarelli, M., L., Gervasini, C., Milani, D., Russo, S., Vezzoni, P. & Larizza, L. (2006). X-linked Cornelia de Lange syndrome owing to SMC1L1 mutations. *Nature Genetics*, 38, 528-530.
- Oliver, C., Arron, K., Hall, S. and Sloneem, J. (2008). The behavioural phenotype of Cornelia de Lange syndrome. *British Journal of Psychiatry*, **193**, 466-470.
- Oliver, C., Jephcott, L., Seri, S., Freiss, S., Kilne, A. & Moss, J. (2007). Self-injurious behavior in Cornelia de Lange and cri du chat syndromes: Associations with peripheral sensory neuropathy and social interactions. *American Journal of Medical Genetics Part A*, 143A, 12, 1399-1400.
- Overhauser, J., Huang, X., Gersch, M., Wilson, W., McMahon, J., Bengtsson, U., Rojas, K., Meyer, M., & Wasmuth, J.J. (1994). Molecular and phenotypic mapping of the short arm of chromosome 5: Sublocalisation of the critical region of the Cri du Chat syndrome. *Human Molecular Genetics*, 3, 247-252.

- Sarimski, K. (2002). Analysis of intentional communication in severely handicapped children with Cornelia-de-Lange syndrome. *Journal of Communication Disorders*, 35, 483-500
- Simonoff, E., Pickles, A., Charman, T., Chandler, S., Loucas, T. & Baird, G. (2008). Psychiatric disorders in children with autism spectrum disorders: Prevalence, comorbidity and associated factors in a population-derived sample. *Journal of the American Academy of Child and Adolescent Psychiatry*, 47, 921-929.
- Sparrow, S. S., Balla, D. A. & Cicchetti, D. V. (1984). The Vineland Adaptive Behavior Scales: Interview Edition-Manual. *American Guidance Service, Inc., Circle Pines*.
- Tonkin, E. T, Wang, T-J., Lisgo, S., Bamshad, M. J. & Strachan, T. (2004). NIPBL, encoding a homolog of fungal Scc2-type sister chromatid cohesion proteins and fly Nipped-B, is mutated in Cornelia de Lange Syndrome. *Nature Genetics*, 36, 636-641.

Acknowledgements

The authors would like to thank all of the individuals who participated in the study, the CdLS Foundation (UK and Ireland) for supporting the research and Lesley Wilkie for her contribution to data collection.

Correspondence concerning this article should be addressed to:

Professor C. Oliver
Centre for Neurodevelopmental Disorders
School of Psychology
University of Birmingham
Edgbaston
Birmingham
B15 2TT
UK
0121 4144909
C.Oliver@Bham.ac.uk

Footnotes

1. The recruitment procedure for the larger study sample is described in detail in Moss *et al.* (2008).

Tables

Table 1: Participant characteristics and comparative analysis

		CdLS (N=12)	CdCS (N=12)	Independent t-test value	P Value
Chronological Age (Years)	Mean (SD)	11:00 (5.15)	8:20 (2.86)	17.17	.11
	<i>range</i>	<i>5.00-18.00</i>	<i>5.00-14.00</i>		
Gender	N Male	4	4		
VABS ABC standard score	Mean (SD)	45.83 (12.50)	36.83 (9.10)	2.01	.12
	<i>range</i>	<i>31.00 – 68.00</i>	<i>23.00 – 54.00</i>		
VABS Classification	N Mild	3	1		
	N Moderate	5	3		
	N Severe	4	8		

Table 2: Operational Definitions of Behaviors and Cohen's Kappa values.

Behavior	Operational Definition	Kappa
Behaviors Indicative of Social Anxiety		
Eye contact	Participant looks up/or at the examiner and fixates on the examiners eyes or face.	.80
Participant – communication	Any verbal communication or use of formal signs directed towards the examiner. This includes prompting, offering information and response to a question.	.90
Moving of hands	Moving of hands to face, head, or another part of the body. For example scratching or touching face, hair, arm, which has no obvious function. Excluding any forms of self-injury, or any communicative gestures.	.86
Examiner Behavior		
Examiner – communication	Any verbal communication directed towards the participant. This includes prompting, offering information and response to a question.	.96

Table 3. Mean frequencies, durations and comparative analyses for behaviors indicative of social anxiety for the CdLS group and CdCS group.

Behavior	CdLS	CdCS	Mann	P Value
			Whitney U	
Eye Contact duration	19.33	15.73	68.00	.82
Participant Communication duration	13.06	9.70	57.00	.39
Moving Hand frequency (per minute)	1.03	.39	52.50	.26

Figure Captions

Figure 1. Mean unconditional probability (filled squares) of the child engaging in the target behavior and conditional probability (unfilled squares) of the child engaging in the target behavior given that they are engaging in the criterion behavior, for 21s before and after the target behavior occurs for the CdLS group and the CdCS group (* = $p < .00001$, $z > 4.5$).

Figure 1 Top