Multiplex proximity extension assay of 425 candidate biomarkers in the sera of bladder cancer patients: Correlation with stage and outcome

Ward, Douglas; Gordon, Naheema; Abbotts, Ben; James, Nicholas; Zeegers, Maurice; Cheng, Kar; Bryan, Richard

DOI:

[10.1016/S1569-9056\(17\)30187-2](https://doi.org/10.1016/S1569-9056(17)30187-2)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Ward, D, Gordon, N, Abbotts, B, James, N, Zeegers, M, Cheng, K & Bryan, R 2017, 'Multiplex proximity extension assay of 425 candidate biomarkers in the sera of bladder cancer patients: Correlation with stage and outcome', *European Urology Supplements*, vol. 16, no. 3. [https://doi.org/10.1016/S1569-9056\(17\)30187-2](https://doi.org/10.1016/S1569-9056(17)30187-2)

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Abstract ID
Number: **AM17-4923**

New Number: **121**

Abstract Title: **Multiplex proximity extension assay of 425 candidate biomarkers in the sera of bladder cancer patients: Correlation with stage and outcome**

Authors: **D. Ward**¹, **N. Gordon**¹, **B. Abbotts**², **N. James**³, **M. Zeegers**⁴, **K.K. Cheng**⁵, **R. Bryan**¹

(1) University of Birmingham, Institute of Cancer and Genomic Sciences, Birmingham, United Kingdom, (2) University of Birmingham, Institute Of Cancer And Genomic Sciences, Birmingham, United Kingdom, (3) University of Warwick, Clinical Trials Unit, Warwick, United Kingdom, (4) University of Maastricht, Dept. of Complex Genetics, Birmingham, United Kingdom, (5) University of Birmingham, School of Health and Population Sciences, Birmingham, United Kingdom

Body:

Introduction & Objectives Serum protein biomarkers that inform on UBC stage and outcome could accelerate and improve clinical decision making but this requires a thoroughly validated high-performance biomarker or panel of biomarkers. Unbiased discovery of serum biomarkers by shotgun proteomics is challenging due to their low abundance in a protein-rich milieu and candidate based discovery is slow and expensive due to the number of individual immunoassays required. In this study we make use of a recently developed multiplex assay platform to measure 425 proteins in the serum of bladder cancer patients with the aim of identifying novel staging and prognostic biomarkers.

Material & Methods All sera were collected as part of the Bladder Cancer Prognosis Programme (ethics approval 06/MRE04/65). Patients whose diagnostic cystoscopy indicated primary UBC were recruited to the study and blood collected prior to TURBT. Ultimately, some of the patients were diagnosed with non-malignant conditions and these serve as non-cancer controls. All sera were prepared following a standard operating procedure and stored at -80°C. The patients used in this study comprised 10 non-cancer controls, 10 G1pTa, 10 G3pTa, 30 G3T1 and 30 G3T2+. There were no significant differences in age and gender between the groups. Proseek multiplex immunoassays (OLINK Proteomics) were used to measure 444 proteins of which 425 passed QC. The statistical analysis was done using R statistical software 3.2.5. Five samples were excluded from the analysis using PCA analysis. Multiple testing correction was done with the Benjamini-Hochberg method and an adjusted p-value<0.05 was considered significant. We used parametric tests and examined any significant hits for heteroscedasticity by evaluation of QQ and box-plots.

Results 425 proteins were successfully measured in the serum of 9 non-cancer controls and 76 UBC patients. 10 proteins were significantly associated with UBC and are listed in order of p-value (smallest first): nectin-4, syndecan-1, kidney injury molecule 1, macrophage colony-stimulating factor 1, matrilysin, thrombopoietin, latency-associated peptide transforming growth factor beta-1, C-C motif chemokine 23, heat shock protein beta-1 and protein AMBP. The serum levels of these proteins all showed a positive association with increasing stage although none showed absolute discrimination between stages. High levels of nectin-4, syndecan-1, macrophage colony-stimulating factor 1 were significantly associated with shorter UBC specific survival times.

Conclusions We have studied the relationship between bladder cancer and the serum concentrations of 425 proteins. Those which reach statistical significance include known biomarkers and new candidates that may warrant further investigation. Bladder cancer causes many biologically plausible changes in the serum proteome which may aid in UBC staging and prognosis.