Influence of sampling approach on concentrations of legacy and "novel" brominated flame retardants in indoor dust

Al-Omran, Layla Salih; Harrad, Stuart

DOI:

[10.1016/j.chemosphere.2017.02.096](https://doi.org/10.1016/j.chemosphere.2017.02.096)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Al-Omran, LS & Harrad, S 2017, 'Influence of sampling approach on concentrations of legacy and "novel" brominated flame retardants in indoor dust', *Chemosphere*, vol. 178, pp. 51-58.

<https://doi.org/10.1016/j.chemosphere.2017.02.096>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

INFLUENCE OF SAMPLING APPROACH ON CONCENTRATIONS OF LEGACY AND “NOVEL” BROMINATED FLAME RETARDANTS IN INDOOR DUST

Layla Salih Al-Omran ^{a,b} and Stuart Harrad^a

^aSchool of Geography, Earth, and Environmental Sciences, University of Birmingham,
Birmingham, B15 2TT, UK

^bDivision of Ecology, College of Science, University of Basrah, Basrah, Iraq

ABSTRACT

The study investigates the impact of sampling method on the concentrations of PBDEs (BDE-28, BDE-47, BDE-99, BDE-100, BDE-153, BDE-154, BDE-183, and BDE-209) and NBFRs (PBEB, EH-TBB, BEH-TEBP, BTBPE and DBDPE) in indoor dust. A total of 36 dust samples were collected from 12 homes in Birmingham, UK (3 samples per home comprising researcher collected dust – both RCD from the living room (RCDL) and bedroom (RCDB), with an additional householder vacuum dust sample - HHVD). BDE-209 was the predominant compound, with average concentrations of 2642, 2336 and 2634 ng/g in RCDL, RCDB and HHVD respectively. The next most abundant BFR was BEH-TEBP, followed by DBDPE, with average concentrations of 306, 339 and 233 ng/g for BEH-TEBP and 155, 91 and 152 ng/g for DBDPE in RCDL, RCDB and HHVD respectively. Average concentrations of Σ_6 tri-hexa-BDEs were 47, 41, and 24 ng/g in RCDL, RCDB and HHVD respectively. With the exception of Σ_6 tri-hexa-BDEs, BDE-153, BDE-99 and to some extent BEH-TEBP, no significant differences were found between BFR concentrations in RCD and HHVD. Statistically significant correlations were observed between concentrations of Σ_6 tri-hexa-BDEs, BEH-TEBP and DBDPE in HHVD and in both RCDL and RCDB. However, comparison of estimates of exposure via dust ingestion based on these two sampling methods revealed that using householder vacuum dust underestimates exposure, particularly for Σ_6 tri-hexa-BDEs, and to some extent for BEH-TEBP. In contrast, HHVD could be a viable alternative to RCD as a metric of exposure for higher brominated BFRs.

HIGHLIGHTS

- BFRs measured in researcher-collected (RCD) and house holder vacuum dust (HHVD).
- Concentrations of more volatile BFRs and BEH-TEBP in HHVD **lower** than those in RCD.
- Concentrations of less volatile do not vary significantly between the two sampling methods.
- Σ_6 tri-hexa-BDEs, BEH-TEBP and DBDPE in HHVD and RCD significantly correlated.
- Using HHVD may underestimate exposure for Σ_6 tri-hexa-BDEs and BEH-TEBP.

1. INTRODUCTION

Polybrominated diphenyl ethers (PBDEs) and “novel” brominated flame retardants (NBFRs) are chemicals added to a wide range of consumer products (electrical and electronic equipment, textiles, polyurethane and polystyrene foams) to meet flame retardancy standards set by various jurisdictions worldwide (Danish EPA, 2013; USEPA, 2014). Since in most applications these chemicals are used additively, they can transfer from such products into the environment (WHO, 1997; Alaei et al., 2003; USEPA, 2010). Evidence of their persistence and capacity for bioaccumulation, coupled with their adverse health effects have led to concern about human exposure (USEPA, 2006; 2008a; 2008b; 2008c; NICNAS, 2007; Noyes et al., 2010; Chevrier et al., 2010; EFSA, 2012; European Commission, 2012; Johnson et al., 2013; Li et al., 2014; Mankidy et al., 2014; Mariani, et al., 2015). Ingestion of indoor settled dust appears to represent a major pathway of exposure to BFRs particularly for young children (Jones-Otazo et al., 2005; Wilford et al., 2005; Harrad et al., 2008a; 2008b; 2010; Abdallah et al. 2008; Lorber, 2008; Roosens et al., 2009; Wang et al., 2010; Basis and Samara 2012; Stapleton et al., 2012; Qi et al., 2014; Hoffman et al., 2015).

To date, there are few studies that have investigated the association between indoor dust sampling method and the concentration of pollutants. For determining exposure of children to lead-contaminated household dust, early studies investigated different house dust sampling methods, indicating that the HVS3 (high-volume small surface sampler) had the highest level of precision among different standardised vacuuming and wipe sampling methods (Sterling et al., 1999), due to the small particles that can be retained by the HVS3 (Lioy et al., 2002). This method has subsequently been widely used, although it can be expensive, complicated and time-consuming (Mercier et al., 2011; USEPA, 2008d). Thus, commercial household vacuum cleaners are widely used as an alternative to the HVS3. By using a household vacuum cleaner, two approaches for dust collection are commonly used in studies of indoor contaminants. One of these approaches involves householders providing the contents of their vacuum cleaners to the researchers (Harrad et al., 2006; Suzuki et al., 2006; Kopp et al., 2012; Shen et al., 2015; Cristale et al., 2016). The principal advantages of the householder vacuum cleaner approach are that: it reflects indoor contamination from all rooms, is cost-effective, provides a large quantity of dust in a short time, and enhances donor compliance, by obviating the need for researchers to enter the home (Harrad et al., 2010). However, dust collected by this approach may be contaminated by the inner part of the vacuum cleaner, thereby reducing the accuracy of this method. Moreover, spatial variability, temporal variability and dust loading cannot be

assessed by this method, as the time and locations covered by the sample are unknown. In addition, vacuum cleaner sampling rates are variable (Harrad et al., 2010). Another approach involves the use of a commercial vacuum cleaner by the researchers themselves by using standardized procedures and specific accessories such as socks inserted in the sampling train to retain dust (Brommer et al., 2012; Ali et al., 2013; Harrad et al., 2016), Soxhlet thimbles (Allen et al., 2008; Stapleton et al., 2012) and filters (Björklund et al., 2012; Thuresson et al., 2012; Newton et al., 2015). The main advantages of such researcher-collected dust approaches are that: it minimises contamination of the sample due to specific accessories which are replaced or cleaned between taking each sample, and that it provides information about the specific time and location of collection of each dust sample, thereby facilitating study of within-room and within-home spatial and temporal variations in BFR concentrations. However, in comparison with the householder vacuum approach, this method is expensive and time-consuming, and may possibly hinder donor compliance as it requires entry of the researcher to the sampled microenvironment (Harrad et al., 2010). Only two studies (Allen et al., 2008; Björklund et al., 2012) have investigated the variation between researcher-collected and household vacuum approaches for analysing PBDE in indoor dust. Overall, the two studies reported PBDE concentrations in researcher-collected dust exceeded significantly those in householder-donated vacuum cleaner dust.

The aim of this study is to investigate the impact of two commonly-employed dust sampling methods on the concentrations of eight PBDEs (BDE-28, BDE-47, BDE-99, BDE-100, BDE-153, BDE-154, BDE-183 and BDE-209) and five NBFRs: pentabromoethylbenzene (PBEB), 2-ethylhexyl-2,3,4,5-tetrabromobenzoate (EH-TBB), bis (2-ethylhexyl) 3,4,5,6-tetrabromophthalate (BEH-TEBP), 2-bis (2,4,6-tribromophenoxy) ethane (BTBPE), and decabromodiphenylethane (DBDPE) in indoor dust and to evaluate the extent to which these sampling methods influence exposure assessments to BFRs via dust ingestion. To our knowledge, this study is the first investigation of the influence of sampling approach on concentrations of NBFRs in indoor dust.

2. MATERIALS AND METHODS

2.1. Sampling and sample preparation

Dust samples (n = 36) were collected from 12 homes between September 2014 and April 2015. In each home, two floor dust samples were collected by the researcher (researcher-collected dust- RCD) with the householder additionally providing the contents of their vacuum cleaner

(household vacuum dust- HHVD). RCD samples were obtained from the living room (RCDL) and bedroom (RCDB) of each house according to a clearly defined standard protocol (Harrad et al., 2008a). Briefly, by using a handheld vacuum cleaner (DIRT DEVIL-DDMHH1-1100W), 1 m² of carpeted floor area was vacuumed for 2 min, using 25 µm pore size nylon sample socks mounted in the furniture attachment tube of the vacuum cleaner. After sampling, socks were closed with a twist tie, sealed in plastic bags. Before sampling, the furniture attachment and the vacuum tubing were cleaned thoroughly using isopropanol-impregnated disposable wipes and dried between collections. HHVD samples were collected at the same time. The dust bag from the householder's own vacuum cleaner was wrapped in aluminium foil and sealed in a plastic bag. All samples were stored at -20 °C until analysis. Prior to analysis, dust samples were passed through a pre-cleaned, *n*-hexane rinsed 250 µm mesh steel sieve covered with the lid and shaken for 3-5 min. Field blanks were conducted which consisted of sodium sulfate that spread on aluminium foil, collected using the vacuum cleaner used to collect RCD and treated as a sample.

2.2. Sample extraction, clean up and instrumental analysis

PBDEs and NBFRs in dust samples were analysed following the same extraction method as reported elsewhere (Ali et al., 2011; Van den Eede et al., 2012, Al-Omran and Harrad 2016a; 2016b) with minor modifications. Accurately weighted aliquots of dust (~ 0.1 g) were spiked with a mixture of internal standards (20 ng of BDE-77, BDE-128, ¹³CBTBPE, ¹³CBEH-TEBP, and 40 ng of ¹³CBDE-209) in isooctane. Dust samples were extracted with 2 mL *n*-hexane: acetone (3:1 v/v), 2× (vortexed for 2 min, sonicated for 5 min) and centrifuged at 3500 rev/min for 5 min. The extraction process was repeated three times and the combined extracts were evaporated to incipient dryness under a gentle nitrogen stream, resolubilised in 1 mL of *n*-hexane.

Concentrated crude sample extracts were purified according to a previously reported method (Al-Omran and Harrad 2016a; 2016b) involving two steps. Briefly, in the first step, the extract was fractionated into two fractions (F1 and F2) using a 2 g Florisil SPE cartridge. F1 (containing PBDEs, DBDPE and PBEB) was eluted with 12 mL of hexane and F2 (containing the rest of the targeted NBFRs) was eluted with 15 mL ethyl acetate. After evaporation to 1 mL, a second purification step for F1 was conducted on 2 g acid silica (44% w/w) eluted with

15 mL *n*-hexane/DCM (1:1, v/v). F2 was evaporated to dryness, resolubilised in 3-5 mL of hexane, then evaporated to 1 mL, and eluted with 12 mL *n*-hexane/DCM (1:1, v/v) using an aminopropyl functionalised silica column (0.5 g). F1 and F2 were combined and evaporated to incipient dryness, before resolubilisation in 100 µL of iso-octane containing PCB-129 at 250 pg/µL ready for GC/MS analysis.

Target PBDEs and NBFRs were quantified using a gas chromatograph (GC) (Trace 1310 Gas Chromatograph) coupled to a mass spectrometer (MS) (ISQ Quadrupole MS); both (Thermo Fisher Scientific, USA) according to our previous study of BFRs in indoor dust (Al-Omran and Harrad 2016b). The GC was equipped with a programmable temperature vaporiser (PTV) injector and fitted with a capillary fused silica column (RESTEK, USA, 15 m x 0.25 mm inner diameter, 0.25 µm film thickness). The MS was operated in the electron capture negative ion (ECNI) mode. Table S1 shows quantification ions, qualification ions and retention times monitored for target compounds, internal standards (IS) and the recovery determination standard (RDS).

2.3. Quality assurance/Quality control

All glassware was cleaned by soaking them overnight in a detergent solution. After washing, glassware and Pasteur pipettes were heated to 470 °C for 5 h. Before use, glassware was rinsed with acetone and hexane. To avoid any degradation that may occur via exposure to light, glassware and the Turbopap instrument were covered with aluminium foil. To assess any possible contamination during sample preparation and analysis method, one laboratory blank was processed in parallel with every set of 6 dust samples and one quality control sample (NIST SRM 2585, organics in indoor dust) was processed with every 12 real dust samples. Limits of detection (LOD) were estimated based on a signal to noise ratio 3:1 and limits of quantification (LOQ) were estimated based on signal to noise ratio 10:1. Where a target compound (as was the case in some instances for BEH-TEBP and BDE-209) was detected a blank, the LOQ for that analyte was calculated as the mean plus 3 times the standard deviation of the concentrations detected in 10 blank samples. Field blanks (n = 5) were also collected to assess any contamination contributed as a result of sampling, transport and storage of samples, in addition to any introduced as a result of extraction and clean-up. The average of internal standard recoveries in dust samples ranged from 75-93%. PBDE and NBFR concentrations in SRM2585

detected in this study were in good agreement with the certified values and those reported in other studies. Tables S3, S4 and S5 report internal standard recovery values, along with PBDE and NBFR concentrations in SRM2585 detected in this study.

2.4. Statistical analysis

Statistical analysis of our data was performed using Microsoft Excel 2013 and IBM SPSS statistics software (V. 20). To test any differences in mean BFR dust concentrations between the two collection methods (researcher-collected and household vacuum), and between the two researcher-collected rooms (living room and bedroom) one way repeated measures ANOVA was performed. After testing for normality using the Shapiro–Wilk test confirmed our data to display a skewed distribution, data were transformed using the natural logarithm of concentrations (ng/g dw). For the purposes of statistical evaluation, all concentrations below LOQ were assigned a value of 0.5 LOQ. A p value < 0.05 was taken to indicate statistical significance. A Pearson correlation was used to test the relationship between BFR concentrations in dust collected via the two sampling methods.

3. RESULTS AND DISCUSSION

3.1. Influence of dust sampling approach on BFR concentrations

3.1.1 Detection frequencies and the relationship between BFRs

In all dust samples ($n = 36$), the detection frequency of PBDEs (BDE-28, BDE-47, BDE-99, BDE-100, BDE-153, BDE-154, BDE-183 and BDE-209) and NBFRs (PBEB, EH-TBB, BTBPE, BEH-TEBP, and DBDPE) ranged from 50% to 100%. The detection frequencies of BDE-209, BEH-TEBP and DBDPE were 100% in both researcher-collected dust (RCD) and household vacuum dust (HHVD). BDE-28, BDE-100, BDE-154, and PBEB were in the lowest detection frequencies. They were thus not accounted for individual statistical comparison, but were instead included in Σ_7 tri-hexa-BDEs (BDE-28, BDE-47, BDE-99, BDE-100, BDE-153 and BDE-154) and Σ_5 NBFRs (PBEB, EH-TBB, BTBPE, BEH-TEBP, and DBDPE). Table S6 lists detection frequencies of PBDEs and NBFRs in this study.

Among our target BFRs, BDE-209 was predominant, with average percentage contributions to Σ BFRs (sum of Σ_7 tri-hepta-BDEs, BDE-209 and Σ_5 NBFRs) of 83.2%, 82.7% and 85.9% in RCDL, RCDB and HHVD respectively. Σ_5 NBFRs were the next most abundant parameter,

with average contributions of 15.1%, 15.8% and 13% in RCDL, RCDB and HHVD respectively. Σ_7 tri-hepta-BDEs displayed the lowest average percentage contributions of our target BFRs. The average percentage contributions of BDE-99, BDE-47 and BDE-183 to Σ_7 tri-hepta-BDEs were 44%, 45% and 38% for BDE-99, 29%, 27% and 23% for BDE-47, and 11%, 5.4% and 23% for BDE-183 in RCDL, RCDB and HHVD respectively. Of our target NBFRs, BEH-TEBP predominated, making mean percentage contributions to Σ_5 NBFRs of 64%, 76% and 58%, followed by DBDPE which contributed 32%, 20% and 38% of Σ_5 NBFRs in RCDL, RCDB and HHVD respectively. Figure 1 depicts the average percentage contributions and congener/compound profiles of target BFRs, tri-hepta-BDEs and NBFRs in RCDL, RCDB and HHVD.

3.1.2. Concentrations of PBDEs and NBFRs in indoor dust obtained via two different sampling methods

The three main commercial PBDE formulations (Penta-BDE, Octa-BDE and Deca-BDE) are represented in this study by Σ_6 tri-hexa-BDEs (BDE-28, BDE-47, BDE-99, BDE-100, BDE-153 and BDE-154) as an indicator of Penta-BDE, BDE-183 as an indicator of Octa-BDE and BDE-209 as an indicator of Deca-BDE. In all dust samples, the highest concentrations of total target Σ PBDEs and Σ_5 NBFRs were found in researcher-collected dust samples from living rooms (RCDL), with values of 4321 and 1450 ng/g for Σ PBDEs and Σ_5 NBFRs respectively. BDE-209 was present at average concentrations of 2642, 2336 and 2634 ng/g, while those of BEH-TEBP were 309, 339 and 233 ng/g in RCDL, RCDB and HHVD respectively. Average concentrations of DBDPE were comparable in both RCDL and HHVD samples, with values of 155 and 152 respectively, while in RCDB it was 91 ng/g. Σ_6 tri-hexa-BDEs average concentrations were 47.3 and 41.3 and 24.4 ng/g in RCDL, RCDB and HHVD respectively. For the rest of our target NBFRs, BTBPE was found in comparable average concentrations in RCDL and HHVD, with values of 11.0 and 11.2 respectively, while in RCDB, it was 9.5 ng/g. Average concentrations of EH-TBB in RCDL and RCDB were comparable (6.9 and 6.4 ng/g) exceeding those in HHVD samples (4.9 ng/g). Tables 1 and 2 provide statistical summaries of concentrations of PBDEs and NBFRs in RCDL, RCDB, and HHVD samples.

Comparison with previous studies (Harrad et al., 2008a; 2008b) in Birmingham, UK, revealed median concentrations of PBDEs in this study to be lower than in the earlier studies by a factor of 2.7 for Σ_8 PBDEs, while DBDPE increased by a factor of 3.6. While our study is based on

too few homes to provide a definitive temporal trend; these data are not inconsistent with restrictions on PBDE use and possible concomitant increased use of NBFR alternatives.

3.1.3. Comparison of BFR concentrations in dust samples from two sampling methods

Tables 1 and 2 show concentrations of BDE-47, BDE-99, BDE-153, Σ_6 tri-hexa-BDEs and EH-TBB in researcher-collected dust from both living room (RCDL) and bedroom (RCDB) exceed those in household vacuum dust (HHVD). In contrast, BDE-183 in HHVD was higher than in RCDL and RCDB. Moreover, BDE-209, BTBPE, and DBDPE concentrations in HHVD were only higher than in RCDB and were comparable in both RCDL and HHVD. Table S7 lists the average concentration ratios for RCDL/ HHVD and RCDB/ HHVD for BDE-47, BDE-99, BDE-153, BDE-183, BDE-209, EH-TBB, BEH-TEBP and DBDPE.

One way repeated measures ANOVA tests were applied to compare means of BFR concentrations in RCDL, RCDB and HHVD. This revealed that, with the exception of Σ_6 tri-hexa-BDEs, BDE-153, BDE-99 and to a moderate extent BEH-TEBP, these differences were not statistically significant ($p > 0.05$). Concentrations of Σ_6 tri-hexa-BDEs and BDE-153 in researcher-collected dust (RCDL and RCDB combined) exceeded significantly those in the household vacuum dust with p values of 0.012 and 0.038 for Σ_6 tri-hexa-BDEs, and 0.025 and 0.016 for BDE-153 in RCDL and RCDB respectively. BDE-99 concentrations in RCDL exceeded significantly those in HHVD with a p value of 0.015. Moreover, BEH-TEBP concentrations in RCDL exceeded those in HHVD at a moderate level of significance ($p = 0.077$). ANOVA tests revealed, with the exception of BDE-183, no significant differences ($p > 0.05$) in BFR concentrations between the living room and bedroom. With respect to BDE-183, concentrations in the living room exceeded significantly those in the bedroom ($p = 0.001$). Based on our results, Penta-BDE and to a lesser extent BEH-TEBP displayed important differences between the two sampling methods, while concentrations of Deca-BDE, Octa-BDE, EH-TBB, BTBPE and DBDPE did not appear significantly impacted by the sampling method employed.

To our knowledge, only two studies (Allen et al., 2008; Björklund et al., 2012) have compared PBDE and HBCDD concentrations in house dust collected via different sampling methods. In a comprehensive study of indoor dust from 20 homes in Boston, USA; Allen et al., (2008) compared concentrations of PBDEs in dust collected using household vacuum cleaner and

researcher-collected (from living room and bedroom) methods. The study reported that Penta-BDE (Σ BDE-17, 28/33, 47, 49, 66, 75, 85/155, 99, 100, 183, 153 and 154) concentrations in researcher-collected dust samples exceeded significantly those in the household vacuum dust for both living rooms ($p = 0.001$) and bedrooms ($p = 0.002$). In addition, the concentrations of Deca-BDE formulation congeners (BDE-206, 207, 208 and 209) in the researcher-collected dust from the living room exceeded significantly ($p = 0.02$) those in the household vacuum dust, with such significant differences to the household vacuum dust not observed for bedroom researcher-collected samples. Moreover, the same study found no significant difference between Octa-BDE concentrations in dust obtained via the two sampling methods. With the exception of the concentrations of Deca-BDE formulation congeners in researcher-collected dust from the living room, our outcomes are consistent with the study of Allen et al. (2008), despite the differences in PBDE distribution profiles between the UK and USA, different sampling accessories (nylon sock and cellulose extraction thimble), different vacuum cleaner brands and different dust particle size fractions ($< 500 \mu\text{m}$ and $< 250 \mu\text{m}$).

Based on dust samples from 19 Swedish homes, Björklund et al., 2012, investigated the differences between PBDE and HBCDD concentrations in samples collected via researcher-collected and household vacuum methods. The researcher-collected method employed involved collection of settled house dust from elevated surfaces (1 m above the floor). Concentrations of all targeted PBDE congeners (BDE-28, BDE-47, BDE-49, BDE-66, BDE-85, BDE-99, BDE-100, BDE-153, BDE-154, BDE-183, BDE-197, BDE-203, BDE-206, BDE-207, BDE-208, and BDE-209) detected in researcher-collected dust exceeded significantly ($P < 0.001$ - 0.003) those collected using the household vacuum method. The significant differences observed by Björklund et al. (2012) between researcher-collected method and household vacuum method dust, exceeded those observed both in our study and that of Allen et al. (2008). This implies that, in addition to the different sampling methods, sampling of different surfaces (floor dust and elevated surface dust) exert an important influence on the findings of the Swedish study. This is consistent with findings reported in our previous studies (Al-Omran and Harrad 2016a; 2016b), that BFR concentrations in elevated surface dust samples exceed significantly those in floor dust.

3.1.4. Correlation between dust sampling methods

Pearson correlation analysis was performed on log-transformed data to determine the relationship between BFR concentrations in household vacuum dust (HHVD) and researcher-collected dust (RCD) from the living room (RCDL) and bedroom (RCDB). The strongest correlations observed between HHVD and RCD methods were for BEH-TEBP concentrations in the living room ($R = 0.793$, $p = 0.002$) and bedroom ($R = 0.883$, $p < 0.001$). Likewise, concentrations of Σ_6 tri-hexa-BDEs in HHVD correlated with those in RCDL ($R = 0.583$, $p = 0.047$) and RCDB ($R = 0.588$, $p = 0.044$), as well as those of DBDPE for which the corresponding correlations with HHVD concentrations were ($R = 0.643$, $p = 0.024$) and ($R = 0.634$, $p = 0.027$) for RCDL and RCDB respectively. Moreover, HHVD concentrations were moderately ($R = 0.532$, $p = 0.075$) associated with RCDL concentrations for BDE-209, and with RCDB for EH-TBB ($R = 0.557$, $p = 0.060$). Figure 2 shows scatter plots and Pearson correlation coefficients obtained when plotting log-transformed concentrations of Σ_6 tri-hexa-BDEs, BEH-TEBP and DBDPE in household vacuum dust against concentrations in both RCDL and RCDB. Table S8 shows Pearson correlation results describing the relationship between BFR concentrations in dust samples collected by a researcher (RCDL and RCDB) and household vacuum (HHVD) approaches.

In general, our findings reveal that BFR concentrations in dust collected via the two sampling methods were highly correlated for BEH-TEBP, DBDPE, Σ_6 tri-hexa-BDEs, and BDE-99, and moderately correlated for BDE-209 and EH-TBB. In contrast, concentrations of BDE-47, BDE-153, BDE-183, and BTBPE were not significantly correlated between researcher-collected and household vacuum dust. For PBDEs, with the exception of BDE-209, these findings are consistent with previous studies (Allen et al. 2008; Björklund et al., 2012). However, Björklund et al., concluded that, when a single high value of BDE-209 was removed from their data analysis, the correlation they observed between concentrations obtained via the 2 dust collection methods was no longer significant (Björklund et al., 2012).

3.1.5 The impact of sampling approach on human exposure assessments to BFRs

To evaluate the extent to which human exposure to our target contaminants via dust ingestion is affected by the choice of sampling approach, we compared the median concentration (for typical exposure) and 95th percentile (for high end exposure) in dust samples collected via the two sampling approaches; researcher collected from the living room and bedroom and household vacuum contents. This comparison revealed that the impact of sampling method on

estimates of exposure to BDE-99 and Σ tri-hexa-BDEs was more important than for other target compounds. Concentrations of (and thus exposure to) BDE-99 and Σ tri-hexa-BDEs in RCD exceeded substantially those in HHVD by factors of 1.5 and 3.5 for BDE-99 and 1.8 and 2.7 for Σ tri-hexa-BDEs for the median and 95th percentile respectively. In addition, concentrations of BEH-TEBP based on analysis of RCD exceeded those for HHVD by factors of 1.4 and 1.5 for the median and 95th percentile respectively. This implies that exposure assessments for these compounds based on analysis of HHVD may be underestimates, particularly when making high-end exposure assessments. In contrast, concentrations of DBDPE in HHVD exceeded those in RCDB by factors of 1.1 and 2.3 for median and 95th percentile concentrations respectively, which implies that analysing HHVD may overestimate exposure to DBDPE. Table 3 illustrates RCDL/HHVD and RCDB/HHVD median and 95th percentile concentration ratios for BDE-99, Σ_6 tri-hexa-BDEs, BDE-209, BEH-TEBP and DBDPE, which illustrate the impact of sampling method on typical and high-end exposure assessments.

4. CONCLUSIONS

This study found that concentrations of BDE-99, BDE-153, Σ_6 tri-hexa-BDEs and – to some extent - BEH-TEBP were significantly lower in HHVD (household vacuum dust) than those in RCD (researcher-collected dust) from both living rooms and bedrooms. This might be due to volatilisation of BFRs as a result of the long residence times of dust in the household vacuum.

In addition, RCD samples were collected from bedrooms and living rooms where large quantities of Penta-BDE and BEH-TEBP may have been used (in articles such as beds, chairs, and sofas), whereas the household vacuum cleaner would contain a complex integral of dust from the entire house, which would include rooms containing fewer products containing Penta-BDE, such as kitchens (Kuang et al., 2016) and hallways. Moreover, small particles (which we have shown previously to contain higher concentrations of some BFRs – Al-Omran and Harrad, 2016b) may have been lost through collecting and transferring processes from the vacuum bag. Our findings indicate that exposure assessments using HHVD may be underestimated for Σ_6 tri-hexa-BDEs and BEH-TEBP, which suggest that this approach is a less suitable method for assessing human exposure to these compounds. However, it could be a viable alternative to RCD for higher brominated BFRs such as BDE-209.

Due to the different particle size distribution pattern of BFRs in indoor dust (Wei et al. 2009; Cao et al., 2013, 2014, 2015; Kefeni et al., 2014), future studies are recommended using particle size analyser to examine the particle size distribution pattern of BFR concentrations in obtained via

the two sampling methods, to test the hypothesis that a greater proportion of fine particles in RCD account for the higher BFR concentrations observed in such dust compared to HHVD. This is because the same compounds (BDE-99, BDE-153, Σ_6 tri-hexa-BDEs and BEH-TEBP) that were significantly elevated in researcher collected compared to household vacuum collected dust, are also significantly higher in the finest particle size fractions of indoor dust (Al-Omran and Harrad, 2016b).

ACKNOWLEDGEMENTS

The authors express their thanks to all the dust donors from Birmingham, UK. Layla Salih Al-Omran acknowledges gratefully the Iraqi government for a PhD Scholarship, the Iraqi Establishment of Martyrs for financial support and the Ministry of Higher Education and Scientific Research for administrative support.

REFERENCES

- Abdallah, M. A., S. Harrad and A. Covaci (2008). "Hexabromocyclododecanes and tetrabromobisphenol-A in indoor air and dust in Birmingham, U.K: implications for human exposure." *Environmental Science & Technology* **42**(18): 6855-6861.
- Alaee, M. (2003). "An overview of commercially used brominated flame retardants, their applications, their use patterns in different countries/regions and possible modes of release." *Environment International* **29**(6): 683-689.
- Ali, N., S. Harrad, D. Muenhor, H. Neels and A. Covaci (2011). "Analytical characteristics and determination of major novel brominated flame retardants (NBFRs) in indoor dust." *Analytical and Bioanalytical Chemistry* **400**(9): 3073-3083
- Ali, N., L. Ali, T. Mehdi, A. C. Dirtu, F. Al-Shammari, H. Neels and A. Covaci (2013). "Levels and profiles of organochlorines and flame retardants in car and house dust from Kuwait and Pakistan: Implication for human exposure via dust ingestion." *Environment International* **55**: 62-70.
- Allen, J. G., M. D. McClean, H. M. Stapleton and T. F. Webster (2008). "Critical factors in assessing exposure to PBDEs via house dust." *Environment International* **34**(8): 1085-1091.
- Al-Omran, L. S. and S. Harrad (2016a). "Polybrominated diphenyl ethers and "novel" brominated flame retardants in floor and elevated surface house dust from Iraq: Implications for human exposure assessment." *Emerging Contaminants* **2**(1): 7-13.
- Al-Omran, L. S. and S. Harrad (2016b). "Distribution pattern of legacy and "novel" brominated flame retardants in different particle size fractions of indoor dust in Birmingham, United Kingdom." *Chemosphere* **157**: 124-131.
- Besis, A. and C. Samara (2012). "Polybrominated diphenyl ethers (PBDEs) in the indoor and outdoor environments--a review on occurrence and human exposure." *Environmental Pollution* **169**: 217-229.

435 Björklund, J. A., U. Sellstrom, C. A. de Wit, M. Aune, S. Lignell and P. O. Darnerud (2012).
 436 "Comparisons of polybrominated diphenyl ether and hexabromocyclododecane
 437 concentrations in dust collected with two sampling methods and matched breast milk
 438 samples." Indoor Air **22**(4): 279-288.

439 Brommer, S., S. Harrad, N. Van den Eede and A. Covaci (2012). "Concentrations of
 440 organophosphate esters and brominated flame retardants in German indoor dust
 441 samples." Journal of Environmental Monitoring **14**(9): 2482-2487.

442 Chevrier, J., K. G. Harley, A. Bradman, M. Gharbi, A. Sjödin and B. Eskenazi (2010).
 443 "Polybrominated Diphenyl Ether (PBDE) Flame Retardants and Thyroid Hormone
 444 during Pregnancy." Environmental Health Perspectives **118**(10): 1444-1449.

445 Cristale, J., A. Hurtado, C. Gómez-Canela and S. Lacorte (2016). "Occurrence and sources
 446 of brominated and organophosphorus flame retardants in dust from different indoor
 447 environments in Barcelona, Spain." Environmental Research **149**: 66-76.

448 Cao, Z., F. Xu, A. Covaci, M. Wu, H. Wang, G. Yu, B. Wang, S. Deng, J. Huang and X.
 449 Wang (2014). "Distribution patterns of brominated, chlorinated, and phosphorus flame
 450 retardants with particle size in indoor and outdoor dust and implications for human
 451 exposure." Environmental Science & Technology **48**(15): 8839-8846.

452 Cao, Z., G. Yu, Y. Chen, C. Liu, K. Liu, T. Zhang, B. Wang, S. Deng and J. Huang (2013).
 453 "Mechanisms influencing the BFR distribution patterns in office dust and implications
 454 for estimating human exposure." Journal of Hazardous Materials **252-253**: 11-18.

455 Cao, Z., F. Xu, W. Li, J. Sun, M. Shen, X. Su, J. Feng, G. Yu, and A. Covaci (2015)."
 456 Seasonal and particle size-dependent variations of hexabromocyclododecanes in settled
 457 dust: Implications for sampling." Environmental Science & Technology **49** (18): 11151-
 458 11157.

459 Danish EPA. Danish Ministry of Environment, Environmental Protection Agency (2013).
 460 "Survey of Brominated Flame Retardants (online). Available from:
 461 <http://www2.mst.dk/Udgiv/publications/2014/01/978-87-93026-90-2.pdf> (Accessed May
 462 2016)

463 EFSA (European Food Safety Authority), 2012. "Scientific Opinion on Emerging and Novel
 464 Brominated Flame Retardants (BFRs) in Food." European Food Safety Authority
 465 Journal **10**(10):2908.

466 European Commission (2012). "Recommendation from the Scientific Committee on
 467 Occupational Exposure Limits for Diphenyl ether, Octabromoderivative (commercial
 468 mixture)" (online). Available from:
 469 <http://ec.europa.eu/social/BlobServlet?docId=7721&langId=en>. (Accessed May 2016).

470 Harrad, S., S. Hazrati and C. Ibarra (2006). "Concentrations of polychlorinated biphenyls in
 471 indoor air and polybrominated diphenyl ethers in indoor air and dust in Birmingham,
 472 United Kingdom: implications for human exposure." Environmental Science &
 473 Technology **40**(15): 4633-4638.

474 Harrad, S., C. Ibarra, M. A.-E. Abdallah, R. Boon, H. Neels and A. Covaci (2008 a).
 475 "Concentrations of brominated flame retardants in dust from United Kingdom cars,
 476 homes, and offices: Causes of variability and implications for human exposure."
 477 Environment International **34**(8): 1170-1175.

478 Harrad, S., C. Ibarra, M. Diamond, L. Melymuk, M. Robson, J. Douwes, L. Roosens, A. C.
 479 Dirtu and A. Covaci (2008b). "Polybrominated diphenyl ethers in domestic indoor dust
 480 from Canada, New Zealand, United Kingdom and United States." Environment
 481 international **34**(2): 232-238.

482 Harrad, S., M. A.-E. Abdallah and A. Covaci (2009). "Causes of variability in concentrations
 483 and diastereomer patterns of hexabromocyclododecanes in indoor dust." Environment
 484 international **35**(3): 573-579.

485 Harrad, S., C. A. de Wit, M. A.-E. Abdallah, C. Bergh, J. A. Björklund, A. Covaci, P. O.
 486 Darnerud, J. de Boer, M. Diamond and S. Huber (2010). "Indoor contamination with
 487 hexabromocyclododecanes, polybrominated diphenyl ethers, and perfluoroalkyl
 488 compounds: an important exposure pathway for people?" Environmental science &
 489 technology **44**(9): 3221-3231.

490 Harrad, S., M. A.-E. Abdallah and T. Oluseyi (2016). "Polybrominated diphenyl ethers and
 491 polychlorinated biphenyls in dust from cars, homes, and offices in Lagos, Nigeria."
 492 Chemosphere **146**: 346-353.

493 Hoffman, K., S. Garantziotis, L. S. Birnbaum and H. M. Stapleton (2015). "Monitoring
 494 Indoor Exposure to Organophosphate Flame Retardants: Hand Wipes and House Dust."
 495 Environmental Health Perspectives **123**(2): 160-165.

496 Johnson, P. I., H. M. Stapleton, B. Mukherjee, R. Hauser and J. D. Meeker (2013).
 497 "Associations between brominated flame retardants in house dust and hormone levels in
 498 men." Science of the total environment **445**: 177-184.

499 Jones-Otazo, H. A., J. P. Clarke, M. L. Diamond, J. A. Archbold, G. Ferguson, T. Harner, G.
 500 M. Richardson, J. J. Ryan and B. Wilford (2005). "Is house dust the missing exposure
 501 pathway for PBDEs? An analysis of the urban fate and human exposure to PBDEs."
 502 Environmental Science & Technology **39**(14): 5121-5130.

503 Kefeni, K. K. and J. O. Okonkwo (2014). "Distribution of polybrominated diphenyl ethers
 504 and dust particle size fractions adherent to skin in indoor dust, Pretoria, South Africa."
 505 Environmental Science And Pollution Research **21**(6): 4376-4386.

506 Kopp, E. K., H. Fromme and W. Völkel (2012). "Analysis of common and emerging
 507 brominated flame retardants in house dust using ultrasonic assisted solvent extraction
 508 and on-line sample preparation via column switching with liquid chromatography–mass
 509 spectrometry." Journal of Chromatography A **1241**: 28-36.

510 Kuang, J., Y. Ma and S. Harrad (2016). "Concentrations of “legacy” and novel brominated
 511 flame retardants in matched samples of UK kitchen and living room/bedroom dust."
 512 Chemosphere **149**: 224-230.

513 Lioy, P. J., N. C. G. Freeman and J. R. Millette (2002). "Dust: a metric for use in residential
 514 and building exposure assessment and source characterization." Environmental Health
 515 Perspectives **110**(10): 969-983.

516 Li, M., Z. Liu, L. Gu, R. Yin, H. Li, X. Zhang, T. Cao and C. Jiang (2014). "Toxic effects of
 517 decabromodiphenyl ether (BDE-209) on human embryonic kidney cells." Frontiers in
 518 Genetics **5**: 118.

519 Lorber, M. (2008). Exposure of Americans to polybrominated diphenyl ethers. Journal of
 520 Exposure Science and Environmental Epidemiology **18**(1), 2-19.

521 Mankidy, R., Ranjan, B., Honaramooz, A., & Giesy, J. P. (2014). Effects of novel
522 brominated flame retardants on steroidogenesis in primary porcine testicular
523 cells. Toxicology letters **224**(1), 141-146.

524 Mariani, A., Fanelli, R., Depaolini, A., & Paola, M. (2015). Decabrominated diphenyl ether
525 and methylmercury impair fetal nervous system development in mice at documented
526 human exposure levels. Developmental neurobiology **75**(1), 23-38.

527 Mercier, F., P. Glorennec, O. Thomas and B. Le Bot (2011). "Organic contamination of
528 settled house dust, a review for exposure assessment purposes." Environmental science
529 & technology **45**(16): 6716-6727.

530 Muenhor, D. and S. Harrad (2012). "Within-room and within-building temporal and spatial
531 variations in concentrations of polybrominated diphenyl ethers (PBDEs) in indoor dust."
532 Environment international **47**: 23-27.

533 Newton, S., U. Sellström and C. A. de Wit (2015). "Emerging Flame Retardants, PBDEs,
534 and HBCDDs in Indoor and Outdoor Media in Stockholm, Sweden." Environmental
535 Science & Technology **49**(5): 2912-2920.

536 NICNAS, 2007, National Industrial Chemicals Notification and Assessment Scheme
537 (online). Available from:
538 [https://www.nicnas.gov.au/data/assets/pdf_file/0003/4944/Final-Interim-Report-](https://www.nicnas.gov.au/data/assets/pdf_file/0003/4944/Final-Interim-Report-PBDE-March.pdf)
539 [PBDE-March.pdf](https://www.nicnas.gov.au/data/assets/pdf_file/0003/4944/Final-Interim-Report-PBDE-March.pdf). Accessed January 2016.

540 Noyes P. D., S. M. Kelly, C. L. Mitchelmore, H. M. Stapleton. (2010) "Characterizing the in
541 vitro hepatic biotransformation of the flame retardant BDE 99 by common carp".
542 Aquatic Toxicology **97**(2): 142-150.

543 Qi, H., W.-L. Li, L.-Y. Liu, Z.-F. Zhang, N.-Z. Zhu, W.-W. Song, W.-L. Ma and Y.-F. Li
544 (2014). "Levels, distribution and human exposure of new non-BDE brominated flame
545 retardants in the indoor dust of China." Environmental Pollution **195**: 1-8.

546 Roosens, L., M. A. Abdallah, S. Harrad, H. Neels and A. Covaci (2009). "Factors
547 influencing concentrations of polybrominated diphenyl ethers (PBDEs) in students from
548 Antwerp, Belgium." Environmental science & technology **43**(10): 3535-3541.

549 Shen, B., T. P. Whitehead, S. McNeel, F. R. Brown, J. Dhaliwal, R. Das, L. Israel, J.-S. Park
550 and M. Petreas (2015). "High Levels of Polybrominated Diphenyl Ethers in Vacuum
551 Cleaner Dust from California Fire Stations." Environmental Science & Technology
552 **49**(8): 4988-4994.

553 Stapleton, H. M., Eagle, S., Sjödin, A., & Webster, T. F. (2012). Serum PBDEs in a North
554 Carolina toddler cohort: associations with handwipes, house dust, and socioeconomic
555 variables. Environmental health perspectives **120**(7): 1049.

556 Sterling, D. A., Roegner, K. C., Lewis, R. D., Luke, D. A., Wilder, L. C., & Burchette, S. M.
557 (1999). Evaluation of four sampling methods for determining exposure of children to
558 lead-contaminated household dust. Environmental research **81**(2): 130-141.

559 Suzuki, G., K. Nose, H. Takigami, S. Takahashi and S. I. Sakai (2006). "PBDEs and
560 PBDD/Fs in house and office dust from Japan". Organohalogen Compounds **141**: 30-41.

561 Thuresson, K., J. A. Björklund and C. A. de Wit (2012). "Tri-decabrominated diphenyl
562 ethers and hexabromocyclododecane in indoor air and dust from Stockholm

- microenvironments 1: Levels and profiles." *Science of The Total Environment* **414**: 713-721.
- USEPA. US Environmental Protection Agency (2006). "Toxicological review of 2,2', 4,4', 5-Pentabromodiphenyl ether (BDE-99)" (online). Available from: https://ofmpub.epa.gov/eims/eimscomm.getfile?p_download_id=474496 (Accessed February 2016).
- USEPA. US Environmental Protection Agency (2008a). "Toxicological Review of 2,2',4,4'-Tetrabromodiphenyl ether (BDE-47)" (online). Available from: <http://www.epa.gov/iris/toxreviews/1010tr.pdf> (Accessed May 2016).
- USEPA. US Environmental Protection Agency (2008b). "Toxicological Review of 2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE-153)" (online). Available from: <http://www.epa.gov/iris/toxreviews/1009tr.pdf> (Accessed May 2016).
- USEPA. US Environmental Protection Agency (2008c). "Toxicological Review of Decabromodiphenyl ether (BDE-209)" (online). Available from: <http://www.epa.gov/iris/toxreviews/0035tr.pdf> (Accessed April 2016).
- USEPA. US Environmental Protection Agency. (2008d) "Guidance for the Sampling and Analysis of Lead in Indoor Residential Dust for Use in the Integrated Exposure Uptake Biokinetic (IEUBK) Model" (online). Available from: <http://nepis.epa.gov> (Accessed April 2016).
- USEPA. US Environmental Protection Agency. (2010) an exposure assessment of polybrominated diphenyl ethers (online). Available from: <http://www.epa.gov/ncea>. (Accessed November 2015).
- USEPA. US Environmental Protection Agency (2014). "An Alternatives Assessment for the Flame Retardant Decabromodiphenyl ether (deca BDE)" (online). Available from: https://www.epa.gov/sites/production/files/2014-05/documents/decabde_final.pdf (Accessed April 2016).
- Van den Eede, N., A. C. Dirtu, N. Ali, H. Neels and A. Covaci (2012). "Multi-residue method for the determination of brominated and organophosphate flame retardants in indoor dust." *Talanta* **89**: 292-300.
- Wang, J., Y. J. Ma, S. J. Chen, M. Tian, X. J. Luo and B. X. Mai (2010). "Brominated flame retardants in house dust from e-waste recycling and urban areas in South China: implications on human exposure." *Environment International* **36**(6): 535-541.
- Wei, H., M. Turyk, S. Cali, S. Dorevitch, S. Erdal and A. Li (2009). "Particle size fractionation and human exposure of polybrominated diphenyl ethers in indoor dust from Chicago." *Journal of environmental science and health. Part A, Toxic/hazardous substances & environmental* **44**(13): 1353-1361.
- WHO. World Health Organization, (1997). Environmental Health Criteria 192 (online). Available from: <http://www.inchem.org/documents/ehc/ehc/ehc192.htm> (Accessed February 2015).
- Wilford, B. H., M. Shoeib, T. Harner, J. Zhu and K. C. Jones (2005). "Polybrominated diphenyl ethers in indoor dust in Ottawa, Canada: implications for sources and exposure." *Environmental science & technology* **39**(18): 7027-7035.

Figure 1: Average percentage contributions and congener/compound profiles of target BFRs, tri-hepta-BDEs and NBFRs in RCDL, RCDB (researcher-collected dust from the living room and bedroom) and HHVD (household vacuum dust)

Figure 2: Correlations between log-transformed concentrations of Σ_6 tri-hexa-BDEs, BEH-TEBP and DBDPE in household vacuum dust and researcher-collected dust from the living room and bedroom

Table 1: Summary statistics for PBDE concentrations (ng/g) in RCDL, RCDB (researcher collected dust from the living room bedroom) and dust HHVD (household vacuum dust)

Target compound	Sampling method	Average	Median	Minimum	Maximum	Standard deviation
BDE-47	RCDL	15.3	13.0	< 0.1	44.8	11.9
	RCDB	12.0	11.1	< 0.1	31.9	7.8
	HHVD	7.4	6.8	< 0.1	14.3	5.1
BDE-99	RCDL	23.3	17.9	4.2	77.1	18.7
	RCDB	19.5	14.5	< 0.2	88.6	22.7
	HHVD	11.8	12.0	7.7	16.1	2.6
BDE-153	RCDL	4.3	4.1	< 0.2	7.3	1.8
	RCDB	4.8	4.0	1.4	14.7	3.6
	HHVD	2.5	2.9	< 0.2	5.9	2.1
Σ_6 tri-hexa-BDEs	RCDL	47.3	40.7	6.8	135	32.6
	RCDB	41.4	33.5	8.9	147	35.4
	HHVD	24.4	22.4	12.3	41.5	9.1
BDE-183	RCDL	5.7	6.0	< 0.2	11.3	3.4
	RCDB	2.4	2.8	< 0.2	5.1	1.7
	HHVD	7.2	2.5	< 0.2	61.2	17.1
BDE-209	RCDL	2642	3066	466	4184	1354
	RCDB	2336	2232	1175	3944	780
	HHVD	2634	2462	1534	3779	802
Σ_8 PBDEs	RCDL	2695	3112	474	4321	1363
	RCDB	2380	2272	1233	3985	775
	HHVD	2666	2519	1568	3795	797

Table 2: Summary statistics for NBFR concentrations (ng/g) in RCDL, RCDB (researcher collected dust from the living room bedroom) and dust HHVD (household vacuum dust)

Target compound	Sampling method	Average	Median	Minimum	Maximum	Standard deviation
EH-TBB	RCDL	6.9	6.4	< 0.5	21.2	6.0
	RCDB	6.4	3.1	< 0.5	24.2	7.9
	HHVD	4.9	3.9	< 0.5	13.5	4.9
BTBPE	RCDL	11.0	11.2	< 2.8	21.4	7.4
	RCDB	9.5	9.8	< 2.8	15.8	4.9
	HHVD	11.2	8.0	< 2.8	35.7	11.5
BEH-TEBP	RCDL	306	175	64	1299	348
	RCDB	339	131	43	1139	380
	HHVD	233	121	33	890	256
DBDPE	RCDL	155	87	14	679	184
	RCDB	91	76	11	236	65
	HHVD	152	85	16	575	170
Σ_5 NBFRs	RCDL	479	394	127	1450	382
	RCDB	446	225	104	1412	420
	HHVD	402	272	129	1302	345

Table 3: Median and 95th percentile concentration ratios of BDE-99, Σ_6 tri-hexa-BDEs, BDE-209 and DBDPE between researcher collected-dust from the living room and bedroom (RCDL and RCDB) and household vacuum dust

Compound	Sampling approach	Median	95 th percentile
BDE-99	RCDL/HHVD	1.5	3.5
	RCDB/HHVD	1.2	3.4
Σ_6 tri-hexa-BDEs	RCDL/HHVD	1.8	2.7
	RCDB/HHVD	1.5	2.6
BDE-209	RCDL/HHVD	1.2	1.1
	RCDB/HHVD	0.9	1.0
BEH-TEBP	RCDL/HHVD	1.4	1.3
	RCDB/HHVD	1.1	1.5
DBDPE	RCDL/HHVD	1.0	1.0
	RCDB/HHVD	0.9	0.4