Pneumolysin mediates heterotypic aggregation of neutrophils and platelets *in vitro*

Nel, Jan G.; Durandt, Chrisna; Theron, Annette J.; Tintinger, Gregory R.; Pool, Roger; Richards, Guy A.; Mitchell, Timothy J.; Feldman, Charles; Anderson, Ronald

DOI:

[10.1016/j.jinf.2017.02.010](https://doi.org/10.1016/j.jinf.2017.02.010)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Nel, JG, Durandt, C, Theron, AJ, Tintinger, GR, Pool, R, Richards, GA, Mitchell, TJ, Feldman, C & Anderson, R 2017, 'Pneumolysin mediates heterotypic aggregation of neutrophils and platelets *in vitro*', *Journal of Infection*, vol. 74, no. 6, pp. 599-608. <https://doi.org/10.1016/j.jinf.2017.02.010>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Pneumolysin mediates heterotypic aggregation of neutrophils and platelets *in vitro*

Jan G. Nel, Chrisna Durandt, Annette J. Theron, Gregory R. Tintinger, Roger Pool, Guy A. Richards, Timothy J. Mitchell, Charles Feldman, Ronald Anderson

PII: S0163-4453(17)30076-2

DOI: [10.1016/j.jinf.2017.02.010](https://doi.org/10.1016/j.jinf.2017.02.010)

Reference: YJINF 3893

To appear in: *Journal of Infection*

Received Date: 30 November 2016

Revised Date: 17 February 2017

Accepted Date: 24 February 2017

Please cite this article as: Nel JG, Durandt C, Theron AJ, Tintinger GR, Pool R, Richards GA, Mitchell TJ, Feldman C, Anderson R, Pneumolysin mediates heterotypic aggregation of neutrophils and platelets *in vitro*, *Journal of Infection* (2017), doi: 10.1016/j.jinf.2017.02.010.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Pneumolysin mediates heterotypic aggregation of neutrophils and platelets *in vitro*

Jan G. Nel*, Chrisna Durandt⁺, Annette J. Theron^o, Gregory R. Tintinger[◇], Roger Pool*, Guy A. Richards[~], Timothy J. Mitchell[▲], Charles Feldman[#], Ronald Anderson^{o+}

^{*}Department of Haematology, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa and Tshwane Academic Division of the National Health Laboratory Service, Pretoria, South Africa

⁺Institute for Cellular and Molecular Medicine, South African Medical Research Council Unit for Stem Cell Research, Department of Immunology, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

^oDepartment of Immunology, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa and Tshwane Academic Division of the National Health Laboratory Service, Pretoria, South Africa

[◇]Department of Internal Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

[~]Department of Critical Care, Charlotte Maxeke Johannesburg Academic Hospital and Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

[▲]Institute of Microbiology and Infection, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

[#]Division of Pulmonology, Department of Internal Medicine, Charlotte Maxeke Johannesburg Academic Hospital and Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Running title Pneumolysin promotes neutrophil:platelet adhesion

Correspondence:

Dr J.G. Nel

Department of Haematology

PO Box 2034

Pretoria

0001

South Africa

Telephone:

+27-12-319-2641

Telefax:

+27-12-323-0732

E-mail:

jan.nel@up.ac.za

Abstract

Objectives. Platelets orchestrate the inflammatory activities of neutrophils, possibly contributing to pulmonary and myocardial damage during severe pneumococcal infection. This study tested the hypothesis that the pneumococcal toxin, pneumolysin (Ply), activates production of platelet-activating factor (PAF) and thromboxane A₂ (TxA₂) by neutrophils, these bioactive lipids being potential mediators of neutrophil:platelet (NP) networking.

Methods. The effects of recombinant Ply (10–80 ng.mL⁻¹) on the production of PAF and TxA₂ by isolated neutrophils were measured using ELISA procedures, and NP aggregation by flow cytometry.

Results. Exposure of neutrophils to Ply induced production of PAF and, to a lesser extent, TxA₂, achieving statistical significance at ≥ 20 ng.mL⁻¹ of the toxin. In the case of NP interactions, Ply promoted heterotypic aggregation which was dependent on upregulation of P-selectin (CD62P) and activation of protease-activated receptor 1 (PAR1), attaining statistical significance at ≥ 10 ng.mL⁻¹ of the toxin, but did not involve either PAF or TxA₂.

Conclusion. Ply induces synthesis of PAF and TxA₂ by human neutrophils, neither of which appears to contribute to the formation of NP heterotypic aggregates *in vitro*, a process which is seemingly dependent on CD62P and PAR1. These pro-inflammatory activities of Ply may contribute to the pathogenesis of pulmonary and myocardial injury during severe pneumococcal infection.

Keywords. Calcium, platelet-activating factor, pneumolysin, P-selectin (CD62P), severe pneumococcal disease.

Pneumolysin (Ply), the cholesterol-binding, pore-forming toxin of *Streptococcus pneumoniae*, is recognised as being the major protein virulence factor of this intransigent respiratory pathogen, the most common bacterial cause of community-acquired pneumonia (CAP) and associated organ damage (1-4). Importantly, Ply has been identified as being a key mediator of both acute lung injury (ALI) (5-8) and myocardial damage (9, 10) in murine models of severe pneumococcal disease. In one such model of ALI, exposure of isolated, perfused lungs to recombinant Ply resulted in the development of pulmonary hypertension and microvascular barrier dysfunction, both of which are key features of this condition in humans (6, 7). The underlying mechanisms appeared to involve increased pulmonary production of the bioactive lipid, platelet-activating factor (PAF), which, in turn, was proposed to activate production of the more potent platelet activator *viz.* the prostanoid, thromboxane A₂ (TxA₂), with resultant vasoconstriction and platelet activation (7). Although the authors speculated that PAF may have originated from Ply-exposed endothelial cells (7), infiltrating neutrophils represent an alternative source of the bioactive lipid. Unlike macrophages, neutrophils express high levels of the PAF-generating enzyme, PAF acetylhydrolase (11). However, to our knowledge a possible link between Ply, neutrophils, PAF and platelet activation has not been described.

In the context of acute cardiovascular events associated with invasive pneumococcal disease, Ply, via its pore-forming activity, has been reported to inflict injury on myocardium through the formation of cardiac microlesions (9, 10). However, the existence of alternative mechanisms of Ply-mediated cardiotoxicity, possibly related to the pro-inflammatory/pro-thrombotic activities of the toxin are largely unexplored (12-14).

To probe the existence of such mechanisms in the pathogenesis of Ply-mediated ALI and myocardial injury, we have investigated the effects of recombinant Ply on the production of PAF and TxA₂ by isolated, human blood neutrophils *in vitro*. In addition, we have also explored the effects of Ply on the formation of potentially, pro-thrombotic, heterotypic aggregates of neutrophils and platelets (15-19), focusing on the involvement of PAF, TxA₂ and other potent platelet activators, as well as the adhesion molecule, P-selectin (CD62P), in this process.

MATERIALS AND METHODS

Permission to draw blood from healthy, adult human volunteers was granted by the Research Ethics Committee, Faculty of Health Sciences, University of Pretoria.

Pneumolysin

Recombinant Ply and the pneumolysoid, delta6Ply, attenuated in respect of pore-forming activity, were prepared as described previously (20, 21). The possible influence of contaminating endotoxin was excluded in both Ply preparations using the Endosafe[®]-PTS[™] system (Charles River Laboratories, Wilmington, MA, USA) which is based on the *Limulus* amoebocyte lysate kinetic chromogenic method. Both active Ply and delta6Ply contained <1 endotoxin unit (EU)/ μ g of protein after purification (1 EU is the lower limit of detection).

Chemicals

PSB 0739, WEB 2086, S 18886 and SCH 79797 antagonists of the platelet purinergic receptor, P2Y₁₂, the PAF receptor, the TxA₂ TP prostanoid receptor, and the protease-activated receptor 1 (PAR1, thrombin activated) respectively were purchased from TOCRIS Bioscience, Bristol, UK. The oral thrombin inhibitor, dabigatran, was provided to one of us (GAR) by Boehringer-Ingelheim Pharma GmbH, Germany. All of these were dissolved to stock solutions of 10 mM in dimethylsulphoxide (DMSO) and used at final concentrations of 10 μ M (final DMSO concentrations of 0.1%) in the assays described below. Appropriate DMSO control systems were included in all of the assays in which these receptor antagonists were used.

Mouse anti-human C62P blocking antibody (P-selectin, non-fluorochrome-labelled), was purchased from Biolegend, London, UK. Unless stated, all other chemicals and reagents were purchased from the Sigma Chemical Co., St. Louis, MO, USA.

Preparation of neutrophils

Neutrophils were prepared from heparinised venous blood (5 units preservative-free heparin.mL⁻¹) as described previously (22). Briefly, neutrophil/erythrocyte pellets obtained following centrifugation of whole blood on Histopaque-1077 (Sigma Diagnostics) were resuspended and sedimented in 3% gelatin to remove most of the erythrocytes. Residual erythrocytes were then removed by differential lysis (brief treatment with 0.83% ammonium chloride) and the resultant neutrophil populations of high purity and viability (>90% and >95% respectively) suspended to a concentration of 1×10^7 cells.mL⁻¹ in Hanks' balanced salt solution (HBSS, indicator-free, pH7.4).

PAF and TxA₂

Neutrophils (2×10^6) suspended in HBSS were prewarmed for 10 min at 37°C followed by addition of one of the following: i) HBSS (negative control); ii) recombinant Ply at final concentrations of 10, 20, 40 and 80 nanograms (ng).mL⁻¹; iii) the pneumolysoid, delta6Ply, which is attenuated with respect to pore-forming activity, at a fixed, final concentration of 80 ng.mL⁻¹; or iv) the calcium ionophore, A23187 at 2 μ M (final) as a positive control system. The final volume in each test tube was 2 mL. After a further 5 min period of incubation at 37°C, the tubes were transferred to an ice-bath to stop the reactions. Following removal of the cells by centrifugation, the concentrations of PAF and TxA₂ in the cell-free supernatants were measured using commercial sandwich ELISA procedures (Cusabio® Life Science, Wuhan, P.R. China and Abnova GmbH, Heidelberg, Germany respectively) and the results expressed as ng.mL⁻¹ and picograms (pg).mL⁻¹ respectively. Cell viability was measured using a propidium iodide-based flow cytometric procedure.

Neutrophil:platelet (NP) aggregate formation

In order to minimise spontaneous activation of platelets, NP-enriched buffy coat suspensions, enumerated for both cell types by standard haematological procedures, were used for these studies in keeping with earlier reports which used whole blood (23-25). These cell suspensions were prepared from the heparinised

blood of healthy, adult humans by sedimentation at 37°C and diluted 1:50 in HBSS to give a final volume of 1 ml. Following 5 min of preincubation at 37°C, recombinant Ply (10–80 ng.mL⁻¹), delta6Ply (80 ng.mL⁻¹), or adenosine 5'-diphosphate (ADP, 100 µM final, agonist of platelet P2Y12 receptors as a positive control) were added to the cell suspensions which were incubated for a further 5 min at 37°C. Following incubation, the cell suspensions were stained with 5 µl of each of the following murine, anti-human, fluorochrome-labelled monoclonal antibodies to detect neutrophils, platelets and total leukocytes: CD16-allophycocyanin (Biolegend, San Diego, CA, USA), CD42a-phycoerythrin (Becton Dickinson, San Jose, CA, USA) and CD45-Krome Orange (Beckman Coulter, Marseille, France) and incubated for 15 min at room temperature in the dark. This was followed by analysis of the various cell suspensions at a slow flow rate using a Gallios flow cytometer (Beckman Coulter, Miami, USA). NP interactions were determined according to the CD16⁺/CD42a⁺ co-expression profiles of CD45⁺ leukocytes and the results expressed as the relative median fluorescence intensities of CD42a expression of these NP aggregates. Platelet aggregates were excluded prior to the aforementioned analysis as indicated in Figure 1 which depicts this gating strategy. Note that residual erythrocytes [confirmed by staining with an anti-CD 235a (glycophorin) monoclonal antibody, Becton Dickinson] in the cell suspensions were not lysed prior to flow cytometric analysis to minimise non-specific activation of platelets.

In a limited series of experiments (2 in the series) undertaken to ensure the veracity of the various antagonists of platelet P2Y12 and PAF receptors and PAR1 (PSB 0739, WEB 2086, and SCH 79797 respectively), these agents were added to platelet-rich plasma which was incubated for 5 min at 37°C prior to the addition of the respective receptor agonists, ADP (100 µM), PAF (400 nM), or thrombin (from human plasma, 1.25 NIH units, final). After a further period of incubation for 5 min at 37°C platelet activation was measured flow cytometrically as described previously according to upregulated expression of the adhesion molecule, CD62P (P-selectin) (14, 25).

The following series of experiments explored the effects of the various platelet receptor (P2Y12, PAF, PAR1, TxA₂) antagonists (all at 10 µM), as well as those of

indomethacin (5 μM) and a mouse anti-human CD62 P blocking monoclonal antibody (5 μL per mL of cell suspension), all added prior to preincubation, on NP aggregation activated by Ply (40 ng.mL^{-1} , final) measured as described above. The following were also investigated: i) the requirement for extracellular Ca^{2+} in the pro-aggregation activity of Ply; ii) the specificity of the PAR1 receptor antagonist, SCH 79797, which was assessed by measuring the effects of this agent on NP aggregation induced by purified thrombin (1.25 NIH units), as well as on spontaneous aggregation and that activated by ADP (100 μM); and iii) the effects of dabigatran (10 μM) on NP aggregation induced by either thrombin or Ply (40 ng.mL^{-1} , final)

Expression and statistical analysis of results

The results of each series of experiments are expressed as median values with interquartile ranges with numbers of different donors and experiments indicated in the text or figure legends. Statistical analyses were performed using GraphPad Prism5 (GraphPad Software, San Diego, USA) using a one-way ANOVA with a Bonferroni correction for multiple comparisons.

RESULTS

Production of PAF and TxA_2 by Ply-activated neutrophils

These results are shown in figures 2A and 2B for PAF and TxA_2 respectively. As shown in figure 2A, exposure of neutrophils to Ply resulted in dose-related activation of generation of PAF which achieved statistical significance at concentrations $\geq 20 \text{ ng.mL}^{-1}$ of the toxin, while the non-physiological positive control, A23187, as expected was extremely potent, and delta6Ply ineffective. The corresponding data for TxA_2 production by neutrophils are shown in figure 2B, which demonstrate similar, albeit lesser, effects.

Neutrophil viability

These results are shown in Figure 3. Exposure of neutrophils to the highest concentrations of Ply (40 and 80 ng.mL⁻¹) or to A23187 caused modest, but nevertheless statistically significant, loss of viability. The median viability values for the control, untreated system and for systems treated with Ply at 40 and 80 ng.mL⁻¹ or A23187 were 98.9% (IQR 98.5-99.4%), 93.9% (p<0,005), 91.0% (p<0,001), and 95.4% (p<0.01) respectively (data from 4 experiments using cells from 4 different donors).

Effect of Ply on the formation of NP heterotypic aggregates

The median neutrophil and platelet counts of the buffy coats used in these and subsequent experiments were $4.67 \times 10^3 \cdot \mu\text{L}^{-1}$ and $451 \times 10^3 \cdot \mu\text{L}^{-1}$ respectively, and the results are shown in Figure 3. Addition of Ply to mixed NP suspensions resulted in dose-related formation of NP aggregates which achieved statistical significance at concentrations of $\geq 20 \text{ ng.mL}^{-1}$ of the toxin and was greater than that observed with ADP, while delta6Ply was ineffective (Figure 4).

Assessment of the veracity of the various platelet receptor antagonists

Prior to assessing their effects on Ply-mediated NP aggregate formation, the efficacy of the various platelet-receptor antagonists (PSB 0739, WEB 2086, SCH 79797, all at 10 μM) was measured in a series of preliminary experiments, using platelet-rich plasma. Following addition of the corresponding, respective receptor agonists ADP (100 μM), PAF (400 nM) or thrombin (1.25 NIH units), platelet activation was measured flow cytometrically according to the level of expression of the adhesion molecule, CD62P. The results, which are shown in Figure 5, demonstrate the activities of the receptor agonists, with PAF being the least potent, as well as the inhibitory activities of the various receptor antagonists. Importantly, upregulation of expression of CD62P by ADP-, PAF- or thrombin-treated platelets

was significantly attenuated by PSB 0739, WEB 2086, and SCH 79797, confirming receptor antagonism.

Effects of the various platelet receptor antagonists, indomethacin, Ca^{2+} depletion, and an anti-CD62P monoclonal antibody on Ply-mediated NP aggregation

The effects of the various platelet receptor antagonists and indomethacin, as well as those of suspension of the cells in Ca^{2+} -free HBSS, on Ply (40 ng.mL⁻¹)-activated formation of NP aggregates are shown in Figure 6A, while those of inclusion of the anti-human CD62P blocking monoclonal antibody are shown in Figure 6B. Exposure of the cells to the PAR1 antagonist, SCH 79797, as well as suspension of the cells in Ca^{2+} -free medium resulted in significant attenuation of Ply-activated formation of NP aggregates, while the other receptor antagonists and indomethacin were ineffective (Figure 6A). Inclusion of the anti-CD62P antibody caused almost complete attenuation of Ply-mediated NP aggregate formation (Figure 6B). Depletion of Ca^{2+} , as well as inclusion of the anti-CD62P antibody also caused significant reductions in basal NP aggregation, underscoring the involvement of both Ca^{2+} and CD62P in basal aggregation. These observations demonstrate significant involvement of CD62P, as well as PAR1, but not the P2Y₁₂, PAF or TxA₂ receptors in Ply-mediated NP aggregate formation.

Effects of SCH 79797 on spontaneous, ADP- and thrombin-activated NP aggregation

To probe the receptor-targeted veracity of SCH 79797 (10 μ M) in the context of NP aggregate formation, the effects of this agent on spontaneous, ADP- or thrombin-activated NP aggregation were investigated and these results are shown in Figure 7. Addition of SCH 79797 to the mixed neutrophil and platelet suspensions during pre-incubation resulted in statistically significant formation of thrombin-activated heterotypic aggregates, but had no effect on either spontaneous or ADP-activated formation of NP aggregates. These findings confirm the selectivity of SCH 79797 for PAR1 and the probable involvement of this receptor in Ply-mediated NP aggregate formation, possibly via thrombin activation.

Effect of dabigatran on Ply-mediated NP aggregation

The thrombin inhibitor, dabigatran, was used to explore the possible involvement of thrombin in Ply-mediated activation of PAR1 and these results are shown in Figure 8. Dabigatran was found to attenuate thrombin-, but not Ply-mediated formation of NP aggregates, apparently excluding the involvement of thrombin derived from either the plasma or cellular elements of the buffy coat preparations in Ply-activated NP aggregation.

Effect of SCH 79797 on the pore-forming activity of Ply

An erythrocyte haemolysis assay was used to exclude possible interference of SCH 79797 with the pore-forming activity of Ply. Erythrocytes are particularly vulnerable to the lytic action of Ply. Briefly, the toxin (20 ng.mL^{-1}) was pre-incubated with SCH 79797 ($10 \text{ }\mu\text{M}$) for 5 min at 37°C followed by the addition of a 0.5% suspension of human erythrocytes in a final volume of 1 mL HBSS. Following 5 min incubation, the remaining erythrocytes were pelleted by centrifugation and haemoglobin in the supernatant fluids measured spectrophotometrically at a wavelength of 490 nm. The mean percentages haemolysis of Ply-treated erythrocytes in the absence or presence of SCH 79797 were 24% and 25% respectively (NS), clearly indicating lack of interference of the PAR1 antagonist with the pore-forming activity of Ply.

DISCUSSION

The results of the current study demonstrate that exposure of neutrophils to Ply, at concentrations representative of those measured in the cerebrospinal fluid of patients with pneumococcal meningitis (26), caused dose-related generation of production of PAF, reaching levels which were about 3-fold higher than those of the untreated control system at the highest concentrations tested (40 and 80 ng.mL⁻¹). The pneumolysoid, delta6Ply, was ineffective, while the calcium ionophore, A23187, included as a positive control system, was more potent than Ply. Similar, but less impressive trends were observed in the case of TxA₂, possibly indicative of intense competition for arachidonic acid by the range of prostanoid/eicosanoid/PAF-generating enzymes present in activated neutrophils. Although not shown, similar effects were observed with TxB₂, excluding conversion of TxA₂ to TxB₂ as a possible cause of the lesser effect of Ply on production of TxA₂ by neutrophils relative to PAF. Although exposure of neutrophils to the toxin at concentrations of 40 and 80 ng.mL⁻¹ resulted in loss of viability, these effects were modest and unlikely to have contributed to the observed activation of production of PAF and TxA₂. In this context, it is noteworthy that some types of mammalian cell can withstand the cytotoxic actions of Ply due to the existence of a mechanism which promotes microvesicle shedding of toxin pores (27). Ply-mediated pore formation in the plasma membrane of inflammatory cells does, however, result in an influx of extracellular Ca²⁺ which either activates or sensitises the cells for increased pro-inflammatory activity (12-14).

A possible association between the production of PAF and TxA₂ by Ply-treated neutrophils and activation of neighbouring platelets was explored by investigating the effects of the toxin on the heterotypic aggregation of these cells in the absence and presence of a PAF or TP receptor antagonist, as well as antagonists of other types of receptor which mediate platelet activation, these being P2Y₁₂ and PAR1. Exposure of mixed NP suspensions to Ply resulted in significant dose-related heterotypic aggregation of these cells which was maximal at 40-80 ng.mL⁻¹ of the toxin, exceeding that observed with ADP, and dependent on the presence of extracellular Ca²⁺, while delta6Ply was ineffective. With respect to the effects of the various receptor antagonists, only SCH 79797, somewhat surprisingly, was found to attenuate Ply-mediated NP heterotypic aggregation, while blockade of

the PAF, P2Y₁₂, and TP receptors, as well as inhibition of cyclooxygenases with indomethacin, were all ineffective. The selectivity of SCH 79797 for the PAR₁ was confirmed by the absence of effects of this agent on either spontaneous or ADP-activated NP aggregation. In addition, SCH 79797 did not interfere with the pore-forming activity of Ply, excluding non-specific inactivation of the toxin as a possible mechanism of interference with NP aggregation. In this context it is noteworthy that antagonism of PAR₁ has recently been reported to decrease the levels of pulmonary, pro-inflammatory cytokines/chemokines and to attenuate alveolar leak in a murine model of experimental pneumococcal pneumonia (28).

The apparent involvement of triggering of PAR₁ on platelets in NP heterotypic aggregation was an unexpected finding, which is most likely a secondary, albeit important, amplification mechanism resulting from interaction of the receptor with putative activators derived from Ply-activated platelets and/or neutrophils, reinforcing and sustaining NP adhesion. Possible contenders include prothrombin released from platelet α -granules (29), which may be converted to thrombin by the action of pro-thrombinase expressed on neighbouring monocytes (30). Activation of PAR₄ which is also expressed on platelets and activated by thrombin, albeit at a slower rate than PAR₁, may also contribute to NP aggregation (31). However, the lack of an effect of the thrombin inhibitor, dabigatran, on Ply-mediated NP aggregation appears to exclude any meaningful involvement of thrombin activation of PARs. An alternative, albeit unexplored mechanism, implicates the serine proteinases, elastase and proteinase 3, as well as the matrix metalloproteinases 8 and 9 expressed by Ply-exposed adherent neutrophils (13, 32) all of which are known activators of PAR₁ (33, 34), while cathepsin G has been reported to activate PAR₄ (35). Addressing this issue is, however, beyond the scope of the current study given the spectrum of neutrophil-derived proteinases and their probable interactions, compounded by the requisite large number of enzyme inhibitors.

Together with the observation that inclusion of an anti-CD62P monoclonal antibody caused almost complete attenuation of Ply-mediated NP aggregate formation, the aforementioned observations appear to be consistent with a sequence of events whereby exposure of platelets to Ply results in influx of extracellular Ca²⁺,

as described previously (14), Ca^{2+} -dependent mobilisation of α -granules, upregulated surface expression of CD62P, and adhesion of neighbouring neutrophils. In this context, interactions between CD62P expressed on platelets and its counter ligand, P-selectin glycoprotein ligand-1 (PSGL-1) expressed on platelets and other cell types, are considered to be the primary mediators of platelet homotypic and heterotypic aggregation (15, 17, 24, 36). Although platelet-derived CD40 ligand has also been reported to mediate this type of interaction, the results of the current study appear to implicate CD62P as being the major player in the pro-adhesive actions of Ply (37, 38). Although speculative, initial CD62P-dependent NP adhesion is then reinforced by neutrophil proteinase-mediated activation of platelet PAR1, resulting in the formation of more stable NP aggregates. Given that endothelial cells also express PAR1 (39), it is likely, albeit unexplored that exposure of endothelium to Ply also results in Ca^{2+} influx and upregulation of endothelial CD62P. This, in turn, may promote the binding of neutrophils and NP aggregates favouring activation of endothelial PAR1 and endothelial dysfunction (39).

In agreement with the findings of the current study, CD62P-dependent formation of NP aggregates following exposure of whole blood to the bacterial pore-forming toxins, streptolysin-O or *Staphylococcus aureus* α -hemolysin, has been described previously (23, 24). In the case of the former, the authors proposed a link between streptolysin-O production, formation of NP aggregates, and vascular occlusions and tissue damage during infection with group A *streptococci* (23). In the case of α -hemolysin, heterotypic aggregate formation was linked to alveolar capillary destruction in haemorrhagic/necrotising pneumonia caused by community-associated, methicillin-resistant *S. aureus* (24). However, unlike the current study, neither of these earlier studies, investigated the pathophysiological mechanisms underpinning toxin-mediated aggregate formation. Very recently, Zhang *et al.* in a study focused primarily on the *S. suis* pore-forming toxin, suilysin, reported that this toxin, as well as Ply, promoted NP aggregation *in vitro* by a Ca^{2+} - and P-selectin-dependent mechanism as described in the current study (40). However, in the study reported by Zhang *et al.* Ply was used at a concentration considerably higher (800 ng.mL^{-1} , fixed) than those used in the current study (10-80 ng.mL^{-1}), while these

446 authors did not investigate the involvement of platelet-activating receptors in either
447 suilysin- or Ply-mediated NP aggregation (40)

448
449 While the exact clinical significance of the findings of the current study await
450 clarification, they do, however, imply a multifaceted role for Ply in the pathogenesis
451 of lung, heart and other types of organ damage during severe pneumococcal
452 disease. Notwithstanding direct Ply-mediated organ damage (6-10), the effects of the
453 toxin described here are also consistent with a pathogenic role for Ply-mediated
454 formation of large, intravascular NP aggregates with resultant microvascular
455 occlusion. Importantly, activated platelets and NP aggregates may also promote
456 tissue injury by amplifying the inflammatory response. On a cautionary note,
457 however, should these harmful activities of the toxin be evident in the clinical setting,
458 therapeutic targeting may prove difficult given the drawbacks and side effect profile
459 of a commercially available PAR1 antagonist, vorapaxar (41), as well as the current
460 lack of pharmacological agents which directly inhibit Ply. In this context, inhibitors of
461 bacterial protein synthesis, especially macrolide antibiotics, may offer the best
462 therapeutic option (42).

REFERENCES

1. Mitchell TJ, Dalziel CE. The biology of pneumolysin. *Subcell Biochem.* 2014;80:145-60. PubMed PMID: 24798011. Epub 2014/05/07. eng.
2. File TM, Jr., Marrie TJ. Burden of community-acquired pneumonia in North American adults. *Postgrad Med.* 2010 Mar;122(2):130-41. PubMed PMID: 20203464. Epub 2010/03/06. eng.
3. Welte T, Torres A, Nathwani D. Clinical and economic burden of community-acquired pneumonia among adults in Europe. *Thorax.* 2012 Jan;67(1):71-9. PubMed PMID: 20729232. Epub 2010/08/24. eng.
4. Musher DM, Rueda AM, Kaka AS, Mapara SM. The association between pneumococcal pneumonia and acute cardiac events. *Clin Infect Dis.* 2007 Jul 15;45(2):158-65. PubMed PMID: 17578773. Epub 2007/06/21. eng.
5. Feldman C, Mitchell TJ, Andrew PW, Boulnois GJ, Read RC, Todd HC, et al. The effect of *Streptococcus pneumoniae* pneumolysin on human respiratory epithelium in vitro. *Microb Pathog.* 1990 Oct;9(4):275-84. PubMed PMID: 2097494. Epub 1990/10/01. eng.
6. Witzernath M, Gutbier B, Hocke AC, Schmeck B, Hippenstiel S, Berger K, et al. Role of pneumolysin for the development of acute lung injury in pneumococcal pneumonia. *Crit Care Med.* 2006 Jul;34(7):1947-54. PubMed PMID: 16715037. Epub 2006/05/23. eng.
7. Witzernath M, Gutbier B, Owen JS, Schmeck B, Mitchell TJ, Mayer K, et al. Role of platelet-activating factor in pneumolysin-induced acute lung injury. *Crit Care Med.* 2007 Jul;35(7):1756-62. PubMed PMID: 17522574. Epub 2007/05/25. eng.
8. Garcia-Suarez Mdel M, Florez N, Astudillo A, Vazquez F, Villaverde R, Fabrizio K, et al. The role of pneumolysin in mediating lung damage in a lethal pneumococcal pneumonia murine model. *Respir Res.* 2007;8:3. PubMed PMID: 17257395. Pubmed Central PMCID: 1790890. Epub 2007/01/30. eng.
9. Brown AO, Mann B, Gao G, Hankins JS, Humann J, Giardina J, et al. *Streptococcus pneumoniae* translocates into the myocardium and forms unique microlesions that disrupt cardiac function. *PLoS Pathog.* 2014 Sep;10(9):e1004383. PubMed PMID: 25232870. Pubmed Central PMCID: 4169480.
10. Alhamdi Y, Neill DR, Abrams ST, Malak HA, Yahya R, Barrett-Jolley R, et al. Circulating Pneumolysin Is a Potent Inducer of Cardiac Injury during Pneumococcal Infection. *PLoS Pathog.* 2015 May;11(5):e1004836. PubMed PMID: 25973949. Pubmed Central PMCID: 4431880. Epub 2015/05/15. eng.
11. Howard KM. Differential expression of platelet-activating factor acetylhydrolase in lung macrophages. *Am J Physiol Lung Cell Mol Physiol.* 2009 Dec;297(6):L1141-50. PubMed PMID: 19837851. Pubmed Central PMCID: Pmc2793189. Epub 2009/10/20. eng.
12. Cockeran R, Durandt C, Feldman C, Mitchell TJ, Anderson R. Pneumolysin activates the synthesis and release of interleukin-8 by human neutrophils in vitro. *J Infect Dis.* 2002 Aug 15;186(4):562-5. PubMed PMID: 12195386. Epub 2002/08/27. eng.
13. Cockeran R, Theron AJ, Steel HC, Matlola NM, Mitchell TJ, Feldman C, et al. Proinflammatory interactions of pneumolysin with human neutrophils. *J Infect Dis.* 2001 Feb 15;183(4):604-11. PubMed PMID: 11170986. Epub 2001/02/15. eng.
14. Nel JG, Durandt C, Mitchell TJ, Feldman C, Anderson R, Tintinger GR. Pneumolysin Mediates Platelet Activation In Vitro. *Lung.* 2016 Aug;194(4):589-93. PubMed PMID: 27192991.
15. Yokoyama S, Ikeda H, Haramaki N, Yasukawa H, Murohara T, Imaizumi T. Platelet P-selectin plays an important role in arterial thrombogenesis by forming large stable platelet-leukocyte aggregates. *J Am Coll Cardiol.* 2005 4/19;45(8):1280-6.
16. Ed Rainger G, Chimen M, Harrison MJ, Yates CM, Harrison P, Watson SP, et al. The role of platelets in the recruitment of leukocytes during vascular disease. *Platelets.* 2015;26(6):507-20. PubMed PMID: 26196409. Epub 2015/07/22. eng.
17. Sreeramkumar V, Adrover JM, Ballesteros I, Cuartero MI, Rossaint J, Bilbao I, et al. Neutrophils scan for activated platelets to initiate inflammation. *Science.* 2014 Dec

- 516 5;346(6214):1234-8. PubMed PMID: 25477463. Pubmed Central PMCID: Pmc4280847. Epub
517 2014/12/06. eng.
- 518 18. McMahon TJ. Platelets promote pulmonary pull of polys. *Blood*. 2015 Nov 5;126(19):2174-5.
519 PubMed PMID: 26542252. Pubmed Central PMCID: PMC4635113. Epub 2015/11/07. eng.
- 520 19. Guo Y, Mishra A, Howland E, Zhao C, Shukla D, Weng T, et al. Platelet-derived Wnt
521 antagonist Dickkopf-1 is implicated in ICAM-1/VCAM-1-mediated neutrophilic acute lung
522 inflammation. *Blood*. 2015 Nov 5;126(19):2220-9. PubMed PMID: 26351298. Pubmed Central
523 PMCID: PMC4635118. Epub 2015/09/10. eng.
- 524 20. Mitchell TJ, Walker JA, Saunders FK, Andrew PW, Boulnois GJ. Expression of the pneumolysin
525 gene in *Escherichia coli*: rapid purification and biological properties. *Biochim Biophys Acta*. 1989 Jan
526 23;1007(1):67-72. PubMed PMID: 2642385. Epub 1989/01/23. eng.
- 527 21. Kirkham LA, Kerr AR, Douce GR, Paterson GK, Dilts DA, Liu DF, et al. Construction and
528 immunological characterization of a novel nontoxic protective pneumolysin mutant for use in future
529 pneumococcal vaccines. *Infect Immun*. 2006 Jan;74(1):586-93. PubMed PMID: 16369015. Pubmed
530 Central PMCID: 1346677. Epub 2005/12/22. eng.
- 531 22. Gravett CM, Theron AJ, Steel HC, Tintinger GR, Cockeran R, Feldman C, et al. Interactive
532 inhibitory effects of formoterol and montelukast on activated human neutrophils. *Eur Respir J*. 2010
533 Dec;36(6):1417-24. PubMed PMID: 20413544. Epub 2010/04/24. eng.
- 534 23. Bryant AE, Bayer CR, Chen RY, Guth PH, Wallace RJ, Stevens DL. Vascular dysfunction and
535 ischemic destruction of tissue in *Streptococcus pyogenes* infection: the role of streptolysin O-
536 induced platelet/neutrophil complexes. *J Infect Dis*. 2005 Sep 15;192(6):1014-22. PubMed PMID:
537 16107954. Epub 2005/08/19. eng.
- 538 24. Parimon T, Li Z, Bolz DD, McIndoo ER, Bayer CR, Stevens DL, et al. *Staphylococcus aureus*
539 alpha-hemolysin promotes platelet-neutrophil aggregate formation. *J Infect Dis*. 2013 Sep
540 1;208(5):761-70. PubMed PMID: 23698812. Pubmed Central PMCID: PMC3733505. Epub
541 2013/05/24. eng.
- 542 25. Mauler M, Seyfert J, Haenel D, Seeba H, Guenther J, Stallmann D, et al. Platelet-neutrophil
543 complex formation-a detailed in vitro analysis of murine and human blood samples. *J Leukoc Biol*.
544 2015 Nov 17. PubMed PMID: 26578648. Epub 2015/11/19. Eng.
- 545 26. Wall EC, Gordon SB, Hussain S, Goonetilleke UR, Gritzfeld J, Scarborough M, et al.
546 Persistence of pneumolysin in the cerebrospinal fluid of patients with pneumococcal meningitis is
547 associated with mortality. *Clin Infect Dis*. 2012 Mar 1;54(5):701-5. PubMed PMID: 22238165.
548 Pubmed Central PMCID: Pmc3275762. Epub 2012/01/13. eng.
- 549 27. Wolfmeier H, Schoenauer R, Atanassoff AP, Neill DR, Kadioglu A, Draeger A, et al. Ca²⁺(+)-
550 dependent repair of pneumolysin pores: A new paradigm for host cellular defense against bacterial
551 pore-forming toxins. *Biochim Biophys Acta*. 2015 Sep;1853(9):2045-54. PubMed PMID: 25219550.
552 Epub 2014/09/16. Eng.
- 553 28. Jose RJ, Williams AE, Mercer PF, Sulikowski MG, Brown JS, Chambers RC. Regulation of
554 neutrophilic inflammation by proteinase-activated receptor 1 during bacterial pulmonary infection. *J*
555 *Immunol*. 2015 Jun 15;194(12):6024-34. PubMed PMID: 25948816. Pubmed Central PMCID:
556 PMC4456635. Epub 2015/05/08. Eng.
- 557 29. Blair P, Flaumenhaft R. Platelet alpha-granules: basic biology and clinical correlates. *Blood*
558 *Rev*. 2009 Jul;23(4):177-89. PubMed PMID: 19450911. Pubmed Central PMCID: PMC2720568.
- 559 30. Robinson RA, Worfolk L, Tracy PB. Endotoxin enhances the expression of monocyte
560 prothrombinase activity. *Blood*. 1992 Jan 15;79(2):406-16. PubMed PMID: 1730086. Epub
561 1992/01/15. eng.
- 562 31. Nieman MT. Protease-activated receptors in hemostasis. *Blood*. 2016 Jul 14;128(2):169-77.
563 PubMed PMID: 27127302. Pubmed Central PMCID: PMC4946198.
- 564 32. Cockeran R, Mitchell TJ, Feldman C, Anderson R. Pneumolysin induces release of matrix
565 metalloproteinase-8 and -9 from human neutrophils. *Eur Respir J*. 2009 Nov;34(5):1167-70. PubMed
566 PMID: 19880617. Epub 2009/11/03. eng.

33. Mihara K, Ramachandran R, Renaux B, Saifeddine M, Hollenberg MD. Neutrophil elastase and proteinase-3 trigger G protein-biased signaling through proteinase-activated receptor-1 (PAR1). *J Biol Chem*. 2013 Nov 15;288(46):32979-90. PubMed PMID: 24052258. Pubmed Central PMCID: PMC3829148. Epub 2013/09/21. eng.
34. Lee EJ, Woo MS, Moon PG, Baek MC, Choi IY, Kim WK, et al. Alpha-synuclein activates microglia by inducing the expressions of matrix metalloproteinases and the subsequent activation of protease-activated receptor-1. *J Immunol*. 2010 Jul 1;185(1):615-23. PubMed PMID: 20511551. Epub 2010/06/01. Eng.
35. Sambrano GR, Huang W, Faruqi T, Mahrus S, Craik C, Coughlin SR. Cathepsin G activates protease-activated receptor-4 in human platelets. *J Biol Chem*. 2000 Mar 10;275(10):6819-23. PubMed PMID: 10702240. Epub 2000/03/04. eng.
36. de Bruijne-Admiraal LG, Modderman PW, Von dem Borne AE, Sonnenberg A. P-selectin mediates Ca(2+)-dependent adhesion of activated platelets to many different types of leukocytes: detection by flow cytometry. *Blood*. 1992 Jul 1;80(1):134-42. PubMed PMID: 1377047. Epub 1992/07/01. eng.
37. Andre P, Nannizzi-Alaimo L, Prasad SK, Phillips DR. Platelet-derived CD40L: the switch-hitting player of cardiovascular disease. *Circulation*. 2002 Aug 20;106(8):896-9. PubMed PMID: 12186789. Epub 2002/08/21. eng.
38. Gerdes N, Seijkens T, Lievens D, Kuijpers MJ, Winkels H, Projahn D, et al. Platelet CD40 Exacerbates Atherosclerosis by Transcellular Activation of Endothelial Cells and Leukocytes. *Arterioscler Thromb Vasc Biol*. 2016 Mar;36(3):482-90. PubMed PMID: 26821950. Epub 2016/01/30. eng.
39. Jose R, Williams A, Sulikowski M, Brealey D, Brown J, Chambers R. Regulation of neutrophilic inflammation in lung injury induced by community-acquired pneumonia. *Lancet*. 2015 Feb 26;385 Suppl 1:S52. PubMed PMID: 26312874. Epub 2015/08/28. eng.
40. Zhang S, Zheng Y, Chen S, Huang S, Liu K, Lv Q, et al. Suilysin-induced Platelet-Neutrophil Complexes Formation is Triggered by Pore Formation-dependent Calcium Influx. *Sci Rep*. 2016 Nov 10;6:36787. PubMed PMID: 27830834. Epub 2016/11/11. Eng.
41. Sharma A, Helft G, Garg A, Agrawal S, Chatterjee S, Lavie CJ, et al. Safety and efficacy of vorapaxar in secondary prevention of atherosclerotic disease: A meta-analysis of randomized control trials. *Int J Cardiol*. 2017 Jan 15;227:617-24. PubMed PMID: 27810296. Epub 2016/11/05. eng.
42. Anderson R, Steel HC, Cockran R, von Gottberg A, de Gouveia L, Klugman KP, et al. Comparison of the effects of macrolides, amoxicillin, ceftriaxone, doxycycline, tobramycin and fluoroquinolones, on the production of pneumolysin by *Streptococcus pneumoniae* in vitro. *J Antimicrob Chemother*. 2007 Nov;60(5):1155-8. PubMed PMID: 17848373. Epub 2007/09/13. eng.

Figure legends:

Figure 1.

Illustration of the gating strategy followed to identify platelet:neutrophil aggregates. Large aggregates were excluded using a Forward Scatter Area vs Forward Scatter Height plot. This was followed by the identification of CD45⁺ leukocytes, after which CD16⁺ neutrophils were identified. The expression intensity of CD42a (marker for platelets) was then measured on the CD45⁺/CD16⁺/CD42a⁺ cells. An erythrocyte lysis step was not included in order to minimise non-specific activation of platelets.

Figure 2.

The effects of addition of pneumolysin (Ply) at concentrations of 10-80 ng.ml⁻¹, as well as those of delta 6 pneumolysin (delta6Ply, 80ng.ml⁻¹) and the calcium ionophore A23187 (2 μM), to neutrophils on the production of platelet-activating factor (PAF) and thromboxane A₂ (TxA₂) are shown in figures 2A and 2B respectively. The data from 5 different experiments, using cells from 5 different individuals, are expressed as the median values with interquartile ranges. BG=background value for unstimulated cells.

*p<0.05-p<0.002

Figure 3.

The effects of the addition of pneumolysin (Ply), at concentrations of 5-80 ng.ml⁻¹, as well as those of delta 6 pneumolysin (delta6Ply, 80ng.ml⁻¹) and the calcium ionophore A23187 (2 μM, positive control) on neutrophil viability.

*p<0.01-p<0.001

Figure 4.

The effects of the addition of ADP (100 μM, positive control) or pneumolysin (Ply, 10-80 ng.ml⁻¹) or delta 6 pneumolysin (delta6Ply 80 ng.ml⁻¹) on the formation of heterotypic neutrophil:platelet (NP) aggregates. The results of 35 experiments, using cell suspensions from 13 different donors are expressed as the CD42a median fluorescence intensity (MFI) with interquartile ranges. The aggregates assessed were positive for co-expression of CD16, CD42a and CD45.

*p<0.001-p<0.0001

Figure 5.

The effects of addition of ADP (100 μM), platelet-activating factor (PAF, 400 nM) or thrombin (1.25 NIH units. ml^{-1}) in the absence and presence of their respective receptor antagonists (PSB 0739, WEB 2086, SCH 79797 all at 10 μM) to platelet-rich plasma on expression levels of the adhesion molecule CD62P. The results of 6 experiments, using platelet rich plasma from 2 different donors are expressed as the median CD16⁺/CD42a⁺/CD45⁺ fluorescence intensities with interquartile ranges.

Figure 6.

The results in the upper figure (5A) show the effects of pneumolysin (Ply 40 ng. ml^{-1}) only or in the presence of WEB 20186, SCH 79797, indomethacin, PSB 0739 or S 18886, all at 10 μM), as well as the effect of calcium depletion from the cell suspending medium, on the formation of neutrophil:platelet (NP) heterotypic aggregates. The results of 18 experiments using cells from 7 donors are expressed as the median CD16⁺/CD42a⁺/CD45⁺ fluorescence intensities with interquartile ranges.

The results in the lower figure (5B) show the effects of addition of an anti-CD62P monoclonal antibody to buffy coat suspensions on the spontaneous (BG) and pneumolysin (Ply 40 ng. ml^{-1})-activated formation of neutrophil:platelet heterotypic aggregates.

* $p < 0.006$ For comparison with the corresponding Ply-treated, drug-free control system.

* $p < 0.0004$ For comparison of the control and corresponding anti-CD62P-treated systems.

Figure 7.

Measurement of the effects of addition of the PAR1 receptor antagonist, SCH 79797 (10 μM), to buffy coat suspensions on the spontaneous (background) and ADP (100 μM)- or thrombin (1.25 NIH units. ml^{-1})-activated formation of neutrophil platelet (NP) heterotypic aggregates. The results of 4 experiments using cells from 4 donors are expressed as the CD42a median fluorescence intensity (MFI) with interquartile ranges. The aggregates assessed were positive for co-expression of CD16, CD42a and CD45.

* $p < 0.001$ For comparison of the thrombin-activated systems without and with SCH 79797

Figure 8.

Measurement of the effects of dabigatran (10 μ M) added to buffy coat suspensions on the formation of heterotypic neutrophil:platelet (NP) aggregates, activated by either thrombin (1.25 NIH units.ml⁻¹) or pneumolysin (Ply 40 ng.ml⁻¹). The results of 6 experiments using cells from 2 donors are expressed as the median CD16⁺/CD42⁺/CD45⁺ fluorescence intensities with interquartile ranges.

*p<0.02 For comparison of the thrombin-activated systems in the absence or presence of dabigatran.

Foot note comments:

None of the authors has any conflict of interest to declare. JGN and CF were funded by research grants awarded by the South African National Research Foundation (NRF).

A

B

Effects of the addition of Ply, delta6Ply or ADP on neutrophil:platelet aggregation

Effects of platelet receptor antagonists on ADP-, PAF- or thrombin-activated upregulation of expression of CD62P

Effects of platelet receptor antagonists, indomethacin or calcium depletion on Ply-induced neutrophil:platelet aggregation

A

B

Effects of the PAR1 receptor antagonist, SCH 79797 on basal, ADP-, or thrombin-activated neutrophil:platelet aggregation

Effects of dabigatran on thrombin- or Ply-activated neutrophil:platelet aggregation

Highlights

- Pneumolysin (Ply) activates production of PAF and thromboxane A₂ (TxA₂) by neutrophils.
- Ply also promotes formation of pro-thrombotic neutrophil:platelet (NP) aggregates.
- Ply-mediated aggregate formation is independent of PAF and TxA₂.
- P-selectin (CD62P) and protease-activated receptor 1 are involved in Ply-induced NP aggregation.
- Ply-mediated NP aggregate formation may contribute to pulmonary and myocardial injury.