Assessing the relative importance of isolated Ficus trees to insectivorous birds in an Indian human-modified tropical landscape

Matthews, Thomas; Cottee-Jones, Eden; Bregman, Tom; Whittaker, Robert

DOI:

[10.1007/s10531-017-1387-8](https://doi.org/10.1007/s10531-017-1387-8)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Matthews, T, Cottee-Jones, E, Bregman, T & Whittaker, R 2017, 'Assessing the relative importance of isolated Ficus trees to insectivorous birds in an Indian human-modified tropical landscape', *Biodiversity and Conservation*. <https://doi.org/10.1007/s10531-017-1387-8>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 05/06/2017

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10531-017-1387-8>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 Title: Assessing the relative importance of isolated *Ficus* trees to insectivorous birds
2 in an Indian human-modified tropical landscape

3 Article type: Original research

4 Authors: Thomas J. Matthews^{*+1,2}; H. Eden W. Cottee-Jones⁺³; Tom P. Bregman⁴;
5 Robert J. Whittaker^{3,5}

6 *Corresponding author

7 ⁺These authors contributed equally to this work.

8 Affiliations: ¹GEES (School of Geography, Earth and Environmental Sciences), The
9 University of Birmingham, Birmingham, B15 2TT. ²CE3C – Centre for Ecology,
10 Evolution and Environmental Changes/Azorean Biodiversity Group and Univ. dos
11 Açores –Depto de Ciências Agrárias, PT-9700-042, Angra do Heroísmo, Açores,
12 Portugal. ³Biodiversity Research Group, School of Geography and the Environment,
13 Oxford University Centre for the Environment, University of Oxford, South Parks
14 Road, Oxford, OX1 3QY, UK. ⁴Edward Grey Institute, Department of Zoology,
15 University of Oxford, OX1 3PS, UK. ⁵Centre for Macroecology, Evolution and
16 Climate, Department of Biology, University of Copenhagen, Copenhagen, Denmark.

17 Email addresses: txm676@gmail.com; henry.cottee-jones@seh.ox.ac.uk;
18 thomas.bregman@zoo.ox.ac.uk; robert.whittaker@ouce.ox.ac.uk

19 Correspondence address: T.J. Matthews, GEES (School of Geography, Earth and
20 Environmental Sciences), The University of Birmingham, Birmingham, B15 2TT.
21 Email: txm676@gmail.com

22 Running headline: Insectivores and isolated *Ficus* trees

23

Acknowledgements

The authors wish to thank Maan Barua, Manju Barua, Barry and Susan Jones, A.J. Tours and Travel, and Wild Grass Eco Lodge for help facilitating this study. Valuable field assistance was provided by Biju Hazarika, Gokul Munda, Soano Rajbonsi, Raju Gogoi, Nakib Ali, Polash Bora, Colia Karmakar, and Humnot Borah. We are very grateful for advice and comments provided by François Rigal, Michael Børregaard, Richard Grenyer, Paul Jepson, Catherine Sheard, Jon Sadler, Ross Crates and an anonymous reviewer.

Compliance with ethical standards

This research was conducted with ethical approval from the University of Oxford (Departmental CUREC reference number: SOGE C1A-99). Surveys were conducted with permission from local landowners where necessary, and permission to conduct this field work in India was granted by the High Commission of India, London, under visa number 4246496. Field studies did not impact the welfare of the animals studied, and the authors have no conflicts of interest to declare.

Abstract

The destruction of forest for agricultural expansion has created a vast estate of human-modified land in tropical regions. One group of organisms that are particularly vulnerable to the loss of forest habitat are insectivorous birds. Despite this, few conservation strategies have been identified for this group in human-modified landscapes. We survey the use of 104 isolated trees by insectivorous birds in rural Assam, India. We used an information theoretic model comparison approach to determine the important variables driving insectivorous bird diversity within these isolated trees. Our work demonstrates that the conservation of large trees in human-modified landscapes may play an important role in maintaining bird diversity and ecological function beyond the forest edge. More specifically, we found that isolated *Ficus* trees hold assemblages with particularly high insectivore abundance, richness and functional diversity when compared to other isolated fruit and large trees. We argue that, where present, *Ficus* trees should be actively conserved in human-modified landscapes to maintain the composition of insectivore communities in a “*Ficus* first” strategy.

Keywords Conservation beyond protected areas, birds, ecological function, *Ficus*, India, insectivores, isolated trees, multimodel inference

Introduction

As agricultural expansion continues to fragment the world’s tropical forests and occupy large areas of land (Phalan et al. 2013), it is increasingly important to devise conservation strategies for human-modified landscapes (Chazdon et al. 2009; Gardner et al. 2009; Melo et al. 2013). The conversion of forest has a range of impacts on different taxa. Here our focus is on the largest avian feeding guild, insectivorous

65 birds, an ecological group that is considered particularly vulnerable to habitat loss
66 (Tschardt et al. 2008).

67 Several studies have demonstrated lower abundance and species richness of
68 insectivores in human-modified landscapes compared to intact forest habitats (Harvey
69 et al. 2006; Şekercioğlu 2012). Studies in habitat fragments have found similar
70 results; especially when there is limited tree cover surrounding fragments (Stouffer
71 and Bierregaard 1995). Not only does this cause deterioration in the conservation
72 status of insectivorous birds (Şekercioğlu et al. 2002), but it also diminishes their
73 beneficial pest control services (Van Bael et al. 2008; Karp and Daily 2013).

74 One possible strategy to mitigate the decline of insectivorous birds in human-
75 modified landscapes might be the conservation of isolated trees (Manning et al. 2006;
76 Cottee-Jones et al. 2015a). These trees can increase the abundance and richness of
77 insectivores by providing connectivity between forest remnants for forest-dependent
78 species (Harvey et al. 2006), along with feeding and nesting sites for matrix-tolerant
79 taxa (Şekercioğlu et al. 2007), thereby moderating the impact of habitat loss
80 (Şekercioğlu 2012). However, our understanding of conservation tools that are
81 appropriate in human-modified landscapes represents a critical frontier in tropical
82 conservation biology (Tschardt et al. 2008; Melo et al. 2013). Indeed, we have very
83 limited experimental evidence to help guide conservation practitioners working in the
84 2.5 billion ha of tropical land area that has been modified by humans (see Supporting
85 Information 1; Fischer et al. 2006; Melo et al. 2013).

86 *Ficus* trees have long been recognised as important food resources for frugivores
87 (Terborgh 1986; Cottee-Jones et al. 2015b), with almost 1,000 frugivorous bird
88 species recorded consuming *Ficus* fruit (Shanahan et al. 2001). They may also be
89 overlooked but important foraging sites for insectivores. Their co-evolved mutualism

with pollinating fig wasps (Hymenoptera, Agaonidae, Agaoninae) means that millions of fig wasps (and non-pollinating fig wasps; Hymenoptera, Chalcidoidea) are found in association with fruiting figs (Harrison 2003; Bain et al. 2013). As well as being a food resource for insectivorous birds themselves, these fig wasps are also the prey of other invertebrates that inhabit or visit *Ficus* trees (Schatz et al. 2008). Numerous other insects, including heteropterans, nematodes, coleopterans, and fruit flies, also exploit the extremely large fruit crop, which may number as many as one million syconia per tree (Cushman et al. 1998). Indeed, notwithstanding chronic under-sampling, there are published records of 1,875 species of arthropod feeding on *Ficus*, including 742 species feeding on syconia, 481 feeding on sap, 369 leaf-chewers, and 283 wood borers (Basset et al. 1997). Ants (Hymenoptera: Formicidae) for example, have been found to specialise in predating non-pollinating fig wasps in some dioecious *Ficus* species, predate on fig dwelling herbivorous insects, consume partially eaten ripe syconia, and nest in figs (Schatz et al. 2008; Harrison 2013). The complex trunk morphology of many strangler *Ficus* trees provides further habitat for arthropods, all of which suggests that *Ficus* trees may be attractive feeding sites for insectivorous birds.

In this study, we sought to test whether isolated *Ficus* trees were particularly important foraging sites for insectivorous birds in human-modified landscapes, relative to other fruit-bearing and large non-fruit trees found in open habitats, as indicated by higher levels of insectivore richness, abundance, and functional diversity.

Materials and methods

Study area

The study took place from April 2012 to June 2013 in the Golaghat District of Assam, North-east India (see Fig. 1). The study site is a $\approx 250 \text{ km}^2$ area bounded by the Western Range of Kaziranga National Park at N26 34.394 E93 15.433, the city of Jorhat at N26 46.198 E94 12.678, and the town of Golaghat at N26 27.819 E93 54.978. The elevation of the study area ranges between 30 and 100 m above sea level, and the mean annual rainfall for the region is 1,500–2,500 mm, most of which falls in the June to September monsoon (Shrivastava and Heinen 2007). The annual temperature range varies from an average minimum of 5°C to an average maximum of 35°C (Barua and Sharma 1999).

The original habitat of moist subtropical deciduous forest (Champion and Seth 1968) was largely cleared following the local commercialisation of tea production in 1840 (Shrivastava and Heinen 2007). Remnants of the original forest remain in the 7.65 km^2 Panbari Forest Reserve on the edge of the Karbi Hills, and in the 430 km^2 Kaziranga National Park (Barua and Sharma 1999). Other small areas of forest regrowth exist, but are typically less than 1 ha. Agriculture is the dominant land-use, with a mixture of small-holder rice cultivation, village home gardens, and large commercial tea estates.

Focal tree sampling

To compare the insectivore assemblages visiting *Ficus* trees to other isolated trees in this human-modified landscape, we surveyed three categories of focal tree: 1) isolated *Ficus* trees, which we surveyed when in fruit; 2) isolated fruit trees that did not belong to the *Ficus* genus, which we surveyed when in fruit; 3) isolated large, non-fruiting trees (i.e. trees that did not produce fruit during the study). We surveyed a total of 40 *Ficus* trees, 33 fruit trees, and 31 large non-fruiting trees in the study area (Table 1). The selection of focal *Ficus* and other non-*Ficus* fruit trees (herein in “fruit

trees”) was determined by their fruiting cycle; only trees with crops of ripe fruit were surveyed. Large non-fruiting trees (“large trees”) were selected from the largest trees in the landscape, exclusive of the two previous groups, with a minimum circumference at breast height of over 1 m, so that we had a dataset that was comparable in tree stature to the *Ficus* trees.

For each tree, we measured the diameter at breast height (DBH), estimated the maximum tree height with a clinometer, and estimated the canopy area by measuring the canopy diameter at ground level along two axes, deriving canopy area using the formula for an ellipse. To obtain a single estimate for tree size, a Principal Components Analysis (PCA) with Kaiser stopping criterion extraction (eigenvalues >1) and oblique rotation was conducted using DBH, maximum height, and canopy area in IBM SPSS Statistics 22 (IBM 2013). The first axis of this PCA explained over 80% of the variance in the three variables and was significantly correlated with all three tree size variables (Pearson's product-moment correlation coefficient greater than 0.8 in each case). Thus, the first axis was used as our tree size variable. The intensity of human land-use within a 100 m radius of each focal tree was recorded using a three-point scale (where 0 is very little human land use; 1 is some human land use, such as cultivation; and 2 is intense human land use, in cases where a road, house, or paddy field were present).

We measured the distance to the nearest protected area with intact forest by marking the focal trees with a GPSmap 62s device, and then overlaying the GPS markers on Landsat 8 satellite images of the region in ArcGIS 10.2.1 (ESRI 2014). We digitised the protected area borders through an on-screen visual interpretation, and then measured the distance (in km) of each focal tree to the nearest protected area. In all cases, protected areas held the only high-quality forest habitat left in the study

landscape. In addition to protected areas, small (≤ 1 ha), low-quality wooded areas were located through consultation with local landholders and marked with a GPS device. We then recorded a second distance measurement: the distance to the nearest wooded area of any quality (whether a protected area or small wooded area).

Insectivore surveys

Each focal tree was surveyed once, for three hours, from first light. Surveys were only conducted in fair weather conditions. During the survey, a single observer would watch the tree from a concealed position with a good view, typically about 20 m from the trunk. Each individual bird that landed in the tree was recorded, and birds that made repeated visits to and from the tree were denoted with an asterisk to avoid double counting. Trees in the *Ficus* and fruit categories were only surveyed when the crop was ripe. Deforestation in the study area occurred over 150 years ago, and was largely uniform in timing, so differential rates of matrix assemblage relaxation were not considered to have an effect on bird communities in the study area. As all trees were surveyed in the same Assamese season, the presence or absence of migratory species was consistent across the three tree groups.

Each species was classified into primary dietary guilds (frugivore, nectivore, insectivore, granivore, or carnivore; omnivores were classified according to their main food type, and were not included in any further analyses on insectivorous birds as they may have been attracted to *Ficus* and fruit trees by the fruit present) following del Hoyo et al. (1992–2002; 2003–2011). Nomenclature also followed del Hoyo et al. (1992–2002; 2003–2011).

Ecomorphological data collection

To calculate functional diversity scores, we preferred to use ecomorphological trait data from the insectivores recorded in the surveys rather than guild classifications extracted from the literature or assessed using our field experience. This is because continuous traits have been argued to produce more accurate representations of species' functional roles in ecosystems (McGill et al. 2006), and the use of continuous trait data removes the need to arbitrarily assign species into different categories. We defined a “trait” as a measurable aspect of an organism, which determines its interaction with the environment (Flynn et al. 2009). Here we were interested in the foraging and dispersal capacity of insectivorous birds, so we measured traits associated with locomotive behaviour, dispersal ability, gape size, bill structure, and body size (Derryberry et al. 2011; Claramunt et al. 2012). In order to obtain ecomorphological trait data, each species recorded in the surveys was measured following Edward Grey Institute protocols at the British Natural History Museum's ornithological collections (Supporting information 2).

Functional diversity calculation

We follow the definition of functional diversity as the distribution of functional traits within multidimensional niche space (Petchey and Gaston 2006), and used Laliberté and Legendre's functional dispersion (FDis) index to measure functional diversity in our dataset (Laliberté and Legendre 2010; please see Supporting information 3 for a justification of the method selected). We calculated FDis for each of our focal trees using “package FD” in Programme R (Laliberté and Shipley 2013; R Core Team 2014).

As our trait data were measured on a continuous scale, rather than classified into nominal groups, a species–species uncorrected distance matrix was computed. A Principal Co-ordinates Analysis (PCoA) was performed after the distance matrix was

corrected for negative eigenvalues. Each trait axis was standardised to have a mean of zero and a standard deviation of one (Petchey and Gaston 2006). These corrected PCoA trait axes were used to calculate the FDis scores for our focal trees. FDis could not be computed for trees with no insectivore records, but these trees were included in further analyses with index scores of 0. Similarly, trees with only one species of insectivore were given a score of 0 following Laliberté and Shipley (2013). We checked the functional relationships between the sampled insectivorous bird species by constructing a dendrogram: we transformed the species–trait data into a distance matrix (Euclidean distance), and applied the UPGMA clustering algorithm. The resulting dendrogram was subsequently converted into a tree object and plotted (Fig. 2).

Statistical analysis

The effect of tree size, land-use intensity, distance to the nearest protected area and nearest forest of any type, and tree type on insectivore abundance, richness, and FDis (the three response variables) were examined using an information-theoretic approach (Burnham and Anderson 2002). For each response variable, we fitted a full generalized linear model (GLM), i.e. a model with all predictor variables included. Abundance and richness are count data and thus for these response variables we used GLMs with the Poisson family and a log link function. As the Poisson distribution assumes that the mean is equal to the variance, a quasi-Poisson model was fitted with each response variable to assess for over-dispersion. In both cases, the data were found to be over-dispersed and thus quasi-AIC_c (QAIC_c; Richards 2008; Bolker 2016) was used for subsequent model comparisons using models with these two response variables. Functional diversity was found to be normally distributed following a log transformation; as the logarithm of zero is not defined, a constant of 0.1 was added to

236 all FDis values. Thus, for models in which FDis was used as the response variable, we
237 fitted GLMs with the Gaussian family and identity link function.

238 Multicollinearity between the continuous predictor variables was assessed using
239 variance inflation factors using the 'car' R package (Fox and Weisberg 2011) and a
240 threshold of five. All continuous predictors had variance inflation factors of less than
241 five and so were included in the model comparisons. Outliers were assessed using
242 Cook's distance and a threshold of one. Two data points were removed prior to the
243 model comparisons based on this criterion. Continuous predictors were assessed for
244 normality: tree size and both distance measures were log transformed to induce
245 normality. A constant of 1.5 was added to tree size values as the raw values (PCA
246 axis values) contained negative numbers.

247 Model comparison was undertaken using an information theoretic approach. When
248 FDis was used as the response variable, we fitted a complete set of models
249 considering all predictor variables using the dredge function in the MuMIn R package
250 (Bartoń 2016). Models were ranked according to AIC_c values (Burnham and
251 Anderson 2002) and we also recorded the ΔAIC_c values and the AIC_c weights for
252 each model.

253 As the dispersion parameter in the Poisson GLM is taken to be one, and model fits
254 using abundance and richness were found to be over-dispersed, we compared models
255 with these two response variables using the quasi-Poisson family and $QAIC_c$
256 (Richards 2008). Thus, instead of using maximum likelihood estimation, we focused
257 on maximising the "quasi-likelihood." For each set of model comparisons, we
258 extracted the dispersion parameter from the full model (i.e. with all predictors) using
259 functions provided by Bolker (2016). We then, separately for each response variable
260 (i.e. abundance and richness), fitted a full set of models considering all predictors and

compared models based on their QAIC_c values; again, also storing the Δ QAIC_c values and the QAIC_c weights. Thus, in total we had three model comparison tables, one for each of the three response variables.

To evaluate the importance of individual variables, for each model comparison table separately we calculated the weight of evidence (WoE) of each predictor by summing the AIC_c weights (or QAIC_c weights) for each model in which a predictor variable was included (Burnham and Anderson 2002; see also Giam and Olden 2016). As the AIC_c and QAIC_c weights sum to one for a given model comparison, the WoE values are constrained to be between 0 and 1. However, WoE values are not expected to be zero even in cases where a predictor variable has no predictive value (see Burnham and Anderson 2002, p. 345). Thus, we followed Burnham and Anderson (2002, p. 345 onwards) and used a bootstrap methodology to compute a baseline WoE value for each predictor variable in each model comparison table. This worked by creating an algorithm that took the i^{th} predictor and randomised the values whilst holding the values of the other predictor variables constant. The model comparison was then repeated and the WoE values calculated in the standard manner. The algorithm then re-arranged the values in the i^{th} predictor back to their original order and moved onto the $i+1^{\text{th}}$ predictor, and so on, until all predictor variables had been randomised. This process was then repeated 500 times and the median value (the bootstrap distribution of WoE values is occasionally skewed and thus the median is a preferable metric; Burnham and Anderson 2002) taken. This approach was only undertaken for the model comparison using FDis as the response variable, as it is straightforward to implement in the context of Gaussian GLMs. In the model comparisons using abundance and richness as the response variables, the models were fitted using the Poisson/quasi-Poisson families and it was found that randomising the predictor values

frequently resulted in very high degrees of over-dispersion and the failure of models to converge. Based on the bootstrap WoE results using FDis, we tentatively used a baseline of 0.3 when analysing WoE results from model comparisons using abundance and richness.

For each model comparison table, we took the full model and best model (i.e. lowest AIC_c or QAIC_c value) and examined the residual plots (e.g. residuals against fitted values, standardised residual values etc.) for any patterns. We also tested for spatial autocorrelation in the residuals of the best model fits using the ‘spdep’ R package (Bivand and Paris 2015), the nb2listw function and row standardised weights. When FDis was used as the response variable, examination of the residuals revealed some sort of pattern; potentially indicating that a variable was missing from the model (Zuur et al. 2009). To account for this, we re-ran the model selection whilst including an interaction term between tree size and tree type in the full model. This resulted in a much more normal distribution of residuals in the best model fit. As such, we re-ran the FDis model selection using the interaction as a fixed term within the ‘dredge’ function in MuMIn. As the interaction term is fixed, it means that the individual variables ‘tree size’ and ‘tree type’ are also fixed. We also re-ran the abundance and richness model selection analyses with this interaction term to assess whether the interaction was important in regards to these response variables. As it was found to improve the distributions of errors in the abundance and richness models, we also fixed the interaction term in this model selection. Finally, for the FDis model comparison we looked at the R² value of the best model, whilst for the abundance and richness model comparisons we computed pseudo R² values for the best models using the formula: 1 - (model deviance / null deviance).

Results

311 Over the 104 surveys, 33 species of insectivorous bird were recorded. The most
312 frequently recorded species were the Oriental white-eye (*Zosterops palpebrosus*) with
313 55 records, common tailorbird (*Orthotomus sutorius*) with 54, and the Oriental
314 magpie robin (*Copsychus saularis*) with 53. The Oriental white-eye was also the most
315 abundant species, with 146 individual records, followed by the great tit (*Parus major*)
316 with 86, and the Oriental magpie robin with 84.

317 *Ficus* trees had higher mean abundance (12.0, standard error = 0.96), richness (7.0,
318 SE = 0.38) and FDis (1.3, SE = 0.07) values compared to the other tree categories,
319 followed by large non-fruit trees (mean richness = 3.7, 2.5 and 0.8, respectively; SE =
320 0.74, 0.45 and 0.15) and then non-*Ficus* fruit trees (2.7, 1.9 and 0.6, SE = 0.37, 0.23
321 and 0.11; Fig. 3).

322 The results of the multimodel comparison analyses are described below for each of
323 the three response variables in turn.

324 Abundance

325 The inclusion of the interaction term resulted in a more normal distribution of errors
326 in the full and best models (for both the abundance and richness models), although
327 there was still a degree of spread towards the extreme tails of the distribution;
328 however, it is known that residuals in Poisson regression models are only
329 approximately normal, and there is expected to be a degree of spread towards the
330 extreme tails of the distribution. Thus, the model selections based on both the
331 abundance and richness (results presented below) response variables were run with
332 the interaction term included. When abundance was used as the response variable,
333 there was one model within 2 ΔQAIC_c values of the best model. The best model
334 contained tree size, tree type and the interaction between them (i.e. the fixed

parameters in the model selection) and the distance to the nearest protected area with intact forest (Table 2a), and had a pseudo- R^2 value of 0.62. The distance to a protected area variable also had a relatively high WoE value (0.59), which was larger than the baseline of 0.3 that we employed for the quasi-Poisson model selections in this study. The parameter estimates for the best model (using the quasi-Poisson family) are included in Table S1 in Supporting information 4. Using the quasi-Poisson family results in the same parameter estimates as the standard Poisson family. We did not look at the significance of parameter estimates for the best model (including for the best richness and FDis models, below), as this is not advised within information theoretic model comparison approaches (Burnham and Anderson 2002). There was no significant spatial autocorrelation in the residuals of the best model (Moran's $I = 0.03$; $P = 0.31$). In regards to the tree type variable, the parameter estimates were negative and relatively large (Table S1) and thus indicated that both fruit trees and large non-fruited trees supported lower abundance than *Ficus* trees. The effect of tree size was positive, indicating abundance increased with the size of tree. Consideration of the interaction term parameter estimates (Table S1) indicates that this is primarily driven by the large non-fruited tree category, i.e. the slope between abundance and tree size is steeper for this tree category relative to the other two. Interestingly, the effect of distance was positive, which implies that the abundance of insectivorous birds increased with distance to the nearest protected area with intact forest. The WoE values for the other distance variable and land use were both below the baseline value (Table 2a), and thus these variables can be considered relatively unimportant in determining abundance in this system.

Richness

When richness was used as the response variable, there were two models within 2 ΔQAIC_c values of the best model. The best model contained tree size, tree type and the interaction between them (i.e. the fixed parameters in the model selection) (Table 2b), and had a pseudo- R^2 value of 0.60. The parameter estimates for the best model (using the quasi-Poisson family) are included in Table S2 in Supporting information 4. In regards to the tree type variable, the parameter estimates were again negative and relatively large (Table S2) and thus indicated that both fruit trees and large non-fruiting trees supported lower richness than *Ficus* trees. As with the best model using abundance, the effect of tree size was positive and the interaction term parameter estimates (Table S2) indicate that this effect is primarily driven by the large non-fruiting tree category. Whilst land use and the distance to the nearest protected area with intact forest were included in the models within 2 ΔQAIC_c of the best model, both variables had WoE values below the baseline of 0.3. The second distance variable also had a WoE value below 0.3 (Table 2b). There was no significant spatial autocorrelation in the residuals of the best model (Moran's $I = 0.03$; $P = 0.29$).

Functional Dispersion

When FDis was used as a response variable the best model had an adjusted R^2 value of 0.29 (see Table S3 in Supporting information 4 for parameter estimates). There were no additional models within 2 ΔAIC_c values of the best model (Table 3). As described above, the FDis model selection analyses included an interaction term between tree size and tree type as a fixed term in the model selection, to ensure a more normal distribution of errors. Thus, tree type, tree size and the interaction term were all included in the best model by default and the WoE values for these variables are not interpretable (see Table S3). In regards to the tree type variable, the parameter estimates were negative and relatively large (Table S3) and thus indicated that both

fruit trees and large non-fruiting trees supported lower FDis than *Ficus* trees. The effect of tree size in the best model was small (-0.06) and negative, indicating that FDis actually decreased with increasing tree size. However, closer inspection of the best model's parameter estimates indicated that this was probably driven by the interaction between tree type and tree size; the slope of the FDis – tree size relationship was steeper for both fruit trees and large non-fruiting trees relative to *Ficus* trees. Land use was also included in the best model and had a relatively high WoE value, which was considerably larger than the bootstrapped baseline value. Inspection of the best model's parameter estimates indicated that increasing land use intensity resulted in a decrease in FDis. Neither of the distance variables were included in the best model, and both had WoE values lower than the baseline values (Table 3). There was no significant spatial autocorrelation in the residuals of the best model (Moran's $I = -0.02$; $P = 0.46$).

Discussion

The conversion of tropical forest to agricultural production causes changes in insectivorous species composition and functional diversity (Tscharntke et al. 2008; Azhar et al. 2013; Edwards et al. 2013). However, we found that isolated *Ficus* trees provide important micro-site level habitat for insectivores in the human-modified Assamese landscape. Compared to the other tree categories, *Ficus* trees had consistently higher insectivore richness, abundance and functional diversity, suggesting that these trees may be more valuable from a conservation perspective in these modified landscapes than other isolated trees.

407 In addition to tree type, our model comparisons also indicated that tree size and the
408 interaction between tree size and tree type were important variables. Although these
409 variables were fixed in the model comparisons, the fact that they were needed to be
410 fixed to improve the error distribution indicates that they are important. In addition,
411 exploratory analysis indicated, once the assumptions of GLMs were temporarily
412 relaxed, that even when the model comparisons were run without fixing these
413 variables they consistently had high WoE values and were included in the best models
414 (results not shown). The interaction terms in the best models indicated that each of the
415 slopes between abundance, richness and FDis, and tree size were steeper in the large
416 non-fruit tree category relative to the other two categories. Figure 4 illustrates these
417 relationships for richness using scaled size (i.e. re-scaling each tree size such that the
418 data cover the same range for each tree type) for each of the three tree types. It can be
419 seen that for *Ficus* trees and the large non-fruit trees category, species richness
420 generally increases with tree size; although the relationship appears to flatten out at
421 large sizes for *Ficus* and the relationship is thus steeper for the large non-fruit tree
422 category. However, there does not appear to be any relationship between tree size and
423 richness for non-*Ficus* fruit trees. It should also be noted that if the tree sizes are
424 simply standardised by the largest tree in the dataset (i.e. not accounting for tree type)
425 the *Ficus* line still lies above the other two lines in Fig. 4 and thus, whilst the
426 relationship is steeper for large non-fruit trees, for any given tree size there is a higher
427 species richness in *Ficus* trees relative to the other tree types. The fact that the
428 relationship is steeper for large non-fruit trees relative to *Ficus* trees is likely due in
429 part to the fact the *Ficus* trees are generally larger than trees in the other two
430 categories and thus there are fewer small *Ficus* trees in our dataset, and in tropical
431 landscapes more generally (partly due to the “strangler” life history of many species).

Nonetheless, in regards to conservation actions and assuming that the retention of ecological services such as pest control is a priority (and assuming that the birds recorded in isolated trees are also foraging in agricultural crops), the conservation of large trees *per se* may be the most effective strategy. In this case, isolated *Ficus* trees would again be a conservation priority, as they were generally the largest trees in the study area.

When abundance was used as the response variable, the model comparison results indicated that abundance increased with increasing distance from a protected area with intact forest. This is an interesting and counter-intuitive finding and is possibly due to the effect of matrix specialist birds (Şekercioğlu 2012). Some of the species with the highest number of recorded individuals were matrix specialist species, such as great tit (*Parus major*), common iora (*Aegithina tiphia*), and crimson sunbird (*Aethopyga siparaja*). These birds showed a distinct preference for non-forest habitats, with increasing occurrence as the distance from the forest increased.

Interestingly, the effect of land use was only important in the model comparisons based on FDis. It was found that increasing land use intensity resulted in a decrease in FDis. This is to be expected and several previous studies have shown that land use change and intensification lead to a reduction in functional diversity (e.g. Schweiger et al. 2007; Flynn et al. 2009; Laliberté et al. 2010). The reason why land use was not an important variable in the abundance and richness model comparisons is unknown. However, again, it may be due to the presence of matrix specialists in the landscape. The inclusion of matrix specialist bird species in analyses such as those in this study have sometimes been found to mask the effects of land use change on abundance and richness (Matthews et al. 2014). If these matrix specialist species are relatively immune to land use change and increase in abundance and richness in more disturbed

environments, but also possess similar trait values to one another, they may compensate the loss of forest specialist abundance and richness with increasing land use intensity whilst simultaneously resulting in a reduction in FDis, as we observed. It is also possible that our coarse three-level ordinal land use scale did not contain enough information to uncover relationships between abundance and richness, and land use.

Several studies have argued for the need to conserve isolated trees in human-modified habitat (Manning et al. 2006; Şekercioğlu et al. 2007; Fischer et al. 2010). Our results build upon these arguments, adding that isolated trees can be important resources for insectivores as well as frugivores (Luck and Daily 2003), and tree-hole nesters (Manning et al. 2004). We consider this a valuable finding, as there are few conservation strategies focused on this vulnerable group in modified landscapes. We also add that *Ficus* trees may be particularly important for insectivorous birds in human-modified landscapes. In our study area at least, this implies that conserving *Ficus* trees ahead of other tree types may be a more effective conservation strategy than conserving isolated trees at random. If *Ficus* trees are found to be similarly important to insectivorous birds in modified landscapes on a wider spatial scale, a “*Ficus* first” approach to isolated tree conservation may be effective across the tropics (see Cottee-Jones & Whittaker 2015; Cottee-Jones et al. 2016). Although additional studies on *Ficus* trees and insectivores are lacking, *Ficus* have been found to support rich and abundant insect communities wherever they have been studied (Basset and Novotny 1999; Pereira et al. 2000; Bain et al. 2013).

The conservation of isolated trees in modified landscapes presents significant challenges, however. It requires a long-term vision with extensive commitment from landowners (Manning et al. 2004), and can incur substantial costs if natural

regeneration is insufficient (Fischer et al. 2010). The regeneration of *Ficus* trees may be particularly challenging, as many species are epiphytic in their early life stages, and so depend upon the presence of large host trees (Leighton and Leighton 1983). Fortunately, evidence from the study area indicates that, in this region at least, *Ficus* trees are regenerating faster in modified habitats than are isolated trees in other parts of the world (Gibbons et al. 2008; Cottee-Jones et al. 2016). If the conservation of isolated *Ficus* trees was adopted in legislation, there is scope for future work to focus on thresholds for the tree size required to qualify for protection, and the size of buffer zones around isolated trees where land-use practices may be restricted.

References

- Azhar B, Lindenmayer DB, Wood J, Fischer J, Manning A, McElhinny C, Zakaria M (2013) The influence of agricultural system, stand structural complexity and landscape context on foraging birds in oil palm landscapes. *Ibis* 155:297–312.
- Bain A, Harrison RD, Schatz B (2013) How to be an ant on figs. *Acta Oecologica* 57:97–108.
- Bartoń K (2016) Package 'MuMIn': multi-model inference. R package version 1.15.6.
- Barua M, Sharma P (1999) Birds of Kaziranga National Park, India. *Forktail* 15:47–60.
- Basset Y, Novotny V (1999) Species richness of insect herbivore communities on *Ficus* in Papua New Guinea. *Biological Journal of the Linnean Society* 67:477–499.
- Basset Y, Novotny V, Weiblen G (1997) *Ficus*: a resource for arthropods in the tropics, with particular reference to New Guinea. In: Watt AD, Stork NE, Hunter MD (eds) *Forests and insects*. Chapman and Hall, London, pp 341–361.
- Bivand R, Piras G (2015) Comparing implementations of estimation methods for spatial econometrics. *Journal of Statistical Software* 63: 1-36.
- Bolker B (2016) Dealing with quasi-models in R. *Compare* 1: 5-452305.
- Burnham KP, Anderson DR (2002) *Model selection and multimodel inference. A practical information-theoretic approach*. Springer, London.
- Champion HG , Seth SK (1968) *A revised survey of forest types of India*. Government of India Press, New Delhi.
- Chazdon RL, Harvey CA, Komar O, Griffith DM, Ferguson BG, Martínez-Ramos M, Morales H, Nigh R, Soto-Pinto L, van Breugel M, Philpott SM (2009) Beyond

515 reserves: a research agenda for conserving biodiversity in human-modified tropical
516 landscapes. *Biotropica* 41:142–153.

517 Claramunt S, Derryberry EP, Remsen JV Jr, Brumfield RT (2012) High dispersal
518 ability inhibits speciation in a continental radiation of passerine birds. *Proceedings of*
519 *the Royal Society B: Biological Sciences* 279:1567–1574.

520 Cottee-Jones HEW, Whittaker RJ (2015) Felling *Ficus*: the cultural status of fig trees
521 in a rural Assamese community, India. *Ethnobiology Letters* 6: 89-98.

522 Cottee-Jones HEW, Bajpai O, Chaudhary LB, Whittaker RJ (2015a) Isolated *Ficus*
523 trees deliver dual conservation and development benefits in a rural landscape. *Ambio*
524 44: 678–684.

525 Cottee-Jones HEW, Matthews TJ, Bregman TP, Barua M, Tamuly J, Whittaker RJ
526 (2015b) Are protected areas required to maintain functional diversity in human-
527 modified landscapes? *PLoS ONE* 10: e0123952.

528 Cottee-Jones HEW, Bajpai O, Chaudhary LB, Whittaker RJ (2016) The importance of
529 *Ficus* (Moraceae) trees for tropical forest restoration. *Biotropica* 48: 413–419.

530 Cushman JH, Compton SG, Zachariades C, Ware AB, Nefdt RJC, Rashbrook VK
531 (1998) Geographic and taxonomic distribution of a positive interaction: ant-tended
532 homopterans indirectly benefit figs across southern Africa. *Oecologia* 116:373–380.

533 del Hoyo J, Elliott A Sargatal J (1992–2002) *Handbook of the Birds of the World*.
534 Vol. 1–7. Lynx Edicions, Barcelona, Spain.

535 del Hoyo J, Elliott A, Christie DA (2003–2011) *Handbook of the Birds of the World*.
536 Vol. 8–16. Lynx Edicions, Barcelona, Spain.

537 Derryberry EP, Claramunt S, Derryberry G, Chesser RT, Cracraft J, Alexio A, Pérez-
 538 Emán J, Remsen JV Jr, Brumfield RT (2011) Lineage diversification and
 539 morphological evolution in a large-scale continental radiation: the Neotropical
 540 ovenbirds and woodcreepers (Aves: Furnariidae). *Evolution* 65:2973–2986.

541 Edwards FA, Edwards DP, Hamer KC, Davies RG (2013) Impacts of logging and
 542 conversion of rainforest to oil palm on the functional diversity of birds in Sundaland.
 543 *Ibis* 155:313–326.

544 ESRI (2014) ArcGIS Desktop: release 10.2.1. Environmental Systems Research
 545 Institute, Redlands, USA.

546 Fischer J, Lindenmayer DB, Manning AD (2006) Biodiversity, ecosystem function,
 547 and resilience: ten guiding principles for commodity production landscapes. *Frontiers*
 548 *in Ecology and the Environment* 4:80–86.

549 Fischer J, Stott J, Law BS (2010) The disproportionate value of scattered trees.
 550 *Biological Conservation* 143:1564–1567.

551 Fox J, Weisberg S (2011) An R companion to applied regression. Second edition.
 552 Sage, California.

553 Flynn FB, Gogol-Prokurat M, Nogeire T, Molinari N, Trautman Richers B, Lin BB,
 554 Simpson N, Mayfield MM, DeClerck F (2009) Loss of functional diversity under land
 555 use intensification across multiple taxa. *Ecology Letters* 12:22–33.

556 Gardner TA, Barlow J, Chazdon RL, Ewers RM, Harvey CA, Peres CA, Sodhi NS
 557 (2009) Prospects for tropical forest biodiversity in a human-modified world. *Ecology*
 558 *Letters* 12:561–582.

559 Giam X, Olden JD (2016) Quantifying variable importance in a multimodel inference
 560 framework. *Methods in Ecology and Evolution* 7: 388-397.

561 Gibbons P, Lindenmayer DB, Fischer J, Manning AD, Weinberg A, Seddon J, Ryan
 562 P, Barrett G (2008) The future of scattered trees in agricultural landscapes.
 563 Conservation Biology 22:1309–1319.

564 Harrison RD (2003) Fig wasp dispersal and the stability of a keystone plant resource
 565 in Borneo. Proceedings of the Royal Society of London B 270:S76–S79.

566 Harrison RD (2013) Ecology of a fig ant-plant. Acta Oecologica 57:88–96.

567 Harvey CA, Medina A, Sánchez DM, Vílchez S, Hernández B, Saenz JC, Maes JM,
 568 Casanoves F, Sinclair FL (2006) Patterns of animal diversity in different forms of tree
 569 cover in agricultural landscapes. Ecological Applications 16:1986–1999.

570 IBM (2013) IBM SPSS Statistics for Windows, Version 22.0. IBM Corp., Armonk,
 571 NY, USA.

572 Karp DS, Daily GC (2013) Cascading effects of insectivorous birds and bats in
 573 tropical coffee plantations. Ecology 95:1065–1074.

574 Laliberté E, Legendre P (2010) A distance-based framework for measuring functional
 575 diversity from multiple traits. Ecology 91:299–305.

576 Laliberté E, Shipley B (2013) FD-package. Measuring functional diversity from
 577 multiple traits and other tools for functional ecology. Version 1.0-11.

578 Laliberté E, Wells JA, DeClerck F, Metcalfe DJ, Catterall CP, Queiroz C, Aubin I,
 579 Bonser SP, Ding Y, Fraterrigo JM, McNamara S, Morgan JW, Merlos DS, Vesik PA
 580 & Mayfield MM (2010) Land-use intensification reduces functional redundancy and
 581 response diversity in plant communities. Ecology Letters 13: 76–86.

582 Leighton M Leighton DR (1983) Vertebrate responses to fruiting seasonality within a
 583 Bornean rain forest. In: Sutton SL, Whitmore TC, Chadwick AC (eds) Tropical rain
 584 forest: ecology and management. Blackwell, Oxford, UK, pp 181–196.

585 Luck GW Daily GC (2003) Tropical countryside bird assemblages: richness,
 586 composition, and foraging differ by landscape context. *Ecological Applications*
 587 13:235–247.

588 Manning AD, Lindenmayer DB, Barry SC (2004) The conservation implications of
 589 bird reproduction in the agricultural “matrix”: a case study of the vulnerable superb
 590 parrot of south-eastern Australia. *Biological Conservation* 120:363–374.

591 Manning AD, Fischer J, Lindenmayer DB (2006) Scattered trees are keystone
 592 structures – implications for conservation. *Biological Conservation* 132:311–321.

593 Matthews TJ, Cottee-Jones HEW, Whittaker RJ (2014) Habitat fragmentation and the
 594 species–area relationship: a focus on total species richness obscures the impact of
 595 habitat loss on habitat specialists. *Diversity and Distributions* 20: 1136–1146.

596 McGill BJ, Enquist BJ, Weiher E, Westoby M (2006) Rebuilding community ecology
 597 from functional traits. *Trends in Ecology & Evolution* 21: 178–185.

598 Melo FPL, Arroyo-Rodríguez V, Fahrig L, Martínez-Ramos M, Tabarelli M (2013)
 599 On the hope for biodiversity-friendly tropical landscapes. *Trends in Ecology and*
 600 *Evolution* 28:462–468.

601 Pereira RAS, Semir J, de Oliveira Menezes A Jr (2000) Pollination and other biotic
 602 interactions in figs of *Ficus eximia* Schott (Moraceae). *Revista Brasileira de Botanica*,
 603 Sao Paulo 23:217–224.

604 Petchey OL, Gaston KJ (2006) Functional diversity: back to basics and looking
 605 forward. *Ecology Letters* 9:741–758.

606 Phalan B, Bertzky M, Butchart SHM, Donald PF, Scharlemann JPW, Stattersfield AJ,
 607 Balmford A (2013) Crop expansion and conservation priorities in tropical countries.
 608 PLoS ONE, 8:e51759.

609 R Core Team (2014) R: A Language and Environment for Statistical Computing.
 610 Vienna: R Foundation for Statistical Computing. Version 3.0.3.

611 Richards SA (2008) Dealing with overdispersed count data in applied ecology.
 612 Journal of Applied Ecology 45:218–227.

613 Schatz B, Kjellberg F, Nyawa S, Hossaert-McKey M (2008) Fig wasps: a staple food
 614 for ants on *Ficus*. Biotropica 40:190–195.

615 Schweiger O, Musche M, Bailey D, Billeter R, Diekötter T, Hendrickx F, Herzog F,
 616 Liira J, Maelfait J-P, Speelmans M, Dziöck F (2007) Functional richness of local
 617 hoverfly communities (Diptera, Syrphidae) in response to land use across temperate
 618 Europe. Oikos 116: 461–472.

619 Şekercioğlu CH (2012) Bird functional diversity and ecosystem services in tropical
 620 forests, agroforests and agricultural areas. Journal of Ornithology 153: S1 53–S1 61.

621 Şekercioğlu CH, Ehrlich PR, Daily GC, Aygen D, Goehring D, Sandi R (2002)
 622 Disappearance of insectivorous birds from tropical forest fragments. Proceedings of
 623 the National Academy of Sciences of the United States of America 99:263–267.

624 Şekercioğlu CH, Loarie SR, Brenes FO, Ehrlich PR, Daily GC (2007) Persistence of
 625 forest birds in the Costa Rican agricultural countryside. Conservation Biology
 626 21:482–494.

627 Shanahan M, So S, Compton SG, Corlett R (2001) Fig-eating by vertebrate
 628 frugivores: a global review. Biological Reviews of the Cambridge Philosophical
 629 Society 76:529–572.

630 Shrivastava RJ, Heinen J (2007) A microsite analysis of resource use around
631 Kaziranga National Park, India. *Journal of Environment and Development* 16:207–
632 226.

633 Stouffer PC, Bierregaard RO Jr (1995) Use of Amazonian forest fragments by
634 understory insectivorous birds. *Ecology* 76:2429–2445.

635 Terborgh J (1986) Keystone plant resources in the tropical forest. In: Soulé ME (ed)
636 *Conservation biology, the science of scarcity and diversity*. Sinauer, Sunderland, MA,
637 USA, pp 330–344.

638 Tschardt T, Şekercioğlu CH, Dietsch TV, Sodhi NS, Hoehn P, Tylianakis JM
639 (2008) Landscape constraints on functional diversity of birds and insects in tropical
640 agroecosystems. *Ecology* 89:944–951.

641 Van Bael SA, Philpott SM, Greenberg R, Bichier P, Barber NA, Mooney KA, Gruner
642 DS (2008) Birds as predators in tropical agroforestry systems. *Ecology* 89:928–934.

643 Wickham H (2009) *ggplot2: elegant graphics for data analysis*. Springer-Verlag, New
644 York.

645 Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009) *Mixed effects models*
646 *and extensions in ecology with R*. New York, Springer.

647

648 **Tables**

649 **Table 1:** Characteristics of the three isolated tree groups surveyed in Assam, India.
 650 DBH is diameter at breast height. Values for DBH, height, and canopy area are mean
 651 $\pm$ standard error. The five most surveyed species are listed in order of decreasing
 652 number of surveys. Fruit refers to large fruit-trees other than *Ficus* and Large to the
 653 category of large trees that did not bear fruit during the study.

Characteristic	<i>Ficus</i>	Fruit	Large
Total no. of individuals surveyed	40	33	31
Total no. of species surveyed	6	12	15
DBH (m)	1.51 $\pm$ 0.13	0.45 $\pm$ 0.02	0.61 $\pm$ 0.05
Height (m)	27.29 $\pm$ 1.40	18.86 $\pm$ 1.03	20.91 $\pm$ 0.89
Canopy area (m ²)	489.32 $\pm$ 67.29	74.01 $\pm$ 7.16	130.11 $\pm$ 21.43
Five most surveyed species (in order of decreasing abundance)	<i>F. religiosa</i> , <i>F. benghalensis</i> , <i>F. rumphii</i> , <i>F. microcarpa</i> , <i>F. benjamina</i>	<i>Artocarpus heterophyllus</i> , <i>Tectona grandis</i> , <i>Artocarpus lakoocha</i> , <i>Syzgium cumini</i> , <i>Toona ciliata</i>	<i>Syzgium cumini</i> (non-fruiting), <i>Albizia lucidor</i> , <i>Albizia procera</i> , <i>Mangifera indica</i> (non-fruiting), <i>Neolamarckia cadamba</i>

654

655

Table 2 Model selection results for a set of the most parsimonious generalized linear models (quasi-Poisson family), modelling the (a) abundance and (b) richness of insectivorous birds in 102 isolated trees, in Assam. The predictor variables included the land use surrounding the trees, tree type, tree size, the distance to the nearest protected area with intact forest (Dist1) and the distance to any forest (Dist2). An interaction between tree type and tree size (Int.) was also included as a term in the model selection as a fixed term. The best model (i.e. lowest QAIC_c) and all models within ΔQAIC_c of < 2 of the best model are given for both (a) and (b). The weight of evidence of each variable, calculated by summing the quasi-Akaike weights of all the models in which a variable was included is also given. + indicates a significant effect of a categorical variable. A blank space indicates that a variable was not included in a model. (L) indicates predictor variables that were log transformed. The ΔQAIC_c (ΔQAIC_c) and QAIC_c weights (wQAIC_c) for each model selection are also presented. *these variables were fixed in the model selection procedure and thus the WoE values are constrained.

a) Abundance

Model Number	Land use	Tree type	Tree size (L)	Dist1 (L)	Dist2 (L)	Int.	ΔQAIC_c	wQAIC _c
1		+	0.39	0.33		+	0	0.38
2		+	0.36			+	0.9	0.24
Weight of evidence	0.18	1*	1*	0.59	0.23	1*		

b) Richness

Model Number	Land use	Tree type	Tree size (L)	Dist1 (L)	Dist2 (L)	Int.	ΔQAIC_c	wQAIC _c
1		+	0.32			+	0	0.39
2		+	0.34	0.13		+	1.83	0.16
3	+	+	0.17			+	1.84	0.16
Weight of evidence	0.27	1*	1*	0.28	0.24	1*		

Table 3 Model selection results for a set of the most parsimonious linear models, modelling the functional dispersion of insectivorous birds in 102 isolated trees, in Assam. The predictor variables included the land use surrounding the trees, tree type, tree size, the distance to the nearest protected area with intact forest (Dist1) and the distance to any forest (Dist2). An interaction between tree type and tree size (Int.) was also included as a term in the model selection as a fixed term. The best model (i.e. lowest AIC_c) and all models within ΔAIC_c of < 2 of the best model are given. The weight of evidence of each variable, calculated by summing the Akaike weights of all the models in which a variable was included, and a bootstrapped baseline (see Materials and methods) with which to compare these values to are also given. + indicates a significant effect of a categorical variable. A blank space indicates that a variable was not included in a model. (L) indicates predictor variables that were log transformed. The ΔAIC_c (ΔAIC_c) and AIC_c weights ($wAIC_c$) for each model selection are also presented. *these variables were fixed in the model selection procedure and thus the WoE values are constrained.

Model Number	Land use	Tree type	Tree size (L)	Dist1 (L)	Dist2 (L)	Int.	ΔAIC_c	$wAIC_c$
1	+	+	-0.06			+	0	0.5
Weight of evidence	0.87	1*	1*	0.26	0.24	1*		
Weight of evidence baseline	0.16	0.23	0.36	0.29	0.29	NA		

Figure legends

Figure 1: A map highlighting the study site location. The inlay shows India and the location of the study landscape within Assam.

Figure 2: The functional dendrogram converted into a tree object, for 33 insectivorous bird species sampled in isolated trees in Assam, India. The dendrogram was constructed by first transforming the species – trait data into a distance matrix (Euclidean distance), and then using the UPGMA clustering algorithm.

Figure 3: Mean insectivore abundance, richness, and FDis (functional dispersion) recorded in isolated tree assemblages in Golaghat District, Assam, India. For *Ficus* trees: n=40; fruit: n=33; and large: n=31. Error bars are standard error.

Figure 4: The relationship between tree size and insectivorous bird species richness for 104 isolated trees in Assam. The data have been split according to the three tree types analysed in the study: fig trees (type 1; red line), non-fig fruiting trees (type 2; green line) and non-fruiting trees (type 3; blue line). The tree size data have been scaled in order for each tree type to cover the same range of tree size; this was achieved by first adding a constant (1.5) to each tree size value (PCA axis; see Methods and material) and then dividing each tree size value by the maximum tree size within that tree type. As the data are not normally distributed we simply fitted loess best fit lines for each of the three tree types, in order to get a rough idea of the patterns.

739

742

743

744

745

746

**Assessing the relative importance of isolated *Ficus* trees to insectivorous birds in
an Indian human-modified tropical landscape – Supporting information**

Biodiversity and Conservation

Thomas J. Matthews^{*+}; H. Eden W. Cottee-Jones⁺; Tom P. Bregman; Robert J.
Whittaker

*Corresponding author

+These authors contributed equally to this work.

Correspondence address: T.J. Matthews, GEES (School of Geography, Earth and
Environmental Sciences), The University of Birmingham, Birmingham, B15 2TT.
Email: txm676@gmail.com.

Supporting information 1:

To calculate the area of land under agricultural production in the world's tropics, we
followed the methodology of Phalan et al. 2013. We defined tropical countries as
those that have at least one-third of their land area between the Tropics of Cancer and
Capricorn, producing a list of 129 countries. We downloaded data on the extent of
total agricultural area in each of these countries from FAO STAT (2014) for the most
recent year (2011). We summed these figures to produce an estimate of the total land
area under agricultural production in tropical countries (2,455,649,900 ha, rounded to
2.5 billion ha).

Supporting information 2:

To compute functional diversity scores, we first collected ecomorphological trait data
on the 34 insectivore species recorded in the study. We measured traits associated
with locomotive behaviour (tarsus length, wing chord, and tail length), dispersal

ability (Kipp's distance, hand-wing index), gape size (gape width), bill structure (culmen length, bill length from nares, bill width, bill depth), and body size (via a Principal Components Analysis, PCA).

Specifically, we measured four specimens of the local population for each species recorded. In almost all cases we were able to measure specimens collected within 150 km of the study area. Two adult males and two adult females of each species were measured with 150 mm outside diameter dial callipers (accurate to 0.1 mm), wing rulers, and tail rulers (accurate to 0.5 mm). The measurements taken were: culmen length (from the base of the skull to the tip of the bill), bill length from nares (from the anterior edge of the nares to the tip of the bill), bill width (the width of the bill at the anterior of the nares), bill depth (the depth of the bill at the anterior of the nares), gape width, tarsus length (the length from the inner bend of the tibiotarsal articulation to the base of the toes, where the scalation pattern changes), wing chord (from the bend in the wing to the unflattened longest primary), Kipp's distance (the distance from the longest primary to the first secondary), and tail length (to the tip of the longest rectrix).

As weight data for birds are often variable (Clark 1979), we preferred to measure body size through a PCA. We initially conducted a pair of PCA analyses, one for locomotive ability (with input measurements of tail length, wing chord, and tarsus length) and one for bill shape (with bill depth, width, and length from nares) using oblique rotation with Kaiser stopping criterion extraction (eigenvalues >1). Each of these PCAs produced two components. In both cases, the first related to size, while the second components were taken as indices for locomotive ability and bill shape, respectively. To produce one index for body size, we ran an additional PCA using the first components from the original analyses (Trisos et al. *in press*). To create an index

for dispersal ability that standardizes for bird size, we calculated the hand-wing index (Claramunt et al. 2012), which is a surrogate for flight performance, migratory behaviour, and natal dispersion in birds.

Supporting information 3:

We used Laliberté and Legendre's functional dispersion (FDis) index to measure functional diversity in our dataset (Laliberté & Legendre 2010). This represents the spread of the species in quantitative trait space by calculating a multidimensional index of the mean distance of an individual species to the centroid of all species in the community (Laliberté & Legendre 2010). A major advantage of FDis over other measures, such as FRic, FEve, and FDiv (Villéger et al. 2008; Mouchet et al. 2010) is that it can be calculated for communities composed of only two species, rather than a minimum of three, which was important for the species-poor insectivore assemblages in the isolated trees. It is also independent of species richness, and can be weighted by abundance, both of which were important considerations for our study.

Supporting information 4:

Table S1: Parameter estimates and standard errors for all terms within the best model, modelling the abundance of insectivorous birds in 102 isolated trees, in Assam. The best model was selected based on comparing QAIC_c values of a complete set of models after fixing the interaction term between tree type and tree size. The predictors included in the best model are tree type (a categorical variable with three levels: 1=*Ficus* trees, 2=non-*Ficus* fruit trees and 3=large non-fruiting trees; see Materials and methods), distance between the tree and the nearest protected area with intact forest (Distance), tree size (the first axis of a PCA using three tree size variables; measured on a log scale) and an interaction between tree size and tree type.

824

Model term	Estimate	Std. error
Intercept	1.20	0.58
Distance	0.33	0.19
Tree type 2	-0.48	0.47
Tree type 3	-0.20	0.45
Tree size	0.39	0.14
Tree type 2 * tree size	-0.37	0.35
Tree type 3 * tree size	0.75	0.39

825

826 **Table S2:** Parameter estimates and standard errors for all terms within the best model,
827 modelling the richness of insectivorous birds in 102 isolated trees, in Assam. The best
828 model was selected based on comparing QAIC_c values of a complete set of models
829 after fixing the interaction term between tree type and tree size. The predictors
830 included in the best model are tree type (a categorical variable with three levels:
831 1=*Ficus* trees, 2=non-*Ficus* fruit trees and 3=large non-fruiting trees; see Materials
832 and methods), tree size (the first axis of a PCA using three tree size variables;
833 measured on a log scale) and an interaction between tree size and tree type.

834

Model term	Estimate	Std. error
Intercept	1.72	0.12
Tree type 2	-1.00	0.21
Tree type 3	-0.79	0.17
Tree size	0.32	0.14
Tree type 2 * tree size	-0.13	0.32

Tree type 3 * tree size	0.65	0.35
-------------------------	------	------

Table S3: Parameter estimates and standard errors for all terms within the best model, modelling the functional dispersion of insectivorous birds in 102 isolated trees, in Assam. The best model was selected based on comparing AIC_c values of a complete set of models after fixing the interaction term between tree type and tree size. The predictors included in the best model are land use (on an ordinal scale: 1=low, 2=medium and 3=high land use intensity), tree type (a categorical variable with three levels: 1=*Ficus* trees, 2=non-*Ficus* fruit trees and 3=large non-fruiting trees; see Materials and methods), tree size (the first axis of a PCA using three tree size variables; measured on a log scale) and an interaction between tree size and tree type.

Model term	Estimate	Std. error
Intercept	1.17	0.49
Land use 2	-0.64	0.34
Land use 3	-1.18	0.41
Tree type 2	-1.43	0.38
Tree type 3	-1.27	0.38
Tree size	-0.06	0.36
Tree type 2 * tree size	0.08	0.49
Tree type 3 * tree size	1.42	0.6

References

Clark GA Jr (1979) Body weights of birds: a review. *The Condor* 81:193–202.

851 FAO STAT (2014) FAO, Rome. <http://faostat3.fao.org/faostat-gateway/go/to/home/E>.
852 Accessed 4 March 2014.

853 Laliberté E, Legendre P (2010) A distance-based framework for measuring functional
854 diversity from multiple traits. *Ecology* 91:299–305.

855 Mouchet MA, Villéger S, Mason NWH, Mouillot D (2010) Functional diversity
856 measures: an overview of their redundancy and their ability to discriminate
857 community assembly rules. *Functional Ecology* 24:867–876.

858 Phalan B, Bertzky M, Butchart SHM, Donald PF, Scharlemann JPW, Stattersfield AJ,
859 Balmford A (2013) Crop expansion and conservation priorities in tropical countries.
860 *PLoS ONE* 8:e51759.

861 Trisos CH, Petchey OL, Tobias JA (2014) Unraveling the interplay of community
862 assembly processes acting on multiple niche axes across spatial scales. *The American*
863 *Naturalist* 184:593–608.

864 Villéger S, Mason NWH, Mouillot D (2008) New multidimensional functional
865 diversity indices for a multifaceted framework in functional ecology. *Ecology*
866 89:2290–2301.

867 Wickham H (2009) *ggplot2: elegant graphics for data analysis*. Springer, New York,
868 USA.

869