Impact of LULCC on the emission of BVOCs during the 21st century

Szogs, Sebastian; Arneth, Almut; Anthoni, Peter; Doelman, Jonathan C.; Humpenöder, Florian; Popp, Alexander; Pugh, Thomas; Stehfest, Elke

DOI:

[10.1016/j.atmosenv.2017.06.025](https://doi.org/10.1016/j.atmosenv.2017.06.025)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Szogs, S, Arneth, A, Anthoni, P, Doelman, JC, Humpenöder, F, Popp, A, Pugh, T & Stehfest, E 2017, 'Impact of LULCC on the emission of BVOCs during the 21st century', *Atmospheric Environment*, vol. 165, pp. 73-87.
<https://doi.org/10.1016/j.atmosenv.2017.06.025>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Impact of LULCC on the emission of BVOCs during the 21st century

Sebastian Szogs, Almut Arneth, Peter Anthoni, Jonathan C. Doelman, Florian Humpenöder, Alexander Popp, Thomas A.M. Pugh, Elke Stehfest

PII: S1352-2310(17)30406-5

DOI: [10.1016/j.atmosenv.2017.06.025](https://doi.org/10.1016/j.atmosenv.2017.06.025)

Reference: AEA 15387

To appear in: *Atmospheric Environment*

Received Date: 13 November 2016

Revised Date: 12 June 2017

Accepted Date: 14 June 2017

Please cite this article as: Szogs, S., Arneth, A., Anthoni, P., Doelman, J.C., Humpenöder, F., Popp, A., Pugh, T.A.M., Stehfest, E., Impact of LULCC on the emission of BVOCs during the 21st century, *Atmospheric Environment* (2017), doi: 10.1016/j.atmosenv.2017.06.025.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Impact of LULCC on the emission of BVOCs during the 21st century

Authors:

Szogs Sebastian^[1,X], Arneth Almut^[1], Anthoni Peter^[1], Doelman Jonathan C.^[2], Humpeöder Florian^[3], Popp Alexander^[3], Pugh Thomas A. M.^[1,4], Stehfest Elke^[2]

^[1] Karlsruhe Institute of Technology (KIT), Institute of Meteorology and Climate Research – Atmospheric Environmental Research (IMK-IFU), Kreuzeckbahnstraße 19, 82467 Garmisch-Partenkirchen, Germany.

^[2] PBL, Netherlands Environmental Assessment Agency, Postbus 30314, 2500 GH The Hague, Netherlands.

^[3] Potsdam Institute for Climate Impact Research (PIK), Telegraphenberg A31, 14473 Potsdam, Germany.

^[4] School of Geography, Earth & Environmental Sciences Birmingham and Institute of Forest Research, University of Birmingham, Edgbaston, Birmingham, B15, 2TT, UK.

Corresponding author: sebastian.szogs@web.de

Abstract

Land-use and land-cover change (LULCC) is one of the key drivers of anthropogenic climate change. In addition to greenhouse gases such as CO₂ or CH₄, LULCC affects also the emission of other carbon trace gases such as biogenic volatile organic compounds (BVOCs). We investigate the impact of changing LULCC on the emission of isoprene and monoterpenes during the 21st century using seven different land-use projections, applying the dynamic vegetation modelling framework LPJ-GUESS. Climate change, and atmospheric CO₂-concentration are based on the RCP2.6 scenario. The different LULCC-scenarios explore the impact of different land-based climate change mitigation strategies (such as afforestation and avoided deforestation, or bioenergy). We show that the increase of land area under crops or grassland would lead to a significant decrease of BVOC emissions, with a strong negative correlation between the fraction of managed global land area and the emission of isoprene and monoterpenes. But the choice of crops is important, especially for the bioenergy scenarios in which increasing fractional cover leads to decreasing BVOC emissions in our simulations; use of woody bioenergy crops can reverse this decrease. The strong impact of LULCC on the global and regional emission of BVOCs implies the need to include the impact of these changes in projections of atmospheric composition and air quality.

Keywords:

- BVOC
- LULCC
- Isoprene
- Monoterpene

1. Introduction

While in the year 1700 an estimated 95% of the ice-free land area was in a (semi) natural state, by 2000 only about 45% remained relatively unaffected by human activities (Ellis et al., 2010). Between 1990 and 2005 alone, nearly 9.000.000 ha of forest area were cleared for cropland and pasture expansion, and human settlements (FAO and JRC, 2012). The rate of deforestation has weakened over recent years, but a further growing population will still need more cropland to meet its energy and food demand (Lambin and Meyfroidt, 2011, Levis, 2010), even though the total area needed depends on many highly uncertain socio-economic factors (Eitelberg et al., 2015). In addition, the need for land-based climate change mitigation by providing biomass for bioenergy, conserving carbon-rich ecosystems from agricultural expansion or enhancing terrestrial carbon sequestration by afforestation will be an additional important driver of land-use change in future (Smith et al. 2014, Popp et al. 2014a).

Land-use and land-cover change (LULCC) is one of the key drivers of anthropogenic greenhouse gas (GHG) emissions. LULCC-related emissions include direct and indirect greenhouse gas emissions from agricultural activities such as livestock farming, manure management, fertiliser use and paddy rice that contribute around 12% of today's total GHG emissions and an additional ca. 10% arise as CO₂ emissions from deforestation (Smith et al., 2008, Smith et al. 2014, Tubiello et al., 2015). Over the period from 1870 to 2014, for example, emissions due to land-cover change were equivalent to nearly 30% of total anthropogenic CO₂ emissions (Le Quéré et al., 2015, 2016). To evaluate future emission from land clearance or the potential of mitigating climate change through land-ecosystem carbon uptake, or through avoided future emissions from deforestation, future land-use scenarios explore different pathways that are based on different assumptions such as technological progress, population growth, or need for energy (e.g., O'Neill et al., 2015, Hurtt et al, 2011). But not only the long-lived GHGs such as carbon dioxide (or methane and nitrous oxide) are affected by LULCC, also atmospherically short-lived species are impacted. Among these are biogenic volatile organic compounds (BVOC) (Rosenkranz et al., 2015). Atmospherically short-lived, their high reactivity and large terrestrial source makes them a crucial component

of atmospheric chemistry. From a mass perspective, isoprene (2-methyl-1,3-butadiene, C_5H_8) is the most important compound, with an estimated emission strength of around 400-600 Tg $C a^{-1}$ (Guenther et al., 2012 Lathière et al., 2006, Arneth et al., 2008). A second important compound group, monoterpenes, consist of two isoprene units ($C_{10}H_{16}$) and have considerably lower (but equally uncertain to isoprene) global total emissions (Arneth et al., 2008).

Once emitted, isoprene is primarily lost through oxidation by the hydroxyl radical, and in the presence of sufficient NO_x leads to the increased formation of tropospheric ozone (O_3), a greenhouse-gas which is also toxic to organisms (Atkinson, 2000, Atkinson and Arey, 2003, Jerret 2009, Ainsworth 2012). Through its atmospheric oxidation by hydroxyl, isoprene is believed to reduce the capacity of the atmosphere to oxidize methane, thus enhancing methane lifetime (Poisson et al., 2000). However, uncertainty as to the extent to which isoprene is a net sink for hydroxyl radicals means that the extent of any reductions is not well understood (Lelieveld et al., 2008; Fuchs et al., 2013). Monoterpenes, but also isoprene, contribute to the growth of secondary organic aerosol particles (SOA) (Carslaw et al., 2010, Kulmala et al., 2005). As aerosols scatter and reflect radiation but also function as cloud condensation nuclei (CCN), SOA (and hence BVOC) affect climate in various, complex ways (direct and indirect effects) related to albedo, cloud lifetime, and surface-atmosphere feedbacks (Spracklen et al., 2008, Carslaw et al., 2010, 2013, Paasonen et al., 2013, Kulmala et al., 2014).

The BVOC total emission strength and underlying regional patterns are still largely unknown, since large-scale observations are difficult to obtain or only indirectly possible. Isoprene emissions, for instance, have been inferred from satellite remote sensing estimates of formaldehyde (HCHO) in the air column. Formaldehyde is an isoprene oxidation product, but interpretation of the signal is complicated by the need to combine such observations with chemistry transport models and because of other processes, such as biomass burning, also affecting HCHO levels (Barkley et al., 2012, Palmer et al., 2006). Ecosystem-atmosphere flux measurements that have become standard over recent decades for monitoring and upscaling of CO_2 exchange (Jung et al., 2011, Reichstein et al., 2007) still pose methodological challenges for BVOC, due to the lack of appropriate fast and easy-maintenance sensors that would allow long-term measurements at different locations. From the available data today it appears that the main source of isoprene emissions is in the tropics, temperate forests emit in a medium range, while boreal forests emit less isoprene, but comparatively large amounts of monoterpene (Guenther et al., 1995, Monson et al., 2007, Arneth et al., 2008a, Harrison et al., 2013). Besides this very general state of knowledge on relative source strengths, fine-

scaled emission patterns, interannual variability and/or seasonality of emissions, and response to underlying biotic or abiotic drivers remain poorly understood.

Models of ecosystem BVOC emissions typically are based on algorithms that account for the known strong leaf-temperature and, in case of isoprene and some monoterpene emitters, radiation dependence, in combination with a number of other factors such as leaf age or weather history (Niinemets et al., 2010). These algorithms are parameterized either for the leaf- or the canopy scale, and are combined with vegetation maps or dynamic global vegetation models to simulate the necessary ecosystem properties (Arneth et al., 2008, Guenther et al., 2012 Niinemets et al. 2010). While using fixed vegetation maps allows to project BVOC emissions into the future, accounting for effects of climate change, coupling emissions algorithms with dynamic global vegetation models (DGVMs) is required to examine combined effects of climate and atmospheric CO₂ concentration on emissions, for instance through enhanced vegetation productivity and leaf area index, and to examine emission response due to changes in vegetation cover (e.g. Arneth et al., 2007a, Schurgers et al., 2011, Lathière et al., 2005).

Simulated BVOC emission potentials are greatly affected by the emission capacity (i.e. emissions under standardized environmental conditions) for a given species or vegetation type (Niinemets et al, 2010). Given that there are typically large differences between woody and herbaceous vegetation both for isoprene, but in particular also for monoterpene emissions, it seems surprising that so far only a relatively small number of studies has focused on land-use change effects on historical and future BVOC emissions (e.g., Lathière et al., 2006, 2010, Squire et al., 2014, Wu et al., 2012, Chen et al., 2009, Ganzeveld et al., 2010, Heald and Spracklen, 2015, Unger, 2013, Unger, 2014, Hantson et al., 2017). From these studies a consensus seems to emerge such that LULCC (in particular when expressed as converting woody into crop or pasture vegetation) decreases global emissions of isoprene and monoterpenes. For instance, Lathière et al. (2010) estimated that by the end of the 20th century anthropogenic cropland expansion reduced isoprene emissions by 15% compared to the beginning of the century. By the year 2095, Squire et al. (2014) estimated a 40% decrease in isoprene emissions due to the effects of land-use change compared to the year 2000. These projections were based on the Sheffield Dynamic Global Vegetation model with a scenario of cropland expansion in the tropics and the SRES B2 emission and climate scenario. Lathière et al. (2006) simulated effects of tropical deforestation over the period from 1983 to 1995 with the ORCHIDEE model, using the ISLSCP-II (International Satellite Land-Surface Climatology Project) satellite-based climate archive and showed a decrease of 41% in monoterpene emissions in the tropics, or 29% decrease globally.

Despite of the large effects of on emissions of isoprene and monoterpenes so far little focus has been on comparing different future LULCC scenarios. In particular, future LULCC, especially in the light of ambitious mitigation targets, will include not only deforestation but also reforestation and/or expansion of bioenergy plantations. To study this further, we apply the dynamic global vegetation model LPJ-GUESS (Smith et al., 2014) to examine how emissions of isoprene and monoterpene compounds respond to changed global vegetation cover, focusing especially on stylized scenarios of maximizing afforestation and reforestation (jointly with avoided deforestation), as well as maximizing land used for growing dedicated 2nd generation bioenergy plants.

2. Methods

2.1 LPJ-GUESS

We used the LPJ-GUESS modelling framework (Smith et al., 2014), which simulates global natural vegetation patterns as well as carbon and water cycles, on basis of climate data input, atmospheric carbon dioxide concentration, nitrogen deposition and soil physical properties. LPJ-GUESS is also able to simulate the emission of BVOCs in response to changing climate and CO₂-concentrations in the atmosphere. Isoprene production is calculated adopting the algorithm from Niinemets et al. (1999), based on the temperature and light dependence as a function of the electron transport required for photosynthesis, and temperature response adopted for rates seen for isoprene. The algorithm was supplemented to include the frequently observed inhibition of emissions, seen at leaf scale, under increasing atmospheric CO₂ (Arneth et al. 2007a). Monoterpene emissions are calculated using similar algorithms for their production rates, but accounting also for storage, as described in Schurgers et al (2009).

DGVMs represent vegetation through plant functional types (PFT), which – in case of LPJ-GUESS - compete at each grid location for water, light and nutrients. Every gridcell has area fractions prescribed for natural and agricultural land. For natural vegetation we used 11 plant functional types as in Smith et al. (2014), 9 PFTs representing boreal, temperate and tropical woody vegetation, while two PFTs represent C3 and C4 herbaceous species. Six different land-use categories were specified: C3 and C4 pastures, and irrigated and non-irrigated crops that were also simplified here by C3 and C4 vegetation. C4 plants are limited by a minimum temperature of 15.5°C for establishment and can thus concur in some regions with C3 plants. Representing harvest and grazing, 50% of the aboveground biomass on crops and pasture areas, respectively, are removed each year and not returned into the soil through litter input. Bioenergy crops were also assumed to be C3 or C4 crops, no woody bioenergy is considered here, which for BVOC emissions is a conservative approach (Rosenkranz et al., 2015). A PFT-

specific standardized leaf-level BVOC emission capacity that corresponds to emission factors (at 30°C temperatures and photosynthetically-active radiation flux of $1000\mu\text{mol m}^{-2} \text{s}^{-1}$ (Guenther et al, 1995, Arneth et al., 2007, Schurgers et al., 2009) was set for every PFT, with values as in Schurgers et al. (2011). For crops we applied the values of C3 and C4 grasses. The model was run with a spatial resolution of $0.5^\circ \times 0.5^\circ$. Since LPJ-GUESS contains some stochastic features in its representation of growth dynamics (Smith et al., 2001, 2014), a number of replicated patches (here set to 10) are specified for each gridcell. In each patch PFTs are competing for resources, and patches are subsequently averaged to represent the output per grid-cell. Atmospheric CO_2 concentration and N deposition for model spin-up and historical simulations over the 20th century were as described in Smith et al (2014). For the pre-industrial spin-up to 1901, we recycled the detrended climate data of 1901-1930 and ran the model for 500 years until the carbon pools were in equilibrium. During the spin-up period atmospheric CO_2 was kept at a constant mixing ratio of 296 ppmv (value in year 1900) and the land-cover was held constant at the historical distribution in year 1901 as given in Hyde 3.1 (Goldewijk et al., 2011). As climate information, we used the IPSL-CM5A-LR Earth System Model (ESM) simulation output (Dufresne et al., 2013), at daily resolution and bias-corrected in the ISI-MIP project (Warszawski et al., 2014). The IPSL-CM5A-LR climate provides data for the years 1950 to 2099. To provide climate data over the whole time period of the LU scenarios (1901 to 2099) we extended the climate to the period 1901 to 1949 by randomly choosing climate data of the years 1950 to 1959. Simulations over the 21st century are based on the RCP2.6-emission scenario concerning evolution of CO_2 -concentration in the atmosphere (van Vuuren et al. 2011).

2.2 Land-use scenarios

The land-use data over the 21st century is derived from two models, the Model of Agricultural Production and its Impact on the Environment (MAGPIE) (Lotze-Campen et al., 2008, Popp et al. 2014b) and the integrated model to assess the global environment (IMAGE) (van Vuuren et al., 2011, Stehfest et al., 2014). Those models calculate future land use and agriculture dynamics for a set of economic regions, considering population growth, economic development, and technological progress affecting potential crop yields, as well as climate change. Land-use changes over the 20th century were based on Hyde 3.1 (Goldewijk et al., 2011) and the projected future output from MAGPIE and IMAGE were mapped onto Hyde 3.1 directly in order to create a harmonized historical-to-future time series. In addition to these scenarios, we also used projections from Hurtt et al. (2011) for RCP 2.6, which are also based on IMAGE.

Three scenarios from MAgPIE and IMAGE were implemented to investigate the impact of different land-use projections on BVOC emissions. First, a scenario based on SSP2 (REF) that does not include land-based climate mitigation (Sitch et al. in prep, see also Humpenöder et al. 2015) was used as a reference. Second, ambitious land-based mitigation targets are achieved through bioenergy in combination with carbon capture and storage (BECCS). In a third scenario a high land-based mitigation target is achieved by means of avoided deforestation and afforestation/reforestation (ADAFF). In each scenario, a number of basic constraints had to be met (e.g. food supply and a minimum area set aside for conservation). For comparison, we also calculated BVOC emission changes using the “standard” LULCC historical-to-future estimates provided for RCP2.6 as described in Hurtt et al. (2011; H11). The managed area fraction used in LPJ-GUESS (pasture, crops and irrigated crops) are derived from the scenarios. Natural land, especially woody vegetation cover was not considered to be managed for forest products in this version of the model.

2.3 Differences between the MAgPIE, IMAGE and H11 model frameworks

The IMAGE and MAgPIE mitigation scenarios are based on the SSP framework and attempt to maximize land-based mitigation but no explicit mitigation target was set, resulting in different land requirements. The variation in land requirements also reflects different modelling approaches. In IMAGE, land-use allocation follows a rule-based approach according to sustainability criteria, which implies that bioenergy production or afforestation can only take place on land that is not needed for food production. In contrast, competition for land between food production and land-based mitigation is explicitly modeled in MAgPIE based on a cost minimization procedure.

The H11 scenario is based on the IMAGE RCP 2.6 scenario (Hurtt et al., 2011, van Vuuren et al., 2011). The largest part of the differences between H11 and the IMAGE REF, BECCS and ADAFF scenarios arises from different socio-economic assumptions. In contrast to the BECCS and ADAFF scenarios the H11 scenario also has a particular climate target (radiative forcing of 2.6 Wm^{-2} in 2100) which considers a broad portfolio of mitigation measures across different sectors. However, the H11 scenario did not include afforestation as a mitigation measure. Lastly, the RCP 2.6 scenario which was the basis for the H11 scenario was produced with IMAGE 2.4 (Bouwman et al., 2006) while the IMAGE mitigation scenarios were produced with IMAGE 3.0 (Stehfest, 2014). Model developments such as the coupling to LPJmL (Bondeau et al., 2007) also explain part of the differences.

3. Results

Maps of the different vegetation cover patterns in the LU scenarios, translated into the LPJ-GUESS dominant PFTs, for the end of 2100 are shown in Figure 1. As expected, the distribution differs strongly both between the scenarios, but also between the different LU models. The reference scenario in MAgPIE and H11 are fairly similar in terms of cropland and pasture distribution, but have some obvious differences e.g., in the assumption of where crops are irrigated (for instance in India). By contrast, the reference IMAGE case shows a larger area dominated by pastures, especially in the northern hemisphere. In both ADAFF stylized scenarios the tropical woody PFTs have a larger areal extent compared to REF, BECCS and also the H11 land cover, a similar observation can be made for parts of the temperate zone (e.g., in the central US and parts of Europe and Asia), where temperate woody vegetation expands. In the BECCS scenario the tropical woody PFTs are reduced by expansion of C4 cropland and C4 pasture, especially in southeastern Brazil and parts of central Africa. In temperate regions such as central USA, Europe or central Asia C3 croplands expand.

Figure 1: Maps showing dominant PFTs per gridcell (PFTs with the highest LAI within each gridcell). Mean values for the period 2081-2100 for all seven different model projections. Abbreviations are: TrBR: Tropical broadleaved raingreen; TrIBE: Tropical shade intolerant broadleaved evergreen; TrBE: Tropical broadleaved evergreen; TeBE: Temperate broadleaved evergreen; IBS: Shade intolerant broadleaved summergreen; TeBS: Temperate broadleaved summergreen; BNS: Boreal needleleaved summergreen; BINE: Boreal shade intolerant needleleaved evergreen; BNE: Boreal needleleaved evergreen; C4-CI: C4 Crops Irrigated; C3-CI: C3 Crops Irrigated; C4-C: C4 Crops; C3-C: C3 Crops; C4-P: C4 Pasture; C3-P: C3 Pasture; C4-G: C4 Grass; C3-G: C3 Grass. (Crops and Pasture simulated as grasses, see methods).

Figure 2 shows the corresponding development of the total fraction of area under some form of crop or pasture in all LU scenarios. At the end of the 21st century, the total land area under agriculture is smallest for the ADAFF scenarios and highest for the MAgPIE BECCS scenario, the latter being more or less similar to the H11 land cover estimates. For only cropland area the development of almost all the scenarios is equivalent to the development of the total fraction of area under some form of agriculture, except for MAgPIE BECCS, where the cropland area fraction shows a strong increase from 2050 on and stays at an even higher value than H11.

Figure 2: Global area fraction under agriculture: total (Pasture and Crops), and cropland-only area fraction (Crops) for all seven LULLC-scenarios.

The mean value of isoprene emission for the REF scenario of both IAMs and the H11 land cover for present day (2000-2010) is $\sim 480 \text{ Tg C a}^{-1}$ (Figure 3), which is well within the range of $412\text{--}601 \text{ Tg C a}^{-1}$ given in Arneth et al. (2008). The mean value of monoterpene emission for the REF scenarios and the H11 land cover is $\sim 39 \text{ Tg C a}^{-1}$ (compared with $32\text{--}127 \text{ Tg C a}^{-1}$; Arneth et al., 2008). Figure 3 shows the change of the BVOC emissions in response to all simulated scenarios. Monoterpene as well as isoprene emissions follow a declining trajectory from 1901 until 2000. This decline is partially due to the CO_2 inhibition of isoprene production and reflecting the increase in the atmospheric CO_2 concentration over the 20th century (Le Quéré et al., 2015). Historical land-cover change enhances

1 the decline in emissions, as woody vegetation with relatively high emission factors is replaced by
 2 herbaceous vegetation cover. In the historical period the difference between H11 and the two IAMs
 3 differ more for monoterpenes than isoprene due to small differences in vegetation cover in the
 4 boreal biome (note also the different scales on the y-axes in Figure 3).

Figure 3: Global sums of emissions for all seven LULCC-scenarios from 1900-2100 [Tg C a⁻¹]. Top panel: isoprene, bottom panel: monoterpene emissions.

Over the 21st century, emissions in the two REF scenarios remain close to present-day levels, with around 470-500 Tg C a⁻¹ for isoprene and 38-40 Tg C a⁻¹ for monoterpenes. Lowest isoprene and monoterpene emissions overall are computed from the H11 land cover (Figure 3, 4, Table 1 & 2). In the ADAFF scenario emissions increase very strongly, especially for isoprene, until the end of the 21st century (Tables 1, 2) in both IAM stylized scenarios, with large differences arising from the differences in projected total vegetation cover changes in MAgPIE and IMAGE. Not surprisingly, the increase in BVOC emissions is dominated by woody PFTs. Emissions in the BECCS scenario show very small changes in different directions, with total isoprene emissions increasing by 1% for IMAGE, but decreasing by -1% in MAgPIE. Monoterpene emissions decrease by -2% (MAgPIE) and do not show any change for IMAGE. In the same period the area fraction under agriculture increases by ~1.5%. When considering cropped areas only (not irrigated) in the MAgPIE BECCS scenario emissions increase by 89% for isoprenes and by 100% for monoterpenes. This increase in emissions is closely linked to the area of cropland (Figure 2), which increases from 10% to 17% in MAgPIE BECCS for the same time period, but the effect on global emissions is small because of the low crop-PFT emission factors. For IMAGE, the increase in emissions from crops is 45% for isoprene and 10% for monoterpenes, while cropland increases by 5% compared to present day. The H11 land cover shows globally the highest increase in area fraction under management (+3% compared to present-day) and associated emissions decrease by -5% for isoprene and by -4% for monoterpene. Cropland increases in H11 are about +5% compared to present day.

Table 1: Mean values of isoprene emission for the 2091-2100 period and the change in percentage compared to the mean value of a present day period (2001-2010) in brackets for each model projection. Regions are split in latitudinal bands (Boreal: 60°N-90°N, Temperate North: 30°N-60°N, Tropics: 30°S-30°N and Temperate South: 30°S-60°S).

	MAgPIE Ref	MAgPIE Beccs	MAgPIE Adaff	IMAGE Ref	IMAGE Beccs	IMAGE Adaff	H11
BNE	5.8 (-26%)	5.7 (-27%)	6.2 (-21%)	6.2 (-23%)	5.6 (-28%)	6.3 (-19%)	5.4 (-31%)
BINE	3.3 (-13%)	3.2 (-16%)	4.6 (+21%)	3.7 (-5%)	3.1 (-18%)	4 (+5%)	2.9 (-22%)
BNS	1.4 (+27%)	1.4 (+27%)	1.4 (+27%)	1.4 (+27%)	1.4 (27%)	1.4 (+27%)	1.4 (+27%)
TeBS	10.9 (+8%)	10.7 (+6%)	15.2 (+50%)	12.4 (+19%)	9.4 (-7%)	14.3 (+42%)	9.8 (-10%)
			47.9				
IBS	30.7 (+41%)	29.9 (+38%)	(+121%)	35.4 (+60%)	28.8 (+33%)	41.2 (+90%)	29 (+31%)
TeBE	35.6 (+5%)	31.7 (-7%)	56.2 (+65%)	32.3 (+12%)	28.6 (-16%)	41.1 (+21%)	34.4 (-5%)
			243.9			214.4	
TrBE	199 (+4%)	187.8 (-2%)	(+27%)	204.4 (+4%)	196.6 (+2%)	(+12%)	179.6 (-9%)
			117.6				
TriBE	78.7 (-3%)	74.1 (-9%)	(+44%)	80 (0%)	78.3 (-4%)	91 (+12%)	65.5 (-18%)
			94.2				
TrBR	51.5 (+20%)	48.5 (+13%)	(+119%)	54.8 (+27%)	52.4 (+22%)	66.1 (+54%)	47.7 (+17%)
C3-G	12.5 (-21%)	11.8 (-26%)	18.5 (+16%)	13.6 (-15%)	12.4 (-22%)	15.1 (-5%)	11.3 (-28%)
C4-G	5.2 (+6%)	4.9 (0%)	7.2 (+47%)	6 (+15%)	5.7 (+16%)	6.8 (+39%)	4.5 (-10%)
C3-P	21.3 (-19%)	18 (-31%)	11.4 (-57%)	24.9 (-13%)	24.9 (-5%)	19.5 (-26%)	21.6 (-15%)
C4-P	15.6 (-5%)	13.1 (-20%)	6.5 (-60%)	16.3 (+1%)	16.3 (-1%)	13.2 (-20%)	15.7 (+1%)

C3-C	13.7 (+18%)	20.3 (+75%) 12.6	4 (-66%)	10.9 (-18%)	16.9 (+46%)	9.5 (-18%)	19.3 (+18%)
C4-C	6.5 (+12%)	(+117%)	2.1 (-66%)	6.7 (-4%)	8.3 (+43%)	5.8 (-5%)	12.7 (+55%)
C3-CI	11.8 (-6%)	11.5 (-9%)	9.7 (-24%)	7.2 (-3%)	7.2 (-43%)	6.8 (-46%)	NA
C4-CI	6.8 (+33%)	5.2 (+2%)	3.5 (-27%) 612.9	2.7 (+23%)	2.7 (-47%)	2.5 (-48%)	NA
Natural	434.8 (+5%)	409.8 (-1%)	(+47%)	450 (+8%)	422.3 (+2%)	(+21%)	391.5 (-7%)
Pasture	37 (-13%)	31.1 (-27%)	17.9 (-58%)	41.2 (-8%)	41.2 (-3%)	32.7 (-23%)	37.3 (-9%)
Total	510.5 (+3%)	490.5 (-1%)	650 (+32%)	518.7 (+6%)	498.5 (+1%)	(+13%)	460.8 (-5%)
Crop							
Rainfed	20.2 (+17%)	32.9 (+89%)	6 (-66%)	17.7 (-13%)	25.2 (+45%)	15.3 (-14%)	32 (+30%)
Crop							
Irrigated	18.6 (+4%)	16.7 (-6%)	13.2 (-24%) 587.2	9.9 (+4%)	9.9 (-44%)	9.3 (-47%)	NA
Woody	417.1 (+6%)	393.1 (0%)	(+49%) 504.8	430.5 (+9%)	404.2 (+3%)	(+24%) 424.3	375.7 (-6%)
Tropics	389.1 (+1%)	372.2 (-3%)	(+32%)	394.2 (+3%)	382.9 (0%)	(+12%)	349.1 (-8%)
Boreal	7.4 (+45%)	7.4 (+45%)	(+547%)	7.4 (+45%)	7.4 (+45%)	7.5 (+47%)	7.4 (+45%)
Temperate	101.1		121.9	104.2		113.4	
N	(+13%)	97.8 (+9%)	(+36%)	(+18%)	95.6 (+9%)	(+29%)	93 (+7%)
Temperate							
S	12.9 (-16%)	13.1 (-15%)	15.8 (+3%)	12.9 (-13%)	12.7 (-14%)	13.7 (-6%)	11.3 (-22%)

Table 2: Mean values of monoterpene emission for the 2091-2100 period and the change in percentage compared to the mean value of a present day period (2001-2010) in brackets for each model projection.

	MAGPIE Ref	MAGPIE Beccs	MAGPIE Adaff	IMAGE Ref	IMAGE Beccs	IMAGE Adaff	H11
BNE	3.4 (-26%)	3.4 (-26%)	3.7 (-20%)	3.7 (-23%)	3.3 (-18%)	3.7 (-23%)	3.2 (-30%)
BINE	1.9 (-17%)	1.9 (-17%)	2.7 (+17%)	2.2 (-4%)	1.9 (-14%)	2.4 (+4%)	1.7 (-23%)
BNS	0.9 (+29%)	0.9 (+29%)	0.9 (+29%)	0.9 (+29%)	0.9 (0%)	0.9 (+29%)	0.9 (+29%)
TeBS	0.4 (0%)	0.4 (0%)	0.6 (+50%)	0.5 (+25%)	0.4 (0%)	0.5 (+25%)	0.4 (0%)
IBS	1.2 (+50%)	1.2 (+50%)	1.9 (+137%)	1.4 (+56%)	1.1 (0%)	1.6 (+100%)	1.1 (+22%)
TeBE	2.4 (+4%)	2.1 (-9%)	3.8 (+65%)	2.2 (+16%)	1.9 (0%)	2.7 (+42%)	2.3 (-4%)
TrBE	6.6 (+3%)	6.2 (-3%)	8.1 (+27%)	6.8 (+5%)	6.5 (+3%)	7.1 (+9%)	6 (-9%)
TrIBE	2.6 (-4%)	2.5 (-7%)	3.9 (+44%)	2.7 (0%)	2.6 (0%)	3 (+15%)	2.2 (-19%)
TrBR	2.8 (+22%)	2.6 (+13%)	5 (+117%)	2.9 (+26%)	2.8 (+8%)	3.5 (+59%)	2.6 (+18%)
C3-G	1.4 (-22%)	1.4 (-22%)	2.1 (+17%)	1.5 (-17%)	1.4 (-7%)	1.7 (-6%)	1.3 (-28%)
C4-G	1.6 (+7%)	1.5 (0%)	2.2 (+47%)	1.8 (+12%)	1.8 (+20%)	2.1 (+40%)	1.4 (-7%)
C3-P	2.3 (-21%)	2 (-31%)	1.3 (-55%)	2.7 (-13%)	2.7 (-4%)	2.1 (-32%)	2.4 (-14%)
C4-P	4.7 (-4%)	4 (-18%)	2 (-59%)	4.9 (0%)	4.9 (+4%)	4 (-20%)	4.7 (0%)
C3-C	1.5 (+25%)	2.2 (+69%)	0.4 (-69%)	1.2 (-4%)	1.8 (+20%)	1 (-33%)	2.1 (+17%)
C4-C	2 (+18%)	3.8 (+124%)	0.6 (-67%)	2 (-5%)	2.5 (+4%)	1.7 (-23%)	3.8 (+52%)
C3-CI	1.3 (0%)	1.2 (-8%)	1 (-23%)	0.8 (0%)	0.8 (0%)	0.7 (-13%)	NA
C4-CI	2 (+33%)	1.6 (+7%)	1.1 (-21%)	0.8 (+14%)	0.8 (+14%)	0.7 (0%)	NA
Natural	25.3 (-2%)	24.1 (-7%)	34.9 (+35%)	26.6 (+2%)	24.6 (-2%)	29.4 (+15%)	23 (-12%)
Pasture	7.1 (-9%)	5.9 (-24%)	3.2 (-59%)	7.6 (-5%)	7.6 (+1%)	6.1 (-26%)	7.1 (-5%)
Total	39.1 (-1%)	38.8 (-2%)	41.2 (+4%)	39 (0%)	38.1 (0%)	39.7 (+2%)	36.1 (-4%)
Crop							
Rainfed	3.4 (+13%)	6 (+100%)	1.1 (-65%)	3.2 (-11%)	4.3 (+10%)	2.8 (-22%)	5.9 (+40%)
Crop							
Irrigated	3.3 (+14%)	2.8 (-3%)	2.1 (-25%)	1.6 (+14%)	1.6 (+7%)	1.5 (+7%)	NA
Woody	22.3 (-1%)	21.2 (-6%)	30.6 (+36%)	23.2 (+3%)	21.5 (-4%)	25.6 (+15%)	20.4 (-10%)
Tropics	24.8 (+1%)	24.7 (0%)	25.7 (+4%)	24.4 (+2%)	24.3 (+1%)	24.7 (+3%)	22.9 (-2%)
Boreal	2.1 (+17%)	2.1 (+17%)	2.1 (+17%)	2.1 (+17%)	2 (+11%)	2.1 (+17%)	2 (+5%)
Temperate							
N	11 (-6%)	10.8 (-8%)	12.2 (+4%)	11.4 (-3%)	10.7 (-9%)	11.8 (+1%)	10 (-11%)
Temperate							
S	1.2 (-14%)	1.2 (-14%)	1.3 (-7%)	1.2 (-8%)	1.1 (-15%)	1.2 (-8%)	1.1 (-15%)

Figures 4 and 5 show isoprene and monoterpene emission difference maps for all scenarios. For isoprene, the clearest difference emerges between the ADAFF and the REF scenario, in particular for MAGPIE. Especially in the tropics the emission differences can be large, with up to $15 \text{ g C m}^{-2} \text{ a}^{-1}$ higher emissions in the ADAFF scenario and reflecting the high emissions simulated for tropical regions (Guenther et al., 1995, Arneth et al., 2011). Differences between the ADAFF scenario and the reference cases for monoterpenes were lower and more broadly distributed between tropical and temperate regions for the MAGPIE case in which differences were up to $0.6 \text{ g C m}^{-2} \text{ a}^{-1}$. By contrast to ADAFF, differences between BECCS and REF were small for both isoprene and monoterpenes, and for both MAGPIE and IMAGE. This may be due to the fact that in this study bioenergy crops are only simulated as grasses, while also woody bioenergy crops could be used in reality. Those areas that show up in Figures 4 and 5 tend to show small decreases for the BECCS scenarios. When comparing H11 and REF landcovers, differences were visible over larger areas compared to the BECCS-Ref case, especially for monoterpenes.

Figure 4: Isoprene emission differences (Δ) between the scenarios Adaff-Ref, Beccs-Ref and H11 - Ref of both MAGPIE and IMAGE. Mean values for the period 2081-2100 [$\text{g C m}^{-2} \text{ a}^{-1}$].

Figure 5: Monoterpene emission differences (Δ) between the scenarios Adaff-Ref, Beccs-Ref and H11 - Ref of both MAgPIE and IMAGE. Mean values for the period 2081-2100 [$\text{g C m}^{-2} \text{a}^{-1}$].

Overall, the larger the amount of agricultural area fraction on global scale, the lower are the emissions of BVOCs. The difference between managed area fractions of the scenarios and the differences in global emissions between the scenarios have a strong negative correlation as indicated by the Pearson coefficient (Table 3), with values between -0.92 and -0.98 for isoprene and -0.8 and -0.94 for monoterpene emissions. This correlation is stronger for isoprene than for monoterpene emissions. Regarding regional differences, the correlations are strong for the tropical region, with values $\sim$ -0.98 for both isoprene and monoterpenes, compared to values in the boreal region where correlations for both isoprene and monoterpenes are between -0.6 and -0.64 (not shown). The impact of LULCC on the emission of BVOCs seems bigger in the tropics than in other regions.

Table 3 Pearson Correlation Coefficient between the global extent of area under agriculture and the global sum of Isoprene and Monoterpene emissions.

	MAgPIE Ref	MAgPIE Beccs	MAgPIE Adaff	IMAGE Ref	IMAGE Beccs	IMAGE Adaff	H11
Isoprene	-0.97	-0.98	-0.92	-0.97	-0.97	-0.97	-0.98
Monoterpene	-0.91	-0.91	-0.80	-0.89	-0.91	-0.85	-0.94

4. Discussion

So far, the number of studies that focused on the emission change of BVOCs due to LULCC, with or without also considering climate change is relatively limited. Lathière et al. (2010) worked with historical land-cover change data based on maps by Loveland et al. (2000), Ramankutty and Foley (1999) and Goldewijk (2001), on which basis emission changes were calculated using the Sheffield Dynamic Global Vegetation Model (SDGVM). Combining the effect of CO₂ inhibition, climate dynamics and LULCC, they showed a decline in isoprene emissions by ~24% from 1901-2002 (Lathière et al. 2010), and similar to our study, they found the highest impact of cropland expansion on isoprene emission in tropical regions, such as the Amazonian region, central Africa and south-east Asia, but also to some extent in central Europe and parts of the eastern USA. Unger (2013) used a biochemical model of isoprene emission within a global chemistry-climate framework to quantify the impact of LULCC, increasing CO₂-concentration and climate change on the emission of isoprene. She found a global decrease of isoprene emissions between 1880 and 2000 of 20% (from 534 to 449 TgCa⁻²), which is of similar magnitude to the earlier studies. Later Unger (2014a, 2014b) shows the same effects for BVOCs in general. In a literature review of the impact of LULCC on atmospheric composition and climate, Heald and Spracklen (2015) estimate the effect of LULCC from the preindustrial until today as a decrease of 15-36% in global isoprene emissions. In earlier work, Lathiere et al. (2006) used the DGVM ORCHIDEE combined with the Guenther et al (1995) BVOC emissions model, and satellite-based present-day climate from the International Satellite Land-Surface Climatology Project, ISLSCP-II. They found under otherwise similar conditions a decline in global isoprene and monoterpene emissions each by ~-30% in response to a tropical deforestation scenario, and very small changes only (+4% and -0.1% for isoprene and monoterpene, respectively) in response to an European afforestation scenario (although in case of the latter, changes when only the European simulation domain was considered were massive). Our results of a ~21% decline during the 20th century fall well within the previously published range.

For future land-use change effects, Wu et al. (2012) and Squire et al. (2014) used the IPCC A1B emission scenario. For land-use change, both studies were based on the IMAGE IAM, specified by Squire et al. (2014) to assume an increase in cropland area by 135% between 2001 and 2095. Despite of the land-use change, Wu et al. (2012) found an increase of isoprene emission by 8% for the period from 2000-2100, while our results for scenarios considering cropland expansion show either little change, or decreasing emissions (+/- 1% for BECCS and -5% for H11). For their simulations Wu et al. (2012) used climate data from the

GISS GCM3, combined with vegetation dynamics from LPJ DGVM (Sitch et al., 2003), and using the MEGAN 2.0 framework for simulating the resulting BVOC emissions (Guenther et al., 2006). The A1B CO₂ emissions pathway is higher than the RCP2.6 scenario used here, however Wu et al. (2012) did not account for the CO₂ inhibition effect, which would act to strongly reduce isoprene emissions relative to considering climate change alone (Arneth et al., 2007).

By contrast, a strong decrease in isoprene emissions by -41% over the 21st century was found by Squire et al. (2014), who used the SDGVM to calculate the vegetation distribution, MEGAN to calculate isoprene emissions, and climate projections from HadGEM1. Their simulation setup combined SRES A1B LULCC and CO₂ emissions, and climate change as in the SRES B2 scenario. The large decline in isoprene emissions could be related to the fact that the authors accounted for CO₂ inhibition in leaf-emissions. The large cropland expansion in their simulations, together with the higher CO₂ concentrations in SRES B2 (comparable to a RCP4.5 pathway; van Vuuren et al. 2011) than used here are likely the chief reasons for the larger emission reduction compared to our study. Likewise, Ganzeveld et al. (2010), applying the EMAC chemistry/climate model with the Guenther et al. (1995) emission algorithms, simulated a 12% decline in isoprene until 2050 when a SRES A2 land cover change scenario was applied, without considering direct CO₂ inhibition or vegetation interacting dynamically with climate change.

In a recent study, Hantson et al. (2017) also used the LPJ-GUESS modelling framework but focusing on the effects of LULCC on the emission of BVOCs using land-use scenarios from Hurtt et al. (2011). Across a broad range of LULCC (using the RCP4.5 climate and CO₂ concentrations) they identified large effects of LULCC on the emission of BVOCs, especially for isoprene compared to a simulation with natural vegetation dynamics. Their results for RCP2.6 LULCC show relatively similar dynamics to those we found here for H11 but with the decline in isoprene emissions even bigger, likely due to the higher CO₂ concentration in RCP4.5. Heald and Spracklen (2015) estimate a decline in isoprene emissions for the 21st century due to the combined effects of increasing CO₂ concentration in the atmosphere and anthropogenic LULCC of 24-27%.

Clearly, the studies available to date on future land-cover change effects on BVOC emissions vary notably in their simulation set-up but unanimously show that BVOC emissions are closely linked not only to climate and CO₂, but also to LULCC. BVOC emissions in regions under cropland expansion decline, and all studies emphasize the crucial role tropical forests play as main emitters of BVOCs on the one hand and as highly threatened regions by agricultural expansion on the other.

Not surprisingly, effects of avoiding deforestation combined with afforestation had largest effects on BVOC trajectories, because of the high emission potentials of woody vegetation and because the difference in vegetation cover (compared to e.g., REF and BECCS) is quite pronounced in tropical regions. As part of the current political debate, in particular after the Paris COP21 agreement it is to be expected that land-based mitigation will become an even larger contributor to land-use change than so-far assumed. While the scenarios we used here are stylized and not calculated within a full Integrated Assessment framework they still serve to highlight the potential effect of avoided deforestation and reforestation vs. BECCS on non-greenhouse trace gas exchanges. In all scenarios we can demonstrate large effects on the amount of BVOCs emitted, related to the total area under management, with less managed area fraction on global scale relatively increasing BVOC-emissions globally and vice versa. But especially for the bioenergy crops it is important to consider which type of vegetation is used. In our scenario we assumed that bioenergy would be crop-based and used therefore a plant functional type with low BVOC emission potential (Guenther et al, 1995, Arneth et al., 2007, Schurgers et al., 2009, 2011). However, in tropical regions, for instance, oil palm is a widely used bioenergy plant, which has an emission potential as high, or even higher than tropical rainforests (Hewitt et al., 2009, Misztal et al., 2011, Silva et al., 2016). Likewise, Ashworth et al. (2012, 2013) simulated for Europe the emissions from poplar plantations used as bioenergy. Poplar is a strong isoprene emitter, and the authors found that the planting of ca. 70 Mha of cropland with bioenergy crops such as poplar (or eucalyptus) would result in an increase in isoprene emissions by roughly 4.5 Tg C a^{-1} . These studies demonstrate that the total area undergoing change towards bioenergy (or other land uses) is a large source of uncertainty for future BVOC emissions – and their effect on atmospheric chemistry - but also that the type of vegetation grown in a certain region will affect emissions substantially. For instance, if we compare our emission rates for C3 grasses, which we used to calculate emissions from bioenergy crops to emission rates of bioenergy crops like short rotation coppice (SRC) willow, the emissions in both BECCS scenarios would be significantly higher. A recent study by Morrison et al. (2016) found isoprene emission factors of willow growing at different locations in England between 0.1 and $16 \mu\text{g g}^{-1} \text{ h}^{-1}$, the upper value being identical to emission factors we used for C3 crops ($16 \mu\text{g g}^{-1} \text{ h}^{-1}$). However, willow emission factors well above $100 \mu\text{g g}^{-1} \text{ h}^{-1}$ were found in other studies (see Table 6, Morrison et al., 2016). Since the calculated emissions scale linearly with the applied emission factor (Niinemets et al., 2010) our calculated rates would, for instance, double if we had applied an emission factor of 32 rather than $16 \mu\text{g g}^{-1} \text{ h}^{-1}$. A more detailed analysis of how bioenergy

plantations affect BVOC emissions would require estimates of the areas of woody and non-woody plants used as bioenergy, and where these are located.

The emission hotspot for isoprene is located in the tropics (Guenther et al., 1995), and it is thus not surprising that also the emission difference between the different LULCC scenarios is largest there. It can be expected that different future trajectories would also impact on tropical atmospheric chemistry (Young et al., 2009, Makkonen et al., 2012, Barkley et al., 2012). However, the form of these impacts is not as clear as was once believed; work over the last decade has revealed significant gaps our understanding of isoprene oxidation and secondary organic aerosol formation (e.g. Lelieveld et al., 2008; Kiendler-Scharr et al., 2009; Fuchs et al., 2013), as well as background chemical conditions over remote areas (Liu et al., 2016). What can be expected with reasonable confidence is that increased BVOC emissions in or near areas with relatively high background levels of NO_x will lead to increased ozone formation. Such increases have been simulated previously for enhancements in isoprene emissions due to land-use change (Chen et al., 2009; Ashworth et al., 2012, 2013; Wu et al., 2012) and the chemistry is relatively well understood (Sillman, 1999). In environments with low NO_x concentrations, such as the remote tropical rainforest, increased BVOC emissions are expected to have the opposite effect, leading to decreased ozone concentrations (Hewitt et al., 2009). Uncertainties in the isoprene oxidation pathway under low- NO_x do not appear to have a large impact on projected changes in ozone concentration (Pugh et al., 2010), but recent evidence suggests that there may be higher levels of NO_x in some of these remote environments than previously thought (Liu et al., 2016), altering the effect of changed BVOC emissions on ozone. Further, in our simulations we did not yet simulate NO_x emissions from changing land-use, which we argue would be an important additional factor in assessing land-use change/air quality interactions, as NO_x emissions will be enhanced in fertilized soils. Combined BVOC/soil NO_x analyses thus will need to be done when assessing the potential ozone formation effects of LULCC. For instance, Ganzeveld et al. (2010), in their study with EMAC, investigated not only changes in isoprene but also altered soil NO_x emissions, resulting in an increase in boundary layer ozone up to 9 ppbv over deforested areas in Africa between present day and 2050 due to LULCC.

Methane lifetime has previously been strongly linked to BVOC emissions. As mentioned in the introduction, however, the uncertainty in the level to which the hydroxyl radical is regenerated in the isoprene oxidation chain (Lelieveld et al., 2008; Fuchs et al., 2013) precludes clear statements as to the influence of changed BVOC emissions on methane lifetime (e.g. Archibald et al., 2011). Further, the formation of secondary organic aerosol (SOA) that takes place from oxidation of monoterpenes (Spracklen et al., 2008, 2011, Carslaw

et al., 2010, 2013) and isoprene (Claeys et al., 2004, Carlton et al., 2009) will also be affected by LULCC, in particular if afforestation or avoided deforestation takes effect in conifer-dominated vegetation of boreal and temperate regions. But very large uncertainties remain regarding formation rates of SOA, for instance, isoprene has been found to inhibit new particle formation in some studies (Kiender-Scharr et al., 2009, 2012). Moreover, climate change has the potential to change the spectrum of different monoterpenes emitted towards chemical species with higher (or lower) atmospheric reactivity (Jardine et al., 2017), and the same can be speculated for land-cover change. Such a change in monoterpene speciation would affect e.g., SOA formation and growth but is not considered in our monoterpene emission algorithm. We therefore refrain from speculating on the implications of our results for methane lifetime and SOA formation, recommending that dedicated chemistry-transport studies are carried out to assess these important implications.

In summary, future pathways of LULCC highly affect the emission of BVOCs. As BVOCs play a crucial role in atmospheric composition, a broader research focus is necessary, compared to the few studies available to date. It seems evident that cropland expansion leads to decreasing emissions, while afforestation leads to increasing emissions. However, not only the impact of LULCC on the emissions itself is crucial, but also the choice of agricultural crop-types, as well as dynamic vegetation response of natural vegetation to changing environmental conditions. Future BVOC emissions and their impacts on chemistry-climate interactions can only be assessed conjointly with dynamic vegetation responses.

Acknowledgements

The authors acknowledge support from the EU FP7 project LUC4C (grant 603542). AA acknowledges also support from the Helmholtz Association ATMO programme and the Innovation and Networking fund. This is paper number 29 of the Birmingham Institute of Forest Research.

Literature:

Ainsworth, E. A., Yendrek, C. R., Sitch, S., Collins, W. J., Emberson, L. D.: The Effects of Tropospheric Ozone on Net Primary Productivity and Implications for Climate Change, *Annu. Rev. Plant Biol.*, 63, 637-661, doi: 10.1146/annurev-arplant-042110-103829, 2012.

Arneth, A., Miller, P. A., Scholze, M., Hickler, T., Schurgers, G., Smith, B., Prentice, I.C.: CO₂ inhibition of global terrestrial isoprene emissions: Potential implications for atmospheric chemistry, *Geophys. Res. Lett.*, 34, L18813, doi:10.11029/12007GL030615, 2007a.

Arneth, A., Niinemets, Ü., Pressley, S., Bäck, J., Hari, P., Karl, T., Noe, S., Prentice, I. C., Serca, D., Hickler, T., Wolf, A., Smith, B.: Process-based estimates of terrestrial ecosystem isoprene emissions:

- 1 incorporating the effects of a direct CO₂-isoprene interaction, *Atmos. Chem. Phys.*, 7, 31-53, doi:
2 10.5194/acp-7-31-2007, 2007b.
- 3 Arneth, A., Schurgers, G., Hickler, T., Miller, P. A.: Effects of species composition, land surface cover,
4 CO₂ concentration and climate on isoprene emissions from European forests, *Plant Biology*, 10, 150-
5 162, doi: 10.5194/acp-8-4605-2008, 2008.
- 6 Arneth, a., Monson, R.K., Schurgers, G., Niinemets, Ü., Palmer, P.I., 2008. Why are estimates of
7 global terrestrial isoprene emissions so similar (and why is this not so for monoterpenes)? *Atmos.*
8 *Chem. Phys.* 8, 4605–4620. doi:10.5194/acp-8-4605-2008
- 9 Arneth, A., Sitch, S., Bondeau, A., Butterbach-Bahl, K., Foster, P., Gedney, N., de Noblet-Ducoudrè, N.,
10 Prentice, I. C., Sanderson, M., Thonicke, K., Wania, R., Zaehle, S.: From biota to chemistry and
11 climate: towards a comprehensive description of trace gas exchange between the biosphere and
12 atmosphere, *Biogeosciences*, 7, 121-149, doi: 10.5194/bg-7-121-2010, 2010.
- 13 Arneth, A., Schurgers, G., Lathière, J., Duhl, T., Beerling, D. J., Hewitt, C. N., Martin, M., Guenther, A.:
14 Global terrestrial isoprene emission models: sensitivity to variability in climate and vegetation,
15 *Atmos. Chem. Phys.*, 11, 8037-8052, doi: 10.5194/acp-11-8037-2011, 2011.
- 16 Ashworth, K., Folberth, G., Hewitt, C. N., Wild, O.: Impacts of near-future cultivation of biofuel
17 feedstocks on atmospheric composition and local air quality, *Atmos. Chem. Phys.*, 12, 919-939, doi:
18 10.5194/acp-12-919-2012, 2012.
- 19 Ashworth, K., Wild, O., Hewitt, C. N.: Impacts of biofuel cultivation on mortality and crop yields,
20 *Nature Climate Change*, 3, S. 492-496, doi: 10.1038/NCLIMATE1788, 2013.
- 21 Atkinson, R.: Atmospheric chemistry of VOCs and NO_x, *Atmos. Environ.*, 34, 2063-2101, doi:
22 10.1016/S1352-2310(99)00460-4, 2000.
- 23 Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a
24 review, *Atmos. Environ.*, 37, 197-219, doi: 10.1016/S1352-2310(03)00391-1, 2003.
- 25 Barkley, M. P., Kurosu, T. P., Chance, K., De Smedt, I., Van Roozendaal, M., Arneth, A., Hagberg, D.,
26 Guenther, A.: Assessing sources of uncertainty in formaldehyde air mass factors over tropical South
27 America: Implications for top-down isoprene emission estimates, *Jour. Geophys. Res.*, 117, D13304,
28 doi:10.1029/2011JD016827, 2012.
- 29 Bondeau, A., Smith, P. C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-Campen,
30 H., Müller, C., Reichstein, M., Smith, B.: Modelling the role of agriculture for the 20th century global
31 terrestrial carbon balance. *Global Change Biology*, 13, 679-706, 2007.
- 32 Bouwman, A. F., Kram, T., Klein Goldewijk, C. G. M.: Integrated modelling of global environmental
33 change. An overview of IMAGE 2.4. PBL, 2006.
- 34 Carlton, A. G., Wiedinmyer, C., Kroll, J. H.: A review of secondary organic aerosol (SOA) formation
35 from isoprene, *Atmos. Chem. Phys.*, 9, 4987-5005, doi: 10.5194/acp-9-4987-2009, 2009.
- 36 Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S., Kulmala, M.: A
37 review of natural aerosol interactions and feedbacks within the Earth system, *Atmos. Chem. Phys.*,
38 10, 1701-1737, doi: 10.5194/acp-10-1701-2010, 2010.

- 1 Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W.,
2 Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., Pierce, J. R.: Large contribution of natural aerosols
3 to uncertainty in indirect forcing, *Nature*, 503, 67-71, doi: 10.1038/nature12674, 2013.
- 4 Chen, J., Avise, J., Guenther, A., Wiedinmyer, C., Salathe, E., Jackson, R. B., Lamb, B.: Future land use
5 and land cover influences on regional biogenic emissions and air quality in the United States, *Atmos.*
6 *Environ.*, 43, 5771-5780, 2009.
- 7 Chini, L.P., G.C. Hurtt, and S. Frolking. 2014. Harmonized Global Land Use for Years 1500 – 2100, V1.
8 Data set. Available on-line [<http://daac.ornl.gov>] from Oak Ridge National Laboratory Distributed
9 Active Archive Center, Oak Ridge, Tennessee, USA. <http://dx.doi.org/10.3334/ORNLDAAAC/1248>
- 10 Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P.,
11 Andreae, M. O., Artaxo, P., Willy, M.: Formation of secondary organic aerosols through
12 photooxidation of isoprene, *Science*, 303, 1173-1176, doi: 10.1126/science.1092805, 2004.
- 13 Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S.,
14 Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F.,
15 Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ehté, C., Fairhead, L., Fichefet, T., Flavoni,
16 S., Friedlingstein P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A.,
17 Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P.,
18 Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S.,
19 Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D.,
20 Szopa, S., Talandier, C., Terray, P., Viovy, N., Vuichard, N.: Climate change projections using the IPSL-
21 CM5A Earth System Model: from CMIP3 to CMIP5, *Clim. Dyn.*, 40, 2123-2165, doi: 10.1007/s00382-
22 012-1636-1, 2013.
- 23 Eitelberg, D. A., van Vliet, J., Verburg, P. H.: A review of global potentially available cropland
24 estimates and their consequences for model-based assessments, *Global Change Biol.*, 21, 1236-1248,
25 doi: 10.1111/gcb.12733, 2015.
- 26 Ellis, E.C., Goldewijk, K.K., Siebert, S., Lightman, D., Ramankutty, N.: Anthropogenic transformation of
27 the biomes, 1700 to 2000, *Global Ecol. Biogeogr.*, 19, 589-606, doi: 10.1111/j.1466-
28 8238.2010.00540.x, 2010.
- 29 Fuchs, H., Hofzumahaus, A., Rohrer, F., Bohn, B., Brauers, T., Dorn, H., Häsel, R., Holland, F.,
30 Kaminski, M., Li, X., Lu, K., Nehr, S., Tillmann, R., Wegener, R., Wahner, A.: Experimental evidence for
31 efficient hydroxyl radical regeneration in isoprene oxidation. *Nat. Geosci.* 6, 1023–1026.
32 doi:10.1038/ngeo1964, 2013.
- 33 Ganzeveld, L., Bouwman, L., Stehfest, E., van Vuuren, D. P., Eickhout, B., Lelieveld, J.: Impact of future
34 land use and land cover changes on atmospheric chemistry-climate interactions, *J. Geophys. Res.*,
35 115, D23301, doi:10.1029/2010JD014041, 2010.
- 36 Goldewijk, K.K.: Estimating global land use change over the past 300 years: THE HYDE Database,
37 *Global Biogeochemical Cycles*, 15, 417-433, doi: 10.1029/1999GB001232, 2001.
- 38 Goldewijk, K. K., Beusen, A., van Drech, G., de Vos, M.: The HYDE 3.1 spatially explicit database of
39 human-induced global land-use change over the past 12.000 years, *Global Ecol. Biogeogr.*, 20, 73-86,
40 doi: 10.1111/j.1466-8238.2010.00587.x, 2011.

- 1 Guenther, A., Hewitt, C.N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau,
2 M., McKay, W.A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., Zimmerman, P.: A
3 global model of natural volatile organic compound emissions, *J. Geophys. Res.*, 100, 8873-8892, doi:
4 10.1029/94JD02950, 1995.
- 5 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., Geron, C.: Estimates of global
6 terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature),
7 *Atmos. Chem. Phys.*, 6, 3181-3210, doi: 10.5194/acp-6-3181-2006, 2006.
- 8 Guenther, A., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., Wang, X.: The
9 Model of Emission of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and
10 updated framework for modeling biogenic emissions, *Geosc. Model Dev.*, 5, 1471-1492, doi:
11 10.5194/gmd-5-1471-2012, 2012.
- 12 Hantson, S., Knorr, W., Schurgers, G., Pugh, T.A.M., Arneth, A.: Global isoprene and monoterpene
13 emissions under changing climate, vegetation, CO₂ and land use, *Atmos. Environ.*, 155, 35-45, doi:
14 10.1016/j.atmosenv.2017.02.010, 2017.
- 15 Harrison, S. P., Morfopoulos, C., Srikanta Dani, K. G., Prentice, I. C., Arneth, A., Atwell, B. J., Barkley,
16 M. P., Leishman, M. R., Loreto, F., Medlyn, B. E., Niinemets, Ü., Possell, M., Penuelas, J., Wright, I. J.:
17 Volatile isoprenoid emissions from plastid to planet, *New Phytologist*, 197, 49-57, doi:
18 10.1111/nph.12021, 2013.
- 19 Hewitt, C.N., MacKenzie, A.R., Di Carlo, P., Di Marco, C.F., Dorsey, J.R., Evans, M., Fowler, D.,
20 Gallagher, M.W., Hopkins, J.R., Jones, C.E., Langford, B., Lee, J.D., Lewis, a C., Lim, S.F., McQuaid, J.,
21 Misztal, P., Moller, S.J., Monks, P.S., Nemitz, E., Oram, D.E., Owen, S.M., Phillips, G.J., Pugh, T. a M.,
22 Pyle, J.A., Reeves, C.E., Ryder, J., Siong, J., Skiba, U., Stewart, D.J.: Nitrogen management is essential
23 to prevent tropical oil palm plantations from causing ground-level ozone pollution. *Proc. Natl. Acad.*
24 *Sci. U. S. A.* 106, 18447–51. doi:10.1073/pnas.0907541106, 2009.
- 25 Heald, C.L., Spracklen, D.V.: Land Use Change Impacts on Air Quality and Climate, *Chem. Rev.*, 115,
26 4476-4496, doi: 10.1021/cr500446g, 2015.
- 27 Humpenoder, F., Popp, A., Stevanovic, M., Müller, C., Bodirsky, B. L., Bonsch, M., Dietrich, J. P., Lotze-
28 Campen, H., Weindl, I., Biewald, A., Rolinski, S.: Land-Use and Carbon Cycle Responses to Moderate
29 Climate Change: Implications for Land-Based Mitigation?, *Environmental Science & Technology*, 49,
30 6731-6739, doi: 10.1021/es506201r, 2015..
- 31 Hurtt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K.,
32 Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Goldewijk, K. K., Riahi, K.,
33 Shevlikova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., Wang, Y. P.:
34 Harmonization of land-use scenarios for the period 1500-2100: 600 years of global gridded annual
35 land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117-161,
36 doi: 10.1007/s10584-011-0153-2, 2011.
- 37 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth
38 Assessment Report of the Intergovernmental Panel on Climate Change, 2013.
- 39 Jardine, K. J., Jardine, A. B., Holm, J. A., Lombardozzi, D. L., Negron-Juarez, R. I., Martin, S. T., Beller,
40 H. R., Gimenez, B. O., Higuchi, N., Chambers, J. Q.: Monoterpene ‘thermometer’ of tropical forest-

- 1 atmosphere response to climate warming, *Plant, Cell and Environ.*, 40, 441-452, doi:
2 10.1111/pce.12879, 2017.
- 3 Jerrett, M., Burnett, R. T., Arden Pope III, C., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle, E., Thun,
4 M.: Long-Term Ozone Exposure and Mortality, *N Engl J Med*, 360, 1085-1095, doi:
5 10.1056/NEJMoa0803894, 2009.
- 6 Jung, M., Reichstein, M., Margolis, H. A., Cescatti, A., Richardson, A. D., Atlatf Arain, M., Arneth, A.,
7 Bernhofer C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E.,
8 Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F.,
9 Williams, C.: Global patterns of landatmosphere fluxes of carbon dioxide, latent heat, and sensible
10 heat derived from eddy covariance, satellite, and meteorological observations, *Jour. Geophys. Res.*,
11 116, G00J07, doi:10.1029/2010JG001566, 2011.
- 12 Kanakidou, M., Myriokefalitakis, S., Daskalakis, N., Fanourgakis, G.: Past, present and future
13 atmospheric nitrogen deposition, *Amer. Meteor. Soc.*, doi:10.1175/JAS-D-15-0278.1, 2016.
- 14 Kiendler-Scharr, A., Wildt, J., Dal Maso, M., Hohaus, T., Kleist, E., Mentel, T.F., Tillmann, R., Uerlings,
15 R., Schurr, U., Wahner, A.: New particle formation in forests inhibited by isoprene emissions. *Nature*
16 461, 381–4. doi:10.1038/nature08292, 2009.
- 17 Kulmala, M., Petäjä, T., Mönkkönen, P., Koponen, I. K., Dal Maso, M., Aalto, P. P., Lehtinen, K. E. J.,
18 Kerminen, V.-M.: On the growth of nucleation mode particles: source rates of condensable vapor in
19 polluted and clean environments, *Atmos. Chem. Phys.*, 5, 409-416, doi: 10.5194/acp-5-409-2005,
20 2005.
- 21 Kulmala, M., Nieminen, T., Nikandrova, A., Lehtipalo, K., Manninen, H. E., Kajos, M. K., Kolari, P.,
22 Lauri, A., Petäjä, T., Krejci, R., Hansson, H.-C., Swietlicki, E., Lindroth, A., Christensen, T. R., Arneth, A.,
23 Hari, P., Bäck, J., Vesala, T., Kerminen, V.-M.: CO₂-induced terrestrial climate feedback mechanism:
24 From carbon sink to aerosol source and back, *Bor. Environ. Res.*, 19, 122-131, ISSN 1797-2469, 2014.
- 25 Lambin E. F., Meyfroidt P.: Global land use change, economic globalization and the looming land
26 scarcity, *PNAS*, 108, 3465-3472, doi: 10.1073/pnas.1100480108, 2011.
- 27 Lathière, J., Hauglustaine, D. A., De Noblet-Ducoudrè, N.: Past and future changes in biogenic volatile
28 organic compound emissions simulated with a global dynamic vegetation model, *Geophys. Res. Lett.*,
29 32, L20818, doi:10.1029/2005GL024164, 2005.
- 30 Lathière J., Hauglustaine, D. A., Friend, A. D., De Noblet-Ducoudrè, N., Viovy, N., Folberth, G. A.:
31 Impact of climate variability and land use changes on global biogenic volatile organic compound
32 emissions, *Atmos. Chem. Phys.*, 6, 2129-2146, doi: 10.5194/acp-6-2129-2006, 2006.
- 33 Lathière, J., Hewitt, C. N., Beerling, D. J.: Sensivity of isoprene emissions from the terrestrial
34 biosphere to 20th century changes in atmospheric CO₂ concentration, climate, and land use, *Global*
35 *Biogeochem. Cycles*, 24, GB1004, doi:10.1029/2009GB003548, 2010.
- 36 Lelieveld, J., Butler, T.M., Crowley, J.N., Dillon, T.J., Fischer, H., Ganzeveld, L., Harder, H., Lawrence,
37 M.G., Martinez, M., Taraborrelli, D., Williams, J.: Atmospheric oxidation capacity sustained by a
38 tropical forest. *Nature* 452, 737–40. doi:10.1038/nature06870, 2008.

- 1 Le Quéré C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P.,
2 Peters, G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J. I., Keeling, R. F., Tans, P., Arneth,
3 A., Bakker, D. C. E., Barbero, L., Bopp, L., Chang, J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M.,
4 Feely, R. A., Gkritzalis, T., Harris, I., Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Goldewijk, K. K.,
5 Koven, C., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N., Millero, F.,
6 Munro, D. R., Murata, A., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T.,
7 Pérez, F. F., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Rödenbeck, C., Saito, S., Schuster, U.,
8 Schwinger, J., Séférian, R., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van
9 der Laan-Luijkx, I. T., van der Werf, G. R., van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A.,
10 Zaehle, S., Zeng, N.: Global carbon budget 2015, *Earth Syst. Sci. Data*, 7, 349-396, doi: 10.5194/essd-
11 7-349-2015, 2015.
- 12 Levis, S.: Modeling vegetation and land use in models of the Earth System, *WIREs Clim Change*, 1,
13 840-856, doi: 10.1002/wcc.83, 2010.
- 14 Lindquist, E.J., D'Annunzio, R., Gerrand, A., MacDicken, K., Achard, F., Beuchle, R., Brink, A., Eva, H.D.,
15 Mayaux, P., San-Miguel-Ayanz, J., Stibig, H-J.: Global forest land-use change 1990:2005, *FAO Forestry*
16 *Paper*, 169, 2012.
- 17 Lotze-Campen, H., Müller, C., Bondeau, A., Rost, S., Popp, A., Lucht, W.: Global food demand,
18 productivity growth, and the scarcity of land and water resources: a spatially explicit mathematical
19 programming approach, *Agric. Econ.*, 39, 325-338, doi: 10.1111/j.1574-0862.2008.00336.x, 2008.
- 20 Loveland, T.R., Reed, B.C., Brown, J.F., Oohlen, D.O., Zhu, Z., Yang, L., Merchant, J.W.: Development
21 of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data, *Int. Jour.*
22 *Of Remote Sensing*, 21, 1303-1330, doi: 10.1080/014311600210191, 2000.
- 23 Makkonen, R., Asmi, A., Kerminen, V.-M., Boy, M., Arneth, A., Guenther, A., Kulmala, M.: BVOC-
24 aerosol-climate interactions in the global aerosol-climate model ECHAM5.5-HAM2, *Atmos. Chem.*
25 *Phys.*, 12, 10077-10096, doi: 10.5194/acp-12-10077-2012, 2012.
- 26 Miralles, D. G., Holmes, T. R. H., De Jeu, R. A. M., Gash, J. H., Meesters, A. G. C. A., Dolman, A. J.:
27 Global land-surface evaporation estimated from satellite-based observations, *Hydrol. Earth Syst. Sci.*,
28 15, 453-469, doi: 10.5194/hess-15-453-2011, 2011.
- 29 Misztal, P. K., Nemitz, E., Langford, B., Di Marco, C. F., Phillips, G. J., Hewitt, C. N., MacKenzie, A. R.,
30 Owen, S. M., Fowler, D., Heal, M. R., Cape, J. N.: Direct ecosystem fluxes of volatile organic
31 compounds from oil palms in South-East Asia, *Atmos. Chem. Phys.*, 11, 8995-9017, doi: 10.5194/acp-
32 11-8995-2011, 2011.
- 33 Monson, R. K., Trahan, N., Rosenstiel, T. N., Veres, P., Moore, D., Wilkinson, M., Norby, R. J., Volder,
34 A., Tjoelker, M. G., Briske, D. D., Karnosky, D. F., Fall, R.: Isoprene emission from terrestrial
35 ecosystems in response to global change: minding the gap between models and observations, *Phil.*
36 *Trans. R. Soc.*, 365, 1677-1695, doi: 10.1098/rsta.2007.2038, 2007.
- 37 Morrison, E. C., Drewer, J., Heal, M. R.: A comparison of isoprene and monoterpene emission rates
38 from the perennial bioenergy crops short rotation coppice willow and *Miscanthus* and the annual
39 arable crops wheat and oilseed rape, *GCB Bioenergy*, 8, 211-225, doi: 10.1111/gcb.12257, 2016.

- 1 Niinemets, Ü., Tenhunen, J. D., Harley, P. C., Steinbrecher, R.: A model of isoprene emission based on
2 energetic requirements for isoprene synthesis and leaf photosynthetic properties for Liquidambar
3 and Quercus, *Plant, Cell and Envir.*, 22, 1319-1335, doi: 10.1046/j.1365-3040.1999.00505.x, 1999.
- 4 Niinemets, Ü., Monson, R. K., Arneth, A., Ciccioli, P., Kesselmeier, J., Kuhn, U., Noe, S. M., Penuelas, J.,
5 Staudt, M.: The leaf-level emission factor of volatile isoprenoids: caveats, model algorithms, response
6 shapes and scaling, *Biogeosciences*, 7, 1809-1832, doi: 10.5194/bg-7-1809-2010, 2010.
- 7 Paasonen, P., Asmi, A., Petäjä, T., Kajos, M.K., Äijälä, M., Junninen, H., Holst, T., Abbatt, J.P.D.,
8 Arneth, A., Birmili, W., van der Gon, H.D., Hamed, A., Hoffer, A., Laakso, L., Laaksonen, A., Richard
9 Leaitch, W., Plass-Dülmer, C., Pryor, S.C., Räisänen, P., Swietlicki, E., Wiedensohler, A., Worsnop, D.R.,
10 Kerminen, V.-M., Kulmala, M.: Warming-induced increase in aerosol number concentration likely to
11 moderate climate change. *Nat. Geosci.* 6, 438–442. doi:10.1038/ngeo1800, 2013.
- 12 Palmer, P. I., Abbot, D. S., Fu, T.-M., Jacob, D. J., Chance, K., Kurosu, T., Guenther, A., Wiedinmyer, C.,
13 Stanton, J. C., Pilling, M. J., Pressley, S. N., Lamb, B., Sumner, A. L.: Quantifying the seasonal and
14 interannual variability of North American isoprene emissions using satellite observations of the
15 formaldehyde column, *Jour. Geophys. Res.*, 111, D12315, doi:10.1029/2005JD006689, 2006.
- 16 Poisson, N., Kanakidou, M., Crutzen, P. J.: Impact of non-methane hydrocarbons on tropospheric
17 chemistry and the oxidizing power of the global troposphere: 3-dimensional modelling results, *J.*
18 *Atmos. Chem.*, 36, 157-230, doi: 10.1023/A:1006300616544, 2000.
- 19 Popp, A., Dietrich, J. P., Lotze-Campen, H., Klein, D., Bauer, N., Krause, M., Beringer, T., Gerten, D.,
20 Edenhofer, O.: The economic potential of bioenergy for climate change mitigation with special
21 attention given to implications for the land system, *Environ. Res. Lett.*, 6, 034017, doi:10.1088/1748-
22 9326/6/3/034017, 2011.
- 23 Popp, A., Rose, S. K., Calvin, K., Van Vuuren, D. P., Dietrich, J. P., Wise, M., Stehfest, E., Humpenöder,
24 F., Kyle, P., Van Vliet, J., Bauer, N., Lotze-Campen, H., Klein, D., Kriegler, E.: Land-use transition for
25 bioenergy and climate stabilization: model comparison of drivers, impacts and interactions with
26 other land use based mitigation options, *Climatic Change*, 123, 459-509, doi: 10.1007/s10584-013-
27 0926-x, 2014a.
- 28 Popp, A., Humpenöder, F., Weindl, I., Bodirsky, B. L., Bonsch, M., Lotze-Campen, H., Müller, C.,
29 Biewald, A., Rolinski, S., Stevanovic, M., Dietrich, J. P.: Land-use protection for climate change
30 mitigation, *Nat. Clim. Change*, doi: 10.1038/NCLIMATE2444, 2014b.
- 31 Pugh, T. A. M., MacKenzie, A. R., Hewitt, C. N., Langford, B., Edwards, P. M., Furneaux, K. L., Heard, D.
32 E., Hopkins, J. R., Jones, C. E., Karunaharan, A., Lee, J., Mills, G., Misztal, P., Moller, S., Monks, P. S.,
33 Whalley, L. K.: Simulation atmospheric composition over a South-East Asian tropical rainforest:
34 performance of a chemistry box model, *Atmos. Chem. Phys.*, 10, 279-298, doi: 10.5194/acp-10-279-
35 2010, 2010.
- 36 Ramankutty, N., Foley, J.A.: Estimating historical changes in global land cover: Croplands from 1700
37 to 1992, *Global Biogeochemical Cycles*, 13, 997-1027, doi: 10.1029/1999GB900046, 1999.
- 38 Reay, D. S., Dentener, F., Smith, P., Grace, J., Feely, R. A.: Global nitrogen deposition and carbon
39 sinks, *Nature Geoscience*, 1, 430-437, doi: 10.1038/ngeo230, 2008.
- 40 Reichstein, M., Papale, D., Valentini, R., Aubinet, M., Bernhofer, C., Knohl, A., Laurila, T., Lindroth, A.,
41 Moors, E., Pilegaard, K., Seufert, G.: Determinants of terrestrial ecosystem carbon balance inferred

- 1 from European eddy covariance flux sites, *Geophys. Res. Lett.*, 34, L01402,
2 doi:10.1029/2006GL027880, 2007.
- 3 Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N., Rafaj, P.:
4 RCP8.5 – A scenario of comparatively high greenhouse gas emissions, *Climatic Change*, 109, 33-57,
5 doi: 10.1007/s10584-011-0149-y, 2011.
- 6 Rosenkranz, M., Pugh, T. A. M., Schnitzler, J.-P., Arneth, A.: Effect of land-use change and
7 management on biogenic volatile organic compound emissions – selecting climate-smart cultivars,
8 *Plant, Cell and Environ.*, 38, 1896-1912, doi: 10.1111/pce.12453, 2015.
- 9 Schurgers, G., Arneth, A., Holzinger, R., Goldstein, A. H.: Process-based modelling of biogenic
10 monoterpene emissions combining production and release from storage, *Atmos. Chem. Phys.*, 9,
11 3409-3423, doi: 10.5194/acp-9-3409-2009, 2009.
- 12 Schurgers, G., Arneth, A., Hickler, T.: Effect of climate driven changes in species composition on
13 regional emission capacities of biogenic compounds, *Jour. Geophys. Res.*, 116, D22304,
14 doi:10.1029/2011JD016278, 2011.
- 15 Sillman, S.: The relation between ozone, NO_x and hydrocarbons in urban and polluted rural
16 environments, *Atmosph. Environ.*, 33, 1821-1845, 1999.
- 17 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht,
18 W., Sykes, M. T., Thonicke, K., Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and
19 terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Glob. Change Biol.*, 9, 161-185,
20 doi: 10.1046/j.1365-2486.2003.00569.x, 2003.
- 21 Silva, S. J., Heald, C. L., Geddes, J. A., Austin, K. G., Kasibhatla, P. S., Marlier, M. E.: Impacts of current
22 and projected oil palm plantation expansion on air quality over southeast asia, *Atmos. Chem. Phys.*
23 *Discuss.*, doi:10.5194/acp-2016-158, 2016.
- 24 Smith, B., Prentice, I. C., Sykes, M. T.: Representation of vegetation dynamics in the modelling of
25 terrestrial ecosystems: comparing two contrasting approaches within European climate space, *Glob.*
26 *Ecol. Biogeogr.*, 10, 621-637, doi: 10.1046/.1466-822X.2001.t01-1-00256.x, 2001.
- 27 Smith, B., Warlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., Zaehle, S.: Implications of
28 incorporating N cycling and N limitations on primary production in an individual-based dynamic
29 vegetation model, *Biogeosciences*, 11, 2027-2054, doi: 10.5194/bg-11-2027-2014, 2014.
- 30 Smith, P., Haberl, H., Popp, A., Erb, K.-H., Lauk, C., Harper, R., Tubiello, F. N., De Siqueira Pinto, A.,
31 Jafari, M., Sohi, S., Masera, O., Böttcher, H., Berndes, G., Bustamante, M., Ahammad, H., Clark, H.,
32 Dong, H., Elsiddig, E. A., Mbow, C., Ravindranath, N. H., Rice, C. W., Abad, C. R., Romanovskaya, A.,
33 Sperling, F., Herrero, M., House, J. I., Rose, S.: How much land-based greenhouse gas mitigation can
34 be achieved without compromising food security and environmental goals?, *Global Change Biology*,
35 19, 2285-2302, doi: 10.1111/gcb.12160, 2013.
- 36 Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsiddig, E. A., Haberl, H., Harper, R.,
37 House, J., Jafari, M., Masera, O., Mbow, C., Ravindranath, N. H., Rice, C. W., Robledo Abad, C.,
38 Romanovskaya, A., Sperling, F., Tubiello, F.: Agriculture, Forestry and Other Land Use (AFOLU),
39 *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth*

- Assessment Report of the Intergovernmental Panel on Climate Change [Edenhofer, O., Pichs-Madruga, J., Sokona, Y., Farahani, E., Kadner, S., Seyboth, K., Adler, A., Baum, I., Brunner, S., Eickemeier, P., Kriemann, B., Savolainen, J., Schlömer, S., von Stechow, C., Zwickel, T., Minx, J. C. (eds.)] Cambridge University Press, Cambridge, United Kingdom and New York, NY, 2014.
- Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Sihto, S.-L., Riipinen, I., Merikanto, J., Mann, G. W., Chipperfield, M. P., Wiedensohler, A., Birmili, W., Lihavainen, H.: Contribution of particle formation to global cloud condensation nuclei concentrations, *Geophys. Res. Lett.*, 35, L06808, doi: 10.1029/2007GL033036, 2008.
- Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W., Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G. Rap, A., Forster, P.: Aerosol mass spectrometer constraint on the global secondary organic aerosol budget, *Atmos. Chem. Phys.*, 11, 12109-12136, doi: 10.5194/acp-11-12109-2011, 2011.
- Squire, O. J., Archibald, A. T., Abraham, N. L., Beerling, D. J., Hewitt, C. N., Lathière, J., Pike, R. C., Telford, P. J., Pyle, J. A.: Influence of future climate and cropland expansion on isoprene emissions and tropospheric ozone, *Atmos. Chem. Phys.*, 14, 1011-1024, doi: 10.5194/acp-14-1011-2014, 2014.
- Stehfest, E., van Vuuren, D., Kram, T., Bouwman, L.: Integrated Assessment of Global Environmental Change with IMAGE 3.0 – Model description and policy applications, PBL, 2014.
- Tubiello, F., Salvatore, M., Ferrara, A. F., House, J., Federici, S., Rossi, S., Biancalani, R., Condor Golec, R. D., Jacobs, H., Flammini, A., Prosperi, P., Cardenas-Galindo, P., Schmidhuber, J., Sanz Sanchez, M. J., Srivastava, N., Smith, P.: The Contribution of Agriculture, Forestry and other Land Use activities to Global Warming, 1990-2012, *Global Change Biology*, 21, 2655-2660, doi: 10.1111/gcb.12865, 2015.
- Unger, N.: Isoprene emission variability through the twentieth century, *Journ. Geophys. Res.*, 118, 606-613, doi: 10.1002/2013JD020978, 2013.
- Unger, N.: Human land-use-driven reduction of forest volatiles cools global climate, *Nat. Clim. Chang.*, 4, doi: 10.1038/NCLIMATE2347, 2014a.
- Unger N.: On the role of plant volatiles in anthropogenic global climate change, *Geophys. Res. Lett.*, 41, 8563-8569, doi: 10.1002/2014GL061616, 2014b.
- Van Vuuren, D. P., Stehfest, E., den Elzen, M. G. J., Kram, T., van Vliet, J., Deetman, S., Isaac, M., Goldewijk, K. K., Hof, A., Beltran, A. M., Oostenrijk, R., van Rujiven, B.: RCP2.6: Exploring the possibility to keep global mean temperature increase below 2°C, *Climatic Change*, 109, 95-116, doi: 10.1007/s10584-011-0152-3, 2011.
- Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., Schewe, J.: The Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP): Project framework, *PNAS*, 111, 3228-3232, doi: 10.1073/pnas.1312330110, 2014.
- Wu, S., Mickley, L. J., Kaplan, J. O., Jacob, D. J.: Impacts of changes in land use and land cover on atmospheric chemistry and air quality over the 21th century, *Atmos. Chem. Phys.*, 12, 1597-1609, doi: 10.5194/acp-12-1597-2012, 2012.

1 Appendix:

2 A: Supplementary Figures:

3
4 *Figure 6: Maps showing present day distribution of the dominant PFTs per gridcell (gridcells show*
5 *PFTs with maximum LAI). Mean values for the period 1981-2000 for all seven different model*
6 *projections. Abbreviations are as in Figure 1.*

Figure 7: Managed Area Fraction [%] - mean values for the period 2081-2100. Differences between the scenarios ADAFF-REF, BECCS-REF and H11-REF for both IAMs. Example equation for the calculation: Gridcell x ($x:n$) in MAGPIE ADAFF = 50% (0.5) area managed, Gridcell x ($x:n$) in MAGPIE REF = 30% (0.3) area managed. $0.5 - 0.3 = 0.2$ (20%) = higher extent of area managed in ADAFF than in REF for Gridcell x .

B: Method-Description – Land-use Datasets:

IMAGE3.0:

IMAGE is an integrated assessment model framework that simulates global and regional environmental consequences of changes in human activities (Stehfest *et al* 2014). The model is a simulation model that uses a set of linked sub-models describing the energy system, the agricultural economy, land use, natural vegetation and the climate system. Most of the socio-economic parameters are modelled for 26 regions and most of the environmental parameters are modelled on a geographic grid.

The land-use component of the IMAGE framework is driven by demand for agricultural commodities, most importantly food crops, animal products and bio-energy. Food crops and animal product demand are modelled on a regional basis by the agro-economic model MAGNET, which takes into

account scenario-specific land availability. MAGNET determines the level of agricultural intensification due to pressures on land availability, for example from policies for biodiversity protection or REDD (Overmars *et al* 2014). The demand for bio-energy is modelled by the energy model TIMER. The land availability for bio-energy production is determined according to a set of sustainability criteria (Hoogwijk 2004): only land that has been used for agriculture in the past but is not required for food production anymore, or areas with low carbon density are available. The IMAGE-core model implements the demands for the various agricultural products (7 irrigated crop groups, 7 rain-fed crop groups, grassland and sugarcane, grass and wood for bio-energy), on a geographic grid of 5' by 5' arc-minutes according to a set of empirically based allocation rules (Stehfest *et al* 2014). Production of animal products is calculated using a detailed description of the livestock sector (Bouwman *et al* 2005). The expansion of built-up area is a function of scenario-specific population growth and urbanization. Forestry activities are also modelled on the grid, including forest degradation and deforestation that cannot be explained by land use expansion for agriculture. In addition, the IMAGE-core model internally calculates climate change using an adapted version of the MAGICC 6.0 model, and a pattern scaling to derive grid-specific temperature and precipitation. The gridded land use is aggregated to 30' by 30' arc-minute fractions for use in the coupled dynamic vegetation model LPJml (Bondeau *et al* 2007). Using land use and climate data, LPJml models the carbon and water cycles, crop yields and natural vegetation dynamics.

In strict climate mitigation scenarios aiming for a maximum warming of two degrees by the year 2100, bio-energy production in combination with carbon capture and storage (BIO CCS) is an important mitigation measure. IMAGE considers three types of bio-energy: ethanol from sugarcane and methanol from lignocellulosic feedstock (grass or wood). The energy model TIMER uses land availability, cost-based competition with other energy carriers and a carbon price to determine final demand for bio-energy. While bioenergy can also be based on residues (Daiglou *et al* 2015), here the bioenergy demand can only be fulfilled by the three explicit feedstocks to achieve optimal comparability between MAgPIE and IMAGE.

Carbon dioxide removal (CDR) from avoided deforestation and afforestation (ADAFF) is not dynamically modelled in IMAGE 3.0. For this study, a stylized implementation is used to determine the mitigation potential of ADAFF under different agricultural intensification pathways. A stepwise approach systematically increases the potential in a set of scenario runs to identify the combination of intensification measures that achieves a specific CDR target. First, degraded forest areas are afforested. Secondly, the efficiencies of the livestock sector are gradually increased until maximum productivity levels are reached. Thirdly, the yield gap in crop production is closed by gradually increasing the intensity towards the maximum potential. Finally, the intensity of livestock pasture use is gradually increased towards the maximum potential. The intensification pathways result in avoided deforestation and abandonment of agricultural land that can be used for afforestation, which is assumed to take place as natural regrowth of vegetation. Areas with high afforestation potential are abandoned first to maximize cumulative CDR. Only biomes with considerable forest cover are accounted as part of the afforestation flux (boreal forests, temperate forests, tropical forests, savannahs).

MAgPIE:

MAgPIE is a global multi regional partial equilibrium model of the agricultural sector (Lotze-Campen *et al* 2008, Popp *et al* 2010, 2014). Objective function of the model is the minimization of global costs for agricultural production throughout the 21st century (5-year time steps) in recursive dynamic mode (**Error! Reference source not found.**). The model is driven by demand for agricultural commodities, which is calculated based on population and income projections for the 21st century

consistent with SSP2 (O'Neill *et al* 2015). The production of agricultural commodities is associated with costs for labor, capital, fertilizer, technological change, transport, and land conversion. Demand and costs enter the model at the level of 10 world regions. For meeting the demand, the model endogenously decides, based on cost-effectiveness, about the level of intensification (yield-increasing technological change), extensification (land-use change), and production relocation (international trade) (Dietrich *et al* 2014, Schmitz *et al* 2012). The optimization process is subject to various spatially explicit biophysical conditions, which are derived from the global crop growth, vegetation, and hydrology model LPJmL (Müller and Robertson 2014, Bondeau *et al* 2007). For instance, climate-induced changes in crop yields can alter cropland requirements, and hence deforestation. Due to computational constraints, spatially explicit input (0.5 degree resolution) is aggregated to 600 simulation units for the optimization process based on a k-means clustering algorithm (Dietrich *et al* 2013).

Land types in MAgPIE include cropland, pasture, forest, built-up and other natural land. Each grid-cell can hold multiple land types at the same time, which add up to the total cell area. Cropland consists of 15 food/feed crop types (e.g. cereals, oil crops, roots), and two bioenergy crop types (woody and grassy). Pasture holds land needed for animal grazing. Forest includes primary forest, modified natural forest and plantations. Built-up area is static in MAgPIE. Other natural land consists of non-forest natural vegetation, abandoned agricultural land and deserts. The distinction between forest and non-forest natural vegetation is based on a carbon density threshold of 20 tC/ha (Hurtt *et al* 2011). The carbon density on abandoned agricultural land increases over time due to regrowth of natural vegetation (Humpenöder *et al* 2014). As soon as the carbon density of re-growing vegetation passes the threshold of 20 tC/ha, the respective area is shifted to the forest land pool.

Demand for bioenergy is exogenous. In this study, bioenergy demand is based on Popp *et al* (2011) and enters the model as a global number. Given the global demand, MAgPIE endogenously derives spatially explicit bioenergy production patterns based on cost-effectiveness (Popp *et al* 2011, Klein *et al* 2014). In MAgPIE, bioenergy demand can only be fulfilled by the production of dedicated grassy and woody energy crops, i.e. residues from agriculture or forestry are not considered. Woody bioenergy, however, is not deployed by the model in this study because grassy bioenergy features higher yields at the same factor costs.

Putting a price on CO₂ emissions from deforestation increases the relative costs of cropland expansion into forests compared to other land conversions (e.g. pasture to cropland). As the objective function of MAgPIE is cost minimization, the model tries to avoid the conversion of carbon-rich forests under a carbon pricing scheme. The same mechanism can be used to incentivize afforestation if carbon dioxide removal is rewarded besides punishing CO₂ emissions (Humpenöder *et al* 2014, 2015). Afforestation is implemented in MAgPIE as managed regrowth of natural vegetation. In this study, we use the same CO₂ price trajectory as in Humpenöder *et al* (2014).

Literature Appendix:

Bondeau A, Smith P C, Zaehle S, Schaphoff S, Lucht W, Cramer W, Gerten D, Lotze-Campen H, Müller C, Reichstein M and Smith B.: Modelling the role of agriculture for the 20th century global terrestrial carbon balance, *Glob Change Biol* **13** 679–706, doi: 10.1111/j.1365-2486.2006.01305.x, 2007.

Bouwman A F, Van der Hoek K W, Eickhout B and Soenario I.: Exploring changes in world ruminant production systems, *Agric. Syst.* **84** 121–53, doi: 10.1016/j.agsy.2004.05.006, 2005.

Daigoglou V, Stehfest E, Wicke B, Faaij A and van Vuuren D P.: Projections of the availability and cost of residues from agriculture and forestry, *GCB Bioenergy* n/a-n/a, doi: 10.1111/gcbb.12285, 2015.

- 1 Dietrich J P, Popp A and Lotze-Campen H.: Reducing the loss of information and gaining accuracy with
2 clustering methods in a global land-use model, *Ecol. Model.* **263** 233–43, doi:
3 10.1016/j.ecolmodel.2013.05.009, 2013.
- 4 Dietrich J P, Schmitz C, Lotze-Campen H, Popp A and Müller C.: Forecasting technological change in
5 agriculture—An endogenous implementation in a global land use model, *Technol. Forecast. Soc.*
6 *Change* **81** 236–49, doi: 10.1016/j.techfore.2013.02.003, 2014.
- 7 Hoogwijk M.: On the global and regional potential of renewable energy sources, Ph.D. Thesis
8 (Utrecht: Universiteit Utrecht, Faculteit Scheikunde), 2004.
- 9 Humpenöder F, Popp A, Dietrich J P, Klein D, Lotze-Campen H, Bonsch M, Bodirsky B L, Weindl I,
10 Stevanovic M and Müller C.: Investigating afforestation and bioenergy CCS as climate change
11 mitigation strategies, *Environ. Res. Lett.* **9** 64029, doi: 10.1038/NCLIMATE2444, 2014.
- 12 Humpenöder F, Popp A, Stevanovic M, Müller C, Bodirsky B L, Bonsch M, Dietrich J P, Lotze-Campen
13 H, Weindl I, Biewald A and Rolinski S.: Land-Use and Carbon Cycle Responses to Moderate Climate
14 Change: Implications for Land-Based Mitigation? *Environ. Sci. Technol.* Online:
15 <http://dx.doi.org/10.1021/es506201r>, 2015.
- 16 Hurtt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K.,
17 Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Goldewijk, K. K., Riahi, K.,
18 Shevlikova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., Wang, Y. P.:
19 Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual
20 land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117–161,
21 doi: 10.1007/s10584-011-0153-2, 2011.
- 22 Klein D, Humpenöder F, Bauer N, Dietrich J P, Popp A, Bodirsky B L, Bonsch M and Lotze-Campen H.:
23 The global economic long-term potential of modern biomass in a climate-constrained world, *Environ.*
24 *Res. Lett.* **9** 74017, doi: 10.1088/1748-9326/9/7/074017, 2014.
- 25 Lotze-Campen, H., Müller, C., Bondeau, A., Rost, S., Popp, A., Lucht, W.: Global food demand,
26 productivity growth, and the scarcity of land and water resources: a spatially explicit mathematical
27 programming approach, *Agric. Econ.*, 39, 325–338, doi: 10.1111/j.1574-0862.2008.00336.x, 2008.
- 28 Müller C and Robertson R D.: Projecting future crop productivity for global economic modeling, *Agric.*
29 *Econ.* **45** 37–50, doi: 10.1111/agec.12088, 2014.
- 30 O'Neill B C, Kriegler E, Ebi K L, Kemp-Benedict E, Riahi K, Rothman D S, van Ruijven B J, van Vuuren D
31 P, Birkmann J, Kok K, Levy M and Solecki W.: The roads ahead: Narratives for shared socioeconomic
32 pathways describing world futures in the 21st century, *Glob. Environ. Change* Online:
33 <http://www.sciencedirect.com/science/article/pii/S0959378015000060>
- 34 Overmars K P, Stehfest E, Tabeau A, van Meijl H, Beltrán A M and Kram T.: Estimating the opportunity
35 costs of reducing carbon dioxide emissions via avoided deforestation, using integrated assessment
36 modelling, *Land Use Policy* **41** 45–60, doi: 10.1016/j.landusepol.2014.04.015, 2014.
- 37 Popp A, Dietrich J P, Lotze-Campen H, Klein D, Bauer N, Krause M, Beringer T, Gerten D and
38 Edenhofer O.: The economic potential of bioenergy for climate change mitigation with special
39 attention given to implications for the land system, *Environ. Res. Lett.* **6** 34017, doi:10.1088/1748-
40 9326/6/3/034017, 2011.

- 1 Popp A, Humpenöder F, Weindl I, Bodirsky B L, Bonsch M, Lotze-Campen H, Müller C, Biewald A,
2 Rolinski S, Stevanovic M and Dietrich J P.: Land-use protection for climate change mitigation, Nat.
3 Clim. Change **4** 1095–8, doi: 10.1038/NCLIMATE2444, 2014.
- 4 Popp A, Lotze-Campen H and Bodirsky B.: Food consumption, diet shifts and associated non-CO2
5 greenhouse gases from agricultural production, Glob. Environ. Change **20** 451–62, doi:
6 10.1016/j.gloenvcha.2010.02.001, 2010.
- 7 Schmitz C, Biewald A, Lotze-Campen H, Popp A, Dietrich J P, Bodirsky B, Krause M and Weindl I.:
8 Trading more food: Implications for land use, greenhouse gas emissions, and the food system, in:
9 Glob. Environ. Change **22** 189–209, doi: 10.1016/j.gloenvcha.2011.09.013, 2012.
- 10 Stehfest E, van Vuuren D, Bouwman L and Kram T.: Integrated Assessment of Global Environmental
11 Change with IMAGE 3.0 : Model description and policy applications (Netherlands Environmental
12 Assessment Agency (PBL)) Online: <http://dspace.library.uu.nl/handle/1874/308545>, 2014.

Highlights:

- LULCC shows a huge effect on the emission of BVOCs.
- Expanding croplands may lead to decreasing BVOC emissions
- Afforestation may lead to increasing BVOC emissions.