UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Isotopic signatures suggest important contributions from recycled gasoline, road dust and non-exhaust traffic sources for copper, zinc and lead in PM_{10} in London, United Kingdom

Dong, Shuofei; Ochoa Gonzalez, Raquel; Harrison, Roy M.; Green, David; North, Robin; Fowler, Geoff; Weiss, Dominik

DOI: 10.1016/j.atmosenv.2017.06.020

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Peer reviewed version

Citation for published version (Harvard):

Dong, S, Ochoa Gonzalez, R, Harrison, RM, Green, D, North, R, Fowler, G & Weiss, D 2017, 'Isotopic signatures suggest important contributions from recycled gasoline, road dust and non-exhaust traffic sources for copper, zinc and lead in PM, in London, United Kingdom', *Atmospheric Environment*, vol. 165, pp. 88-98. https://doi.org/10.1016/j.atmosenv.2017.06.020

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Isotopic signatures in atmospheric particulate matter suggest important contributions from recycled gasoline for lead and non-exhaust traffic sources for copper and zinc in aerosols in London, United Kingdom

Shuofei Dong, Raquel Ochoa Gonzalez, Roy M. Harrison, David Green, Robin North, Geoff Fowler, Dominik Weiss

PII: S1352-2310(17)30401-6

DOI: 10.1016/j.atmosenv.2017.06.020

Reference: AEA 15382

To appear in: Atmospheric Environment

Received Date: 2 February 2017

Revised Date: 6 June 2017

Accepted Date: 10 June 2017

Please cite this article as: Dong, S., Ochoa Gonzalez, R., Harrison, R.M., Green, D., North, R., Fowler, G., Weiss, D., Isotopic signatures in atmospheric particulate matter suggest important contributions from recycled gasoline for lead and non-exhaust traffic sources for copper and zinc in aerosols in London, United Kingdom, *Atmospheric Environment* (2017), doi: 10.1016/j.atmosenv.2017.06.020.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1	
2	
3	
4	
5	
5	
6	Isotopic signatures in atmospheric particulate matter suggest important
7	contributions from recycled gasoline for lead and non-exhaust traffic
8	sources for copper and zinc in aerosols in London, United Kingdom
9	
10	
11	
12 13	Shuofei Dong ^{1,2} , Raquel Ochoa Gonzalez ^{1*} , Roy M. Harrison ^{3,4} , David Green ⁵ , Robin North ⁶ , Geoff Fowler ¹ and Dominik Weiss ¹
14	
15	¹ Imperial College London, Department of Earth Science and Engineering, London, SW7 2AZ, UK
16	² CRPG, UMR 7358 CNRS-UL, Nancy 54500, France
17	³ National Centre for Atmospheric Sciences, Division of Environmental Health and Risk Management, University of
18	Birmingham, B15 2TT, UK
19	⁴ Department of Environmental Sciences / Center of Excellence in Environmental Studies, King Abdulaziz University, PO
20	Box 80203, Jeddah, 21589, Saudi Arabia
21	⁵ Centre for Environment and Health, King's College London, London SE1 9NH, UK
22	⁶ Transport Systems Catapult, Milton Keynes, MK9 1BP, UK
23	
24	
25	
26	*Corresponding author
27	E-mail: raquelocglez@yahoo.es
28	Tel: +44 (0)20 7594 63 83
29	
30	
31	

32 Abstract

33 The aim of this study was to improve our understanding of what controls the isotope composition of Cu, Zn 34 and Pb in particulate matter (PM) in the urban environment and to develop these isotope systems as possible 35 source tracers. To this end, isotope ratios (Cu, Zn and Pb) and trace element concentrations (Fe, Al, Cu, Zn, Sb, Ba, Pb, Cr, Ni and V) were determined in PM₁₀ collected at two road sites with contrasting traffic densities in 36 37 central London, UK, during two weeks in summer 2010, and in potential sources, including non-combustion 38 traffic emissions (tires and brakes), road furniture (road paint, manhole cover and road tarmac surface) and 39 road dust. The isotope signatures of other important sources (gasoline and exhaust emissions) were taken from 40 previous published data. Iron, Ba and Sb were used as proxies for emissions derived from brake pads, and Ni, 41 and V for emissions derived from fossil fuel oil.

The isotopic composition of Pb (expressed using ²⁰⁶Pb/²⁰⁷Pb) ranged between 1.1137 and 1.1364. The 42 isotope ratios of Cu and Zn expressed as δ^{65} Cu_{NIST976} and δ^{66} Zn_{Lvon} ranged between -0.01 ‰ and +0.51 ‰ and 43 44 between -0.21 ‰ and +0.33 ‰, respectively. We did not find significant differences in the isotope signatures 45 in PM_{10} over the two weeks sampling period and between the two sites, suggesting similar sources for each 46 metal at both sites despite their different traffic densities. The stable isotope composition of Pb suggests 47 significant contribution from road dust resuspension and from recycled leaded gasoline. The Cu and Zn 48 isotope signatures of tires, brakes and road dust overlap with those of PM_{10} . The correlation between the 49 enrichments of Sb, Cu, Ba and Fe in PM_{10} support the previously established hypothesis that Cu isotope ratios 50 are controlled by non-exhaust traffic emission sources in urban environments (Ochoa Gonzalez et al., 2016). 51 Analysis of the Zn isotope signatures in PM_{10} and possible sources at the two sites suggests significant 52 contribution from tire wear. However, temporary additional sources, likely high temperature industrial 53 emissions, need to be invoked to explain the isotopically light Zn found in 3 out of 18 samples of PM_{10} .

54

55 **1. Introduction**

56

57 Metals in atmospheric particulate matter (PM) are a serious concern as they induce adverse health effects. 58 Cardiopulmonary diseases have been attributed to the capacity of inhaled particles and metals to cause 59 inflammation and injury at the air-lung interface (Costa and Dreher, 1997; Chen and Lippmann, 2009; Gamble 60 and Lewis, 1996; Jarup, 2003; Ozkaynak et al., 1996). Lead exposure damages the human central nervous 61 system and kidneys, affects biochemical processes and impairs psychological and neurobehavioral functions 62 (Tong et al., 2000). Zinc is associated with oxidative stress and a contributing factor in many chronic diseases 63 (Prasad et al., 2004; Walsh et al., 1994), and Cu causes damage to proteins and lipids (Brewer, 2010). Thus, 64 developing a full understanding of the major sources of metals in urban PM is critical to control toxic 65 emissions in urban environments.

Previous studies analyzed the sources of multiple elements in urban PM by using conventional chemical 66 67 techniques and receptor modeling methods such as principal component analysis (PCA) and positive matrix 68 factorization (PMF). These suggested that industrial emissions, exhaust and non-exhaust traffic emissions 69 and resuspended road dust are the major sources that contribute to the metal aerosol burden in large cities 70 around the world including Birmingham, Thessaloniki, and Boston (Cooper and Watson, 1980; Harrison et al., 71 1997; Laden et al., 2000; Lin et al., 2010; Manoli et al., 2002; Polissar et al., 2001; Querol et al., 2001; 72 Thurston and Spengler, 1985). Iron and Cr are often associated with the bulk matrix from road dust, Zn from 73 tire and brake wear, and road dust re-suspension, Pb from the past use of leaded gasoline, Fe from brake drum 74 abrasion, wear-off of engines, tires and brakes, and V and Ni with emissions from oil combustion (Garg et al., 75 2000; Hjortenkrans et al., 2007; Manoli et al., 2002; Thorpe et al. 2007; Thurston et al., 2011). Copper, Sb 76 and Ba are abundant in brake lining materials whereas brakes, yellow road paint and road pavement are 77 typically sources of Cr (Adachi and Tainosho, 2004; Garg et al., 2000; Hjortenkrans et al., 2007; Iijima et al., 78 2007; Rodriguez et al., 2004; Salma and Maenhaut, 2006; Thorpe and Harrison, 2008).

79 The success of using metal concentrations in combination with PCA and PMF to identify single sources of 80 trace metals remains limited. PCA does not provide a unique solution because of its simplified approach and 81 the results of PMF vary depending on different uncertainty structures (Samek et al., 2016). Consequently, the 82 use of isotope ratios is widely investigated to improve source tracing in atmospheric particles (Wiederhold, 83 2015, Weiss et al., 2008). This approach was indeed highly successful for Pb, where isotopes identified 84 leaded gasoline as the main source in urban PM in London, and despite the restrictions in the use of leaded 85 gasoline in the 1980s, a contribution of 60% was still found in west London between 1995 and 1996 (Monna et 86 al., 1997; Noble et al., 2008). Stable isotope systems have been tested for source apportionment of trace 87 elements in PM such as Fe, Sr, Zn and Nd (Flament et al., 2008; Geagea et al., 2008; Majestic et al., 2009; 88 Widory et al., 2010). Zinc isotopes in urban PM were studied in Metz (Cloquet et al., 2006), São Paulo (Gioia et al., 2008), and Barcelona and London (Ochoa Gonzalez et al., 2016). In Metz, the δ⁶⁶Zn_{Lvon} in ambient PM 89 90 collected in a bus and an urban waste incinerator showed similar values close to +0.12 ‰. In São Paolo, in 91 contrast, Gioia et al. (2008) found lighter ratios and significantly larger variations in δ^{66} Zn_{Lyon}, ranging between 92 -1.05 and -0.46 ‰ in PM_{2.5-10} and between -1.13 and -0.07 ‰ in PM_{2.5}. In line with these findings, Ochoa 93 Gonzalez et al. (2016) found isotopically light Zn in PM_{10} collected in Barcelona, with $\delta^{66}Zn_{Lvon}$ ranging 94 between -0.45 and -0.83 ‰, and isotopically heavy Cu in coarse PM collected in London during late autumn

and the winter season (δ^{65} Cu_{NIST976} between +0.66 and +0.97 ‰) (Ochoa Gonzalez et al., 2016). The observed 95 isotope signatures in the PM were associated with sources derived from combustion and smelting processes. 96 97 Other studies are in line with this interpretation as they showed that residues and flue gas particles from metal 98 refining and coal combustion are highly fractionated relative to the starting materials, with isotopically lighter 99 Zn emitted in the PM (between -0.73 and -0.63 ‰) and heavy Zn remaining in the residues (up to +1.49 ‰) 100 (Mattielli et al., 2009; Ochoa Gonzalez and Weiss, 2015; Sivry et al., 2008). These signatures seem to 101 partially overlap with the isotope ratios of Zn in anthropogenic materials produced during low temperature processes like galvanisation which show δ^{66} Zn_{Lvon} between +0.12 and +1.15 ‰ (Araújo et al., 2017; John et al, 102 103 2007). In summary, these studies suggested strongly that Cu and Zn isotopes can play an important role in the 104 identification of emission sources from metal refining and combustion due to their larger fractionation, as well 105 as Pb isotopes to assess the contribution from the past use of leaded gasoline.

106 The aim of this study was to improve our present knowledge on what controls the isotopic variability of Cu, 107 Zn and Pb in urban PM and to critically assess the potential of these isotope systems for source tracing. To this 108 end, we determined short-term temporal and spatial isotope variability of Cu, Zn and Pb in PM₁₀ collected in 109 2010 in central London and in potential sources including road furniture (manhole cover, road paint and road 110 tarmac surface), non-combustion vehicle sources (tire and brake wear) and road dust (that represents other 111 potential sources deposited in the road environment). We collected possible sources and PM_{10} samples at a high 112 and a low traffic density site over a 20-day sampling period during summer. The concentrations of trace elements including Fe, Sb, Ba, Cr, Ni, and V, and enrichment factors (EF) were combined with the isotope ratios of Zn, 113 114 Cu and Pb to constrain possible contributions from sources. Finally, we critically compare the isotope data in 115 PM₁₀ and potential sources in London with previously published isotope signatures of other potential sources and PM from major urban cities. 116

117

118 **2. Materials and methods**

119

120 **2.1. Sample collection**

Samples of PM₁₀ were collected for 48 h using a Partisol-Plus Model 2025 Sequential Air Sampler (Thermo 121 Scientific, MA, USA) at two different road sites in central London between 8th July 2010 and 28th July 2010 at a 122 123 height of 3.5 m. Nine samples (MR1-MR9) were collected at the heavily traffic street canyon Marylebone 124 Road (denoted as MR) in London, with more than 80,000 vehicles per day and frequently congested (Charron 125 and Harrison, 2005). Nine samples (NK1-NK9) were collected at a sampling location in North Kensington 126 (denoted as NK) which is a lightly trafficked open urban site situated next to a school playground (Bigi and 127 Harrison, 2010). The sampling site NK has an average daily traffic flow of 8,000 vehicles and the nearest road 128 is more than 30 m away (Abdalmogith and Harrison, 2006). Polytetrafluoroethylene (PTFE) filters were used 129 for PM_{10} collection and leached prior use for 2 days in 1.5 M sub-boiling distilled HNO₃ at 100°C on a hotplate.

Potential ssources for various metals were constrained following a critical literature review and included brake pads for Cu, Sb and Ba; tire wear and road dust for Zn, Fe, Pb and Cr; road paint and road surface for Cr (Adachi and Tainosho, 2004; Hjortenkrans et al., 2007; Thorpe and Harrison, 2008; Thorpe et al., 2007), and manhole covers for Zn (Fry et al., 2005). The isotopic composition of Zn and Pb from gasoline and vehicle exhaust emissions have been taken from the literature (Cloquet et al., 2006; Gioia et al., 2008; Monna et al., 135 1997).

Three samples of road dust were taken at each site, one next to the PM sampling tube and two in the 136 137 opposite directions from each sampling site. Approximately 30 g of road dust were taken using a brush and a 138 pan previously cleaned; samples were sieved at 2 mm and then milled using an agate pestle and mortar for 139 subsequent analysis. The road dust samples were taken during clear days without rainfall 24 h prior sample 140 collection. Road furniture (manhole cover and road paint, two samples at each site; road tarmac surface, two 141 samples at the NK site) were sampled within approximately 10 m of each aerosol sampling site and were 142 stored in plastic bags. There were no natural soils exposed close to the sampling sites. There were manhole 143 covers in the middle of the roads where the PM₁₀ were collected, as well on the side. There were 144 approximately five manhole covers within 50 m.

Non-combustion vehicle sources, i.e., tear-off from tires and brakes, were characterized using a collection of used tires and new brake pads. Twelve individual car tires of common commercial brands were shredded and mixed in three sets (labelled as R43/54-1, R43/54-2 and R56/57-1) before analysis. The brake samples were scratched from the surface of two different brands of metallic brake pads commonly used. These include metallic pads HBP Ferodo for Peugeot and Renault cars, and FSB Halfords for Volkswagen family cars (labelled as HBP and FSB, respectively).

151

152 **2.2. Sample preparation**

The filters were weighed before and after the collection to determine the mass of material. The metals were 153 154 leached from the filter with 5 ml of aqua regia for 24 hours followed by 10 ml of 0.5 M HNO₃ for 48 hours on a 155 hotplate at 80°C. An ultrasonic bath was used for 1 hour every 12 hours to improve the removal efficiency. The filters were then washed with 5 ml 0.5 M HNO₃ three times. The two solutions were mixed and evaporated 156 157 to dryness before further treatment. The standard reference material BCR-723 (road dust) and all the samples 158 were digested using a conc. HNO3:HClO4 mixture (1:4) in sealed PFA vials (Savillex, MN, USA) at 150°C in a 159 hotplate for 24 hours. HClO₄ was used to favor the oxidation of the organic matter. The samples were dried 160 and digested using a mixture of conc. HNO_3 and HF (1:4) on a hotplate at 140°C for another 24 hours. The 161 solutions were taken to dryness again. The samples were re-fluxed in 300 µl of conc. HNO₃ and evaporated twice to remove the excess of fluorides. After re-dissolution in 2.5 ml of 0.5 M HNO₃, 0.5 ml was used for 162 163 concentration measurement using quadrupole ICP-MS and the remaining volume of each solution was prepared 164 for anion exchange column separation of Pb, Cu and Zn.

165 All the experimental work was carried out in a Class 1000 clean lab and under Class 10 laminar flow 166 hoods in the MAGIC laboratories at Imperial College London. Sub-distilled mineral acids (HNO₃ 15 mol l^{-1} , 167 HCl 6 mol l^{-1}), commercial HF (40 % v/v Suprapur, Merck, Germany) and HClO₄ (70 % v/v Suprapur, Merck, 168 Germany), and 18.2 MΩ·cm grade Millipore water (Bedford, MA, USA) were used throughout this study.

169

170 **2.3. Concentration analysis and enrichment factors**

The concentration of Fe, Al, Cu, Zn, Sb, Ba, Pb, Cr, Ni and V were determined using quadrupole ICP-MS
(Varian, CA, USA). Table S1 shows the concentrations of Cu, Zn, Sb, Ba and Pb determined for BCR-723
(road dust) and the indicative values provided by the European Commission Joint Research Centre Institute for

- 174 Reference Materials and Measurements. The analytical uncertainty of the concentration determinations for all
 - sence Materials and Measurements. The analytical differentiative of the concentration determinations for

- elements was better than 6% (n=5, 2SD) and the recoveries, relative to the indicative values were higher than
 89 % with respect to the reference values for all the elements.
- 177 The EFs of the PM_{10} were calculated to assist the source assessment study using Al as as a proxy for the 178 upper continental crust according to Eq. 1,

$$EF_{C} = \frac{C_{sample}/Al_{sample}}{C_{crust}/Al_{crust}}$$
 [Eq. 1]

180 where C is the element of interest and C_{sample} , Al_{sample} , C_{crust} and Al_{crust} are the concentrations of C and Al in the 181 sample and the continental crust. The average chemical composition of the upper continental crust was used as 182 previously reported (Wedepohl, 1995).

183

203

179

184 **2.4. Isotope ratio analysis of Cu, Zn and Pb**

185 Copper, Zn and Pb in all of the samples were separated from other matrix elements using ion exchange 186 chromatography. The details of the Cu, Zn and Pb separation methods and sample treatment are described in 187 detail elsewhere (Dong et al., 2013; Weiss et al., 2004) and summarized in the supplementary material. The 188 isotope ratios were determined using a Nu Plasma multi collector ICP-MS (Nu Instruments Limited, UK) 189 equipped with a Nu DSN-100 Desolvation Nebuliser System. The instrumental mass bias was corrected using 190 standard-sample bracketing for Cu isotope ratio measurement (Peel et al. 2008), Tl doping for Pb isotope ratio measurements (Weiss et al., 2004) and a ⁶⁴Zn-⁶⁷Zn double spike for Zn isotope ratio measurements (Arnold et 191 192 al., 2010, Ochoa Gonzalez et al., 2016).

193 The total procedural blank was <4 ng for Cu and Zn, and <80 pg for Pb. These contributions were less 194 than 0.01% of the total element content in all samples analyzed and had no significant effect on the accuracy of 195 the isotope ratio measurements. The Romil Cu and Romil Zn solutions were processed and measured with the samples and showed values of δ^{65} Cu_{NIST976} = +0.17 ± 0.10 ‰ (2SD, n=24) and δ^{66} Zn_{Lvon} = -9.12 ± 0.08 ‰ (2SD, 196 n=20), respectively. These are within error relative to previous published values for Romil Cu (δ^{65} Cu_{NIST976}= 197 198 $+0.18 \pm 0.06$ ‰, n=19, Moeller et al., 2012; δ^{65} Cu_{NIST976} = $+0.18 \pm 0.05$ ‰, n=4, Ochoa Gonzalez et al., 2016) and Romil Zn (δ^{66} Zn_{Lyon} = -9.14 ± 0.08 ‰, n=21, Chapman et al., 2006). During this study ²⁰⁶Pb/²⁰⁴Pb, 199 207 Pb/ 204 Pb and 208 Pb/ 204 Pb for NIST-SRM 981 Pb (when treated as non-spiked standard) were 16.9372 \pm 200 201 0.0021, 15.4954 ± 0.0021 and 36.7158 ± 0.0057 (n=50), respectively, which are in good agreement with 202 previous published data (Weiss et al., 2004).

204 **3. Results and discussion**

205 3.1. Source assessment based on concentrations and enrichment factors

The concentrations of PM_{10} and of Fe, Cu, Al, Zn, Sb, Ba, Fe, Cr, Pb, Ni and V in PM_{10} and local sources are shown in Table 1 and Table 2. The concentrations of PM_{10} are approximately twice as high at the MR site $(31\pm7 \ \mu g \ m^{-3})$ compared to the NK site $(14\pm3 \ \mu g \ m^{-3})$. According to London Air, average hourly concentrations of PM_{10} at the MR and NK sites over the sampling period during July 2010 are $40\pm15 \ \mu g \ m^{-3}$ and $16\pm7 \ \mu g \ m^{-3}$, respectively (London Air Website, www.londonair.org.uk).

The concentrations and EFs of Fe, Cu, Zn, Sb, Ba and Cr in the PM_{10} collected at the MR site are much higher than those at the NK site (Tables 1 and 3). These elements are highly enriched relative to natural dust represented geochemically by the upper continental crust (Table 3), suggesting that they are derived from traffic related emissions. Iron, Cu, Zn, Sb and Ba are typically high in tires and brakes (McKenzie et al., 2009). The correlations between the EF_{Sb} and those of Fe, Cu, Zn and Ba (Figure 1) in PM₁₀ at the NK and the MR sites show R² between 0.8686 and 0.9972, supporting the hypothesis of similar sources at the sampling sites and of non-combustion traffic emissions as main source. The EF_{Sb} and those of typical tracers of brakes (i.e., Fe, Cu and Ba) show similar and very high R² (R² between 0.9949 and 0.9972), suggesting that non-exhaust traffic emissions are the dominant source for these elements in urban PM₁₀ in London.

Re-suspension of small particles of road dust may have a significant effect on the generation of PM_{10} enriched in metals (Amato et al., 2014; Thorpe and Harrison, 2008). The road dust samples collected at the MR site (MRRD-1, MRRD-2 and MRRD-3) are enriched in Fe, Cu, Zn, Sb, Ba, Cr and Pb compared to road dust at the NK site (NKRD-1, NKRD-2 and NKRD-3) (Table 3), most probably reflecting the higher traffic density. This observation supports the idea that re-suspended road dust is an important source of these elements in urban PM (Amato et al., 2014).

Brake wear is the dominant source of Ba in urban and natural PM (Gietl and Klemm, 2009; Gietl et al., 2010; Iijima et al., 2007; Moreno et al., 2015). We find high concentrations of Ba in brakes $(310 - 1300 \ \mu g \ g^{-1})$ and samples of manhole cover $(120-1100 \ \mu g \ g^{-1})$ (Table 2). The concentrations of Ba in the PM₁₀ at the NK site $(1.6\pm0.4 \ ng \ m^{-3})$ are much lower than at the MR site $(5.7 - 84 \ ng \ m^{-3})$, supporting the dominant contribution of brake wear.

The concentration of Zn in PM₁₀ ranges between 1.4 and 3.1 ng m⁻³ at the NK site, and between 3.7 and 41 231 232 ng m⁻³ at the MR site. The concentration of Cu in PM_{10} ranges between 1.9 and 3.6 ng m⁻³ at the NK site and between 9.4 and 130 ng m⁻³ at the MR site. The element concentrations show that the content of Zn in tires 233 234 and of Cu in brakes, tires and manhole cover are significantly higher than in other sources (Table 2). In addition, 235 samples of road tarmac surface and manhole cover contain significant amounts of Zn. Cu/Sb ratios between 3.3 236 and 9.1 have been used to identify brake wear in PM collected in Cologne, London, Stockholm and Budapest 237 (Hjortenkrans et al., 2007; Salma and Maenhaut, 2006; Weckwerth, 2001). In this study similar Cu/Sb ratios 238 are observed in PM₁₀, although with a smaller range between 5.8 and 8.3 (Table 1). The average Cu/Sb ratios in 239 the PM₁₀ at the NK site are slightly higher than those in the PM₁₀ at the MR site. However, the Cu/Sb ratios in 240 the studied brake pads range between 27 and 39 and are large compared with typical values for brakes in PM 241 (Table 2). This observation is likely explained by the recent restrictions on the use of Sb_2S_3 during the 242 manufacturing process of brake pads due to its potential carcinogenic properties (Hüner et al., 2001).

243 The concentrations of Cr in PM₁₀ collected at the MR site $(0.9 - 8.2 \text{ ng m}^{-3})$ are higher than those at the 244 NK site (0.3±0.1 ng m³). Multiple sources including brakes, manhole cover, yellow road paint (NKRP-2) and 245 road dust are enriched in this element relative to natural dust (Table 3). However, the positive correlation between the EF_{Sb} and the EF_{Cr} ($R^2 = 0.9385$) in PM_{10} (Figure 1), and between the EF_{Sb} and those of other 246 247 elements linked with non-exhaust traffic emissions, i.e., Cu, Ba, and Fe, suggests that not only Cu, Ba and Fe, 248 but also Cr is mainly controlled by non-exhaust traffic emissions in the urban atmosphere. The concentrations 249 of Fe and EF_{Fe} in the PM₁₀ collected at MR are higher than those in the PM₁₀ at NK, and significantly larger 250 concentrations of this element are found in samples of brakes and manhole cover. In addition to these sources, 251 brake discs contribute significantly to PM₁₀ enriched in Fe (Wåhlin et al., 2006).

252 The elemental concentrations and EF of Ni and V are similar in the PM_{10} at both sites, which suggests that

these elements are not directly affected by the density of traffic (Tables 1 and 3). Previous studies suggested that Ni and V are mainly derived from fuel oil, gas refineries, and industrial processes (Moreno et al., 2010; Nriagu and Pacyna, 1988; Peltier and Lippmann, 2010), and less from non-exhaust traffic sources (Thorpe and Harrison, 2008). In a recent receptor modeling study conducted using PM_{10} samples collected at the NK site (Beddows et al., 2015), Ni and V were found to be associated with high concentrations of sulphate in a fuel oil-related factor.

Lead concentrations in PM_{10} are slightly higher at the MR site (1.1-8.8 ng m⁻³) compared to the NK site (1.0±0.3 ng m⁻³) but the EF_{Pb} are similar at both sites. Some studies based on emissions of metals in PM_{10} and PM_{2.5} suggested that following the phasing out of leaded gasoline, Pb is emitted into the urban atmosphere from fuel and motor oil combustion or brake wear (Lough et al., 2005). Significantly high concentrations of Pb relative to other possible sources have been found in tires and brakes (4.0 – 38 µg g⁻¹) and in manhole covers (56 – 690 µg g⁻¹) (Table 2).

265

266 **3.2 Source identification of lead using stable isotope ratios**

The ²⁰⁶Pb/²⁰⁴Pb, ²⁰⁶Pb/²⁰⁷Pb and ²⁰⁸Pb/²⁰⁶Pb isotope ratios determined in the PM₁₀ and the sources are shown in 267 Table 4. Figure 2a shows that the Pb isotope ratios in PM_{10} are similar at the MR and NK sites. This 268 observation, and similarity between the EF_{Pb} in the PM₁₀ at the MR and NK sites (Table 1), suggest that the 269 source(s) at NK and MR sites are the same. The Pb isotope ratios measured in the PM_{10} at NK ($^{206}Pb'^{204}Pb =$ 270 17.446-17.598, ${}^{206}Pb'^{207}Pb = 1.1218 - 1.1304$, ${}^{208}Pb'^{206}Pb = 2.1288 - 2.1382$) fall within the respective ranges at 271 MR ($^{206}Pb'^{204}Pb = 17.305 - 17.701$, $^{206}Pb'^{207}Pb = 1.1137 - 1.1364$, $^{208}Pb'^{206}Pb = 2.1246 - 2.1476$). These Pb 272 signatures are significantly different from those of natural sources in the UK ($^{206}Pb^{/207}Pb= 1.18 - 1.19$. Monna 273 274 et al., 1997), which suggests an ongoing contribution of anthropogenic sources of Pb in atmospheric PM. The ²⁰⁸Pb^{/206}Pb and ²⁰⁶Pb^{/207}Pb isotope ratios determined in our study are similar to those reported in 1995, 1998 and 275 1999 and partly overlap with years 1996, 2000 and 2001 (Monna et al., 1997; Noble et al., 2008) (Figure 2b). 276 The ²⁰⁶Pb^{/207}Pb determined for urban PM₁₀ in our study fall within the range determined in France and the UK 277 278 during mid-90s, that ranged between 1.1008 and 1.1453 (Monna et al., 1997). Noble et al. (2008) showed that 279 since leaded gasoline was banned in the UK in 1998, the ²⁰⁶Pb/²⁰⁷Pb isotope ratio in urban PM in London shifted slightly from 1.1169 - 1.1388 to more radiogenic ratios during 2000 and 2001, and they stabilized 280 thereafter (1.1354 – 1.1507). The isotope ratios of road paint ($^{206}Pb'^{204}Pb = 17.866 - 18.301$, $^{206}Pb'^{207}Pb = 17.866 - 18.301$, $^{206}Pb'^{207}Pb = 17.866 - 18.301$, $^{206}Pb''^{207}Pb = 18.300$, $^{206}Pb''^{207}Pb = 18.300$, $^{206}Pb''^{207}Pb = 18.300$, 206 281 1.1471 - 1.1708, ${}^{208}Pb'^{206}Pb = 2.0843 - 2.1012$) and of road dust (${}^{206}Pb'^{204}Pb = 17.347 - 17.890$, ${}^{206}Pb'^{207}Pb = 17.347 - 17.890$, 282 1.1163 - 1.1417, ²⁰⁸Pb^{/206}Pb = 2.1176 - 2.1446) collected at the two sampling sites show significant Pb isotope 283 284 ratio variability. As illustrated in Figure 2a, the Pb isotope ratios of the PM_{10} fall on a mixing line with 285 gasoline used in the UK before its phasing out as one end member (Monna et al., 1997), and road dust which 286 contains re-mobilized Pb from leaded gasoline and other source deposits such as tires, brakes and road 287 furniture as the other end member. The Pb isotope composition of the PM_{10} collected at NK and MR sites is 288 similar to those of road dust and manhole cover which suggests that re-suspended road dust may contribute to 289 Pb enrichment in urban PM. Previous elemental analyses of PM₁₀ from the NK site showed appreciable 290 concentrations of crustal elements such as Al and Ca, that might have arisen from road dust (Beddows et al., 291 2015). Therefore, it seems likely that road dust with residual contamination from the earlier use of leaded 292 gasoline contribute to PM₁₀ due to resuspension into the atmosphere.

294 **3.3** Copper and zinc isotope ratios variations in source materials and PM₁₀

295 The Cu and Zn isotope ratios of PM₁₀ collected at the sites with high and low traffic densities and the potential 296 local sources are illustrated in Figure 3. Previous published data for PM2.5-80 collected in London between 2014 297 and 2015 and for PM₁₀ collected in Barcelona between 2012 and 2013 (Ochoa Gonzalez et al., 2016) are shown 298 for reference. The δ^{65} Cu_{NIST976} values vary in the range between +0.01±0.13 ‰ and +0.51±0.10 ‰ at the MR site and between -0.01±0.10 ‰ and +0.46±0.07 ‰ at the NK site which suggests same sources of Cu in the 299 PM₁₀ collected at the two sampling sites (Table S2-Sn). This agrees with our hypothesis derived from the 300 301 linear correlations between the EF of typical tracers of non-exhaust emissions (i.e. Cu, Sb, Ba, Zn and Fe) in PM_{10} (Figure 1). The $\delta^{66}Zn_{Lyon}$ values in PM_{10} collected at the MR site range between -0.14±0.04 ‰ and 302 303 $+0.33\pm0.04$ %, and at the NK site between -0.21 ± 0.14 % and $+0.26\pm0.12$ %. We do not observe a significant 304 temporal trend in the isotope composition during the sampling period of 20 days and there is no correlation 305 between the shifts in the isotopic composition of Cu and those of Zn (Table S2-Sn). The latter observation 306 probably reflects different contribution from the non-exhaust traffic sources of Zn and those of Cu. Although 307 higher Cu and Zn concentrations are found at the high traffic site (Table 1), the isotope data set suggests that 308 the dominant sources and/or controlling processes are the same at both sites studied. The similar variability in 309 the Cu and Zn isotope ratios in the PM_{10} collected at the low and high traffic sites during 3 weeks over July 310 2010 (Figure 3) suggests that the sources of Cu and Zn in PM_{10} do not change on that temporal and spatial 311 scale. This observation is in good agreement with previous studies over an entire year that showed only 312 significant Cu isotope variability during the winter season and suggested that traffic sources are the main 313 contributor to Cu and Zn enrichment in PM_{2.5-80} (Ochoa Gonzalez et al., 2016) and PM_{2.5-10} (Visser et al., 2015) 314 in London. The Cu and Zn isotope ratios of PM2.5-80 collected at building height during July and August 2014 and July and August 2015 showed $\delta^{65}Cu_{NIST976}$ between +0.46±0.12 ‰ and +0.55±0.15 ‰ and $\delta^{66}Zn_{Lyon}$ 315 between +0.02±0.04 ‰ and +0.17±0.02 ‰, with one light sample collected during summer 2015 (δ^{66} Zn_{Lvon}= 316 317 -0.29 ± 0.04 ‰) (Ochoa Gonzalez et al., 2016). The Cu and Zn isotope signatures in PM₁₀ collected in London 318 at the NK and MR sites during 2010 in London agree well with the isotope signatures in PM2.5-80 collected in 319 another street canyon during 2014 and 2015 in the same city (Ochoa Gonzalez et al., 2016).

320

293

321 3.3.1 Controls of copper isotope signatures in urban particulate matter in London

322 Brakes, tires, manhole cover, road paint, and road tarmac surface have $\delta^{65}Cu_{NIST976}$ values ranging between 323 -0.18±0.14 ‰ and +0.71±0.09 ‰ (Table S2-Sn). Four out of the six road dust samples show Cu isotope signatures falling within this range and two samples show more negative isotope signatures ($\delta^{65}Cu_{NBS976} =$ 324 325 -0.28±0.08 ‰ and -0.22±0.08 ‰) at the MR site. Isotopically heavy Cu is found in road paint, road tarmac surface and selected samples of brakes and manhole cover, with $\delta^{65}Cu_{NBS976}$ values above +0.55 %. The 326 δ^{65} Cu_{NIST976} of the non-exhaust traffic sources (i.e., tires and brakes) show significant isotope variability, 327 328 ranging between $+0.17\pm0.10$ ‰ and $+0.33\pm0.12$ ‰ for tires and between $+0.28\pm0.10$ ‰ and $+0.63\pm0.14$ ‰ for 329 brakes, most probably reflecting the different Cu species such as metallic Cu, Cu oxides or Cu sulfides 330 (Grigoratos and Martini, 2015; Straffelini et al., 2015), the different geographical location of the ores used 331 and/or possible isotopic fractionation during the manufacturing processes.

332 Cu/Sb ratios in the PM have been used to identify source contribution of brake wear in urban PM

333 (Hjortenkrans et al., 2007; Salma and Maenhaut, 2006; Weckwerth, 2001). The relationship between Cu/Sb 334 ratios and the Cu isotope composition in the samples of PM analysed during this work and a previous study 335 conducted in Barcelona and London (Ochoa Gonzalez et al., 2016) is illustrated in Figure 4. The PM₁₀ samples from Barcelona and London with low Cu/Sb ratios ranging between 5.8 and 12, which fall within the range of 336 Cu/Sb found for brakes, have isotopically lighter Cu with δ^{65} Cu_{NBS976} ranging between -0.01 and +0.51 ‰. 337 Indeed, these isotope signatures largely overlap with the Cu isotope signatures in samples of brakes analyzed 338 during this study (δ^{65} Cu_{NIST976} = +0.28± 10‰ - +0.63±14 ‰) (Table S2-Sn). Coarse particles (PM_{2.5-80}) in 339 London with Cu/Sb ratios larger than 12 and isotopically heavier Cu (δ^{65} Cu_{NIST976} ranging between +0.63 and 340 341 +0.97 ‰) were attributed to particles emitted during fossil fuel combustion emissions which seemed more 342 important during end-autumn and the winter season (Ochoa Gonzalez et al., 2016).

343

344 3.3.2 Controls of zinc isotope signatures in urban particulate matter in London

Brakes, tires, manhole cover, road paint, and road tarmac surface show δ^{66} Zn_{Lyon} ranging between -0.02±0.09 ‰ 345 and +0.58±0.10 ‰ (Table S2-Sn). The Zn isotope signatures of the samples of road dust range between 346 347 +0.21±0.05 ‰ and +0.34±0.10 ‰ at both sites, except one sample with a δ^{66} Zn_{Lvon} of -0.03±0.08 ‰ at the NK 348 site. These ratios overlap with the signatures in PM₁₀ at the NK and MR sites. The variations in the Zn isotope 349 signatures found in the road dust samples at the two sites likely reflect the potential accumulation of different 350 sources. Manhole cover, road paint and road surface show significant isotope variability within each source 351 type, up to 0.51 ‰. This suggests that the source materials and/or processes during the manufacturing of these 352 products have an effect on the isotope variability of each kind of source. The isotopically heaviest Zn has been measured in road paint (δ^{66} Zn_{Lyon} between +0.07 and +0.58 ‰, n=4). The Zn isotope composition of 353 354 brakes and tires vary within a small range with δ^{66} Zn_{Lyon} between +0.15±0.07 ‰ and +0.21±0.08 ‰ (n=6) and 355 are not significantly different relative to an average internal precision of 0.08 ‰. Together with the high 356 levels of Zn in tires (Table 2), we propose that Zn in most of the PM₁₀ sampled at the NK and the MR sites 357 $(\delta^{66}$ Zn_{Lvon} ranging between -0.04±0.14 ‰ and +0.33±0.04 ‰) are derived from tires (Figure 3). However, a 358 source enriched in isotopically light Zn which can explain the Zn isotope signatures found in three samples of 359 PM₁₀, i.e., in NK3, NK7 and MR8 (Figure 3) has not been identified. The isotopic composition of Zn in PM₁₀ in London is similar to that of PM_{2.5-80} collected at a high density traffic site in the same city during 2014 and 360 2015, with δ^{66} Zn_{Lvon} ranging between -0.29 ‰ and +0.17 ‰ (Figure 3). The δ^{66} Zn_{Lvon} in bus air filters 361 362 analysed during previous studies (Cloquet et al., 2006) overlap with the signatures in the PM_{10} collected at the 363 NK and MR sites, whereas those of gasoline (Gioia et al., 2008) only overlap with three samples of PM_{10} out of 364 18 (NK3, NK7 and MR8) (Figure 3, Table S2-Sn). The relatively constant traffic densities and the similar 365 Zn/Sb ratios in PM₁₀ at each sampling site (Table 2) suggest that contribution from gasoline combustion is less 366 likely, in line with the typically low concentrations of Zn found in gasoline (Pignalosa et al., 2005). The light Zn isotope signatures found in PM₁₀ in Barcelona (δ^{66} Zn_{Lvon} = -0.83 - -0.45 ‰) were tentatively attributed to 367 368 metallurgical emissions (Ochoa Gonzalez et al., 2016). This hypothesis agrees with previous observations 369 pointing that industrial processes such as smelting or combustion exert an effect on the fractionation due to 370 evaporation and/or condensation processes (Mattielli 2009; Bigalke 2010; Ochoa Gonzalez and Weiss, 2015; 371 Thapalia et al., 2015). These findings suggest that tire wear largely contribute to Zn in PM₁₀ in London, 372 whereas for some samples the isotope signatures suggest a minor contribution from high temperature industrial

or fossil fuel combustion.

374

375 **4. Conclusions**

376 The aim of this study was to determine the short-term temporal and spatial isotope variability of Cu, Zn and Pb 377 in urban PM₁₀ collected at two sites with different traffic densities in central London, UK, and in possible sources, including non-combustion traffic emissions (tires and brakes), road furniture (road paint, manhole 378 379 cover and road tarmac surface) and road dust. Samples of PM₁₀ and possible sources were also characterized 380 for the EF of other metals (Ni, V, Fe, Sb, Ba and Cr) to assist the source apportionment study. Copper and Zn 381 concentrations and EF in the PM₁₀ were higher at the sampling location with high traffic density, along with 382 other elements commonly associated with traffic related sources including Fe, Sb, Ba and Cr. In contrast, the 383 EF of Pb showed no significant differences between the two sites although was enriched relative to crustal 384 values.

The ²⁰⁶Pb/²⁰⁷Pb in the PM₁₀ ranges between 1.1137 and 1.1364, and the isotope ratios of Cu and Zn 385 expressed as δ^{65} Cu_{NIST976} and δ^{66} Zn_{Lvon} range between -0.01 and +0.51 ‰ and between -0.21 and +0.33 ‰, 386 respectively. The results indicate that traffic density does not exert a significant effect on the observed isotope 387 388 signatures, suggesting similar sources for these toxic metals at both sites. The Pb isotope signatures of the PM₁₀ 389 collected at the NK and MR sites overlap with those of road dust which suggests that re-suspended road dust is 390 an important source. The similarity between the Cu isotope signatures in samples of PM_{10} ($\delta^{65}Cu_{NRS976}$ ranging between -0.01±0.10 ‰ and +0.51±0.10 ‰) and brakes (δ^{65} Cu_{NIST976} ranging between +0.28±0.10 ‰ and 391 392 +0.63±0.14 ‰), and the low Cu/Sb ratios (Cu/Sb ranging between 5.8 and 12) which fall within the range of 393 Cu/Sb found for brakes, suggest that brake wear is the dominant source of Cu. The Zn isotope signatures in 394 PM₁₀ at NK and MR overlap with the signatures of tires and road dust but additional sources such as high 395 temperature industrial emissions that impart isotopically light Zn, need to be invoked to explain the light 396 signatures found in 3 out of 18 samples.

397

398 Acknowledgments

399 We thank Barry Coles and Katharina Kreissig for assistance in the MAGIC laboratories and Stanislav 400 Strekopytov for the analysis by ICP-MS at the Natural History Museum. Johanna Gietl is thanked for her help 401 during sampling collection. Raquel Ochoa Gonzalez thanks the European Commission 402 (FP7-PEOPLE-2012-IEF) for funding the project ISOTRACE (proposal 329878). We acknowledge financial 403 support from the UK-China Scholarships for Excellence for Shuofei Dong.

405 **References**

- 406 Abdalmogith, S.S., Harrison, R.M., 2006. An analysis of spatial and temporal properties of daily sulfate, nitrate and 407 chloride concentrations at UK urban and rural sites. J. Environ. Monit. 8, 691-699.
- 408 Adachi, K., Tainosho, Y., 2004. Characterization of heavy metal particles embedded in tire dust. Environ. Int. 30, 409 1009-1017.
- 410 Amato, F., Cassee, F.R., Denier van der Gon, H.A.C., Gehrig, R., Gustafsson, M., Hafner, W., Harrison, R.M., Jozwicka,
- M., Kelly, F.J., Moreno, T., Prevot, A.S.H., Schaap, M., Sunyer, J., Querol, X., 2014. Urban air quality: The challenge of
 traffic non-exhaust emissions. Journal of Hazardous Materials 275, 31-36.
- 413 Araújo, D.F., Boaventura, G.R., Machado, W., Viers, J., Weiss, D., Patchineelam, S.R., Ruiz, I., Rodrigues, A.P.C.,
- 414 Babinski, M., Dantas, E., 2017. Tracing of anthropogenic zinc sources in coastal environments using stable isotope 415 composition. Chem. Geol. 449, 226-235.
- 416 Arnold, T., Schönbächler, M., Rehkämper, M., Dong, S., Zhao, F.J., Kirk, G.J.D., Coles, B.J., Weiss, D.J., 2010.
- 417 Measurement of zinc stable isotope ratios in biogeochemical matrices by double-spike MC-ICPMS and determination of
- 418 the isotope ratio pool available for plants from soil. Anal. Bioanal. Chem. 398, 3115-3125.
- 419 Beddows, D.C.S., Harrison, R.M., Green, D.C., Fuller, G.W., 2015. Receptor modelling of both particle composition and
- 420 size distribution from a background site in London, UK. Atmos. Chem. Phys. 15, 10107-10125.
- Bigi, A., Harrison, R.M., 2010. Analysis of the air pollution climate at a central urban background site. Atmos. Environ. 44,
 2004-2012.
- 423 Brewer, G.J., 2010. Copper toxicity in the general population. Clin. Neurophysiol. 121, 459-460.
- Chapman, J.B., Mason, T.F.D., Weiss, D.J., Coles, B.J., Wilkinson, J.J., 2006. Chemical Separation and Isotopic Variations
 of Cu and Zn From Five Geological Reference Materials. Geostand. Geoanal. Res. 30, 5-16.
- 426 Cloquet, C., Carignan, J., Libourel, G., 2006. Isotopic composition of Zn and Pb atmospheric depositions in an 427 urban/periurban area of northeastern France. Environ. Sci. Technol. 40, 6594-6600.
- 428 Cooper, J.A., Watson, J.G., 1980. Receptor Oriented Methods of Air Particulate Source Apportionment. J. Air Pollut.
 429 Control Assoc. 30, 1116-1125.
- 430 Costa, D.L., Dreher, K.L., 1997. Bioavailable transition metals in particulate matter mediate cardiopulmonary injury in 431 healthy and compromised animal models. Environ. Health Perspect. 105, 1053-1060.
- Charron, A., Harrison, R.M., 2005. Fine (PM2.5) and coarse (PM2.5-10) particulate matter on a heavily trafficked London
 highway: Sources and processes. Environ. Sci. Technol. 39, 7768-7776.
- Chen, L.C., Lippmann, M., 2009. Effects of Metals within Ambient Air Particulate Matter (PM) on Human Health. Inhal.
 Toxicol. 21, 1-31.
- 436 Dong, S., Weiss, D.J., Strekopytov, S., Kreissig, K., Sun, Y., Baker, A.R., Formenti, P., 2013. Stable isotope ratio
- measurements of Cu and Zn in mineral dust (bulk and size fractions) from the Taklimakan Desert and the Sahel and inaerosols from the eastern tropical North Atlantic Ocean. Talanta 114, 103-109.
- Flament, P., Mattielli, N., Aimoz, L., Choel, M., Deboudt, K., de Jong, J., Rimetz-Planchon, J., Weis, D., 2008. Iron
 isotopic fractionation in industrial emissions and urban aerosols. Chemosphere 73, 1793-1798.
- 441 Fry, C., Jarvis, K.E. and Parry, S.J., 2005. Diagnostic tool for source apportionment of heavy metals around road systems.
- 442 Environment Agency Report.
- 443 Gamble, J.F., Lewis, R.J., 1996. Health and respirable particulate (PM(10)) air pollution: A causal or statistical association?
- 444 Environ. Health Perspect. 104, 838-850.

- 445 Garg, B.D., Cadle, S.H., Mulawa, P.A., Groblicki, P.J., Laroo, C., Parr, G.A., 2000. Brake wear particulate matter 446 emissions. Environ. Sci. Technol. 34, 4463-4469.
- 447 Geagea, M.L., Stille, P., Gauthier-Lafaye, F., Millet, M., 2008. Tracing of industrial aerosol sources in an urban 448 environment using Pb, Sr, and Nd isotopes. Environ. Sci. Technol. 42, 692-698.
- 449 Gietl, J.K., Klemm, O., 2009. Analysis of Traffic and Meteorology on Airborne Particulate Matter in Münster, Northwest
- 450 Germany. J. Air Waste Manage. Assoc. 59, 809-818.
- 451 Gietl, J.K., Lawrence, R., Thorpe, A.J., Harrison, R.M., 2010. Identification of brake wear particles and derivation of a
- 452 quantitative tracer for brake dust at a major road. Atmos. Environ. 44, 141-146.
- 453 Gioia, S., Weiss, D., Coles, B., Arnold, T., Babinski, M., 2008. Accurate and Precise Zinc Isotope Ratio Measurements in
- 454 Urban Aerosols. Anal. Chem. 80, 9776-9780.
- 455 Grigoratos, T., Martini, G., 2015. Brake wear particle emissions: a review. Environ Sci Pollut Res 22, 2491-2504.
- Harrison, R.M., Deacon, A.R., Jones, M.R., Appleby, R.S., 1997. Sources and processes affecting concentrations of PM10
 and PM2.5 particulate matter in Birmingham (UK). Atmos. Environ. 31, 4103-4117.
- and 1 M2.5 particulate matter in Brinningham (OK). Atmos. Environ. 51, 4105-4117.
- Hjortenkrans, D.S.T., Bergbäck, B.G., Häggerud, A.V., 2007. Metal Emissions from Brake Linings and Tires: Case Studies
 of Stockholm, Sweden 1995/1998 and 2005. Environ. Sci. Technol. 41, 5224-5230.
- 460 Hüner, R., Melcher, B., Milczarek, R., Kienleitner, H., 2001. Solid lubricants with a tin sulphide and carbon base. Google461 Patents.
- 462 Iijima, A., Sato, K., Yano, K., Tago, H., Kato, M., Kimura, H., Furuta, N., 2007. Particle size and composition distribution
- analysis of automotive brake abrasion dusts for the evaluation of antimony sources of airborne particulate matter. Atmos.Environ. 41, 4908-4919.
- 465 Jarup, L., 2003. Hazards of heavy metal contamination. British Medical Bulletin 68, 167-182.
- John, S.G., Park, J.G., Zhan, Z.T., Boyle, E.A., 2007. The isotopic composition of some common forms of anthropogenic
 zinc. Chem. Geol. 245, 61-69.
- Laden, F., Neas, L.M., Dockery, D.W., Schwartz, J., 2000. Association of fine particulate matter from different sources
 with daily mortality in six US cities. Environ. Health Perspect. 108, 941-947.
- 470 Lin, L., Lee, M.L., Eatough, D.J., 2010. Review of Recent Advances in Detection of Organic Markers in Fine Particulate
- 471 Matter and Their Use for Source Apportionment. J. Air Waste Manag. Assoc. 60, 3-25.
- Lough, G.C., Schauer, J.J., Park, J.S., Shafer, M.M., Deminter, J.T., Weinstein, J.P., 2005. Emissions of metals associated
 with motor vehicle roadways. Environ. Sci. Technol. 39, 826-836.
- 474 Majestic, B.J., Anbar, A.D., Herckes, P., 2009. Stable Isotopes as a Tool to Apportion Atmospheric Iron. Environ. Sci.
 475 Technol. 43, 4327-4333.
- 476 Manoli, E., Voutsa, D., Samara, C., 2002. Chemical characterization and source identification/apportionment of fine and
 477 coarse air particles in Thessaloniki, Greece. Atmos. Environ. 36, 949-961.
- 478 Mattielli, N., Petit, J.C.J., Deboudt, K., Flament, P., Perdrix, E., Taillez, A., Rimetz-Planchon, J., Weis, D., 2009. Zn
- isotope study of atmospheric emissions and dry depositions within a 5 km radius of a Pb-Zn refinery. Atmos. Environ. 43,1265-1272.
- 481 McKenzie, E.R., Money, J.E., Green, P.G., Young, T.M., 2009. Metals associated with stormwater-relevant brake and tire 482 samples. Sci. Total Environ. 407, 5855-5860.
- 483 Moeller, K., Schoenberg, R., Pedersen, R.B., Weiss, D., Dong, S., 2012. Calibration of the New Certified Reference
- 484 Materials ERM-AE633 and ERM-AE647 for Copper and IRMM-3702 for Zinc Isotope Amount Ratio Determinations.

- 485 Geostand. Geoanal. Research 36, 177-199.
- 486 Monna, F., Lancelot, J., Croudace, I.W., Clundy, A.B., Lewis, J.T., 1997. Pb isotopic composition of airborne particulate
- 487 material from France and the southern United Kingdom: implications for Pb pollution sources in urban areas. Environ. Sci.
- 488 Technol. 31, 2277-2286.
- 489 Moreno, T., Martins, V., Querol, X., Jones, T., BéruBé, K., Minguillón, M.C., Amato, F., Capdevila, M., de Miguel, E.,
- 490 Centelles, S., Gibbons, W., 2015. A new look at inhalable metalliferous airborne particles on rail subway platforms. Sci.
- 491 Total Environ. 505, 367-375.
- 492 Moreno, T., Perez, N., Querol, X., Amato, F., Alastuey, A., Bhatia, R., Spiro, B., Hanvey, M., Gibbons, W., 2010.
- 493 Physicochemical variations in atmospheric aerosols recorded at sea onboard the Atlantic-Mediterranean 2008 Scholar Ship
- 494 cruise (Part II): Natural versus anthropogenic influences revealed by PM10 trace element geochemistry. Atmos. Environ.
 495 44, 2563-2576.
- 496 Noble, S.R., Horstwood, M.S.A., Davy, P., Pashley, V., Spiro, B., Smith, S., 2008. Evolving Pb isotope signatures of
- 497 London airborne particulate matter (PM10) constraints from on-filter and solution-mode MC-ICP-MS. J. Environ. Monit.
 498 10, 830-836.
- 499 Nriagu, J.O., Pacyna, J.M., 1988. Quantitative assessment of worldwide contamination of air, water, and soils by trace
 500 metals. Nature 333, 134-139.
- 501 Ochoa Gonzalez, R., Strekopytov, S., Amato, F., Querol, X., Reche, C., Weiss, D., 2016. New Insights from Zinc and
- 502 Copper Isotopic Compositions into the Sources of Atmospheric Particulate Matter from Two Major European Cities.
 503 Environ. Sci. Technol. 50, 9816-9824.
- Ochoa Gonzalez, R., Weiss, D., 2015. Zinc Isotope Variability in Three Coal-Fired Power Plants: A Predictive Model for
 Determining Isotopic Fractionation during Combustion. Environ. Sci. Technol. 49, 12560-12567.
- 506 Ozkaynak, H., Xue, J., Spengler, J., Wallace, L., Pellizzari, E., Jenkins, P., 1996. Personal exposure to airborne particles 507 and metals: Results from the particle team study in Riverside, California. J. Expo. Anal. Env. Epid. 6, 57-78.
- 508 Peel, K., Weiss, D., Chapman, J., Arnold, T., Coles, B., 2008. A simple combined sample-standard bracketing and
- inter-element correction procedure for accurate mass bias correction and precise Zn and Cu isotope ratio measurements. J.Anal. At. Spectrom. 23, 103-110.
- Peltier, R.E., Lippmann, M., 2010. Residual oil combustion: 2. Distributions of airborne nickel and vanadium within New
 York City. J. Expo. Sci. Env. Epid. 20, 342-350.
- 513 Pignalosa, G., Knochen, M., Cabrera, N., 2005. Determination of Zinc-Based Additives in Lubricating Oils by
- 514 Flow-Injection Analysis with Flame-AAS Detection Exploiting Injection with a Computer-Controlled Syringe. J. Autom.
- 515 Method. Manag. 1, 1-7.
- 516 Polissar, A.V., Hopke, P.K., Poirot, R.L., 2001. Atmospheric aerosol over Vermont: Chemical composition and sources.
- 517 Environ. Sci. Technol. 35, 4604-4621.
- Prasad, A.S., Bao, B., Beck, F.W.J., Kucuk, O., Sarkar, F.H., 2004. Antioxidant effect of zinc in humans. Free Radical Bio.
 Med. 37, 1182-1190.
- Querol, X., Alastuey, A., Rodriguez, S., Plana, F., Ruiz, C.R., Cots, N., Massague, G., Puig, O., 2001. PM10 and PM2.5
 source apportionment in the Barcelona Metropolitan area, Catalonia, Spain. Atmos. Environ. 35, 6407-6419.
- 522 Rodriguez, S., Querol, X., Alastuey, A., Viana, M.M., Alarcon, M., Mantilla, E., Ruiz, C.R., 2004. Comparative
- 523 PM10-PM2.5 source contribution study at rural, urban and industrial sites during PM episodes in Eastern Spain. Sci. Total
- 524 Environ. 328, 95-113.

- 525 Salma, I., Maenhaut, W., 2006. Changes in elemental composition and mass of atmospheric aerosol pollution between
- 526 1996 and 2002 in a Central European city. Environ. Pollution 143, 479-488.
- 527 Samek, L., Stegowski, Z., Furman, L., 2016. Preliminary PM2.5 and PM10 fractions source apportionment complemented
- 528 by statistical accuracy determination. Nukleonika 61, 75-83.
- 529 Sivry, Y., Riotte, J., Sonke, J.E., Audry, S., Schäfer, J., Viers, J., Blanc, G., Freydier, R., Dupré, B., 2008. Zn isotopes as
- tracers of anthropogenic pollution from Zn-ore smelters The Riou Mort–Lot River system. Chem. Geol. 255, 295-304.
- 531 Straffelini, G., Ciudin, R., Ciotti, A., Gialanella, S., 2015. Present knowledge and perspectives on the role of copper in
- 532 brake materials and related environmental issues: A critical assessment. Environ. Pollution 207, 211-219.
- Thorpe, A.J., Harrison, R.M., Boulter, P.G., McCrae, I.S., 2007. Estimation of particle resuspension source strength on a
 major London Road. Atm. Environ. 41, 8007-8020.
- Thorpe, A., Harrison, R.M., 2008. Sources and properties of non-exhaust particulate matter from road traffic: A review. Sci.
 Total Environ. 400, 270-282.
- 537 Thurston, G.D., Spengler, J.D., 1985. A quantitative assessment of source contributions to inhalable particulate matter
- 538 pollution in metropolitan Boston. Atm. Environ. (1967) 19, 9-25.
- Tong, S., von Schirnding, Y.E., Prapamontol, T., 2000. Environmental lead exposure: a public health problem of global
 dimensions. B. World Health Organ. 78, 1068-1077.
- 541 Visser, S., Slowik, J.G., Furger, M., Zotter, P., Bukowiecki, N., Dressler, R., Flechsig, U., Appel, K., Green, D.C., Tremper,
- 542 A.H., Young, D.E., Williams, P.I., Allan, J.D., Herndon, S.C., Williams, L.R., Mohr, C., Xu, L., Ng, N.L., Detournay, A.,
- 543 Barlow, J.F., Halios, C.H., Fleming, Z.L., Baltensperger, U., Prévôt, A.S.H., 2015. Kerb and urban increment of highly
- time-resolved trace elements in PM10, PM2.5 and PM1.0 winter aerosol in London during ClearfLo 2012. Atmos. Chem.
 Phys. 15, 2367-2386.
- 546 Wåhlin, P., Berkowicz, R., Palmgren, F., 2006. Characterisation of traffic-generated particulate matter in Copenhagen.
- 547 Atm. Environ. 40, 2151-2159.
- Walsh, C.T., Sandstead, H.H., Prasad, A.S., Newberne, P.M., Fraker, P.J., 1994. Zinc: health effects and research priorities
 for the 1990s. Environ. Health Persp. 102, 5-46.
- Weckwerth, G., 2001. Verification of traffic emitted aerosol components in the ambient air of Cologne (Germany). Atmos.
 Environ. 35, 5525-5536.
- 552 Wedepohl, K.H., 1995. The composition of the continental crust. Geochim. Cosmochim. Acta 59, 1217-1232.
- 553 Weiss, D.J., Kober, B., Dalgopolova, A., Gallagher, K., Spiro, B., Le Roux, G., Mason, T.F.D., Kylander, M., Coles, B.J.,
- 554 2004. Accurate and precise Pb isotope ratio measurements in environmental samples by MC-ICP-MS. Int. J. Mass 555 Spectrom. 232, 205-215.
- 556 Weiss, D.J., Rehkdmper, M., Schoenberg, R., McLaughlin, M., Kirby, J., Campbell, P.G.C., Arnold, T., Chapman, J., Peel,
- 557 K., Gioia S., 2008. Application of Nontraditional Stable-Isotope Systems to the Study of Sources and Fate of Metals in the
- 558 Environment. Environ. Sci. Technol. 42, 655–664.
- Widory, D., Liu, X.D., Dong, S.P., 2010. Isotopes as tracers of sources of lead and strontium in aerosols (TSP & PM2.5) in
 Beijing. Atmos. Environ. 44, 3679-3687.
- Wiederhold J.G., 2015. Metal Stable Isotope Signatures as Tracers in Environmental Geochemistry. Environ. Sci. Technol.
 49, 2606–2624.
- 563



Figure 1. Relationship between the enrichment factors of Sb (EF_{Sb}) and EF_{Fe} (R^2 =0.9960), EF_{Cu} (R^2 =0.9972), EF_{Zn} (R^2 =0.8686), EF_{Ba} (R^2 =0.9949) and EF_{Cr} (R^2 =0.9385) in the PM₁₀ collected in North Kensington (open symbols) and Marylebone Road (closed symbols).

CER CER



Figure 2. (a) 208 Pb/ 206 Pb versus 206 Pb/ 207 Pb plot of the PM₁₀ samples collected in Marylebone (MR site) and North Kensington (NK site) and possible sources in this study; (b) 208 Pb/ 206 Pb versus 206 Pb/ 207 Pb plot of the PM₁₀ measured in London since the phasing out of leaded gasoline between 1996 and 2001, as well as gasoline and industrial emissions (Noble et al., 2008 and Monna et al., 1997) and during 2010 (this study). 2SD are not displayed in the figure as they are small and fall within the marker dimensions.



Figure 3. Zinc (a) and Cu (b) isotopic composition in PM_{10} collected at the NK site (PM-NK London) and the MR site (PM-MR London) and local sources during this study (summer 2010). Average 2SD for Zn and Cu isotope measurements are 0.08 and 0.10‰, respectively. Also shown are previous published Zn and Cu isotope signatures in atmospheric $PM_{2.5-80}$ collected in South Kensington in London during 18 months between 2014 and 2015 (PM – SK – London), and PM_{10} collected in Barcelona during two sampling campaigns in 2012 and 2013 labelled as PM - Barcelona (Ochoa Gonzalez et al., 2016)^a with average 2SD for Zn and Cu isotope ratios of 0.08 and 0.11‰, respectively. The Zn isotope composition of gasoline (Gioia et al., 2008)^b and that from exhaust emissions represented by previously analyzed bus air filters (Cloquet et al., 2006)^c are also shown. The Cu isotope signatures in PM collected during the winter season in London (PM – SK – London) are highlighted as they shown heavier values, explained by an increasing fossil fuel combustion. Previously published Zn isotope ratios referring to the IRMM-3702 standard were recalculated to JMC 3-0749L Lyon using an isotopic offset of +0.32‰ for Zn (Moeller et al., 2012).



Figure 4. Relationship between Cu/Sb ratios and δ^{65} Cu_{NIST976} in PM₁₀ collected at the NK site and the MR site in this study (London PM₁₀), and previous publish data in PM_{2.5-80} from London (London PM_{2.5-80}) and in PM₁₀ from Barcelona (Ochoa Gonzalez et al., 2016).

1	Table 1. Concentrations of PM ₁₀	, major and trace elements	in the PM_{10} samples	collected in Marylebone	(MR) and North Kensington (NK),
---	---	----------------------------	--------------------------	-------------------------	---------------------------------

2 and Cu/Sb ratios.

						(Concentration	s					
Sample	Reference	PM_{10}	Fe	Al	Cu	Zn	Sb	Ba	Pb	Cr	Ni	V	Cu/Sb
		(µg m ⁻³)	(ng m ⁻³)										
NK - PM ₁₀	NK1	12	79	23	3.1	2.9	0.5	2.0	1.3	0.2	0.5	0.7	6.2
	NK2	12	46	14	2.3	1.4	0.3	1.1	0.6	0.3	0.4	0.5	7.7
	NK3	10	66	13	3.1	2.7	0.5	1.7	1.3	0.2	0.2	0.3	6.2
	NK4	17	58	12	2.5	1.8	0.3	1.6	0.8	0.4	0.3	0.4	8.3
	NK5	15	46	6.1	1.9	1.4	0.3	1.2	0.7	0.1	0.2	0.2	6.3
	NK6	17	89	22	3.5	3.1	0.5	2.0	1.1	0.3	0.4	0.6	7.0
	NK7	15	68	8.2	3.6	2.3	0.5	1.9	0.9	0.3	0.2	0.2	7.2
	NK8	12	58	8.6	2.5	2.2	0.4	1.5	1.2	0.3	0.2	0.1	6.3
	NK9	14	87	29	3.5	2.8	0.6	1.8	1.4	0.4	0.2	0.1	5.8
	$Average \pm SD$	14±3	66±16	15±8	2.9±0.6	2.3±0.6	0.4±0.1	1.6±0.4	1.0±0.3	0.3±0.1	0.3±0.1	0.3±0.2	6.8±0.9
MR - PM ₁₀	MR1	37	2660	150	130	41	22	84	8.8	8.2	3.6	2.6	5.8
	MR2	28	550	20	26	8.0	4.3	18	1.1	1.7	0.7	0.6	6.1
	MR3	29	436	24	21	7.8	3.5	14	1.9	1.5	0.6	0.4	6.1
	MR4	38	481	25	24	8.4	3.7	15	1.4	1.9	0.9	0.4	6.4
	MR5	29	494	16	24	7.3	3.9	16	1.3	1.7	0.7	0.3	6.0
	MR6	41	641	35	33	11	5.2	20	1.8	2.5	1.0	0.8	6.3
	MR7	34	2300	89	110	35	19	72	6.8	7.8	2.7	1.5	6.0
	MR8	27	279	15	13	5.5	2.2	7.9	1.7	1.1	0.3	0.2	6.0
	MR9	18	212	17	9.4	3.7	1.6	5.7	1.2	0.9	0.3	0.1	5.9
	Average \pm SD	31±7	-	-) -	-	-	-	-	-	-	-	6.1±0.2

3 4

Table 2. Concentrations of major and trace elements in local sources close to Marylebone (MR) and North Kensington (NK), including
 Cu/Sb ratios. '<' means below detection limit; '-' means not calculated.

						Concentr	ations					
Sample	Reference	Fe	Al	Cu	Zn	Sb	Ba	Pb	Cr	Ni	V	Cu/Sb
		$(mg g^{-1})$	$(mg g^{-1})$	$(\mu g g^{-1})$	$(\mu g g^{-1})$	$(\mu g g^{-1})$	$(\mu g \ g^{-1})$	$(\mu g g^{-1})$	$(\mu g g^{-1})$	$(\mu g g^{-1})$	$(\mu g g^{-1})$	
Brake	FSB-1	6.8	9.5	6.1	20	0.2	1000	5.3	89	51	9.7	27
	FSB-2	5.2	7.1	4.4	17	0.1	1300	4.0	80	41	7.0	39
	HBP-1	89	5.5	25	46	0.8	310	38	110	28	15	30
Tire	R43/54-1	0.4	0.2	43	8100	<	12	9.9	<	<	<	-
	R43/54-2	0.3	0.3	41	13000	<	19	14	<	1.4	<	-
	R56/57-1	0.2	0.4	26	7400	<	2.1	9.1	<	<	<	-
Manhole cover	NKMHC-1	68	5.7	52	370	7.4	550	690	130	100	26	7.0
	NKMHC-2	25	2.8	20	130	3.6	1100	56	93	86	15	5.4
	MRMHC-1	52	5.2	170	530	17	120	210	2300	1700	76	9.9
	MRMHC-2	43	1.5	89	420	7.5	120	89	120	110	120	12
Road dust	NKRD-1	9.4	4.4	70	220	3.3	93	140	29	11	19	21
	NKRD-2	7.6	3.2	17	80	1.0	34	22	29	100	15	17
	NKRD-3	18	13	25	170	0.7	86	26	31	18	56	35
	MRRD-1	17	6.0	800	560	21	160	410	55	26	23	38
	MRRD-2	14	3.7	74	350	10	97	170	54	13	15	7.3
	MRRD-3	10	4.0	140	350	40	94	54	40	17	14	3.5
Road paint	NKRP-1	0.6	0.3	1.8	44	<	2.4	2.8	3.2	4.0	0.5	-
	NKRP-2	0.4	0.2	<	4.8	93	200	3100	840	3.4	0.7	-
	MRRP-1	0.7	0.3	4.3	14	0.5	6.1	3.8	1.1	2.4	0.7	8.6
	MRRP-2	1.3	2.4	2.6	14	<	5.4	2.9	1.3	6.6	1.9	-
Road tarmac	NKRS-1	8.2	4.9	7.1	40	0.4	64	11	16	15	28	18
surface	NKRS-2	7.3	4.4 ¥	5.2	34	0.3	100	8.8	20	19	21	17

Table 3. Enrichment factors (EFs) in local sources and the PM_{10} samples collected at Marylebone

11	(MR) and North	Kensington	(NK) sites.	'-' means not calculated.
----	----------------	------------	-------------	---------------------------

Sample	Reference	EFs										
Sample	Reference	Fe	Cu	Zn	Sb	Ba	Pb	Cr	Ni	V		
NK - PM ₁₀	NK1	8.7	740	190	5300	10	258	18	83	44		
	NK2	8.2	900	150	4700	9.0	199	47	110	54		
	NK3	12	1300	300	9000	15	458	41	68	31		
	NK4	13	1200	230	7100	16	317	67	100	45		
	NK5	19	1700	340	11000	22	485	26	110	55		
	NK6	10	850	210	5500	11	220	33	72	36		
	NK7	21	2300	430	15000	27	484	81	95	37		
	NK8	17	1600	380	13000	20	663	80	100	25		
	NK9	7.6	650	150	5200	7.3	225	34	28	7.0		
	Average±SD	13±5	1200 ± 600	300±200	8000±4000	15±7	400±200	50 ± 30	90±30	40±20		
$MR - PM_{10}$	MR1	45	4600	410	37000	65	270	120	100	25		
	MR2	69	7100	600	54000	100	245	190	150	44		
	MR3	46	4900	490	37000	66	362	140	110	22		
	MR4	48	5100	490	37000	69	247	170	150	25		
	MR5	78	8000	680	61000	110	366	240	180	31		
	MR6	46	5000	450	37000	66	232	160	120	31		
	MR7	65	7000	580	54000	93	348	190	130	25		
	MR8	47	4800	560	37000	62	518	160	93	19		
	MR9	31	3000	330	23000	39	334	120	64	13		
	Average±SD	50±20	6000±2000	500±200	40000±20000	70±30	330±90	170±40	130±40	26±9		
Brake	FSB-1	1.8	3.5	3.2	5.9	13	2.6	21	22	1.5		
	FSB-2	1.9	3.3	3.5	3.9	21	2.6	25	24	1.4		
	HBP-1	41	25	13	37	6.6	32	46	21	3.9		
Tire	R43/54-1	4.6	1200	60000	-	6.9	230	-	-	-		
	R43/54-2	2.6	780	69000	-	7.5	220	-	21	-		
	R56/57-1	1.5	400	31000	-	0.7	110	-	-	-		
Manhole	NKMHC-1	30	49	97	330	11	550	50	75	6.6		
cover	NKMHC-2	23	38	72	320	44	92	74	130	8.1		
	MRMHC-1	25	170	150	800	2.7	180	970	1400	21		
	MRMHC-2	73	330	420	1300	9.1	280	190	310	120		
Road dust	NKRD-1	5.3	85	73	190	2.4	140	14	10	6.3		
	NKRD-2	6.0	29	37	79	1.2	32	20	131	6.8		
	NKRD-3	3.6	10	19	14	0.8	9.0	5.2	5.8	6.4		
	MRRD-1	7	730	140	880	3.2	300	21	18	5.7		
	MRRD-2	9.5	110	140	680	3.0	200	32	14	5.9		
	MRRD-3	6.3	190	130	2500	2.7	62	22	17	5.3		
Road paint	NKRP-1	4.6	28	190	-	0.8	37	21	48	2.2		
	NKRP-2	5.1	-	39	130000	120	76000	10000	76	5.4		
	MRRP-1	7.0	88	80	450	2.7	67	9.6	39	3.9		
	MRRP-2	1.4	5.8	8.5	-	0.3	5.5	1.2	11	1.1		
Road	NKRS-1	4.2	7.9	12	22	1.5	11	7.3	13	8.3		
surface	NKRS-2	4.2	6.4	12	18	2.8	9.2	10	18	7.1		

12 **Table 4.** Lead isotope ratios of PM_{10} samples collected at Marylebone Road (MR) and North 13 Kensington (NK) and sources analyzed during this study. Each sample has been measured 3 or 4 14 times to calculate the 2SD.

		Pb isotope ratios								
Sample	Sample ID	²⁰⁶ Pb/ ²⁰⁴ Pb	2SD	²⁰⁶ Pb/ ²⁰⁷ Pb	2SD	²⁰⁸ Pb/ ²⁰⁶ Pb	2SD			
NK - PM ₁₀	NK1	17.446	0.004	1.1218	0.0001	2.1382	0.0002			
	NK2	17.552	0.003	1.1277	0.0001	2.1327	0.0002			
	NK3	17.465	0.005	1.1229	0.0001	2.1374	0.0002			
	NK4	17.537	0.002	1.1270	0.0001	2.1352	0.0001			
	NK5	17.554	0.003	1.1277	0.0001	2.1321	0.0001			
	NK6	17.569	0.004	1.1287	0.0001	2.1315	0.0001			
	NK7	17.598	0.004	1.1304	0.0001	2.1291	0.0001			
	NK8	17.487	0.004	1.1239	0.0001	2.1355	0.0001			
	NK9	17.564	0.003	1.1282	0.0001	2.1288	0.0001			
MR - PM ₁₀	MR1	17.512	0.002	1.1256	0.0001	2.1351	0.0001			
	MR2	17.643	0.005	1.1330	0.0001	2.1279	0.0001			
	MR3	17.581	0.005	1.1294	0.0001	2.1313	0.0002			
	MR4	17.677	0.003	1.1349	0.0001	2.1261	0.0001			
	MR5	17.550	0.002	1.1275	0.0001	2.1327	0.0001			
	MR6	17.678	0.002	1.1349	0.0001	2.1257	0.0002			
	MR7	17.701	0.003	1.1364	0.0001	2.1246	0.0001			
	MR8	17.463	0.001	1.1227	0.0001	2.1368	0.0002			
	MR9	17.305	0.002	1.1137	0.0001	2.1476	0.0001			
Brake	FSB-1	19.569	0.003	1.2200	0.0001	1.9598	0.0001			
	FSB-2	20.199	0.002	1.2547	0.0001	1.9151	0.0001			
	HBP-1	18.176	0.004	1.1651	0.0001	2.1052	0.0001			
Tire	R30-1	17.937	0.003	1.1502	0.0001	2.1075	0.0002			
	R43/54-1	18.190	0.002	1.1660	0.0001	2.0973	0.0001			
	R43/54-2	18.181	0.002	1.1653	0.0001	2.0983	0.0001			
	R56/57-1	18.138	0.002	1.1613	0.0001	2.0995	0.0002			
Manhole cover	NKMHC-1	17.883	0.003	1.1450	0.0001	2.1262	0.0001			
	NKMHC-2	17.547	0.004	1.1281	0.0001	2.1355	0.0002			
	MRMHC-1	17.557	0.004	1.1284	0.0001	2.1332	0.0001			
	MRMHC-2	17.550	0.002	1.1280	0.0001	2.1335	0.0002			
Road dust	NKRD-1	17.438	0.002	1.1218	0.0001	2.1403	0.0001			
	NKRD-2	17.762	0.001	1.1405	0.0001	2.1222	0.0002			
	NKRD-3	17.728	0.002	1.1379	0.0001	2.1245	0.0001			
	MRRD-1	17.890	0.003	1.1417	0.0001	2.1242	0.0001			
	MRRD-2	17.777	0.002	1.1412	0.0001	2.1176	0.0002			
	MRRD-3	17.347	0.002	1.1163	0.0001	2.1446	0.0001			
Road paint	NKRP-1	17.930	0.002	1.1509	0.0001	2.0894	0.0001			
r	NKRP-2	18.301	0.002	1.1708	0.0001	2.0843	0.0001			
	MRRP-1	17.866	0.001	1.1471	0.0001	2.1012	0.0001			
	MRRP-2	18.135	0.001	1.1621	0.0001	2.0973	0.0002			
Road tarmac	NKRS-1	18.148	0.001	1.1633	0.0001	2.0953	0.0001			
	NKRS-2	18 252	0.001	1 1695	0.0001	2.0869	0.0001			

Isotopic signatures in atmospheric particulate matter suggest important contributions from recycled gasoline for lead and nonexhaust traffic sources for copper and zinc in aerosols in London, United Kingdom

Highlights

• Isotope ratios of Cu, Zn and Pb were determined in PM₁₀ collected at two road sites with contrasting traffic densities in central London, UK, and in potential sources.

• Despite the different traffic densities, no significant differences in the isotope signatures of Cu, Zn and Pb are found in PM₁₀, suggesting similar sources for each metal at both sites.

• Analysis of the isotopic composition of Cu, Zn and Pb in PM₁₀ and potential sources indicate important contributions from road dust resuspension, recycled leaded gasoline, tire and brake wear.