Promoting physical activity for disabled people who are ready to become physically active:

Jaarsma, Eva; Smith, Brett

DOI:

[10.1016/j.psychsport.2017.08.010](https://doi.org/10.1016/j.psychsport.2017.08.010)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Jaarsma, E & Smith, B 2017, 'Promoting physical activity for disabled people who are ready to become physically active: a systematic review', *Psychology of Sport and Exercise*.
<https://doi.org/10.1016/j.psychsport.2017.08.010>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Table 1 Study characteristics of the included studies

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Arbour-Nicitopoulos, Martin Ginis, & Latimer, 2009)	44 (30/14)	49.70 (12.71)	SCI	10 weeks	RCT	Questionnaire measuring: - intention - self-efficacy (coping, barrier) - PARA-SCI - frequency of action planning - coping planning	-	- Exercise equipment, - Written materials (action plans, PA pamphlet, guidelines and safety tips, log books) - Telephone counselling sessions (3 in total)	- Intention - Coping self-efficacy: General barriers self-efficacy, Facility barriers self-efficacy, Scheduling self-efficacy. - Short version of the PARA-SCI over 7 days. - Frequency of action planning - Coping planning	- LTPA (exp vs con): d=0.71; p<0.03 - Intention (over time): d=0.18; p<0.03 - General barriers self-efficacy (over time): d=0.60; p<0.01 - Coping self-efficacy: Facility barriers (exp vs con): d=-0.65; p<0.04 General barriers (exp vs con): d=0.83; p<0.01 Scheduling (exp vs con): d=0.87; p<0.01 - Scheduling in week 1 predicting LTPA in week 5: β =0.31; p<0.03	- LTPA: Time effect was not significant, nor was the time x condition interaction (ps >0.60). - Intentions: No significant main effect for condition or time x condition interaction (ps <0.10).
(Arbour-Nicitopoulos, Tomasone, Latimer-Cheung, & Martin Ginis, 2014)	65 (37/27)	50.42 (12.78)	SCI	6 months	Cohort	Self-report LTPA Questionnaire for People with SCI	HAPA	Telephone counselling sessions (14 in total)	- Intention - Self-report LTPA Questionnaire for People with SCI for 7 days	- Intentions for regular LTPA at start and after 6 months remained high: ds=0.02-0.20; p = 0.44	- Increase in clients being regularly active at baseline (35%) versus 4 months (48%; p = 0.13) and 6 months (52%, p = 0.09)

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Cardinal, Kosma, & McCubbin, 2004)	322(122/200)	52.5 (13.9)	AMP, CP, joint and connective tissue disease, MS, MD, PP, SB, SCI, CVA, unspecified	-	Cross-sectional	Questionnaire measuring: - Stage of change - Process of change (behaviour and cognitive) - Decision balance - Exercise barriers)	TTM	-	- Stages of change - Behavioural processes of change - Cognitive processes of change - Self-efficacy - Decision balance: Pros for exercise Cons for exercise - Exercise barriers	- Behavioural processes of change: $d=1.63$; $p<0.001$ - Self-efficacy: $d=1.31$; $p<0.001$ - Pros for exercise: $d=0.97$; $p<0.001$ - Cons for exercise: $d=0.87$; $p<0.001$ - Cognitive processes of change: $d=0.87$; $p<0.001$ - Exercise barriers: $d=0.84$; $p<0.001$ Predicting stages of change: - Maintenance (91.3%) - Precontemplation (73.8%) - Contemplation (48.3%) - Preparation (23.8%) - Action (5.3%)	
(Gernigon, Pereira Dias, Riou, Briki, & Ninot, 2015)	18(13/5)	36.0 (16.1)	SCI	16 weeks	Cross-Sectional	- Approach and Avoidance Questionnaire for Sport and Physical Education - Physical Self-Perception Profile	-	-	- Approach and Avoidance Questionnaire for Sport and Physical Education - Physical Self-Perception Profile	Participants vs non-participants: - Mastery avoidance goals: $d=1.06$; $p<0.05$ - Physical self-worth: $d=1.53$; $p<0.01$	- global self-esteem: $d=1.06$; $p=0.07$ Participants vs non-participants: No significant differences were found for mastery-approach goals, performance-approach goals, performance-avoidance goals, physical condition, physical strength, body attractiveness, and sport competence ($p>0.05$).

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Jaarsma, Geertzen, de Jong, Dijkstra, & Dekker, 2014)	76 (30/46)	30.5 (9.7)	AMP, CP, SCI, VI, Les Autres, other Neuro	-	Cross-sectional	Self-constructed questionnaire on barriers and facilitators of sport	TPB, ICF	-	- Self-constructed questionnaire focusing on personal and environmental barriers and facilitators of sport	Wheelchair vs ambulant: - Experienced barriers: d= 0.53; p=0.023 - Lack of sports facilities: p<0.01 - Sports facilities not adjusted: p<0.01 - Health and physical fitness to maintain active: p=0.015 Initiate vs maintain: - Health and physical fitness: d=0.59; p=0.012 - Competition: d=0.62; p=0.009	Wheelchair vs ambulant: - Dependency on others: p=0.055 - transport: p=0.055
(Jaarsma, Dekker, Koopmans, Dijkstra, & Geertzen, 2014)	648 (311/337)	49.1 (18)	VI	-	Cross-sectional	Self-constructed questionnaire on barriers and facilitators of sport	ICF	-	- Self-constructed questionnaire focusing on personal and environmental barriers and facilitators of sport	- Higher education: d=0.24; p=0.039 Disability (experienced as barrier): d=-0.31; p=0.03 - Costs: d=-0.73; p<0.001 - Lack of peers/buddies: d=-1.05; p<0.001 - Use of computer software: d=0.35; p=0.003	- Using a white cane: d=0.029; p=0.801 - Having a guide dog: d=0.23; p=0.170 - Age: d=0.0027; p=0.368 - Gender: d=0.10; 0.361

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Keegan, Chan, Ditchman, & Chiu, 2012)	126 (83/43)	43.5 (13.3) [19-76]	SCI	-	Cross-sectional	<ul style="list-style-type: none"> - Self-Report Functional Independence Measure basic Activities of Daily Living subscale. - TPB physical activity and exercise questionnaire - Friend Support for Exercise Habits Scale/ Family Support for Exercise Habits Scale - Outcome Expectations for Exercise Scale - Barriers to Health Promoting Activities for Disabled Persons Scale - SCI Exercise Self-Efficacy Scale - Revised Planning for Exercise Scale - International Physical Activity Questionnaire - Physical Activity Stages of Change Instrument 	Pender's Health promotion model (SCT & TPB)	-	<ul style="list-style-type: none"> - Pre-injury physical activity - Self-Report Functional Independence Measure - Normative and control beliefs about physical activity and exercise scale. - Friend Support for Exercise Habits Scale/ Family Support for Exercise Habits Scale - Outcome Expectations for Exercise Scale - Barriers to Health Promoting Activities for Disabled Persons Scale - SCI Exercise Self-Efficacy Scale - Revised Planning for Exercise Scale - International Physical Activity Questionnaire - Physical Activity Stages of Change Instrument 	<ul style="list-style-type: none"> - Physical activity/exercise participation: Pre-injury PA: $\beta=0.17$; $p<0.05$ - Severity of SCI: $\beta=0.20$; $p<0.05$ - Commitment to action plan: $\beta=0.41$; $p<0.01$ - Commitment to action plan: Friend/family support: $\beta=0.40$; $p<0.01$ - Perceived benefits: $\beta=0.17$; $p<0.05$ - Perceived self-efficacy: $\beta=0.35$; $p<0.05$ 	-

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Kennedy, Taylor, & Hindson, 2006)	35 (30/5)	31.91 (10.60) [18-61]	SCI	6 weeks	Cohort	<ul style="list-style-type: none"> - Life Satisfaction Questionnaire - Hospital Anxiety and Depression Scale - Perceived Manageability - Generalised Self-Efficacy Scale - Overall gains (via interviews) 	-	Course with multiple or single activity program (1 week)	<ul style="list-style-type: none"> - Life Satisfaction questionnaire - Hospital Anxiety and Depression Scale - Needs Assessment Checklist - Generalised Self-Efficacy Scale - Overall gains of intervention: <ol style="list-style-type: none"> 1) Why would you recommend Back-Up to other people with SCI? 2) Describe how being involved in Back-Up has had a positive effect on your rehabilitation. 3) What do you feel you have gained from Back-Up? 	<ul style="list-style-type: none"> - Life Satisfaction General: $d=0.88$; $p=0.016$ - Leisure Satisfaction: $d=1.021$; $p=0.007$ - Anxiety: $d=0.50$; $p<0.01$ - GSES: $d=0.93$; $p=0.012$ 	<i>Quantitative results:</i> <ul style="list-style-type: none"> - Perceived Manageability $d=0.88$ <i>Qualitative results:</i> <ul style="list-style-type: none"> - Overall gains of intervention: <ol style="list-style-type: none"> 1) 'Meeting people and making friends.' 'Perception of possibilities and capabilities.' 2) 'Self-confidence and sense of achievement.' 'Skills and knowledge' 3) 'Self-confidence and sense of achievement.' 'Meeting people and making friends.'
(Kosma, Cardinal, & McCubbin, 2004)	151 (34/117)	37.9 (8.8)	SCI, CP, MS	-	Cross-sectional	Questionnaire measuring: <ul style="list-style-type: none"> - Stage of change - Process of change - Self-efficacy - Decision balance 	TTM	-	<ul style="list-style-type: none"> - Stages of change - Self-efficacy - Decision balance 	Most important stages of change predictors: <ul style="list-style-type: none"> - Function 1: Behavioural changes: $r=0.94$ Cognitive changes: $r=0.71$ Self-efficacy: $r=0.57$ Decision balance: $r=0.36$ - Function 2: Cognitive processes of change: $r=0.58$ 	-

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Latimer, Martin Ginis, & Arbour, 2006)	54 (26/28)	40.61 (10.89)	SCI	8 weeks	RCT	Questionnaire measuring: - intention - PARA-SCI - Perception of control	TPB	- Exercise equipment - Written materials (action plans, PA pamphlet, guidelines and safety tips, log books)	- Intention - Physical Activity Recall Assessment for Individuals with SCI - PBC - Scheduling self-efficacy - Barrier self-efficacy	- Minutes PA (exp vs con): $d=0.52$, $p=0.04$ - Intention as predictor for PA duration (only exp): $\beta=0.68$, $p=0.05$ - Intention as predictor for PA frequency (only exp): $\beta=0.76$, $p=0.05$ - Treatment effect on intention: $d=0.73$, $p=0.04$ - Treatment effect on scheduling self-efficacy: $d=0.71$, $p=0.04$	- Number of days participants engaged in ≥ 30 min of physical activity. - Intention was no predictor for PA duration and frequency $ps > 0.84$ - PBC not significant for experimental and control group. - No significant treatment effects for the PBC or barrier self-efficacy measures ($ps > 0.05$).
(Latimer, Martin Ginis, & Craven, 2004)	124 (86/38)	43.45* (16.21)*	SCI	-	Cross-sectional	Questionnaire measuring TPB constructs (attitude, subjective norm, PBC) - Godin Leisure Time Exercise Questionnaire	TPB	-	- Self-constructed questionnaire assessing TPB construct: Attitude Subjective norm PBC Intentions - Godin Leisure Time Exercise Questionnaire	- PBC as predictor of intention (only for TP): $\beta=0.59$, $p<0.01$ - PBC predictor of exercise in moderate intensity exercise model (only for TP): $\beta=0.33$, $p=0.03$	- For individuals with paraplegia, none of the TPB constructs predicted intentions. - For TP intentions were not a significant predictor of exercise behaviour at any intensity. - For individuals with paraplegia, the TPB constructs did not predict exercise behaviour at any intensity.

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Martin Ginis et al., 2013)	238 (179/57, 2 missing)	44.14 (12.74)	SCI	-	Cross-sectional	PARA-SCI, Questionnaire measuring LTPA outcome expectancies, self- efficacy (task, maintenance, recovery, scheduling, goal setting), intentions, planning and action control	TPB, HAPA	-	Motivational phase constructs: - LTPA Outcome Expectancies - Task self-efficacy - Intentions Volitional phase constructs: - Planning - Maintenance Recovery, Scheduling, Goal Setting self-efficacy - Action Control	- Actors reported more min/day of moderate and heavy intensity LTPA than intenders and non-intenders, ps < 0.001. - Actors scored significantly higher than both intenders and non-intenders on all constructs, ps ≤ 0.01. - Intenders scored significantly higher than non-intenders on all constructs, ps ≤ 0.01.	Intenders and non-intenders did not differ on min/day of moderate and heavy intensity LTPA.

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Molton, Jensen, Nielson, Cardenas, & Ehde, 2008)	130 (93/37)	45 (14.4) [18-82]	SCI	-	Cross-sectional	<ul style="list-style-type: none"> - Numerical Rating Scale for pain intensity - Multi-dimensional Pain Readiness to Change Questionnaire - Motivational Model of Pain Self-Management 	Motivational Model of Pain Self-Management	-	<ul style="list-style-type: none"> - Numerical Rating Scale - Multidimensional Pain Readiness to Change Questionnaire - Motivational Model of Pain Self-Management 	<ul style="list-style-type: none"> - Effect Perceived importance on exercise behaviour: $\beta = 0.48$, $p < 0.001$ - Effect Perceived importance on Readiness to exercise: $\beta = 0.56$, $p < 0.001$ - Readiness to exercise on exercise behaviour: $\beta = 0.54$, $p < 0.001$ - Self-efficacy on exercise behaviour: $\beta = 0.41$, $p < 0.001$ - Self-efficacy on readiness to exercise: $\beta = 0.56$, $p < 0.001$ - Readiness to exercise on exercise behaviour: $\beta = 0.23$, $p = 0.01$ 	<ul style="list-style-type: none"> - Effect of perceived importance on exercise behaviour (including readiness to exercise) $\beta = 0.18$, $p = 0.04$ (after α correction)

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Pelletier, Latimer-Cheung, Warburton, & Hicks, 2014)	17 (13/4)	42.1 (10.6)	SCI	16 weeks	Cohort	- Exercise beliefs questionnaire	SCT	Telephone counselling sessions (5 in total)	- (Self-reported) Adherence to twice-weekly exercise program for 16-week period - Exercise belief questionnaire	-	- No difference in attendance rates between inpatient compared to outpatient counselling groups. - Effect size: Inpatient compared to outpatient counselling groups: $d=0.63$, $p=0.22$ Outpatient referral only compared to referral plus counselling: $d=0.79$, $p=0.22$. - Exercise belief questionnaire: No significant differences in constructs between groups. $p>0.05$. No significant correlation between adherence and constructs. $p=0.21$

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Perrier, Sweet, Strachan, & Latimer-Cheung, 2012)	201 (119/82)	44.0 (12.8)	SCI, AMP, other (stroke, polio)	2 weeks	Pre-post testing	Questionnaire measuring: - Athletic identity (AIMS) - Outcome expectancies - Risk perceptions - Self-efficacy - Intentions - Planning	HAPA	-	<ul style="list-style-type: none"> - Athletic Identity Measurement Scale - Instrumental expectancies - Affective expectancies - Negative expectancies - Health risk perceptions - Task self-efficacy - Intentions - Scheduling self-efficacy - Barrier self-efficacy - Action planning - Coping planning - Recovery self-efficacy - modified version of the 7 day short form Leisure Time Physical Activity Questionnaire for People with Spinal Cord Injury 	<ul style="list-style-type: none"> - Negative outcome expectancies negatively predicts intention on sport participation: $\beta=-0.24$, $p=0.001$ - Higher task self-efficacy, decrease in planning: $\beta=0.22$, $p=0.015$ - Indirect effect task self-efficacy on planning through intention: $\beta=0.13$, $p=0.002$ - Maintenance self-efficacy on sport participation: $\beta=0.48$, $p=0.003$ - Indirect effect planning on sport participation through self-efficacy: $\beta=0.33$, $p=0.002$ 	<ul style="list-style-type: none"> - Health risks did not predict intentions to participate in sport: $\beta=-0.09$, $p=0.17$ - Relationship between planning and sport participation: $\beta=0.052$, $p=0.65$ - recovery self-efficacy and sport: $\beta=0.19$, $p=0.11$

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration intervention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Perrier, Shirazipour, & Latimer-Cheung, 2015)	201 (119/82)	44.27 (12.08)	Acquired physical disabilities, such as SCI	-	Cross-sectional	Questionnaire measuring: - Intention - Outcome expectancies - Risk perceptions - Self efficacy - Planning	HAPA	-	<ul style="list-style-type: none"> - Staging sport - Outcome expectancies - Risk perceptions - Task self-efficacy - Intentions - Maintenance self-efficacy - Planning - Recovery self-efficacy 	Task self-efficacy, $p < 0.001$: $d(NI, IN) = 0.25$ $d(NI, A) = 1.34$ $d(IN, A) = 1.15$ Intentions, $p < 0.001$: $d(NI, IN) = 0.57$ $d(NI, A) = 2.29$ $d(IN, A) = 1.32$ Scheduling self-efficacy, $p < 0.001$: $d(NI, IN) = 0.51$ $d(NI, A) = 1.49$ $d(IN, A) = 0.76$ Affective outcome expectancies, $p < 0.001$: $d(NI, IN) = 1.10$ $d(NI, A) = 1.20$ $d(IN, A) = 0.0080$ Recovery self-efficacy, $p < 0.001$: $d(NI, IN) = 0.76$ $d(NI, A) = 1.46$ $d(IN, A) = 0.17$ Barrier self-efficacy, $p < 0.001$: $d(NI, IN) = 0.95$ $d(NI, A) = 1.58$ $d(IN, A) = 0.78$ Action plans, $p < 0.001$: $d(NI, IN) = 0.92$ $d(NI, A) = 2.56$ $d(IN, A) = 1.06$	Bonferroni-adjusted $\alpha = 0.006$: - Instrumental outcome expectancies, $p = 0.13$: $d(NI, IN) = 0.43$ $d(NI, A) = 0.46$ $d(IN, A) = 0.0061$ - Negative outcome expectancies, $p = 0.006$: $d(NI, IN) = 0.15$ $d(NI, A) = 0.44$ $d(IN, A) = 0.25$ - Risk perceptions, $p = 0.21$: $d(NI, IN) = 0.18$ $d(NI, A) = 0.42$ $d(IN, A) = 0.68$ - Coping plans, $p = 0.028$ $d(NI, IN) = 0.28$ $d(NI, A) = 0.71$ $d(IN, A) = 0.38$

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Saebu & Sorensen, 2011)	327 (149/178)	24.15 (3.88)	CP, SB, SCI, MD, VI	-	Cross-sectional	- International Physical Activity Questionnaire - Functioning and Disability - Barriers to exercise (environmental factors) - Exercise Self-Regulation Questionnaire	SDT, ICF	-	- International Physical Activity Questionnaire (Short Form) - Self construction questionnaire about function and disability - Self constructed questionnaire about environmental factors - Exercise Self-Regulation Questionnaire - Exercise self-schema - Medical Outcome Study Short Form 12	- Acquired disability $\beta=0.12$, $p<0.05$ - Low need for personal activity equipment $\beta=0.15$, $p<0.01$ - Employed $\beta=0.16$ - Available local activities $\beta=0.11$, $p<0.05$ - High physical component summary (PCS) $\beta=0.12$, $p<0.05$ - Exerciser schematics $\beta=0.27$, $p<0.01$ - High intrinsic motivation $\beta=0.14$, $p<0.01$	- No need for personal aids - High education - Need for more than 3 hours of daily personal care - Functional personal activity equipment - Adapted facilities at site - High level of information of activities - Age - Gender

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Sweet, Martin Ginis, & Latimer-Cheung, 2012)	541 (411/130)	47.6 (13.4)	SCI	18 months	Observational study	- TPB to predict LTPA in persons with SCI - PARA-SCI	TPB	-	- Theory of Planned Behaviour Measures - Physical Activity Recall Assessment for People with SCI	Significant results compared to inactive group: - Intention: Increaser: d=0.16, p=0.02 Decreaser: d= 0.29, p<0.01 Stable active: d=0.43, p<0.01 - Less severe injuries: Decreaser: d=0.27, p=0.01 Stable active: d=0.20, p=0.05 - Stable active Younger: d=0.016, p=0.05 - Fewer years post injury: d=0.03, p<0.01	Non-significant results (p>0.05) compared to inactive group: - Age: Increaser, Decreaser - Gender: Increaser, Decreaser, Stable active - Years post injury: Increaser, Decreaser - Injury severity: Increaser - Subjective norms: Increaser, Decreaser, Stable active - Attitude: Increaser, Decreaser, Stable active - PBC: Increaser, Decreaser, Stable active
(Thomas et al., 2011)**	21(10/11)	43.6 (14.2)	SCI	9 months	RCT	- TTM questionnaire - Borg Rating of Perceived Exertion scale for physical activity intensity	TTM	- Telephone counselling sessions (7 in total), - Written materials (brochures, tailored exercise instructions), - Exercise materials (dvd)	- Stages of Change (TTM) - Self-reported activity log (number of days, total minutes per day, types of activity, and intensity of each activity for one full week per month for each of the 3 months) - Borg Rating of Perceived Exertion scale for physical activity intensity	- CON: PA increase at T2, T3 and T4 compared to T1 (p<0.05, p<0.01, p<0.05 respectively). - EXP: PA increase T3 and T4 compared to T1 (p<0.05 for both).	- EXP: no increase in PA at T2 compared to T1 (p>0.05). - No significant differences in improvement between groups

Authors	Sample size (M/F)	Age (SD) [range]	Disability group(s)	Duration inter-vention	Design	Assessment tool	Theory/ Model	Intervention delivery mode	Outcome measures	Significant Results	Non-significant results
(Warms, Belza, Whitney, Mitchell, & Stiens, 2004)	16 (13/3)	43.2 (11.3) [24-68]	SCI	6 weeks	Cohort	<ul style="list-style-type: none"> - Stage of Readiness for Change in Exercise Behaviour - Barriers to Health Activity Among Disabled Persons - Self-rated Abilities for Health Practices Scale - Self-rated Health Scale 	-	<ul style="list-style-type: none"> - Face to face counselling (motivational interviewing, goal setting, action plan), - Telephone counselling sessions (4 in total), - Written materials (pamphlets, tailored physical activity information) 	<ul style="list-style-type: none"> - Accelerometer and a physical activity record for 4 days. - Stages of change (Trans Theoretical Model) - Barriers to Health Activity Among Disabled Persons Scale. - Self-rated Abilities for Health Practices Scale. - Self-rated Health Scale. - CES-D - Isometric strength of elbow and shoulder flexors/extensors was measured bilaterally using a handheld dynamometer 	<ul style="list-style-type: none"> - Motivational barriers: d=1.5, p=0.01 - Exercise self-efficacy: d=-1.1, p=0.05 - Self-rated Health: d=-1.1, p=0.04 - Muscle Strength: d=-3.6, p<0.001 	<ul style="list-style-type: none"> - Activity score: d=-0.68, p = 0.32 - Total barriers score: d=1.1, p=0.06 - external barriers: d= 0.50, p=0.37 - self-rated abilities for health practices: d=-0.45, p=0.39 - Depression: d=0.64, p=0.24

* Pooled mean and standard deviation

** Effect sizes could not be calculated based on the results provided in the study.

A = Actors, AMP = Amputation, CES-D = Centre for Epidemiologic Studies Depression Scale, CON = control group, CP = Cerebral Palsy, CVA = Cerebral Vascular Accident (which includes both stroke and traumatic brain injury), EXP = Experimental group, F = Female, HAPA = Health Action Process Approach, I = Interview, ICF = International Model of Classification, Functioning and Disability, IN = Intenders, LTPA = Leisure Time Physical Activity, M = Male, MD = Muscular Disease, MS = Multiple Sclerosis, NI = Non-Intenders, PA = Physical Activity, PARA-SCI = Physical Activity Recall Assessment for People with Spinal Cord Injuries, PBC = Perceived Behavioural Control, PP = Post-Polio, Q = Questionnaire, RCT = Randomised Controlled Trial, SB = Spina Bifida, SCI = Spinal Cord Injury, SCT = Social Cognitive Theory, SD = Standard Deviation, SDT = Self-Determination Theory, SOC = Stages of Change, TP = Tetraplegia, TPB = Theory of Planned Behaviour, TTM = Trans Theoretical Model, VI = Visual Impairment