Antibodies against collagen type II are not a general marker of acute arthritis onset

Manivel, Vivek Anand; van der Woude, D; Toes, Rene; Filer, Andrew; Raza, Karim; Rönnelid, Johan

DOI:

[10.1136/annrheumdis-2017-211974](https://doi.org/10.1136/annrheumdis-2017-211974)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Manivel, VA, van der Woude, D, Toes, R, Filer, A, Raza, K & Rönnelid, J 2017, 'Antibodies against collagen type II are not a general marker of acute arthritis onset', *Annals of the Rheumatic Diseases*.
<https://doi.org/10.1136/annrheumdis-2017-211974>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Antibodies against collagen type II are not a general marker of acute arthritis onset.

Vivek Anand Manivel¹, Diane van der Woude², Rene Toes², Andrew Filer^{3,4}, Karim Raza^{3,5}, Johan Rönnelid¹

¹Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden

²Department of Rheumatology, Leiden University Medical Center, Leiden, Netherlands

³Rheumatology Research Group, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

⁴University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

⁵Department of Rheumatology, Sandwell and West Birmingham Hospitals NHS Trust, Birmingham, UK

The fibrillar protein collagen type II (CII) is essentially confined to hyaline cartilage in diarthrodial joints. Antibodies against CII (anti-CII) were previously described in 3-27% of rheumatoid arthritis (RA) patients, and Kim et al. described anti-CII to be associated with elevated levels of CRP and ESR in a heterogeneous group of RA patients with 2-432 months of disease duration. [1] Contrary to anti-citrullinated protein antibodies, anti-CII are not detected before RA onset. [2] We have shown that anti-CII levels are highest at the time of RA diagnosis and thereafter decline, and that elevated anti-CII levels at diagnosis associate with elevated CRP, ESR, swollen joint count, disease activity score and radiological destruction at the time of diagnosis but not later, thus representing an acute onset RA phenotype. [3-5] It is plausible that production of pro-inflammatory cytokines by macrophages is stimulated by anti-CII-containing immune complexes (IC) in RA joints, as anti-CII producing B cells are detectable in synovial fluid but not in the circulation. [6, 7] Analogously, in rodents, collagen antibody-induced arthritis (CAIA) develops soon after injection of a defined blend of anti-CII antibodies. [8] Polymorphonuclear granulocytes (PMN) are central in CAIA pathogenesis, and we have recently shown that PMN induce chemokine production after stimulation with IC containing anti-CII from RA patients. [9]

In patients with very early synovitis with ≤ 3 months duration, anti-CII are not more common in patients developing RA than in patients developing other diagnoses during the following year. [10] Post-publication subgroup analysis showed that 3/11 (27%) of reactive arthritis and 2/7 (29%) of gouty arthritis patients had elevated anti-CII levels at the early synovitis stage (unpublished). As both reactive arthritis and gouty arthritis often have an acute arthritis onset, we hypothesized that high levels of anti-CII might not only be a marker of acute onset RA, but a marker of acute onset arthritis in general.

To investigate the hypothesis that high anti-CII levels might be a general marker of acute onset arthritides we collected first visit sera from 26 patients with gouty arthritis, 3 patients with pseudogout, 44 patients with reactive arthritis, and 33 patients with remitting seronegative symmetrical synovitis with pitting edema (RS3PE), diagnoses that often start with acute onset arthritis. Patient

characteristics are described in table 1. Antibodies against native human CII were measured with ELISA, and levels above the 95th percentile among controls were regarded as positive in agreement with our previous studies. [4-6] Serially followed internal laboratory controls showed that the analysis performance remained unaltered. [3, 4]

Eight of the 106 (7.5%) first visit non-RA patients were anti-CII positive, very similar to our previously published data on early RA (6.6-8.8%). [3, 4] There was no significant difference between any diagnostic groups (table 1). Only moderately elevated anti-CII levels were found; no non-RA patients exhibited very high anti-CII levels comparable to the high outlier group that we have described in early RA patients (figure 1). [4, 5]

We conclude that although high anti-CII levels are associated with acute onset in RA, this cannot be generalized to most patients with other acute onset arthritides.

Competing interests: None declared.

References:

- [1] Kim WU, Yoo WH, Park W, et al. IgG antibodies to type II collagen reflect inflammatory activity in patients with rheumatoid arthritis. *J Rheumatol* 2000;27:575-581.
- [2] Möttönen T, Hannonen P, Oka M, et al. Antibodies against native type II collagen do not precede the clinical onset of rheumatoid arthritis. *Arthritis Rheum* 1988;31:776-779.
- [3] Manivel VA, Mullazehi M, Padyukov L, et al. Anticollagen type II antibodies are associated with an acute onset rheumatoid arthritis phenotype and prognosticate lower degree of inflammation during 5 years follow-up. *Ann Rheum Dis* 2017.
- [4] Mullazehi M, Mathsson L, Lampa J, et al. High anti-collagen type-II antibody levels and induction of proinflammatory cytokines by anti-collagen antibody-containing immune complexes in vitro characterise a distinct rheumatoid arthritis phenotype associated with acute inflammation at the time of disease onset. *Ann Rheum Dis* 2007;66:537-541.
- [5] Mullazehi M, Wick MC, Klareskog L, et al. Anti-type II collagen antibodies are associated with early radiographic destruction in rheumatoid arthritis. *Arthritis Res Ther* 2012;14:R100.
- [6] Mullazehi M, Mathsson L, Lampa J, et al. Surface-bound anti-type II collagen-containing immune complexes induce production of tumor necrosis factor alpha, interleukin-1beta, and interleukin-8 from peripheral blood monocytes via Fc gamma receptor IIA: a potential pathophysiologic mechanism for humoral anti-type II collagen immunity in arthritis. *Arthritis Rheum* 2006;54:1759-1771.
- [7] Rönnelid J, Lysholm J, Engström-Laurent A, et al. Local anti-type II collagen antibody production in rheumatoid arthritis synovial fluid. Evidence for an HLA-DR4-restricted IgG response. *Arthritis Rheum* 1994;37:1023-1029.
- [8] Nandakumar KS, Holmdahl R. Collagen antibody induced arthritis. *Methods Mol Med* 2007;136:215-223.
- [9] Manivel VA, Sohrabian A, Rönnelid J. Granulocyte-augmented chemokine production induced by type II collagen containing immune complexes is

mediated via TLR4 in rheumatoid arthritis patients. *Eur J Immunol* 2016;46:2822-2834.

[10] Raza K, Mullazehi M, Salmon M, et al. Anti-collagen type II antibodies in patients with very early synovitis. *Ann Rheum Dis* 2008;67:1354-1355.

Table 1. Patient characteristics for the investigated patients. Not all patients had data on RF and ACPA.

	Gouty arthritis	Pseudogout\$	Reactive arthritis	RS3PE	All non-RA	RA (from [4])
Number	26	3	44	33	106	274
Female; number (%)	9 (34)	1 (33)	17 (39)	17 (52)	44 (42)	193 (70)
Age, years; median (IQR)	62 (48-69)	80 (74-80)	33 (27-47)	52 (28-72)	46 (30-66)	56 (46-70)
CRP; median (IQR)	7 (3-30)	52 (0-100)	13 (3-38)	5 (3-11)	7 (3-26)	14.5 (7-31)
ESR; median (IQR)	16 (10-37)	60 (17-77)	19 (8-63)	11 (6-26)	16 (8-39)	21 (12-40)
SJC (66 joint count); median (IQR)	3 (1.5-9.5)	11 (1-12)	2 (1-4)	3 (1-6)	2 (1-6)	9 (5-13)
TJC (66 joint count); median (IQR)	4 (1-6)	13 (1-17)	3 (2-5)	5 (2-10.5)	4 (2-6)	7 (4-12)
RF positive/total measured (%)	4/26 (15.4)	0/3 (0)	3/44 (6.8)	4/33 (12.1)	11/106 (10.4)	172/272 (63.2)
ACPA positive/total measured (%)	0/19 (0)	0/3 (0)	1/12 (8.3)	0/6 (0)	1/41 (2.4)	157/274 (57.3)
Anti-CII* positive/total measured (%)	2/26 (7.7)	0/3 (0)	3/44 (6.8)	3/33 (9.1)	8/106 (7.5)	24 (8.8)

* samples yielding > 29 AU/ml of anti-CII and showing lower OD in bovine serum albumin blocked control wells without CII than in anti-CII coated and subsequently blocked wells. \$ patients presenting with an inflammatory arthritis of at least one joint in which calcium pyrophosphate crystals were identified in synovial fluid microscopy and in whom no alternative diagnosis was made.

Figure 1. Levels of antibodies against native human collagen type II. The dotted horizontal line represent the cutoff based on the 95th percentile among healthy controls (29 arbitrary units/ml). Values above the cutoff are divided into true positive samples (triangles) and samples with non-specific binding (stars) reacting with higher OD levels in control wells coated only with bovine serum albumin. Only true positive values are reported as positive in table 1. Solid horizontal bars represent median values in each group. ReA= reactive arthritis, RS3PE= remitting seronegative symmetrical synovitis with pitting edema, RA=rheumatoid arthritis. Data on the 274 RA patients were published previously. [4, 5]