Use of the SAME-TT2R2 score to predict anticoagulation control in atrial fibrillation and venous thromboembolism patients treated with vitamin K antagonists: A review

Zulkifly, Hanis; Lip, Gregory; Lane, Deirdre

DOI:

[10.1016/j.hrthm.2017.11.026](https://doi.org/10.1016/j.hrthm.2017.11.026)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Zulkifly, H, Lip, G & Lane, D 2017, 'Use of the SAME-TT2R2 score to predict anticoagulation control in atrial fibrillation and venous thromboembolism patients treated with vitamin K antagonists: A review', *Heart Rhythm*.
<https://doi.org/10.1016/j.hrthm.2017.11.026>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Use of the SAME-TT₂R₂ score to predict anticoagulation control in atrial fibrillation and venous thromboembolism patients treated with vitamin K antagonists: A review

Running head: The SAME-TT₂R₂ score: a review

Hanis Zulkifly, MPharm^a

Gregory Y.H. Lip, MD^{a, b}

Deirdre A. Lane, PhD^{a, b}

^aInstitute of Cardiovascular Sciences, University of Birmingham, Birmingham, United Kingdom

^bAalborg Thrombosis Research Unit, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

HHZ523@student.bham.ac.uk

g.y.h.lip@bham.ac.uk

deirdrelane@nhs.net

Corresponding Author: Dr Deirdre A Lane

Tel: +44 121-507-5080 Email: deirdrelane@nhs.net

Mailing address: University of Birmingham Institute of Cardiovascular Sciences, City Hospital,
Dudley Road, Birmingham, B18 7QH, UK

Word count: 5957

Article type: Review article

23 **Conflict of interest:**

24 HHZ: Received PhD funding from Malaysian Ministry of Education and Universiti Teknologi MARA
25 but not directly for the work under consideration.

26 GYHL: Consultant for Bayer/Janssen, BMS/Pfizer, Biotronik, Medtronic, Boehringer-Ingelheim,
27 Microlife and Daiichi-Sankyo. Speaker for Bayer, BMS/Pfizer, Medtronic, Boehringer-Ingelheim,
28 Microlife, Roche and Daiichi-Sankyo. No fees are received personally. Co-author of original SAME-
29 TT₂R₂ score.

30 DAL: Institution received educational grants from BMS and Boehringer-Ingelheim. Speaker and
31 consultancy for Boehringer-Ingelheim, Bayer, and BMS/Pfizer.

32

Abstract

Identifying patients who are likely to achieve and maintain a therapeutic INR when prescribed a vitamin K antagonist (VKA) for stroke prevention in atrial fibrillation (AF) and venous thromboembolism (VTE) is challenging. The SAME-TT₂R₂ score was developed based on common clinical factors that can highlight patients who may be unable to achieve and maintain good anticoagulation control and for whom a 'trial of warfarin' would be inadvisable. This review summarises the main published prospective and retrospective studies that have validated the SAME-TT₂R₂ score in AF and VTE patients treated with a VKA and how the SAME-TT₂R₂ score could aid clinical decision-making; 19 studies were included. Taken together validation studies suggest that the SAME-TT₂R₂ score is able to predict good or poor anticoagulation control among AF and VTE patients, although data on VTE patients are limited (3 studies). The available evidence suggests that the SAME-TT₂R₂ score may be a useful tool to aid clinical decision-making for oral anticoagulants (OAC) in AF and VTE patients.

Keywords: SAME-TT₂R₂ score; atrial fibrillation; venous thromboembolism; vitamin K antagonist; decision-making; oral anticoagulation

Introduction

For decades, vitamin K antagonists (VKA, e.g., warfarin) have been the cornerstone of stroke prevention in atrial fibrillation (AF) and prevention of venous thromboembolism (VTE).¹ However, VKA efficacy and safety requires achievement of an international normalised ratio (INR) between 2.0-3.0. Achieving this target INR alone is an inadequate measure of the therapeutic efficacy of VKA.¹

Time in therapeutic range (TTR) is one measure that summarises INR control over time. TTR is an important and independent predictor of thromboembolic and bleeding outcomes in AF patients on VKA.^{2,3} An average individual TTR $\geq$ 65% is recommended by NICE guidelines,¹ while European guidelines⁴ recommend TTR $\geq$ 70% to maximize effectiveness and safety of VKAs.

However, identifying patients who are likely to achieve and maintain a therapeutic INR is more difficult. Based on common clinical factors that influence INR and anticoagulation control in everyday clinical practice, a clinical scoring system, the SAME-TT₂R₂ score⁵ (**Table 1**) was developed in 2013 to identify risk factors highlighting those patients who may be unable to achieve/maintain good anticoagulation control and for whom a 'trial of warfarin' would be inadvisable. The frequency of INR measurements are not factored-in (or intended to be). This score assigns 1 point each to female sex, age <60 years, history of $\geq$ 2 co-morbidities (hypertension, diabetes mellitus, coronary artery disease or myocardial infarction, peripheral artery disease, congestive heart failure, previous stroke, pulmonary, hepatic, or renal disease) and treatment with drugs interacting with VKA (e.g., amiodarone) and 2 points each for current/recent tobacco use (within 2-years) and non-white ethnicity⁵ (**Table 1**). The score can be used to aid decision-making by identifying those patients who would probably do well on VKA (achieving a high TTR, $\geq$ 65%) or conversely, those would need additional interventions to achieve good INR control or to be started on/switched to a non-VKA oral anticoagulant (NOAC). The current review summarises studies which have assessed and/or validated the SAME-TT₂R₂ score in patients treated with VKA for AF or VTE.

Methods

A comprehensive structured literature search was performed using MEDLINE and EMBASE from 2013 until February 2017; the SAME-TT₂R₂ score was first published in 2013. The search strategy included keywords and MeSH terms relating to AF, deep vein thrombosis, VTE, stroke prevention, warfarin, VKAs, oral anticoagulant, inception cohort, adverse effect, poor control, INR and SAME-TT₂R₂ score (without MeSH term) individually and in combination. Primary published research articles and abstracts on prospective or retrospective studies validating the SAME-TT₂R₂ score were included. Studies that did not provide comparative outcomes, information on follow-up time, or were not published in English language were excluded. Manual search of citations was also performed, and discussion with content experts was undertaken to identify any other relevant studies (**Figure 1**).

Results

Searches identified 166 citations. After removal of duplicates and screening of titles and abstracts, 24 full-text articles were assessed for eligibility and 19 studies were included (see **Figure 1**). Current studies assessing the SAME-TT₂R₂ score are summarised in **Table 2** and baseline patient characteristics of these cohorts in **Table 3**. With the exception of three⁶⁻⁸ all were performed in AF patients.^{2,5,9-22} Most studies (n=11)^{5-7,11,14,17,19-22} were performed prospectively, with follow-up duration ranging from six-months¹⁷ to 4.7 years.¹⁵ The number of participants included in VTE cohorts ranged from 135⁶ to 1943⁸ and between 104¹⁴ to 8120²¹ in studies on AF patients.

Fourteen studies were performed in European populations,^{5-7,9-12,14,17-22} two in Asian populations,^{15, 16} (with one reporting a target INR 2.0-3.0¹⁵) and two in North American populations.^{8, 13} Proietti et al¹¹ studied a mixed indication clinical trial cohort including patients from Europe, Asia and Australasia.

Most studies were performed in elderly (mean/median age ranging from 61-76 years) white-Western populations, mainly using warfarin (13 studies)^{5-11,13,15,16,19-21} as the OAC of choice. Most patients had multiple comorbidities with hypertension the most common, except for the study by Lip et al²¹ where congestive heart failure was most prevalent. All studies reported a low prevalence of

smoking status and use of amiodarone for rhythm-control, with the exception of the original derivation study; 35% of patients used amiodarone.²¹ As the SAME-TT₂R₂ score categories increase, the mean TTR of their study population decreases, except for one study by Domelo-Rodriguez⁶ which showed the opposite relationship (Figure 2).

Five studies^{8,12,13,15,18} investigated the relationship between components included in the SAME-TT₂R₂ score and TTR. Three studies^{12,13,18} showed that female sex was associated with poor anticoagulation control; one¹⁸ showed that having ≥ 2 comorbidities was related to poor TTR and one¹³ showed that black ethnicity (as well as NYHA IV) was associated with poorer anticoagulation control. Chan et al¹⁵ also reported that having heart failure and diabetes mellitus independently predicts poor anticoagulation control.

Eight studies^{2,5,7-9,12,18, 21} reported the predictive ability of the SAME-TT₂R₂ score using c-statistics (**Figure 3**). Taken together these validation studies suggest that the SAME-TT₂R₂ score is able to predict good or poor anticoagulation control among AF patients better than chance, with c-statistics ranging from 0.56¹² to 0.72;⁵ the evidence is less robust in VTE patients (c-statistic 0.52-0.65).^{7,8}

Eight studies^{11,15,18,20-22} also examined if the SAME-TT₂R₂ score could discriminate AF patients with clinical events. Five^{11,15,18,21,22} demonstrated some positive associations for SAME-TT₂R₂ score predicting clinical events, with c-statistics ranging from 0.55²¹ to 0.62²² (**Table 4**). Another study,⁸ also examined if the SAME-TT₂R₂ score was associated with clinical outcomes, in particular recurrent VTE and International Society on Thrombosis and Haemostasis (ISTH) major bleeding rates in a VTE cohort; patients with a score > 2 had more overall adverse event rates (composite of recurrent VTE and ISTH major bleeding) than those with a score of 0-2 (7.9 vs. 4.5 overall adverse event rates/100 patient-years respectively).⁸

Discussion

This review of studies assessing and validating the SAME-TT₂R₂ score extends and updates a previous narrative review²³ with the addition of validation studies in VTE populations^{6, 7} and validations in Asian AF populations.^{15,16} Overall, eight studies^{2,5,7-9,12,18,21} suggest that the SAME-TT₂R₂ score is able

to modestly predict quality of anticoagulation control in AF patients receiving VKA therapy, with c-statistics ranging from 0.56¹² to 0.72.⁵ Many risk scores based on clinical factors such as CHADS₂, CHA₂DS₂-VASc, Killip and TIMI scores show broadly similar modest c-indexes (approx. 0.6) when used to predict patients categorised at ‘high risk’ who actually sustain clinical events.^{24,25}

The original purpose of developing the SAME-TT₂R₂ score was to produce a simple clinical schema which could be used routinely in everyday practice to help assess the likelihood of an AF patient being able to achieve and maintain good anticoagulation control on VKA therapy, using patient-related clinical parameters which are readily available. The availability of NOACs worldwide has resulted in increased usage due to their advantages. These include faster onset-of-action (average maximum effect approximately three hours after intake²⁶ compared to VKA (onset 36-72 hours)), greater reduction in stroke/systemic embolism (+19% compared to VKA⁴), avoidance of INR monitoring with NOACs,²⁷ and absence of achieving/maintaining adequate TTR (as with warfarin). Achieving a therapeutic INR can take 2-4 weeks and often longer.³ After termination of study drug in the NOAC trials, of those patients switching to warfarin, <40% achieved a therapeutic INR within 15 days, and <80% after 30 days;²⁸ more strokes occurred during that period in the patients who went from study drug to VKA than from VKA to VKA.^{28,29} This strongly argues for using NOACs over VKAs where possible, however, VKAs are still widely used globally and will not disappear from use especially for AF patients with severe renal impairment, moderate to severe mitral stenosis or mechanical heart valves.⁴

In addition, in low- and middle-income countries where cost plays an important role in options available for OAC treatment VKA is still the first-line antithrombotic agent of choice, therefore the SAME-TT₂R₂ score will remain an important decision-making tool, currently and in the future, to guide physicians choice of anticoagulant treatment.³⁰ Most validation studies included in this review demonstrated good predictive ability except two^{6,19} which demonstrate that the SAME-TT₂R₂ score was unable to predict anticoagulation control well in their populations. Although both studies were prospective, results should be interpreted with care as both included small numbers of participants (135⁶ and 180¹⁹ respectively) and thus may not be adequately powered to test the predictive ability of the SAME-TT₂R₂ score in regard to anticoagulation control.

Importance of good anticoagulation control

Achieving good anticoagulation control ($TTR \geq 65-70\%$) as recommended by guidelines^{1,4} is essential for managing AF and VTE patients treated with VKA. Numerous studies have demonstrated that a high TTR translates into lower risk of stroke and bleeding.³¹⁻³⁵ A systematic review demonstrated that a 7% and 12% improvement in TTR can lead to a reduction in major bleeding and thromboembolic events, respectively, by 1 event per 100 patient years.³⁴ A real-world study³² of 27,458 warfarin-treated AF patients (≥ 3 INR measurements), showed that in patients with good anticoagulation control ($TTR \geq 70\%$), stroke risk was reduced to 79% compared to patients with poor INR control ($TTR \leq 30\%$). However, achieving and maintaining a therapeutic INR can be difficult to accomplish and therefore, NOACs are preferred to VKA in the majority of patients requiring OAC initiation.⁴

SAMe-TT₂R₂ score and clinical events

Evident in most studies included in this review,^{2,5,7,9-18,20-22} increasing SAMe-TT₂R₂ score demonstrated poorer TTR values which might also translate into poorer clinical outcomes. This can be evidenced by studies that showed the SAMe-TT₂R₂ score relating to severe bleeding²² and major bleeding (defined by the Bleeding Academic Research Consortium),²¹ stroke/TE,²¹ adverse cardiovascular events²² and death^{21, 22} during follow-up. In an observational study performed in 911 Spanish AF patients, the SAMe-TT₂R₂ score also successfully predicted the composite outcome of major bleeding, TE complications and death.¹⁸ A Chinese study also demonstrated that a SAMe-TT₂R₂ score of ≤ 2 vs. SAMe-TT₂R₂ of 3 vs. SAMe-TT₂R₂ ≥ 4 is associated with lower annual stroke risk (3.49%/year vs. 4.56% per year vs. 6.41%/year, respectively).¹⁵

Impact of different methods of calculating TTR

Fauchier and colleagues³⁶ have raised concern about the different methods used to calculate TTR, whether to use TTR based on the Rosendaal method, percentage of INRs in range (PINRR) (traditional method) or percentage of visits in range on a given date (cross-sectional method), as these methods are not interchangeable. In this review, 17 studies^{2,5-17,19,20,22} reported TTR using the Rosendaal method, only one¹⁸ calculated time in therapeutic range according to PINRR, while the other reported 'labile INR' as their measure of anticoagulation control.²¹ Currently there is no evidence on the optimal method of calculating percentage of INR in range, as each method has its own unique strengths and weaknesses.³⁷ While TTR via the Rosendaal method calculates the exact percentage of days the INR falls within range, its calculation is more complex than the others and is based on linear extrapolation. In contrast, calculating TTR via the PINRR method is simpler as it only looks at the number of INRs that fall within the therapeutic range divided by the total number of INR tests undertaken. However, the PINRR method does not take into account the actual number of days of anticoagulant treatment and thus might underestimate control in patients with inconsistent INR monitoring, patients who have temporarily discontinued therapy and patients with a long gaps between each INR test, in contrast to the Rosendaal method where these factors will be accounted for, resulting in a lower TTR.

Factors affecting anticoagulation control

In this review, only 5 studies^{8,12,13,15,18} investigated the relationship of individual components of the SAME-TT₂R₂ score with the quality of anticoagulation control. Among these female sex^{12,13,18}, >2 comorbidities,¹⁸ heart failure and diabetes mellitus¹⁵ (individually) and black ethnicity⁵ were associated with poor TTR control, however no studies found any association between age <60 years and smoking with poor TTR.

It is interesting to speculate how some elements of the SAME-TT₂R₂ score could influence anticoagulation control. Some studies^{38,39} investigating predictors of TTR have demonstrated that women have poorer anticoagulation control compared to men (translating into poorer outcomes), although the precise mechanism remains unclear.⁵ Similarly, women are known to be at higher risk of AF-related stroke irrespective of warfarin use.^{40,41} Tobacco use within 2 years scores 2-points in the

SAMe-TT₂R₂ score, however most validation studies reported low prevalence of smoking (6.3%-30%) except in the external validation study by Apostolakis et al⁵ (49% reported as smoker/ex-smoker (within 2 years)). How smoking can influence anticoagulation control is unclear but it may reflect less interest in maintaining good health which may translate into poorer adherence to oral anticoagulants, thus resulting in poor TTR.⁵

The original SAMe-TT₂R₂ score publication suggested that patients who are younger and have more comorbidities probably have adherence issues with VKA therapy which are reflected by poor TTR³. In terms of non-white ethnicity, some studies have shown that African-Americans and Hispanics have poorer anticoagulation control compared to whites and suggest that this may be due to various reasons including socioeconomic status, poor understanding of therapy, adherence issues, genetic predisposition, etc.^{42,43} However, these aspects need to be further investigated as studies in these areas are lacking.

Another editorial⁴⁴ suggests that other factors, not currently within the SAMe-TT₂R₂ score, could be included in the assessment of anticoagulation control, such as distance from home to anticoagulation clinic, which could be the main reason preventing patients attending for regular follow-up. There is clearly the need for a large prospective randomised trial to evaluate the impact of SAMe-TT₂R₂ score-guided therapy with VKA or NOAC not only in relation to anticoagulation control (TTR) but also towards clinical outcomes (stroke and bleeding), which would formalise its utility in clinical practice. Hence, where patients have chosen VKA over a NOAC for stroke prevention or treatment of VTE or where NOACs are contraindicated but a high SAMe-TT₂R₂ score (>2) is present, perhaps more frequent follow-up visits and reviews, educational interventions and counselling⁴⁵ may be required to ensure that INRs are within the therapeutic range in order to achieve the best outcomes and minimise treatment complications.

Limitations

The main limitation of the included studies is study design; none utilised a randomised controlled trial design and most were performed in white populations. Given that one of the risk factors for poorer

anticoagulation control is ethnicity, SAME-TT₂R₂ score in these populations is automatically worse compared to non-whites; thus a lower score predicts better control of VKA therapy. Thus, future studies need to ascertain whether the threshold of the SAME-TT₂R₂ score used to indicate probability of poorer anticoagulation control (SAME-TT₂R₂ score ≥ 2) needs to be modified in non-white populations so that the SAME-TT₂R₂ score is applicable globally. In addition, only three studies have validated the SAME-TT₂R₂ score in VTE cohorts to date, hence more studies are needed specifically in VTE cohorts to enhance its applicability in these patients. Lastly, only 8 studies reported the c-statistic to quantify the predictive ability of the SAME-TT₂R₂ score.

Conclusions

Making decisions when choosing OAC therapy can be challenging. The available evidence suggests that the SAME-TT₂R₂ score is a useful tool to aid decision-making for OAC in AF (and VTE) patients and adequately predicts those who are likely to be able/unable to achieve and maintain good INR control.

References

1. National Institute for Care Excellence (2014) Atrial Fibrillation: Management. NICE Guideline (CG180)2014.
2. Ruiz-Ortiz M, Bertomeu V, Cequier A, Marin F, Anguita M. Validation of the SAME-TT₂R₂ score in a nationwide population of nonvalvular atrial fibrillation patients on vitamin K antagonists. *Thromb Haemost* 2015;114:695-701.
3. De Caterina R, Husted S, Wallentin L, et al. Vitamin K antagonists in heart disease: Current status and perspectives (Section III). Position Paper of the ESC Working Group on Thrombosis – Task Force on Anticoagulants in Heart Disease. *Thromb Haemost* 2013;110:1087-1107.
4. Kirchhof P, Benussi S, Kotecha D, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J* 2016;37:2893-2962.
5. Apostolakis S, Sullivan RM, Olshansky B, Lip GYH. Factors affecting quality of anticoagulation control among patients with atrial fibrillation on warfarin: The same-tt₂r₂ score. *Chest* 2013;144:1555-1563.
6. Demelo-Rodríguez P, Postigo-Esteban A, García-Fernández-Bravo I, Agud M, Calpena S, Camacho M, del Toro-Cervera J. Evaluation of the SAME-TT₂R₂ score to predict the quality of anticoagulation control in a cohort of patients with venous thromboembolism treated with vitamin K antagonists. *Thromb Res* 2016;147:58-60.
7. Palareti G, Antonucci E, Lip GYH, Testa S, Guazzaloca G, Falanga A, Pengo V, Poli D. The SAME-TT₂R₂ score predicts the quality of anticoagulation control in patients with acute VTE. A real-life inception cohort study. *Thromb Haemost* 2016;115:1101-1108.
8. Kataruka A, Kong X, Haymart B, Kline-Rogers E, Almany S, Kozlowski J, Krol GD, Kaatz S, McNamara MW, Froehlich JB. SAME-TT₂R₂ predicts quality of anticoagulation in patients with acute venous thromboembolism: The MAQI2 experience. *Vasc Med* 2017;22:197-203.

- 284 **9.** Pivatto Júnior F, Scheffel R, Ries I, Wolkind R, Marobin R, Barkan S, Amon L, Biolo A.
285 SAME-TT2R2 Score in the Outpatient Anticoagulation Clinic to Predict Time in Therapeutic
286 Range and Adverse Events. *Arq Bras Cardiol* 2017;108:290-296.
- 287 **10.** Szymanski FM, Lip GYH, Filipiak KJ, Platek AE, Karpinski G. Usefulness of the SAME-
288 TT2R2 score to predict anticoagulation control on VKA in patients with atrial fibrillation and
289 obstructive sleep apnea. *Int J Cardiol* 2016;204:200-205.
- 290 **11.** Proietti M, Lane DA, Lip GYH. Relation of the SAME-TT2R2 score to quality of
291 anticoagulation control and thromboembolic events in atrial fibrillation patients: Observations
292 from the SPORTIF trials. *Int J Cardiol* 2016;216:168-172.
- 293 **12.** Lobos-Bejarano JM, Barrios V, Polo-García J, Escobar C, Vargas-Ortega D, Marín-Montañés
294 N, Prieto-Valiente L, Fuentes S, Prieto MA, García-Ortiz L. Evaluation of SAME-TT2R2
295 score and other clinical factors influencing the quality of anticoagulation therapy in non-
296 valvular atrial fibrillation: a nationwide study in Spain. *Curr Med Res Opin* 2016;32:1-7.
- 297 **13.** Lip GY, Waldo AL, Ip J, Martin DT, Bersohn MM, Choucair WK, Akar JG, Wathen M,
298 Halperin JL. Determinants of Time in Therapeutic Range in Patients Receiving Oral
299 Anticoagulants (A Substudy of IMPACT). *Am J Cardiol* 2016;118:1680-1684.
- 300 **14.** Gorzalak-Pabis P, Zyzak S, Krewko L, Broncel M. Assessment of the mean time in the
301 therapeutic INR range and the SAME-TT2R2 score in patients with atrial fibrillation and
302 cognitive impairment. *Pol Arch Med Wewn* 2016;126:494-501.
- 303 **15.** Chan PH, Hai JJ, Chan EW, Li WH, Tse HF, Wong ICK, Lip GYH, Siu CW. Use of the
304 SAME-TT(2)R(2) Score to Predict Good Anticoagulation Control with Warfarin in Chinese
305 Patients with Atrial Fibrillation: Relationship to Ischemic Stroke Incidence. *PLoS One*
306 2016;11:e0150674.
- 307 **16.** Bernaitis N, Ching CK, Chen L, Hon JS, Teo SC, Davey AK, Anoopkumar-Dukie S. The
308 Sex, Age, Medical History, Treatment, Tobacco Use, Race Risk (SAME TT2R2) Score
309 Predicts Warfarin Control in a Singaporean Population. *J Stroke Cerebrovasc Dis* 2016;26:64-
310 69.

17. Roldan V, Cancio S, Galvez J, Valdes M, Vicente V, Marin F, Lip GY. The SAME-TTR Score Predicts Poor Anticoagulation Control in AF Patients: A Prospective 'Real-world' Inception Cohort Study. *Am J Med* 2015;128:1237-1243.
18. Abumuaileq RR, Abu-Assi E, Raposeiras-Roubin S, Lopez-Lopez A, Redondo-Diequez A, Alvarez-Iglesias D, Rodriguez-Manero M, Pena-Gil C, Gonzalez-Juanatey JR. Evaluation of SAME-TT2R2 risk score for predicting the quality of anticoagulation control in a real-world cohort of patients with non-valvular atrial fibrillation on vitamin-K antagonists. *Europace* 2015;17:711-717.
19. Skov J, Bladbjerg E-M, Bor MV, Gram J. SAmett2r2 does not predict time in therapeutic range of the international normalized ratio in patients attending a high-quality anticoagulation clinic. *Chest* 2014;145:187-188.
20. Poli D, Antonucci E, Testa S, Lip GY. A prospective validation of the SAME-TT2R 2 score: how to identify atrial fibrillation patients who will have good anticoagulation control on warfarin. *Intern Emerg Med* 2014;9:443-447.
21. Lip GYH, Haguenoer K, Saint-Etienne C, Fauchier L. RElationship of the same-tt2r2 score to poor-quality anticoagulation, stroke, clinically relevant bleeding, and mortality in patients with atrial fibrillation. *Chest* 2014;146:719-726.
22. Gallego P, Roldan V, Marin F, Galvez J, Valdes M, Vicente V, Lip GY. SAME-TT2R2 score, time in therapeutic range, and outcomes in anticoagulated patients with atrial fibrillation. *Am J Med* 2014;127:1083-1088.
23. Esteve-Pastor MA, Roldan V, Valdes M, Lip GY, Marin F. The SAME-TTR score and decision making between a Vitamin K Antagonist or a Non-Vitamin K Antagonist Oral Anticoagulant in patients with atrial fibrillation. *Expert Rev Cardiovasc Ther* 2015;14:177-187.
24. Senoo K, Lip GY. Predictive abilities of the HAS-BLED and ORBIT bleeding risk scores in non-warfarin anticoagulated atrial fibrillation patients: An ancillary analysis from the AMADEUS trial. *Int J Cardiol* 2016;221:379-382.

- 338 **25.** Lip GY, Lane DA. Bleeding risk assessment in atrial fibrillation: observations on the use and
339 misuse of bleeding risk scores. *J Thromb Haemost* 2016;14:1711-1714.
- 340 **26.** Heidbuchel H, Verhamme P, Alings M, et al. Updated European Heart Rhythm Association
341 practical guide on the use of non-vitamin-K antagonist anticoagulants in patients with non-
342 valvular atrial fibrillation: Executive summary. *Eur Heart J* 2017;38:2137-2149.
- 343 **27.** Mekaj YH, Mekaj AY, Duci SB, Miftari EI. New oral anticoagulants: their advantages and
344 disadvantages compared with vitamin K antagonists in the prevention and treatment of
345 patients with thromboembolic events. *Ther Clin Risk Manag* 2015;11:967-977.
- 346 **28.** Mahaffey KW, Hellkamp AS, Patel MR, et al. End of Study Transition From Study Drug to
347 Open-Label Vitamin K Antagonist Therapy. *Circ Cardiovasc Qual Outcomes* 2013;6:470.
- 348 **29.** Ruff CT, Giugliano RP, Braunwald E, Mercuri M, Curt V, Betcher J, Grip L, Cange AL,
349 Crompton AE, Murphy SA, Deenadayalu N, Antman EM. Transition of Patients From
350 Blinded Study Drug to Open-Label Anticoagulation: The ENGAGE AF-TIMI 48 Trial. *J Am*
351 *Coll Cardiol* 2014;64:576-584.
- 352 **30.** Bista D, Chalmers L, Bereznicki L, Peterson G. Potential use of NOACs in developing
353 countries: pros and cons. *Eur J Clin Pharmacol* 2014;70:817-828.
- 354 **31.** Gallego P, Roldan V, Marín F, Romera M, Valdés M, Vicente V, Lip GYH. Cessation of oral
355 anticoagulation in relation to mortality and the risk of thrombotic events in patients with atrial
356 fibrillation. *Thromb Haemost* 2013;110:1189-1198.
- 357 **32.** Gallagher AM, Setakis E, Plumb JM, Clemens A, van Staa T-P. Risks of stroke and mortality
358 associated with suboptimal anticoagulation in atrial fibrillation patients. *Thromb Haemost*
359 2011;106:968.
- 360 **33.** Morgan CL, McEwan P, Tukiendorf A, Robinson PA, Clemens A, Plumb JM. Warfarin
361 treatment in patients with atrial fibrillation: Observing outcomes associated with varying
362 levels of INR control. *Thromb Res* 2009;124:37-41.
- 363 **34.** Wan Y, Heneghan C, Perera R, Roberts N, Hollowell J, Glasziou P, Bankhead C, Xu Y.
364 Anticoagulation Control and Prediction of Adverse Events in Patients With Atrial
365 FibrillationCLINICAL PERSPECTIVE. *Circ Cardiovasc Qual Outcomes* 2008;1:84-91.

35. White HD, Gruber M, Feyzi J, et al. Comparison of outcomes among patients randomized to warfarin therapy according to anticoagulant control: Results from sportif iii and v. Arch Intern Med 2007;167:239-245.
36. Fauchier L, Poli D, Olshansky B. The SAME-TT₂R₂ score and quality of anticoagulation in AF: Can we predict which patient benefits from anticoagulation? Thromb Haemost 2015;114:657-659.
37. Schmitt L, Speckman J, Ansell J. Quality Assessment of Anticoagulation Dose Management: Comparative Evaluation of Measures of Time-in-Therapeutic Range. J Thromb Thrombolysis 2003;15:213-216.
38. Senoo K, Lip GYH. Female Sex, Time in Therapeutic Range, and Clinical Outcomes in Atrial Fibrillation Patients Taking Warfarin. Stroke 2016;47:1665.
39. Rose AJ, Hylek EM, Ozonoff A, Ash AS, Reisman JI, Berlowitz DR. Patient characteristics associated with oral anticoagulation control: results of the Veterans Affairs Study to Improve Anticoagulation (VARIA). J Thromb Haemost 2010;8:2182-2191.
40. Fang MC, Singer DE, Chang Y, Hylek EM, Henault LE, Jensvold NG, Go AS. Gender Differences in the Risk of Ischemic Stroke and Peripheral Embolism in Atrial Fibrillation. Circulation 2005;112:1687.
41. Tsadok M, Jackevicius CA, Rahme E, Humphries KH, Behloul H, Pilote L. Sex differences in stroke risk among older patients with recently diagnosed atrial fibrillation. JAMA 2012;307:1952-1958.
42. Yong C, Azarbal F, Abnoui F, et al. Racial Differences in Quality of Anticoagulation Therapy for Atrial Fibrillation (from the TREAT-AF Study). Am J Cardiol 2016;117:61-68.
43. Golwala H, Jackson LR, 2nd, Simon DN, et al. Racial/ethnic differences in atrial fibrillation symptoms, treatment patterns, and outcomes: Insights from Outcomes Registry for Better Informed Treatment for Atrial Fibrillation Registry. Am Heart J 2016;174:29-36.
44. Roldan V, Marin F. The importance of excellence in the quality of anticoagulation control whilst taking vitamin K antagonists. Thromb Haemost 2015;113:671-673.

393 **45.** Clarkesmith DE, Pattison HM, Lip GYH, Lane DA. Educational intervention improves
394 anticoagulation control in atrial fibrillation patients: the TREAT randomised trial. PLoS One
395 2013;8:e74037.

396

397

Table 1: The SAME-TT₂R₂ score

Component		Score
S	Sex (female)	1
A	Age (<60 years)	1
Me	Medical history [†]	1
T	Treatment (interacting drugs, e.g., amiodarone)	1
T	Tobacco use (within 2-years)	2
R	Race (non-white ethnicity)	2
Maximum total score		8

[†]≥2 of the following: hypertension, diabetes mellitus, coronary artery disease/myocardial infarction, peripheral arterial disease, congestive heart failure, previous stroke, pulmonary, hepatic, or renal disease.

Table 2: Studies assessing the SAME-TT₂R₂ score in atrial fibrillation and venous thromboembolism cohorts

	a. Study design b. Mean follow-up c. Method INR monitoring	Population a. Number b. Mean (SD)/median (IQR) age (range, years) c. Race/ethnicity d. OAC used	SAME-TT₂R₂ score distribution (%); mean TTR (%) ± SD	Percentage of patients with dichotomised TTR (%)
Pivatto Junior⁹ 2017 Brazil	a. Retrospective b. 1 year c. Hospital OAC clinic	a. 263 AF b. 71.2 (64.1-78.5) c. White d. 97.3% Warfarin	0-1: 138 (52.5); 69.2 ≥2: 125 (47.5); 56.3	-
Kataruka⁸ 2017 USA	a. Retrospective b. Median 0.56 years ± 1.13 c. Hospital OAC clinic	a. 1943 VTE b. 61.8 (15.7) c. White d. Warfarin	0-1: 665; 57±21 2: 432; 55±22 >2: 846; 50±23	TTR<60: 57.6
Bernaitis¹⁶ 2016 Singapore	a. Retrospective b. - c. Hospital	a. 1137 AF b. 71 (63-77) c. Asian d. Warfarin	0-1: 0 2: 339; 63.2±34.1 >2: 798; 55.8±34.1	-
Chan¹⁵ 2016 Hong Kong	a. Retrospective b. 4.7 ± 3.6 years c. Hospital	a. 1428 NVAF b. 76.2 (8.7) c. Chinese d. Warfarin	2: 22(14.3); 70 [†] 3: 80 (51.9); 70 4: 41 (26.6); 70 5: 7 (4.5); 70 6: 4 (2.6); 70	TTR≥70: 11 TTR<70: 89
Demelo-Rodriguez⁶ 2016 Spain	a. Prospective b. 72,668 patient-years c. Primary care	a. 135 VTE b. Median 66 [#] c. White d. Warfarin	0-1: 91; 64.7±19.5 ≥2: 44; 66 ±20.5	-
Gorzalak-Pabis¹⁴ 2016 Poland	a. Prospective b. - c. Hospital	a. 104 AF with cognitive impairment b. 75 (10) c. White d. 61% Acenocoumarol	0-1: 64±26 ≥2: 50±28	-

Lip¹³ 2016 USA	a. Prospective b. 438 days c. Trial setting	a. 229 AF b. 66.7 (11) c. 80.3% White d. Warfarin	0-1:0.571±0.22 ≥2: 0.498±0.24	-
Lobos-Bejarano¹² 2016 Spain	a. Retrospective b. >12 months c. Primary care	a. 1524 NVAF b. 77.4 (8.7) c. White d. 94.8% Acenocoumarol	0-1: 69.6% ± 17.4 ≥2: 66.6% ± 18.5	TTR≥65: 60.6 TTR<65: 39.4
Palareti⁷ 2016 Spain	a. Prospective b. 998 patient-years c. Hospital OAC clinic	a. 1308 VTE b. 68(51-78) c. White d. Warfarin	0-1: 916 (70); 61±22 ≥2: 392 (30); 56±23	TTR≥65: 50.4 TTR<65: 49.6
Proietti¹¹ 2016 Europe, Asia, Australasia	a. Prospective b. Median 563 days c. Trial setting	a. 3665 AF b. 72(66-77) c. Mixed‡ d. Warfarin	0-2: 2914 (80.4); 69.05 (55.63-79.89) >2: 710 (19.6); 66.55 (52.83-77.46)	TTR>70: 46.9 TTR≤70: 53.1
Szymanski¹⁰ 2016 Poland	a. Retrospective b. - c. Hospital	a. 211 AF b. 57.1 (10.2) c. White d. 75.4% warfarin	0-1: 114 (54); 52.3 ≥2: 97 (46); 51.3	TTR>70: 25.2 TTR≤70: 74.8
Abumuaileq¹⁸ 2015 Spain	a. Retrospective b. 10 months c. Hospital OAC clinic	a. 911 NVAF b. 73 (11) c. White d. 93% Acenocoumarol	0-1:672 (74); 59±18 [¶] ≥2: 239 (26); 54±19 [¶]	PINRR>65:39 PINRR≤65:61
Roldán⁴⁴ 2015 Spain	a. Prospective b. 6 months c. Hospital OAC clinic	a. 459 NVAF b. 76 (70-82) c. White d. Acenocoumarol	<2: 253 (55); 67±18 ≥2: 206 (44.8); 61±16	TTR>65:54 TTR≤65:46
Ruiz-Ortiz² 2015 Spain	a. Retrospective b. Median 27 months c. Cardiology clinic	a. 1056 NVAF b. 73.6 (9.8) c. White d. Acenocoumarol	0-1:613 (58); 65.6±26.2 ≥2: 443 (42); 61.3±25.3	TTR≥65:52.7 TTR≥65:47.3

Gallego²² 2014 Spain	a. Prospective b. Median 952 days c. Hospital OAC clinic	a. 972 NVAF b. 76 (70-82) c. White d. Acenocoumarol	0-1:431 (44); 79.67 ±19.46 ≥2: 332 (34); 78.4 ± 20.28 >2:208 (21); 74.25 ± 20.24	-
Lip²¹ 2014 France	a. Prospective b. 1016±1018 days c. Clinicians -hospital	a. 8120 AF b. 70 (15) c. White d. Warfarin	0-1: 4504 (55); 77(1.7) [§] ≥2: 2252 (28); 52(2.3) [§] >2:1364 (17); 43(3.2) [§]	-
Poli²⁰ 2014 Italy	a. Prospective b. 4.6 years c. Hospital OAC clinic	a. 1089 AF b. 75 (30-94) c. White d. Warfarin	0-1:624 (57); 72.3 ± 15.3 2: 288 (26); 72.0 ± 15.6 >2:177 (16); 68.2 ±16.4	-
Skov¹⁹ 2014 Denmark	a. Prospective b. 1 year c. Hospital OAC clinic	a. 182 AF b. 70.2 [#] c. White d. Warfarin	0-1:105 (58); 76 ≥2: 77 (42); 76	-
Apostolakis⁵ 2013 United Kingdom	a. Retrospective and prospective b. 3.5 years c. Clinical trial (internal-validation)/Hospital OAC clinic (external-validation)	a. 1305 AF b. 69(8)/74(10) c. 8.7%, 19.3 % non-white (internal/external-validation) d. Warfarin	(Internal/External validation) 0: 242 (19); 0.66±0.16/0.7±0.13 1: 413 (32); 0.65±0.18/0.66±0.17 2: 303 (23); 0.63±0.17/0.66±0.16 3:185 (14); 0.59±0.22/0.65±0.17	Internal validation TTR>70:35.7 TTR≤70:64.3 External validation TTR>70:44.1 TTR≤70:55.9

AF: atrial fibrillation; CV: cardiovascular; INR: international normalised ratio; IQR: interquartile range; Max: maximum; MI: myocardial infarction; NVAF: non-valvular atrial fibrillation; OAC: oral anticoagulant/anticoagulation; ROC: area under curve; SD: standard deviation; SAME-TT₂R₂ score: sex (female), age (<60 years, medical history (≥2 of the following: hypertension, diabetes, coronary artery disease or myocardial infarction, peripheral arterial disease, congestive heart failure, previous stroke, pulmonary, hepatic, or renal disease), treatment with interacting drugs (e.g. amiodarone[all 1 point], current tobacco use and race (non-white) [2 points]; TTR: time to therapeutic; TE: thromboembolism; VTE: venous thromboembolism

†TTR presented as ≥70% and <70% not mean TTR; ‡mixed population: White, Black, Asian, other; §number of patients with labile INR, (%); ¶PINRR % (mean ± SD); # no SD or IQR reported; - not reported

Table 3: Baseline characteristics of studies assessing SAME-TT₂R₂ score in AF and VTE cohorts

Patient characteristic, N (%)	Sex (female)	Age <60 y	Hypertension	Diabetes mellitus	Heart failure	Prior stroke/TIA	Peripheral arterial disease	Renal disease	Coronary artery disease	COPD	Current smoking habit	Previous bleeding	Treatment: Amiodarone
PivattoJunior ⁹	113 (43.0)	41 (15.6)	231 (87.8)	108 (41.1)	149 (56.7)	96 (36.5)	25 (9.5)	7 (2.7)	76 (28.9)	36 (13.7)	37 (14.1)	24 (9.1)	26 (9.9)
Kataruka ⁸	1017 (52.3)	1060 (54.6)	-	-	-	-	-	-	-	-	575 (29.6)	-	22 (1.1)
Bernaitis ¹⁶	448 (39.4)	172 (15.1)	677 (59.5)	343 (30.2)	88 (7.7)	45 (4.0)	-	156 (13.7)	271 (23.8)	-	84 (7.4)	-	78 (6.9)
Chan ¹⁵	671 (52.5)	48.0 (3.4)	922 (64.6)	387 (27.1)	367 (25.7)	496 (34.7)	102 (7.1)	2.9 (2.0)	407 (28.5)	-	71.0 (5.0)	-	94 (6.6)
Demelo-Rodriguez ⁶	(50.4)	-	(51.9)	(18.5)	(3.7)	(5.2)	(3.0)	(15.6)	-	(17.0)	(18.5)	-	-
Gorzelak-Pabis ¹⁴	63 (60.6)	-	92 (88.5)	30 (28.8)	72 (69.2)	15 (14.0)	-	-	-	-	20 (19.2)	-	8 (7.7)
Lip ¹³	47 (20.5)	57 (24.9)	206 (90.0)	106 (46.3)	126 (55.0)	26 (11.4)/ 14 (6.1)	31 (13.5)	-	178 (77.7)	-	-	-	46 (20.1) [#]
L-Bejarano ¹²	741 (48.6)	66 (4.3)	1223 (80.2)	473 (31.0)	392.0 (25.7)	209.0 (13.7)	99 (6.5)	92 (6.0)	286 (18.8)	-	100 (6.6)	134 (8.8)	100 (6.6)
Palareti ⁷	698.0 (53.4)	446 (34.1)	678 (51.8)	107 (8.2)	36.0 (2.8)	66 (5.0)	54 (4.1)	73 (5.6)	99.0 (7.6)	-	134 (10.0)	-	15 (1.1)
Proietti ¹¹	1116 (30.5)	72 [§] (66-77)	2812 (76.7)	860 (23.5)	1372 (37.4)	753 (20.5)	-	-	1619 (44.2)	-	334 (9.1)	208 (5.7)	-
Szymanski ¹⁰	79 (37.4)	108 (51.2)		27 (12.8)	8.0 (3.8)	16 (7.6)	-	-	-	-	31.0 (14.7)	-	17 (8.1)
Abumuaileq ¹⁸	306 (33.6)	-	678 (74.4)	220 (24.1)	343 (37.7)	103 (11.3)	92 (10.1)	36 [¶] (4)	127 (13.9)	183 (20.1)	77 (8.5)	115 (12.6)	-
Roldán ¹⁷	237 (53.0)	38 (8.0)	368 (80.0)	141 (31.0)	87 (19.0)	67 (15.0)	-	51 (11.0)	70 (15.0)	50 (11.0)	38 (8.0)	37 (8.0)	72 (16.0)
Ruiz-Ortiz ²	443 (42.0)	-	884 (83.7)	321 (30.4)	235 (22.2)	150 (14.2)	-	153 (14.5)	215 (20.3)	176 (16.7)	76 (7.2)	56 (5.3) ^{††}	102 (9.7)

Gallego²²	494 (51.0)	66 (7.0)	796 (82.0)	249 (26.0)	350 (36.0)	182 (19.0)	-	94 (10.0)	182 (19.0)	-	136 (14.0)	79 (8.0)	-
Lip²¹	3,129 (39)	-	3,405 (42.0)	1,244 (15.0)	4,466 (55.0)	674 (8.0)	-	734 (9.0)	2,434 (30.0)	870 (11.0)	1,053 (13.0)	-	1,670 (35.0)
Poli²⁰	412 (37.8)	61 (5.6)	745 (68.7)	216 (19.9)	268 (24.7)	313 (28.8)	143 (13.2)	-	239 (22.1)	-	181 (16.6)	-	200 (18.4)
Skov¹⁹	54 (29.6)	23 (12.6)	-	-	-	-	-	-	-	-	41 (22.5)	-	27 (14.8)
Apostolakis^{5†}	382 (37.5)	147 (14.4)	692 (67.9)	200 (19.6)	197 (19.3)	130 (12.8)	57 (5.6)	53 (5.2) ^{‡‡}	173 (17.0) ^{§§}	-	64.0 (6.3)	-	129 (12.7)
Apostolakis^{5‡}	157 (67.1)	30.0 (10.5)	234 (81.8)	64 (22.4)	45 (15.7)	30.0 (12.8)	8 (2.8)	2.0 (0.7) ^{‡‡}	44 (15.4) ^{§§}	-	140 (49.0)	-	26 (9.1)

COPD: chronic obstructive pulmonary disease; eGFR: estimated glomerular filtration rate; MI: myocardial infarction; TIA: transient ischemic attack

†Internal validation; ‡external validation; §median age (IQR); ¶eGFR 30 ml/min/1.73m²; [#] antiarrhythmic; ††Major bleed; ‡‡ hepatic/renal disease; §§ history of MI

Table 4: Predictive ability (c-statistics) of SAME-TT₂R₂ for anticoagulation control and clinical events

	Anticoagulation control, c-statistics (95% CI)	Clinical events, c-statistics (95% CI)
PivattoJunior ⁹	TTR $\geq$ 65: 0.612 (0.544-0.681; p=0.002)	-
Kataruka ⁸	TTR<60: 0.61(-) TTR<65: 0.65(-) TTR<70: 0.65 (-)	-
Chan ¹⁵	-	Stroke: 0.54 (0.52-0.57)
Lobos-Bejarano ¹²	TTR $\geq$ 65: 0.562 (0.533-0.592; p<0.001)	-
Palareti ⁷	TTR<65: 0.52 (0.48-0.55; p:0.35)	-
Abumuaileq ¹⁸	PINRR $\leq$ 70: 0.60 (0.56-0.64; p<0.001)	Composite major bleeding, thromboembolic complication or death: 0.57 (0.51-0.62)
Ruiz-Ortiz ²	TTR $\geq$ 65: 0.57 (0.53-0.60; p<0.0005)	-
Gallego ²²	-	Adverse CV event: 0.62 (0.57-0.68; p<0.001) Bleeding: 0.55 (0.49-0.62; p=0.117) All-cause mortality: 0.62 (0.55-0.68; p<0.001)
Lip ²¹	Labile INR: 0.589 (0.574-0.603)	Stroke/TE: 0.561 (0.547-0.575) Severe bleeding: 0.552 (0.537-0.566) Major BARC bleeding: 0.574 (0.560-0.589) Death: 0.544 (0.530-0.559)
Apostolakis ⁵	TTR 31% internal 0.72 (0.64-0.795) TTR 36% external 0.70 (0.57-0.82)	-

BARC: Bleeding Academic Research Consortium; CV: cardiovascular; INR: international normalised ratio; PINRR: percentage of INR in range; TE: thromboembolism; TTR: time in therapeutic range; - not reported

Figure legends:

Figure 1: Selection of studies for inclusion – PRISMA flowchart

Figure 2: Mean TTR vs. SAMe-TT₂R₂ categories in validation studies

Legend: SAMe-TT₂R₂ categories: black= score 0-1; grey= score of 2; white= score >2

Figure 3: Predictive ability (c-statistics and 95% confidence intervals) of SAMe-TT₂R₂ and anticoagulation control in validation studies