Mesocosm experiments reveal the direction of groundwater-surface water exchange alters the hyporheic refuge capacity under warming scenarios

Folegot, Silvia; Krause, Stefan; Mons, Raphael; Hannah, David M.; Datry, Thibault

DOI:

[10.1111/fwb.13049](https://doi.org/10.1111/fwb.13049)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Folegot, S, Krause, S, Mons, R, Hannah, DM & Datry, T 2017, 'Mesocosm experiments reveal the direction of groundwater-surface water exchange alters the hyporheic refuge capacity under warming scenarios', *Freshwater Biology*. <https://doi.org/10.1111/fwb.13049>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Folegot S, Krause S, Mons R, Hannah DM, Datry T. Mesocosm experiments reveal the direction of groundwater-surface water exchange alters the hyporheic refuge capacity under warming scenarios. *Freshwater Biol.* 2017;00:1–13. <https://doi.org/10.1111/fwb.13049>, which has been published in final form at 10.1111/fwb.13049. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Mesocosm experiments reveal the direction of groundwater-surface water exchange alters the hyporheic refuge capacity under warming scenarios

Journal:	<i>Freshwater Biology</i>
Manuscript ID	FWB-P-May-17-0222.R3
Manuscript Type:	Standard Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Folegot, Silvia; University of Birmingham, School of Geography, Earth and Environmental Sciences Krause, Stefan; University of Birmingham, School of Geography, Earth and Environmental Sciences Mons, Raphael; IRSTEA, MALY Hannah, David; University of Birmingham, School of Geography, Earth and Environmental Sciences Datry, Thibault; Irstea, MALY
Keywords:	Climate change < Applied Issues, Conservation / biodiversity < Applied Issues, Hyporheic zone < Habitat, Behaviour < Process / Approach / Methods, Hydrology < Process / Approach / Methods, Crustacea < Taxonomic Group / Assemblage

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Mesocosm experiments reveal the direction of groundwater-surface water exchange**
2 **alters the hyporheic refuge capacity under warming scenarios.**

3 Silvia Folegot¹, Stefan Krause¹, Raphael Mons², David M. Hannah¹ & Thibault Datry²

4 ¹University of Birmingham, School of Geography, Earth and Environmental Sciences, B15 2TT Edgbaston,
5 Birmingham, U.K.

6 ²UR MALY, IRSTEA, 5 rue de la Doua, CS70077, F-69626 Villeurbanne, France

7

8 **Correspondence to:** Silvia Folegot, tel. +44 121 414 5535, fax: +44 121 414 3971, email:
9 SXF356@student.bham.ac.uk

10 **Running title:** the hyporheic zone and climate change.

11 **Keywords:** climate change, hyporheic zone, hydrologic exchange, thermal refuge, riverine
12 biodiversity.

Summary

1. Climate change is expected to affect hydrologic and thermal regimes of river ecosystems. During dry periods when river flows decrease and water temperatures increase, the hyporheic zone (HZ) can provide a refuge to surface aquatic organisms and enhance the resilience capacity of riverine ecosystems. However, shifts from up- to downwelling flow conditions in the HZ could jeopardize this capacity.
2. Using laboratory mesocosms and high resolution fiber-optic distributed temperature sensing, we explored the combined effects of 5 different increased surface water temperature treatments (from 15 to 27 °C at 3 °C intervals) and the direction of water exchange on the ability of *Gammarus pulex* (Crustacea, Amphipoda) to migrate into the HZ as a response to warming. We determined the survival rates of this ubiquitous hyporheic dweller and its rates of consumption of *Alnus glutinosa* leaf litter in the HZ.
3. Results showed that at increasing surface water temperature leaf litter breakdown was observed at a greater depth in the sediments under downwelling flow conditions, i.e. *Gammarus pulex* migrated deeper into the HZ compared to upwelling conditions, resulting in greater survival rates (64 ± 11 vs. 44 ± 10 %). However, under both upwelling and downwelling conditions, we found evidence for potential use of the hyporheic zone as a thermal refuge for *G. pulex*. Below sediment depths of 25 cm, temperatures remained low (< 22 °C) even when surface waters were at 27 °C, so temperatures deep in the hyporheic zone never exceeded critical thermal thresholds for *G. pulex*.
4. This study provides evidence that alterations to the direction of groundwater-surface water exchange can alter the capacity of the HZ to provide a refuge for benthic invertebrates, thereby affecting the resilience of river communities to warming under climate change.

38 **Introduction**

39 Significant changes in the hydrologic and thermal regimes of rivers are expected to occur
40 under global warming (Webb & Nobilis, 2007; van Vliet *et al.*, 2013), affecting biodiversity
41 and functioning of freshwater ecosystems (Woodward, Perkins & Brown, 2010; Ledger &
42 Milner, 2015; Leigh *et al.*, 2015). Typical consequences include geographical range shifts in
43 animal and plant communities (Walther *et al.*, 2002; Root *et al.*, 2003; Holzinger *et al.*,
44 2008), habitat loss or fragmentation (Mantyka-Pringle, Martin & Rhodes, 2012) and altered
45 food webs interactions (Woodward, Dybkjær, Ólafsson *et al.*, 2010; Kratina *et al.*, 2012;
46 Ledger *et al.*, 2013). As most aquatic organisms are ectotherms, they are highly sensitive to
47 temperature increases (Sibly & Atkinson, 1994; Daufresne *et al.*, 2004; Daufresne,
48 Lengfellner & Sommer, 2009; Vander Vorste *et al.*, 2016a). So changes to river thermal
49 regimes alter freshwater community diversity and composition (Brown, Hannah & Milner,
50 2007; Datry *et al.*, 2014; Leigh *et al.*, 2016).

51 The hyporheic zone (HZ), defined as the saturated interstices below and adjacent to river
52 channels (White, 1993) in which groundwater and surface water mix (Krause, Hannah,
53 Fleckenstein *et al.*, 2011), can provide a refuge for river organisms (Palmer, Bely & Berg,
54 1992; Stubbington, 2012; Vander Vorste *et al.*, 2016a). Refuges, *sensu* Sedell *et al.* (1990),
55 can favour the survival of many riverine species including invertebrates and fish, particularly
56 in a context of global change (Keppel *et al.*, 2015; Ledger & Milner, 2015). Because the HZ
57 is characterized by reduced daily and annual temperature amplitudes compared to surface
58 water (Hannah, Webb & Nobilis, 2008; Krause, Hannah & Blume, 2011), it is a potential
59 refuge for surface river organisms during adverse thermal conditions (Palmer *et al.*, 1992;
60 Stubbington, 2012; Vander Vorste *et al.*, 2016a). Surface and HZ habitats are vertically
61 interconnected by upwelling (exfiltration) and downwelling (infiltration) fluxes of water,
62 solutes and organisms (Brunke *et al.*, 1997; Boulton, Findlay & Marmonier, 1998).

Upwelling conditions reflect water fluxes from the HZ into the surface, whereas downwelling is the infiltration of surface water into the HZ. Water temperatures in the HZ are generally lower than channel water in summer and higher in winter (Evans, Greenwood & Petts, 1995; Arrigoni *et al.*, 2008; Krause, Hannah & Blume, 2011). Therefore, the HZ represents a potential thermal refuge for surface organisms when surface temperatures become unfavourable. Early signals of *Gammarus pulex* actively using the HZ to avoid exposure to elevated temperatures (Wood *et al.*, 2010; Vander Vorste *et al.*, 2016a) or desiccation (Vadher, Stubbington & Wood, 2015; Vander Vorste *et al.*, 2016b) have been detected in natural systems. Hence, the HZ may mitigate the negative effects of climate warming on organisms resilience and associated ecosystem processes, such as organic matter decomposition (Stubbington, 2012; Kawanishi *et al.*, 2013; Vander Vorste *et al.*, 2016a).

In a climate change context, the capacity of the HZ to provide a thermal refuge may be at risk due to shifts in the direction of groundwater-surface water exchange, potentially reducing the resilience of riverine ecosystems. The combination of reduced runoff and greater demand for water resources increases human reliance upon groundwater causing increased pumping and lower groundwater levels (Green *et al.*, 2011; Treidel, Martin-Bordes & Gurdak, 2012; Taylor *et al.*, 2013). Lower groundwater tables contribute less groundwater to river base flow (Fetter, 2001; Sophocleous, 2002), altering interactions between groundwater and surface waters (Krause & Bronstert, 2007; Kløve *et al.*, 2014) and reversing conditions from upwelling to downwelling (Stanley & Valett, 1992; Dole-Olivier & Marmonier, 1992b; Dahm *et al.*, 2003). The consequences of such complex interacting pressures, (warming under climate change, more frequent and extreme events and increased groundwater abstraction), on the refuge capacity of the HZ are still poorly understood (Dole-Olivier, 2011; Stubbington, 2012). On one hand, enhanced downwelling could increase hyporheic water temperatures with heat being propagated deeper into the HZ by additional heat advection

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

88 (Boulton *et al.*, 1998; Malard *et al.*, 2002; Krause, Hannah & Blume, 2011), and this could
89 preclude the HZ from acting as a thermal refuge during warming. On the other hand,
90 downwelling conditions may favour the passive downward migration of aquatic organisms
91 from the surface into the HZ and promote their survival (Dole-Olivier, Marmonier & Beffy,
92 1997; Stubbington, Wood & Reid, 2011). To accurately predict the response of riverine
93 communities and ecosystem processes to climate change, it is crucial to understand how the
94 direction of groundwater-surface water exchange, heat transport and animal behaviour
95 interact and possibly alter the potential capacity of the HZ to act as a refuge.

96 In this study, we addressed the effects of change in the direction of groundwater-surface
97 water exchange on the capacity of the HZ of gravel-bed rivers to act as a thermal refuge for
98 surface organisms. Using laboratory mesocosms, we simulated real ranges of increased
99 surface water temperatures, representing for instance disconnected standing pools associated
100 with stream channel contraction, and we manipulated the direction of water exchange. We
101 tested the following hypotheses:

102 1) The HZ provides a thermal refuge for river organisms when surface water temperature
103 increases because it will remain cooler with a narrower range of temperatures than surface
104 waters, as predicted under climate change in many riverine systems;

105 2) The direction of groundwater-surface water exchange mediates this refuge capacity, which
106 will be lower in downwelling conditions than in upwelling conditions because warmer
107 surface water flow into the HZ will raise the temperature of the HZ under downwelling
108 conditions, but upwelling water will remain cooler than surface waters.

109 **Methods**

110 **Experimental design**

111 We used a set of 10 experimental mesocosms to mimic gravel-bed river HZs and simulated
112 increased surface water temperature and reversed flow direction due to climate change. We
113 applied 5 temperature treatments, from 15 to 27 °C, and 2 contrasting hydrological
114 conditions, comprising upwelling (exfiltration) and downwelling (infiltration) flow (Table 1),
115 and observed the vertical migration of *G. pulex* in response to these treatments. This
116 amphipod was used as a model organism (see details below).

117 We conducted Fiber-Optic Distributed Temperature Sensing (FO-DTS) high-resolution
118 monitoring of vertical temperature profiles and kept dissolved oxygen levels close to
119 saturation to avoid any possible anoxia. To assess the vertical migration of *G. pulex* into the
120 HZ we used rates of *Alnus glutinosa* leaf litter breakdown (Navel *et al.*, 2010; Vander Vorste
121 *et al.*, 2016a; Foucreau *et al.*, 2016). The experiments ran for 15 days and were repeated 3
122 times (n total = 30) within a 4-month period (see details below).

123 **Mesocosm design**

124 The mesocosms were made of opaque PVC, 120 cm high, 25 cm in diameter and filled to the
125 height of 90 cm with washed gravel (sediment size = 10-14 mm), to provide a substrate not
126 limiting to the vertical migration of *G. pulex* (Navel *et al.*, 2010; Vadher *et al.*, 2015) into the
127 HZ (Fig. 1a,b,c). Each mesocosm was divided into two main parts (Fig. 1c); a 30-cm surface
128 zone: 10 cm at the top were left for gas exchange and 20 cm for surface water; and a 90-cm
129 sediment zone representing the HZ. To analyse physical and chemical pore water properties,
130 mesocosms had lateral tubing outlets every 15 cm from -5 cm from the free water-sediment

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

interface to -80 cm depth (6 in total each), screened with 500 µm mesh to prevent *G. pulex* from escaping the mesocosms.

Temperature treatments

We generated five distinct surface water temperatures of 15, 18, 21, 24 and 27 °C to simulate climate-induced warming of rivers water (Table 2). The chosen temperature values spanned the range of temperatures observed *in-situ* (Zwolsman & van Bokhoven, 2007; van Vliet & Zwolsman, 2008) or projected for rivers in temperate regions under future climate change through modelling approaches (Mantua, Tohver & Hamlet, 2010; van Vliet *et al.*, 2013). To heat the water, we used heating cables (0.5 cm diameter) (Hydrokable, Hydor Inc. Sacramento, CA USA) placed onto the sediment surface in the free water column and coiled around the inner wall of the mesocosms (Vander Vorste *et al.*, 2016a). We controlled surface water temperature using an electronic thermostat (± 0.1 °C) (Hobby, Dohse Aquaristik GmbH & Co., Graftschaff, Germany) and kept surface water temperatures constant until the end of the experiment. A 12:12-h light:dark cycle was applied using Grolux (35 W, 8500 K, Sylvania Inc., Noida, India) aquarium lights above mesocosms. Throughout the experiment, we kept mesocosms in a temperature-controlled room (16.4 ± 0.4 °C).

Hydrological treatments

We used peristaltic pumps to generate up- and downwelling conditions. The resulting infiltration rate in the mesocosms was 1.9 L/h (Darcy velocity: 6.7 cm/h) which generated an interstitial water velocity of 22.3 cm/h. Upwelling flow was simulated by pumping continuously dechlorinated tap water (which was kept aerated by air bubblers) from a 1000-L tank into the bottom of the mesocosms (n = 5). Water drained (1.9 L/h) through a 2-cm diameter hole, screened with 500 µm mesh, located 10 cm below the top of each mesocosm. Downwelling flow was simulated by pumping water from the tank into the top of the

mesocosms ($n = 5$) and forcing water to flow through the sediments by pumping out interstitial water from the bottom of the column (1.85 L/h), while 0.05 L/h drained through a 2-cm diameter hole, screened with 500 μm mesh, located 10 cm below the top of each mesocosm.

In each mesocosm water volume (22.8 L) was renewed for both flow paths every 12 hours to avoid any possible hypoxia, particularly for downwelling treatments. We measured dissolved oxygen and temperature in interstitial water twice during each experimental run. They were measured at 3 depths (5, 35 and 80 cm in the HZ) by drawing interstitial water from the outlets and using a portable multi-parameter meter (HQ40D, Hach, Loveland, USA, DO resolution = 0.01 mg/L, temperature = 0.1 $^{\circ}\text{C}$). An air bubbler kept surface water in each mesocosm aerated, and dissolved oxygen concentrations in interstitial water varied between 6.53 and 9.64 mg/L in the HZ.

High resolution temperature sensing profiles

Raman-backscatter Distributed Temperature Sensing (DTS) is being increasingly used in environmental applications including hydrological processes (Selker *et al.*, 2006) because it provides high-resolution, continuous temperature data collection in space and time (Briggs *et al.*, 2012). DTS systems provide temperature measurements along a fiber-optic cable by analysing the ratio of the amplitudes of the temperature-independent backscatter, Stokes, to temperature-dependent anti-Stokes signal of the light pulse emitted by the instrument (Selker *et al.*, 2006; Tyler *et al.*, 2009). The timing of these backscatter returns yields a measure of location (Briggs *et al.*, 2012). The precision of the measurements depends on the accuracy of Stokes/anti-Stokes ratio, and greater signal strength requires longer integration time (Selker *et al.*, 2006).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We continuously monitored vertical temperature profiles in the mesocosms for each of the 15-day experimental runs at high spatial and temporal resolution using 10 high resolution temperature sensing profiles (HRTS) (Briggs *et al.*, 2012), specifically constructed for the purpose. The DTS instrument applied was a Silixa XT-DTS (Silixa, Elstree, UK), having a sampling resolution of 25 cm, suggesting a minimal spatial resolution of > 50 cm along the fiber based on the Nyquist criterion (van de Giesen *et al.*, 2012). A small armoured bend-insensitive fiber-optic temperature sensing cable with stainless steel loose tube containing 2 optical fibers, 1.6 mm diameter (Brugg Kabel AG, Brugg, Switzerland) was wrapped around a PVC pipe (6-cm external diameter hollow PVC pipe, 1.2 m long), pre-threaded at a specific pitch, to create a HRTS with 0.004 m vertical sampling resolution. The PVC pipes were threaded along the central 1 metre of length, leaving the first and last 10 cm of the pipes unthreaded. Each HRTS was placed vertically in the centre of the 10 mesocosms. Temperature values were taken continuously every 2 minutes for the total length of the experiment. Alternate single-ended monitoring mode was adopted (Krause & Blume, 2013), and a dynamic instrument calibration was used, based on matching the temperatures of two separate sections of the fibre in a control bath. Specifically, at both ends of the fibre-optic cable, sections of > 20 metres were coiled and kept at a constant temperature in a 0 °C ice bath during the experiment (Tyler *et al.*, 2009) and their monitoring temperatures matched during calibration to account for potential drift caused by differential loss along the cable length.

The number of HRTS that could be connected in series by splicing the fiber cable together before signal loss occurred was affected by the total number of splices (Tyler *et al.*, 2009). This limited the number of mesocosms that could be employed in a single experimental run. For this reason, based on published literature (Briggs *et al.*, 2012), we limited the number of mesocosms to 10 and repeated the experiment three times using an identical design.

203 Model organism

204 We used *G. pulex* (Amphipoda: Crustacea, Linnaeus, 1758) as a biological model because of
205 its wide distribution and abundance throughout Europe (Graça, Maltby & Calow, 1994;
206 Macneil, Dick & Elwood, 1997). It is a facultative component of the hyporheos (Dole-Olivier
207 & Marmonier, 1992a), able to burrow up to 2 m into deep sediments during adverse surface
208 conditions (Dole-Olivier *et al.*, 1997; Stubbington *et al.*, 2011), is eurythermic (Foucreau *et*
209 *al.*, 2014) and can tolerate moderate hypoxia for several days (Danielopol, 1989). Its crucial
210 role in leaf litter breakdown has been well documented in streams (Graça *et al.*, 1994; Navel
211 *et al.*, 2010; Piscart *et al.*, 2011). Together, these reasons make *G. pulex* a valuable and
212 widely used model for laboratory and environmental change studies (Navel *et al.*, 2010;
213 Foucreau *et al.*, 2014; Vander Vorste *et al.*, 2016a).

214 During the experiment, we twice collected (early March and mid-May 2016) adult amphipods
215 of similar size (5-7 mm) from a first-order stream near Dijon, France (see Vander Vorste *et*
216 *al.* 2016a;b for details). We kept the amphipods in a temperature-controlled room (16.4 ± 0.4
217 $^{\circ}\text{C}$) and allowed them to acclimatize to temperature, water (collected from the same stream as
218 amphipods, pH = 6.99, T = 10.4°C , EC = $527\ \mu\text{S}/\text{cm}$) and food source in aquaria (40 x 22 x
219 25 cm) for two weeks before the start of the experiment (Navel *et al.*, 2010). A thermostatic
220 water pump (TECO, Ravena, Italy) kept water temperature constant ($16.4 \pm 0.4^{\circ}\text{C}$) and air
221 bubblers kept dissolved oxygen concentration near saturation. We fed the amphipods with
222 conditioned alder leaves (*Alnus glutinosa*), their most preferred food source (Graça, Maltby
223 & Calow, 1993a; Friberg & Jacobsen, 1994; Foucreau *et al.*, 2014). At each experimental
224 run, we introduced 120 *G. pulex* into each mesocosm ($3849\ \text{individuals}/\text{m}^3$), representing a
225 density occurring in natural streams (Stubbington *et al.*, 2011; Vander Vorste *et al.*, 2016a), a
226 couple of hours before starting to warm the surface water.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

227 **Assessing *G. pulex* survival rate**

228 We quantified the percentage of individuals alive after 15 days by elutriating the sediments of
229 each mesocosm. Water was removed from the mesocosms, and amphipods were washed out
230 with the water and collected using sieves (500 µm). Wet sediments were then vacuumed
231 (Kärcher WD6 Premium, 2000 W power, 30 L capacity) and mesocosms carefully washed.
232 Prior experiments showed that vacuuming did not kill amphipods. Mesocosm sediments were
233 then placed into separate large plastic cases and carefully elutriated, taking small sediment
234 portions each time. Amphipods found with eyes intact and with no signs of soft tissue
235 breakdown were counted as alive prior to mesocosm deconstruction. Amphipods that did not
236 meet this criterion were considered dead.

237 **Assessing *G. pulex* vertical migration**

238 For each mesocosm, we assessed the average depth to which *G. pulex* migrated by
239 determining (*G. pulex* mediated) leaf litter breakdown rates at different depths in the HZ (Fig.
240 1c, Table 1). Alder leaves, dried at 60 C for 24 h (0.4317 ± 0.0036 g dry mass) with primary
241 veins removed, were enclosed in 7.5 x 8-cm plastic mesh bags (0.8 cm diameter) (n = 6) and
242 positioned in the HZ at 6 different depths for each mesocosm (Fig. 1c). The mesh size
243 allowed amphipods to enter the bags freely and consume leaf litter. To facilitate colonization
244 by fungi and increase leaf palability (Graça, Maltby & Calow, 1993b; Graça *et al.*, 1994), leaf
245 bags were pre-conditioned in aerated stream water for 7 days (Suberkropp & Chauvet, 1995),
246 before being placed into the sediments. To account for microbial leaf litter decomposition,
247 fine mesh leaf bags (500 µm, 7.5 x 6-cm) pre-conditioned in the same way (n = 6) were
248 placed next to the coarse mesh bags at all depths (Foucreau *et al.*, 2016). The mesh size of
249 fine mesh leaf bags excluded *G. pulex* without limiting microbial colonization (Boulton &
250 Boon, 1991). In the same way, we prepared and pre-conditioned 3 additional coarse and 3

251 fine mesh leaf bags in order to correct the initial weight for loss due to handling and leaching
252 of soluble components within 24-h after immersion (Gessner, Chauvet & Dobson, 1999).

253 After each run, leaves from both coarse and fine leaf bags were dried at 60 °C for 24 h and
254 weighed. At each depth, we calculated the net leaf litter breakdown (net LLB) rate as: (final
255 dry coarse leaf mass - initial dry coarse leaf mass corrected for leaching) – (final dry fine leaf
256 mass - initial dry fine leaf mass corrected for leaching).

257 Statistical analysis

258 Temperature vertical patterns in the HZ for down- and upwelling conditions

259 To test our first hypothesis that the HZ provides a thermal refuge for *G. pulex* when surface
260 water temperature increases, we first explored the vertical temperature profiles for each
261 treatment. Secondly, to evaluate whether flow direction and surface water temperature
262 influenced differences in HZ temperature between depth 1 and depth 6 (hereafter ΔT), we
263 used linear mixed effect models with Gaussian error distribution (LME) (Bolker *et al.*, 2009;
264 Öckinger *et al.*, 2010). Run was considered a random effect to account for variability among
265 runs. Temperatures were log-transformed prior to statistical analysis. Linear regressions were
266 performed to determine the significance of the correlations among variables when interaction
267 effects were statistically significant.

268 Survival rates and vertical migration of *G. pulex*

269 To test our second hypothesis that flow direction can impair the capacity of the HZ to provide
270 a refuge when surface water temperature increases, we tested for differences in *G. pulex*
271 survival rates among temperature and flow direction treatments using an LME. The
272 percentage of *G. pulex* found alive at each run was treated as the response variable, and flow
273 and mean surface water temperature were modelled as fixed effects. Run was considered a

random effect. Percentages of *G. pulex* found alive were arcsin-transformed prior to statistical tests to meet the assumption of normality. Subsequently, we tested for differences in vertical migrations of *G. pulex* among treatments. To do so, we first calculated a leaf litter breakdown averaged depth (D) for each mesocosm, as follows:

$$D = \sum_1^6 (\text{net LLB} * \text{depth}) / \sum_1^6 \text{net LLB}$$

D represents the average depth (m) at which *G. pulex* mediated leaf litter breakdown (LLB) was the highest. We then fit a LME to test for differences in D among treatments. Mean D for each mesocosm calculated for each run was treated as the response variable; mean surface water temperature, flow direction and the percentage of *G. pulex* found were modelled as fixed effects. We included in the model the percentage of *G. pulex* found in each mesocosm to account for the influence of the number of amphipods found at each run on D. All statistical analyses were performed using the nlme package (Pinheiro *et al.*, 2016) in R 3.3.1 (R Core Team, 2016).

Results

Is the HZ a thermal refuge when surface water temperature increases?

For every treatment across the 3 runs, temperature was highest in the shallow sediments of the HZ (depth 1, - 5 cm) and strongly decreased from depth 3 (- 35 cm) (Fig. 2). On average, temperature at depth 3 was below 20 °C and ranged from 15.9 ± 0.1°C (18 °C, upwelling treatment) to 19.6 ± 1.2 °C (27 °C, downwelling treatment).

Temperatures in the HZ under downwelling conditions were on average 1.1 ± 0.3 , 2.0 ± 0.3 , 2.5 ± 0.5 and 3.6 ± 0.5 higher for 18, 21, 24 and 27 °C treatments respectively than under upwelling conditions (one-way ANOVA, flow effect, $P < 0.01$). At 15 °C, mean temperatures in the HZ under downwelling flow conditions were not different from those under upwelling conditions (15.9 ± 0.3 and 16.0 ± 0.3 , respectively). Vertical temperature profiles in the HZ varied with flow direction (Table 3, interaction factor, $P < 0.0001$).

When surface water temperature increased, ΔT increased more under downwelling ($R^2 = 0.98$) than upwelling ($R^2 = 0.60$) conditions (Fig. 3). ΔT values ranged from -0.5 to 8.1 °C (mean value: 3.7 ± 0.4 °C) under downwelling flow, and it varied from 1 to 4.1 °C (mean value: 2.7 ± 1.1 °C) under upwelling conditions.

Does the direction of groundwater-surface water exchange affect the capacity for HZ to provide a refuge?

The percentage of amphipods found alive at the end of each run varied with flow direction, but not with surface temperature (Fig. 4, Table 4). On average, 64 ± 11 % of amphipods survived under downwelling conditions, whereas 44 ± 10 % survived under upwelling conditions (Fig. 4). At the end of the experiment, the mean percentage of amphipods found dead was similar between downwelling and upwelling flow condition, 3 ± 3 and 3 ± 2 % respectively (Table 5). The mean percentage of *G. pulex* not found, presumably consumed by conspecifics (due to *G. pulex* propensity for cannibalism), was 43 ± 15 % (Table 5).

D, (leaf litter breakdown averaged depth), increased with surface water temperature (Table 4, $P = 0.012$) and varied with flow direction ($P < 0.0001$), with no significant interaction (Fig. 5, Table 4). In upwelling conditions, D ranged from 0.13 ± 0.1 (15 °C treatment) to 0.22 ± 0.05 m (27 °C treatment), with a mean value of 0.18 ± 0.1 m. In downwelling conditions, D

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ranged from 0.27 ± 0.0 (15 °C treatment) to 0.49 ± 0.1 cm (27 °C treatment), with a mean value of 0.37 ± 0.1 m.

Discussion

Vulnerability of aquatic organisms to global warming has been demonstrated (Verberk & Bilton, 2013; Pyne & Poff, 2017), particularly for those species occupying habitats near the limits of their thermal tolerance, (e.g. in arid regions Stewart, Close, Cook *et al.*, 2013). The identification and conservation of potential refuges has therefore become a priority (Keppel *et al.*, 2012, 2015). By manipulating surface water temperature and the direction of groundwater-surface water exchange to mimic potential climate change effects on the thermal and hydrological regime of HZs, we showed that hyporheic sediments could be a potential refuge for *G. pulex*. Specifically, we found that the survival of *G. pulex* in the HZ under the range of interstitial flow velocity tested is strongly influenced by the direction of groundwater-surface water exchange when surface temperatures increase. However, our hypothesis that downwelling flow areas provide less effective refuges compared to upwelling zones was not supported. These results challenge the current paradigm that upwelling areas provide better refuges for river invertebrates during disturbance. Although we tested the response of only one species, these results show that a more comprehensive understanding is required of the potential consequences of climate change for riverine biodiversity and ecosystem resilience and how to mitigate these effects.

The HZ acts as a thermal refuge

When the temperature of surface water increased, the resulting vertical temperature patterns in the HZ differed between up- and downwelling flow conditions. However, in all treatments the deeper hyporheic sediments remained a potential thermal refuge for *G. pulex*. It is known

that the downward flow of water transports heat from the surface into hyporheic sediments (Constantz & Stonestrom, 2003), and that higher infiltration rates lead to greater advection, deeper penetration and shorter lags of thermal surface signals at a given depth (Clark, Webb & Ladle, 1999; Arrigoni *et al.*, 2008; Constantz, 2008; Krause, Hannah & Blume, 2011). In our case, hyporheic temperatures were steady and not influenced by increased surface water temperature at a depth of 80 cm. Within the range of temperatures tested, similar vertical temperature patterns under downwelling conditions have been reported from previous field studies (Constantz & Stonestrom, 2003; Vogt *et al.*, 2010; Briggs *et al.*, 2012). Upwelling conditions generally provide more stable and cooler temperatures due to upward advection of groundwater and smaller variations in sediment temperature are produced compared to downwelling conditions (Alexander & Caissie, 2003; Constantz & Stonestrom, 2003; Caissie *et al.*, 2014). Similarly, in our mesocosms, increases in surface water temperature were buffered in the shallow sediments of the HZ even at the highest temperature treatment. The simulated upwelling flow had a mean temperature of 15.3 ± 0.3 °C, which is a frequent hyporheic temperature observed *in-situ*, for example in lowland alluvial rivers in the UK (Evans & Petts, 1997; Krause, Hannah & Blume, 2011), France (Capderrey *et al.*, 2013) and within the range of temperatures observed in an anthropogenic channel in Germany (Schmidt, Bayer-Raich & Schirmer, 2006). Although heat propagated deeper into the HZ under down- than upwelling conditions, for all treatments, at sediment depths below 25 cm the hyporheic temperature was < 22°C, providing a potential thermal refuge for even the most sensitive aquatic invertebrates like Ephemeroptera (mayflies) (Stewart, Close, Cook *et al.*, 2013).

The use of the HZ by *G. pulex* when surface water temperature increases

Even when surface temperature increased up to 27°C, far above the upper limit of the thermal window for *G. pulex* (10 - 20 °C, Maazouzi *et al.*, 2011), there was no significant effect of temperature on *G. pulex* survival rate, suggesting that the HZ successfully provided a thermal

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

refuge. *G. pulex* is known to be an active vertical crawler (Elser, 2001) and it has been found in hyporheic sediments during spates under downwelling conditions (Marmonier & des Châtelliers, 1991; Dole-Olivier & Marmonier, 1992a), low flow (Stubbington *et al.*, 2011) and drying events (Wood *et al.*, 2010). In our mesocosms, we created optimal physical conditions for observing such vertical migration behaviour because of the porous gravel matrix, the absence of fine sediments clogging interstices, sufficient interstitial dissolved oxygen concentrations and food resources available at different depths. Under both up- and downwelling conditions we found evidence that *G. pulex* used the HZ to avoid increased surface water temperatures but no evidence that these increased temperatures led to lower survival rates. This indicates that the HZ acts as a refuge under flow in both directions. In addition to the findings reported in recent laboratory studies (Vadher *et al.*, 2015; Vander Vorste *et al.*, 2016a b), this study shows that the HZ may also act as a thermal refuge under both up- and downwelling conditions and therefore its potential capacity to mitigate the negative effects of climate change on river ecosystems.

The direction of groundwater-surface water exchange flow affects *G. pulex* success in using the HZ as a thermal refuge

The direction of groundwater-surface water exchange influenced the survival of *G. pulex* and its use of the HZ. Across the temperature treatments, *G. pulex* survival rates were always higher under downwelling (64 ± 11 %) compared to upwelling (44 ± 10 %) conditions. For upwelling conditions, survival rates were in the range of those reported by Vander Vorste *et al.* (2016a). Surprisingly, downwelling conditions seemed to better promote the survival of *G. pulex*. This result is in contrast to the assumption that upwelling zones represent thermal refuges during unfavourable surface conditions due to the upwelling of cool groundwater (Malard *et al.*, 2002; Dole-Olivier, 2011; Stubbington, 2012).

Higher survival rates under downwelling conditions corresponded to a deeper migration into the HZ by *G. pulex* compared to upwelling conditions. The average depth at which most of the leaf litter was consumed by *G. pulex* increased with surface water temperature, but was always higher under downwelling than upwelling conditions. While higher temperatures flowing into the HZ with downwelling water triggered the vertical migration of *G. pulex* deeper into the sediments, upwelling flow seemed to constrain habitat availability resulting in more organisms occupying shallow hyporheic sediments. If available habitat was constrained to the shallow hyporheic sediments, biotic interactions might have intensified as competition for food resources (leaf litter) and space increased, and organisms were exposed to high temperatures. These factors could explain the lower survival rates found for upwelling conditions, also corroborated by the fact that the mean percentage of organisms that disappeared at the end of the experiment under upwelling was higher than under downwelling conditions; we assumed that missing amphipods were the victims of cannibalism, commonly observed when *G. pulex* is under stress (Dick, 1995; McGrath *et al.*, 2007; Vander Vorste *et al.*, 2016a).

Potentially, the relatively high hyporheic water velocities used here might have prevented *G. pulex* from moving against the flow direction in upwelling water, whilst favouring downwards migration under downwelling conditions. Interstitial water velocity was ~ 22.3 cm/h, slightly higher than the one generated in previous mesocosms experiments (Mermillod-Blondin, Mauclaire & Montuelle, 2005; Navel *et al.*, 2010; Vander Vorste *et al.*, 2016a), but in the range of those reported from field surveys (Morrice *et al.*, 2000; Gerech *et al.*, 2011). Stubbington *et al.* (2011) hypothesized that the energetic costs for organisms of long-term position maintenance in upwelling flow could be very high and our results corroborate this hypothesis. In contrast, downwelling conditions may facilitate downwards migration. This likely helped *G. pulex* to avoid lethal temperatures at the surface in the mesocosms and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

413 facilitated access to the thermal refuge in the HZ. Consequently, the broader range of depths
414 accessed by *G. pulex* under downwelling condition as indicated by the leaf litter breakdown
415 rates could reveal that organisms were actively moving up and down in the HZ. Organisms
416 might have used the flow to move deeper into the HZ to escape warmer surface temperatures,
417 but could also have moved against the flow (positive rheotaxis) to compensate downstream
418 drift (Hughes, 1970). This seems not to have happened under upwelling conditions, probably
419 because of the higher metabolic costs required. In natural systems upwelling zones are often
420 characterized by depleted dissolved oxygen levels (Dole-Olivier, 2011) which may also
421 decrease the refuge potential of the HZ. Further exploration of the capacity of the HZ to
422 enhance the resilience of riverine biodiversity is needed because the responses by individual
423 species to changing climate vary depending on species traits and interacting drivers of change
424 (Chen *et al.*, 2011), and because these results may not hold in the face of more severe
425 warming that, even in the HZ, exceeds the thermal tolerances of organisms.

426 Determining the complex relationships between groundwater-surface water exchange and
427 organismal behaviour under climate change pressure will require further analysis to advance
428 our understanding of the use of the HZ as a refuge. Indeed, to date, most ecological research
429 seems to have overlooked the eco-hydraulics of HZs, focusing more on how the
430 physicochemistry and biotic interactions shape hyporheic communities. Although laboratory
431 experiments simplify reality (*e.g.* one taxon, controlled conditions), the use of mesocosms
432 provided useful insights for understanding organismal responses to interacting factors linked
433 to climate change which would have been virtually impossible to disentangle in the natural
434 environment. As a next step, the novel experimental design applied in this study can be
435 replicated and refined to recreate more realistic mesocosms conditions (Ledger *et al.*, 2009;
436 Stewart, Dossena, Bohan *et al.*, 2013) where for instance water quality mimics that of natural

437 systems and thus differs between up- and downwelling conditions and sediment grain size
438 distribution is more heterogeneous.

439 HZs could provide thermal refuges for some surface-dwelling organisms when vertical
440 connectivity is efficient, enabling the HZ to contribute strongly to the survival and resilience
441 of surface species in a changing climate. Our results indicate that downwelling conditions
442 might promote the use of different depths of the HZ by *G. pulex* even when surface water
443 temperatures increase up to 27 °C. However, the combination of increased temperature and
444 shifts between up- and downwelling conditions can jeopardize this refuge capacity. These
445 results show the need to develop a landscape perspective of the HZ in rivers (Malard *et al.*,
446 2002) and call for additional field surveys to gain a better understanding of how hydrological
447 conditions, and their temporal shifts, can influence riverine communities and ecosystem
448 resilience. Additional laboratory experiments addressing the effects of altered vertical
449 connectivity in a context of climate change where increased surface temperature, drying
450 events and increased biotic interactions occur represents a promising research avenue for
451 developing efficient tools and guidelines to manage river ecosystems.

452 **Aknowledgments**

453 The research was funded by the European Climate Knowledge and Innovation Community
454 (EIT - Climate KIC) fellowship “A risk assessment framework for quantifying drought
455 impacts on thermal and water quality extremes” and the Leverhulme Trust International
456 Network Grant (IN-2013-042) “Where rivers, groundwater and disciplines meet: a hyporheic
457 research network”. SF wishes to thank Martin Briggs (USGS) for his technical support during
458 the design of the HRTS and Stephen West (Biosciences Workshop, University of
459 Birmingham) for his help while constructing them. Grateful thanks to all UR MALY team,
460 and particularly to Nicolas Lamouroux for his critical input and help with the statistics, Clara

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

461 Mendoza-Lera, Maxence Forcellini, Guillaume Le Goff, Bertrand Launay and Laura Plichard
462 for suggestions about the experimental design and help in the laboratory. Ross Vander Vorste
463 and Paul Wood made insightful and constructive suggestions on an earlier draft of this
464 manuscript.

Copy for Review

References

- Alexander M.D. & Caissie D. (2003) Variability and comparison of hyporheic water temperatures and seepage fluxes in a small Atlantic salmon stream. *Ground Water* **41**, 72–82.
- Arrigoni A.S., Poole G.C., Mertes L.A.K., O'Daniel S.J., Woessner W.W. & Thomas S.A. (2008) Buffered, lagged, or cooled? Disentangling hyporheic influences on temperature cycles in stream channels. *Water Resources Research* **44**, 1–13.
- Bolker B.M., Brooks M.E., Clark C.J., Geange S.W., Poulsen J.R., Stevens M.H.H., *et al.* (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology and Evolution* **24**, 127–135.
- Boulton A. & Boon P. (1991) A review of methodology used to measure leaf litter decomposition in lotic environments: time to turn over an old leaf? *Marine and Freshwater Research* **42**, 1.
- Boulton A., Findlay S. & Marmonier P. (1998) The functional significance of the hyporheic zone in streams and rivers. *Annual Review of Ecology and Systematic* **29**, 59–81.
- Briggs M.A., Lautz L.K., McKenzie J.M., Gordon R.P. & Hare D.K. (2012) Using high-resolution distributed temperature sensing to quantify spatial and temporal variability in vertical hyporheic flux. *Water Resources Research* **48**, 1–16.
- Brown L.E., Hannah D.M. & Milner A.M. (2007) Vulnerability of alpine stream biodiversity to shrinking glaciers and snowpacks. *Global Change Biology* **13**, 958–966.
- Brunke M., Gonser T., Brunke M., Gonser T. & Gonser T. (1997) The ecological significance of exchange processes between rivers and groundwater. *Freshwater biology* **37**, 1–33.
- Caissie D., Kurylyk B.L., St-hilaire A., El-jabi N. & Macquarrie K.T.B. (2014) Streambed

1
2
3 488 temperature dynamics and corresponding heat fluxes in small streams experiencing
4
5 489 seasonal ice cover. *Journal of Hydrology* **519**, 1441–1452.
6
7
8 490 Capderrey C., Datry T., Foulquier A., Claret C. & Malard F. (2013) Invertebrate distribution
9
10 491 across nested geomorphic features in braided-river landscapes. *Freshwater Science* **32**,
11
12 492 1188–1204.
13
14
15 493 Chen I.-C., Hill J.K., Ohlemüller R., Roy D.B. & Thomas C.D. (2011) Rapid range shifts of
16
17 494 species associated with high levels of climate warming. *Science* **333**, 1024 LP-1026.
18
19
20 495 Clark E., Webb B. & Ladle M. (1999) Microthermal gradients and ecological implications in
21
22 496 Dorset rivers. *Hydrological Processes* **13**, 423–438.
23
24
25 497 Constantz J. (2008) Heat as a tracer to determine streambed water exchanges. *Water*
26
27 498 *Resources Research* **44**, W00D10.
28
29
30 499 Constantz J. & Stonestrom D.A. (2003) *Heat as a tracer of water movement near streams, in*
31
32 500 *Heat as a Tool for Studying the Movement of Ground Water Near Streams - Circular*
33
34 501 *1260*. Reston, Virginia.
35
36
37 502 Dahm C.N., Baker M.A., Moore D.I. & Thibault J.R. (2003) Coupled biogeochemical and
38
39 503 hydrological responses of streams and rivers to drought. *Freshwater Biology* **48**, 1219–
40
41 504 1231.
42
43
44 505 Danielopol D.L. (1989) Groundwater Fauna Associated with Riverine Aquifers. *Journal of*
45
46 506 *the North American Benthological Society* **8**, 18–35.
47
48
49 507 Datry T., Larned S.T., Fritz K.M., Bogan M.T., Wood P.J., Meyer E.I., *et al.* (2014) Broad-
50
51 508 scale patterns of invertebrate richness and community composition in temporary rivers:
52
53 509 effects of flow intermittence. *Ecography* **37**, 94–104.
54
55
56 510 Daufresne M., Lengfellner K. & Sommer U. (2009) Global warming benefits the small in
57
58
59
60

- 1
2
3 511 aquatic ecosystems. *Proceedings Of The National Academy Of Sciences Of The United*
4
5 512 *States Of America* **106**, 12788–12793.
6
7
8 513 Daufresne M., Roger M.C., Capra H. & Lamouroux N. (2004) Long-term changes within the
9
10 514 invertebrate and fish communities of the Upper Rhône River: Effects of climatic factors.
11
12 515 *Global Change Biology* **10**, 124–140.
13
14
15 516 Dick J.T.A. (1995) The cannibalistic behaviour of two *Gammarus* species (Crustacea:
16
17 517 Amphipoda). *Journal of Zoology* **236**, 697–706.
18
19
20 518 Dole-Olivier M.-J. (2011) The hyporheic refuge hypothesis reconsidered: A review of
21
22 519 hydrological aspects. *Marine and Freshwater Research* **62**, 1281–1302.
23
24
25 520 Dole-Olivier M.J. & Marmonier P. (1992a) Effects of spates on the vertical distribution of the
26
27 521 interstitial community. *Hydrobiologia* **230**, 49–61.
28
29
30 522 Dole-Olivier M.J. & Marmonier P. (1992b) Patch distribution of interstitial communities:
31
32 523 prevailing factors. *Freshwater Biology* **27**, 177–191.
33
34
35 524 Dole-Olivier M.-J., Marmonier P. & Beffy J.-L. (1997) Response of invertebrates to lotic
36
37 525 disturbance: Is the hyporheic zone a patchy refugium? *Freshwater Biology* **37**, 257–276.
38
39
40 526 Elser P. (2001) Assessing small-scale directional movements of benthic invertebrates in
41
42 527 streams by using a multidirectional cage trap. *Limnologia - Ecology and Management*
43
44 528 *of Inland Waters* **31**, 119–128.
45
46
47 529 Evans E.C., Greenwood M.T. & Petts G.E. (1995) Thermal profiles within river beds.
48
49 530 *Hydrological Processes* **9**, 19–25.
50
51
52 531 Evans E.C. & Petts G.E. (1997) Hyporheic temperature patterns within riffles. *Hydrological*
53
54 532 *Sciences* **42 (2)**, 199–213.
55
56
57 533 Fetter C.W. (2001) *Applied Hydrogeology*. Prentice-Hall, Upper Sadle River, New Jersey.
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

534 Foucreau N., Cottin D., Piscart C. & Hervant F. (2014) Physiological and metabolic
535 responses to rising temperature in *Gammarus pulex* (Crustacea) populations living under
536 continental or Mediterranean climates. *Comparative Biochemistry and Physiology - Part*
537 *A* **168**, 69–75.

538 Foucreau N., Piscart C., Puijalon S. & Hervant F. (2016) Effects of rising temperature on a
539 functional process: consumption and digestion of leaf litter by a freshwater shredder.
540 *Fundamental and Applied Limnology / Archiv für Hydrobiologie* **187**, 295–306.

541 Friberg N. & Jacobsen D. (1994) Feeding plasticity of two detritivore-shredders. *Freshwater*
542 *Biology* **32**, 133–142.

543 Gerecht K.E., Cardenas M.B., Guswa A.J., Sawyer A.H., Nowinski J.D. & Swanson T.E.
544 (2011) Dynamics of hyporheic flow and heat transport across a bed-to-bank continuum
545 in a large regulated river. *Water Resources Research* **47**, 1–12.

546 Gessner M.O., Chauvet E. & Dobson M. (1999) A perspective on leaf litter breakdown in
547 streams. *Oikos* **85**, 377–384.

548 Graça M.A.S., Maltby L. & Calow P. (1994) Comparative ecology of *Gammarus pulex* (L.)
549 and *Asellus aquaticus* (L.) II: population dynamics and microdistribution. *Hydrobiologia*
550 **281**, 155–62.

551 Graça M.A.S., Maltby L. & Calow P. (1993a) Importance of fungi in the diet of *Gammarus*
552 *pulex* (L.) and *Asellus aquaticus* (L.): II Effects on growth, reproduction and physiology.
553 *Oecologia* **96**, 304–309.

554 Graça M.A.S., Maltby L. & Calow P. (1993b) Importance of fungi in the diet of *Gammarus*
555 *pulex* and *Asellus aquaticus* I: feeding strategies. *Oecologia* **93**, 139–144.

556 Green T.R., Taniguchi M., Kooi H., Gurdak J.J., Allen D.M., Hiscock K.M., *et al.* (2011)
557 Beneath the surface of global change: Impacts of climate change on groundwater.

- 558 *Journal of Hydrology* **405**, 532–560.
- 559 Hannah D.M., Webb B.W. & Nobilis F. (2008) River and stream temperature: Dynamics,
560 processes, models and implications. *Hydrological Processes* **22**, 899–901.
- 561 Holzinger B., Hülber K., Camenisch M. & Grabherr G. (2008) Changes in plant species
562 richness over the last century in the eastern Swiss Alps: Elevational gradient, bedrock
563 effects and migration rates. *Plant Ecology* **195**, 179–196.
- 564 Hughes D.A. (1970) Some factors affecting drift and upstream movements of *Gammarus*
565 *pulex*. *Ecology* **51**, 301–305.
- 566 Kawanishi R., Inoue M., Dohi R., Fujii A. & Miyake Y. (2013) The role of the hyporheic
567 zone for a benthic fish in an intermittent river: A refuge, not a graveyard. *Aquatic*
568 *Sciences* **75**, 425–431.
- 569 Keppel G., Mokany K., Wardell-Johnson G.W., Phillips B.L., Welbergen J.A. & Reside A.E.
570 (2015) The capacity of refugia for conservation planning under climate change.
571 *Frontiers in Ecology and the Environment* **13**, 106–112.
- 572 Keppel G., Van Niel K.P., Wardell-Johnson G.W., Yates C.J., Byrne M., Mucina L., *et al.*
573 (2012) Refugia: identifying and understanding safe havens for biodiversity under
574 climate change. *Global Ecology and Biogeography* **21**, 393–404.
- 575 Kløve B., Ala-Aho P., Bertrand G., Gurdak J.J., Kupfersberger H., Kværner J., *et al.* (2014)
576 Climate change impacts on groundwater and dependent ecosystems. *Journal of*
577 *Hydrology* **518**, 250–266.
- 578 Kratina P., Greig H.S., Thompson P.L., Carvalho-Pereira T.S.A. & Shurin J.B. (2012)
579 Warning modifies trophic cascades and eutrophication in experimental freshwater
580 communities. *Ecology* **93**, 1421–1430.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

581 Krause S. & Blume T. (2013) Impact of seasonal variability and monitoring mode on the
582 adequacy of fiber- optic distributed temperature sensing at aquifer- river interfaces.
583 *Water Resources Research* **49**, 2408–2423.

584 Krause S. & Bronstert A. (2007) The impact of groundwater–surface water interactions on
585 the water balance of a mesoscale lowland river catchment in northeastern Germany.
586 *Hydrological Processes* **21**, 169–184.

587 Krause S., Hannah D. & Blume T. (2011) Interstitial pore-water temperature dynamics across
588 a pool-riffle-pool sequence. *Ecohydrology* **4**, 549–563.

589 Krause S., Hannah D.M., Fleckenstein J.H., Heppell C.M., Kaeser D., Pickup R., *et al.* (2011)
590 Inter-disciplinary perspectives on processes in the hyporheic zone. *Ecohydrology* **4**,
591 481–499.

592 Ledger M., Harris R., Armitage P. & Milner A. (2009) Realism of model ecosystems: an
593 evaluation of physicochemistry and macroinvertebrate assemblages in artificial streams.
594 *Hydrobiologia; The International Journal of Aquatic Sciences* **617**, 91–99.

595 Ledger M.E., Brown L.E., Edwards F.K., Hudson L.N., Milner A.M. & Woodward G. (2013)
596 Extreme climatic events alter aquatic food webs. A Synthesis of evidence from a
597 mesocosm drought experiment. *Advances in Ecological Research* **48**, 343–395.

598 Ledger M.E. & Milner A.M. (2015) Extreme events in running waters. *Freshwater Biology*
599 **60**, 2455–2460.

600 Leigh C., Bonada N., Boulton A.J., Hugueny B., Larned S.T., Vander Vorste R., *et al.* (2016)
601 Invertebrate assemblage responses and the dual roles of resistance and resilience to
602 drying in intermittent rivers. *Aquatic Sciences* **78**, 291–301.

603 Leigh C., Bush A., Harrison E.T., Ho S.S., Luke L., Rolls R.J., *et al.* (2015) Ecological
604 effects of extreme climatic events on riverine ecosystems: insights from Australia.

- 605 *Freshwater Biology* **60**, 2620–2638.
- 606 Maazouzi C., Piscart C., Legier F. & Hervant F. (2011) Ecophysiological responses to
607 temperature of the “ killer shrimp” *Dikerogammarus villosus*: Is the invader really
608 stronger than the native *Gammarus pulex*? *Comparative Biochemistry and Physiology*,
609 *Part A* **159**, 268–274.
- 610 Macneil C., Dick J.T.A. & Elwood R.W. (1997) The trophic ecology of freshwater
611 *Gammarus* spp. (Crustacea:Amphipoda): problems and perspectives concerning the
612 functional feeding group concept. *Biological Reviews* **72**, 349–364.
- 613 Malard F., Tockner K., Dole-Olivier M.-J. & Ward J. V. (2002) A landscape perspective of
614 surface-subsurface hydrological exchanges in river corridors. *Freshwater Biology* **47**,
615 621–640.
- 616 Mantua N., Tohver I. & Hamlet A. (2010) Climate change impacts on streamflow extremes
617 and summertime stream temperature and their possible consequences for freshwater
618 salmon habitat in Washington State. *Climatic Change* **102**, 187–223.
- 619 Mantyka-Pringle C.S., Martin T.G. & Rhodes J.R. (2012) Interactions between climate and
620 habitat loss effects on biodiversity: A systematic review and meta-analysis. *Global*
621 *Change Biology* **18**, 1239–1252.
- 622 Marmonier P. & des Châtelliers M.C. (1991) Effects of spates on interstitial assemblages of
623 the Rhône River. Importance of spatial heterogeneity. *Hydrobiologia* **210**, 243–251.
- 624 McGrath K.E., Peeters E.T.H.M., Beijer J.A.J. & Scheffer M. (2007) Habitat-mediated
625 cannibalism and microhabitat restriction in the stream invertebrate *Gammarus pulex*.
626 *Hydrobiologia* **589**, 155–164.
- 627 Mermillod-Blondin F., Mauclaire L. & Montuelle B. (2005) Use of slow filtration columns to
628 assess oxygen respiration, consumption of dissolved organic carbon, nitrogen

1
2
3 629 transformations, and microbial parameters in hyporheic sediments. *Water Research* **39**,
4
5 630 1687–1698.
6
7
8 631 Morrice J.A., Dahm C.N., Valett H.M., Unnikrishna P. V. & Campana M.E. (2000) Terminal
9
10 632 electron accepting processes in the alluvial sediments of a headwater stream. *Journal of*
11
12 633 *the North American Benthological Society* **19**, 593–608.
13
14
15 634 Navel S., Mermillod-Blondin F., Montuelle B., Chauvet E., Simon L., Piscart C., *et al.* (2010)
16
17 635 Interactions between fauna and sediment control the breakdown of plant matter in river
18
19 636 sediments. *Freshwater Biology* **55**, 753–766.
20
21
22 637 Öckinger E., Schweiger O., Crist T.O., Debinski D.M., Krauss J., Kuussaari M., *et al.* (2010)
23
24 638 Life-history traits predict species responses to habitat area and isolation: A cross-
25
26 639 continental synthesis. *Ecology Letters* **13**, 969–979.
27
28
29 640 Palmer M., Bely A.E. & Berg K.E. (1992) Response of invertebrates to lotic disturbance: A
30
31 641 test of the hyporheic refuge. *Oecologia* **89**, 182–194.
32
33
34 642 Pinheiro J., Bates D., DebRoy S., D. S. & R Core Team (2016) nlme: Linear and nonlinear
35
36 643 mixed effects models.
37
38
39 644 Piscart C., Navel S., Maazouzi C., Montuelle B., Cornut J., Mermillod-Blondin F., *et al.*
40
41 645 (2011) Leaf litter recycling in benthic and hyporheic layers in agricultural streams with
42
43 646 different types of land use. *Science of the Total Environment* **409**, 4373–4380.
44
45
46 647 Pyne M.I. & Poff N.L.R. (2017) Vulnerability of stream community composition and
47
48 648 function to projected thermal warming and hydrologic change across ecoregions in the
49
50 649 western United States. *Global Change Biology* **23**, 77–93.
51
52
53 650 R Core Team (2016) R: A language and environment for statistical computing.
54
55
56 651 Root T., Price J., Hall K. & Schneider S. (2003) Fingerprints of global warming on wild
57
58
59
60

- animals and plants. *Nature* **421**, 57–60.
- Schmidt C., Bayer-Raich M. & Schirmer M. (2006) Characterization of spatial heterogeneity of groundwater-stream water interactions using multiple depth streambed temperature measurements at the reach scale. *Hydrology and Earth System Sciences* **10**, 849–859.
- Sedell J.R., Reeves G.H., Hauer F.R., Stanford J.A. & Hawkins C.P. (1990) Role of refugia in recovery from disturbances: Modern fragmented and disconnected river systems. *Environmental Management* **14**, 711–724.
- Selker J.S., Thévenaz L., Huwald H., Mallet A., Luxemburg W., Giesen V. De, *et al.* (2006) Distributed fiber optic temperature sensing for hydrologic systems. *Water Resources Research* **42**.
- Sibly R.M. & Atkinson D. (1994) How rearing temperature affects optimal adult size in ectotherms. *Functional Ecology* **8**, 486–493.
- Stanley E.H. & Valett H.M. (1992) Interactions between drying and the hyporheic zone of a desert stream. In: *Global Climate Change and Freshwater Ecosystems*. (Eds P. Firth & S.G. Fisher), pp. 234–249. Springer New York, New York, NY.
- Stewart B.A., Close P.G., Cook P.A. & Davies P.M. (2013) Upper thermal tolerances of key taxonomic groups of stream invertebrates. *Hydrobiologia* **718**, 131–140.
- Stewart R.I.A., Dossena M., Bohan D.A., Jeppesen E., Kordas R.L., Ledger M.E., *et al.* (2013) *Mesocosm Experiments as a Tool for Ecological Climate-Change Research*, 1st edn. Elsevier Ltd.
- Stubbington R. (2012) The hyporheic zone as an invertebrate refuge: A review of variability in space, time, taxa and behaviour. *Marine and Freshwater Research* **63**, 293–311.
- Stubbington R., Wood P.J. & Reid I. (2011) Spatial variability in the hyporheic zone

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

675 refugium of temporary streams. *Aquatic Sciences* **73**, 499–511.

676 Suberkropp K. & Chauvet E. (1995) Regulation of leaf breakdown by fungi in streams:
677 Influences of water chemistry. *Ecology* **76**, 1433–1445.

678 Taylor R.G., Scanlon B., Döll P., Rodell M., van Beek R., Wada Y., *et al.* (2013) Ground
679 water and climate change. *Nature Climate Change* **3**, 322–329.

680 Treidel H., Martin-Bordes J.L. & Gurdak J.J. (2012) *Climate change effects on groundwater*
681 *resources: a global synthesis of findings and recommendations*. CRC Press/Balkema,
682 Leiden, The Netherlands.

683 Tyler S.W., Selker J.S., Hausner M.B., Hatch C.E., Torgersen T., Thodal C.E., *et al.* (2009)
684 Environmental temperature sensing using Raman spectra DTS fiber-optic methods.
685 *Water Resources Research* **45**, W00D23.

686 Vadher A.N., Stubbington R. & Wood P.J. (2015) Fine sediment reduces vertical migrations
687 of *Gammarus pulex* (Crustacea: Amphipoda) in response to surface water loss.
688 *Hydrobiologia* **753**, 61–71.

689 van de Giesen N.C., Steele Dunne S., Jansen J., Hoes O., Hausner M., Tyler S., *et al.* (2012)
690 Double-ended calibration of fiber-optic Raman spectra distributed temperature sensing
691 data. *Sensors* **12**, 5471–5485.

692 van Vliet M.T.H., Franssen W.H.P., Yearsley J.R., Ludwig F., Haddeland I., Lettenmaier
693 D.P., *et al.* (2013) Global river discharge and water temperature under climate change.
694 *Global Environmental Change* **23**, 450–464.

695 van Vliet M.T.H. & Zwolsman J.J.G. (2008) Impact of summer droughts on the water quality
696 of the Meuse river. *Journal of Hydrology* **353**, 1–17.

697 Vander Vorste R., Mermillod-Blondin F., Hervant F., Mons R. & Datry T. (2016a)

- 698 *Gammarus pulex* (Crustacea: Amphipoda) avoids increasing water temperature and
699 intraspecific competition through vertical migration into the hyporheic zone: a
700 mesocosm experiment. *Aquatic Sciences* **79**, 1–11.
- 701 Vander Vorste R., Mermillod-Blondin F., Hervant F., Mons R., Forcellini M. & Datry T.
702 (2016b) Increased depth to the water table during river drying decreases the resilience of
703 *Gammarus pulex* and alters ecosystem function. *Ecohydrology* **9**, 1177–1186.
- 704 Verberk W.C.E.P. & Bilton D.T. (2013) Respiratory control in aquatic insects dictates their
705 vulnerability to global warming. *Biology Letters* **9**, 20130473–20130473.
- 706 Vogt T., Schneider P., Hahn-Woernle L. & Cirpka O.A. (2010) Estimation of seepage rates in
707 a losing stream by means of fiber-optic high-resolution vertical temperature profiling.
708 *Journal of Hydrology* **380**, 154–164.
- 709 Walther G.R., Post E., Convey P., Menzel A., Parmesan C., Beebee T.J.C., *et al.* (2002)
710 Ecological responses to recent climate change. *Nature* **416**, 389–395.
- 711 Webb B.W. & Nobilis F. (2007) Long-term changes in river temperature and the influence of
712 climatic and hydrological factors. *Hydrological Sciences Journal* **52**, 74–85.
- 713 White D.S. (1993) Perspectives on defining and delineating hyporheic zones. *Journal of the*
714 *North American Benthological Society* **12**, 61–6961.
- 715 Wood P.J., Boulton A.J., Little S. & Stubbington R. (2010) Is the hyporheic zone a refugium
716 for aquatic macroinvertebrates during severe low flow conditions? *Fundamental and*
717 *Applied Limnology* **176**, 377–390.
- 718 Woodward G., Dybkjær J.B., Ólafsson J.S., Gíslason G.M., Hannesdóttir E.R. & Friberg N.
719 (2010) Sentinel systems on the razor's edge: Effects of warming on Arctic geothermal
720 stream ecosystems. *Global Change Biology* **16**, 1979–1991.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Woodward G., Perkins D.M. & Brown L.E. (2010) Climate change and freshwater ecosystems: impacts across multiple levels of organization. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **365**, 2093–2106.

Zwolsman J.J.G. & van Bokhoven A.J. (2007) Impact of summer droughts on water quality of the Rhine River - a preview of climate change? *Water Science and Technology* **56**.

Tables

Treatment	N ^o of levels	Labels
Flow direction	2	up down
Surface water temperature	5	15 °C 18 °C 21 °C 24 °C 27 °C
Leaf litter breakdown (LLB) in the HZ	6	Depth 1 = 5 cm Depth 2 = 20 cm Depth 3 = 35 cm Depth 4 = 50 cm Depth 5 = 65 cm Depth 6 = 80 cm

Table 1. Overview of 2 flow, 5 temperature and 6 leaf litter breakdown (LLB) treatments generated in the experiment. The surface water temperature treatments chosen represented real or projected water temperature values for rivers under global warming.

Surface water T. (°C)	Flow direction	Mean (± SD) surface water T (°C) measured (n=3)
15	up	16.4 ± 0.6
15	down	15.5 ± 0.5
18	up	17.8 ± 0.2
18	down	18.0 ± 0.1
21	up	20.7 ± 0.3
21	down	20.8 ± 0.1
24	up	23.8 ± 0.0
24	down	23.6 ± 0.1
27	up	26.6 ± 0.3
27	down	26.8 ± 0.2

Table 2. Mean value (± SD) for surface water temperature (°C) as given by the high resolution temperature sensing (HRTS) profiles for each temperature and flow direction treatment during the three experimental runs.

Dependent variable	Factor	d.f.	F-value	P-value
ΔT	Flow direction (Flow)	1	14.75	0.1465
	Mean surface water T measured (T)	1	285.16	<.0001
	Flow x T	1	49.69346	<.0001

Table 3. Linear mixed effect model (LME) analysis results for temperature differences between deep and shallow hyporheic sediments (ΔT , °C) associated with flow direction and measured mean surface water temperature and the interaction between these factors.

Dependent variable	Factor	d.f.	F-value	P-value
% <i>G. pulex</i> found alive	Flow direction (Flow)	1	131.88	<.0001
	Mean surface water T measured (T)	1	0.11	0.7407
	Flow x T	1	1.51	0.2304
Leaf litter breakdown averaged depth (D)	Flow direction (Flow)	1	31.34	<.0001
	Mean surface water T measured (T)	1	7.67	0.0118
	% <i>G. pulex</i> found	1	0.03	0.8663
	Flow x T	1	1.77	0.1988
	Flow x % <i>G. pulex</i> found	1	0.16	0.6950
	T x % <i>G. pulex</i> found	1	0.46	0.5052
	Flow x T x % <i>G. pulex</i> found	1	0.05	0.8280

Table 4. Linear mixed effect model (LME) analysis results for *G. pulex* survival rates associated with flow direction and measured mean surface water temperature and the interaction between these factors; LME analysis results for leaf litter breakdown averaged depths associated with flow direction, measured mean surface water temperature and the percentage of organisms found alive and the interactions between these factors.

Surface water T. (°C)	Flow direction	% <i>G. pulex</i> found (n = 3)		% <i>G. pulex</i> not found (mean ± SD, n = 3)
		Alive (mean ± SD)	Dead (mean ± SD)	
15	up	46 ± 5	3 ± 1	50 ± 10
15	down	62 ± 13	6 ± 4	32 ± 5
18	up	47 ± 15	3 ± 2	50 ± 15
18	down	60 ± 8	4 ± 3	36 ± 8
21	up	48 ± 14	2 ± 2	50 ± 17
21	down	69 ± 17	2 ± 1	29 ± 13
24	up	38 ± 3	2 ± 1	61 ± 14
24	down	67 ± 16	3 ± 3	31 ± 2
27	up	41 ± 9	3 ± 2	57 ± 10
27	down	63 ± 8	3 ± 1	34 ± 6

Table 5. Mean (± SD) percentage of *G. pulex* found with distinction between alive and dead organisms and mean (± SD) percentage of organisms not found for each temperature and flow direction treatment.

Figure 1. Mesocosm design with outside view (a), details from the inside (b) and mesocosm schematic representation indicating the dimensions of the surface and HZ, with the infiltration rates for down- and upwelling flow direction and the position of the fine and coarse leaf litter bags in the hyporheic sediments (c).

299x424mm (300 x 300 DPI)

Figure 2. Mean temperature values with standard deviation (n = 3) in the HZ for both up-and downwelling flow treatments at increasing surface water temperature. Dashed horizontal line represents location of the free water-sediment interface.

152x60mm (300 x 300 DPI)

Figure 3. Median of the temperature differences ($\pm$ SD) between deep and shallow hyporheic sediments (ΔT , $^{\circ}C$) for both down-and upwelling flow conditions at increasing surface water temperature.

152x114mm (300 x 300 DPI)

Figure 4. Mean ($\pm$ SD) percentage of *G. pulex* found alive (arcsin-transformed) for each surface water temperature and flow direction treatments.

152x114mm (300 x 300 DPI)

Figure 5. Median of the leaf litter breakdown averaged depths (m) for both down-and upwelling flow conditions at increasing surface water temperature.

152x114mm (300 x 300 DPI)