Some recent progress on sharp Fourier restriction theory

Foschi, Damiano; Silva, Diogo Oliveira e

DOI:

[10.1007/s10476-017-0306-2](https://doi.org/10.1007/s10476-017-0306-2)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Foschi, D & Silva, DOE 2017, 'Some recent progress on sharp Fourier restriction theory', *Analysis Mathematica*, vol. 43, no. 2, pp. 241–265. <https://doi.org/10.1007/s10476-017-0306-2>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is a post-peer-review, pre-copyedit version of an article published in *Analysis Mathematica*. The final authenticated version is available online at: <http://10.1007/s10476-017-0306-2>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

SOME RECENT PROGRESS ON SHARP FOURIER RESTRICTION THEORY

DAMIANO FOSCHI AND DIOGO OLIVEIRA E SILVA

ABSTRACT. The purpose of this note is to discuss several results that have been obtained in the last decade in the context of sharp adjoint Fourier restriction/Strichartz inequalities. Rather than aiming at full generality, we focus on several concrete examples of underlying manifolds with large groups of symmetries, which sometimes allows for simple geometric proofs. We mention several open problems along the way, and include an appendix on integration on manifolds using delta calculus.

1. INTRODUCTION

Curvature causes the Fourier transform to decay. This observation links geometry to analysis, and lies at the base of several topics of modern harmonic analysis. Given the long history of the subject, it is perhaps surprising that the possibility of restricting the Fourier transform to curved submanifolds of Euclidean space was not observed until the late sixties. The Fourier transform of an integrable function is uniformly continuous, and as such, can be restricted to any subset. On the other hand, the Fourier transform of a square-integrable function is again square-integrable, and in view of Plancherel's theorem no better properties can be expected. In particular, restricting the Fourier transform of a square-integrable function to a set of zero Lebesgue measure is meaningless. The question is what happens for intermediate values of $1 < p < 2$.

It is not hard to check that the Fourier transform of a radial function in $L^p(\mathbb{R}^d)$ defines a continuous function away from the origin whenever $1 \leq p < \frac{2d}{d+1}$, see for instance [52]. The corresponding problem for non-radial functions is considerably more delicate. To introduce it, let $\mathcal{M}$ be a smooth compact hypersurface in $\mathbb{R}^d$, endowed with a surface-carried measure $d\mu = \psi d\sigma$. Here σ denotes the surface measure of $\mathcal{M}$, and the function ψ is smooth and non-negative. Given $1 < p < 2$, for which exponents q does the *a priori* inequality

$$(1.1) \quad \left(\int_{\mathcal{M}} |\widehat{f}(\xi)|^q d\mu_{\xi} \right)^{\frac{1}{q}} \leq C \|f\|_{L^p(\mathbb{R}^d)}$$

hold? A complete answer for $q = 2$ is given by the celebrated Tomas–Stein inequality.

Date: January 25, 2017.

2010 Mathematics Subject Classification. 42B10.

Key words and phrases. Fourier restriction theory, Strichartz estimates, optimal constants, extremizers.

The second author was partially supported by the Hausdorff Center for Mathematics.

Theorem 1 ([51, 56]). *Suppose $\mathcal{M}$ has non-zero Gaussian curvature at each point of the support of μ . Then the restriction inequality (1.1) holds for $q = 2$ and $1 \leq p \leq \frac{2d+2}{d+3}$.*

The range of exponents is sharp, since no $L^p \rightarrow L^2(\mu)$ restriction can hold for $\mathcal{M}$ if $p > \frac{2d+2}{d+3}$. This is shown via the famous Knapp example, which basically consists of testing the inequality dual to (1.1) against the characteristic function of a small cap on $\mathcal{M}$. Moreover, some degree of curvature is essential, as there can be no meaningful restriction to a hyperplane except in the trivial case when $p = 1$ and $q = \infty$. An example to keep in mind is that of the unit sphere $\mathbb{S}^{d-1} = \{\xi \in \mathbb{R}^d : |\xi| = 1\}$, with constant positive Gaussian curvature. However, nonvanishing Gaussian curvature is a strong assumption that can be replaced by the nonvanishing of some principal curvatures, at the expense of decreasing the range of admissible exponents p .

The question of what happens for values of $q < 2$ is the starting point for the famous restriction conjecture. One is led by dimensional analysis and Knapp-type examples to guess that the correct range for estimate (1.1) to hold is $1 \leq p < \frac{2d}{d+1}$ and $q \leq (\frac{d-1}{d+1})p'$, where $p' = \frac{p}{p-1}$ denotes the dual exponent. This is depicted in Figure 1. Note that the endpoints of this relation are the trivial case $(p, q) = (1, \infty)$, and $p, q \rightarrow \frac{2d}{d+1}$.

FIGURE 1. Range of exponents for the restriction problem

Despite tremendous effort and very promising partial progress, the restriction conjecture is only known to hold for $d = 2$. The restriction conjecture implies the Keakeya conjecture and is implied by the Bochner–Riesz conjecture. Multilinear versions of the restriction and Keakeya conjectures have been established by Bennett, Carbery and Tao [5], and played a crucial role in the very recent work of Bourgain and Demeter [10] on ℓ^2 decoupling. For more on the restriction problem, and its relation to other prominent problems in modern harmonic analysis, we recommend the works [51, 59] and especially [55].

Tomas–Stein type restriction estimates are very much related to Strichartz estimates for linear partial differential equations of dispersion type. Let us illustrate this point in two cases, that of solutions $u = u(x, t)$ to the homogeneous Schrödinger equation

$$(1.2) \quad i\partial_t u = \Delta u,$$

and that of solutions to the homogeneous wave equation

$$(1.3) \quad \partial_t^2 u = \Delta u.$$

In both situations, $(x, t) \in \mathbb{R}^{d+1}$. The following theorem was originally proved by Strichartz.

Theorem 2 ([54]). *Let $d \geq 1$. Then there exists a constant $S > 0$ such that*

$$(1.4) \quad \|u\|_{L^{2+\frac{4}{d}}(\mathbb{R}^{d+1})} \leq S \|f\|_{L^2(\mathbb{R}^d)},$$

whenever u is the solution of (1.2) with initial data $u(x, 0) = f(x)$. If $d \geq 2$, then there exists a constant $W > 0$ such that

$$(1.5) \quad \|u\|_{L^{2+\frac{4}{d-1}}(\mathbb{R}^{d+1})} \leq W \|(f, g)\|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^d) \times \dot{H}^{-\frac{1}{2}}(\mathbb{R}^d)},$$

whenever u is the solution of (1.3) with initial data $u(x, 0) = f(x)$ and $\partial_t u(x, 0) = g(x)$.

A hint that Theorems 1 and 2 might be related comes from the numerology of the exponents: The Strichartz exponent $2 + \frac{4}{d}$ coincides with the dual of the Tomas–Stein exponent in dimension $d + 1$. It turns out that Strichartz estimates for the Schrödinger equation correspond to restriction estimates on the paraboloid, whereas Strichartz estimates for the wave equation correspond to restriction estimates on the cone. Note that the Gaussian curvature of the cone is identically zero because one of its principal curvatures vanishes. This in turn translates into estimate (1.5) holding for the Strichartz exponent in one lower dimension. Perhaps more significantly, neither of these manifolds is compact. However, they exhibit some scale invariance properties that enable a reduction to the compact setting. We shall return to this important point later in our discussion.

In this note, we are interested in extremizers and optimal constants for sharp variants of restriction and Strichartz-type inequalities. Apart from their intrinsic mathematical interest and elegance, such sharp inequalities often allow for various refinements of existing inequalities. The following are natural questions, which in particular can be posed for inequalities (1.1), (1.4) and (1.5):

- (1) What is the value of the optimal constant?
- (2) Do extremizers exist?
 - (a) If so, are they unique, possibly after applying the symmetries of the problem?
 - (b) If not, what is the mechanism responsible for this lack of compactness?
- (3) How do extremizing sequences behave?
- (4) What are some qualitative properties of extremizers?
- (5) What are necessary and sufficient conditions for a function to be an extremizer?

Questions of this flavor have been asked in a variety of situations, and in the context of classical inequalities from Euclidean harmonic analysis go back at least to the seminal work

of Beckner [1] for the Hausdorff–Young inequality, and Lieb [36] for the Hardy–Littlewood–Sobolev inequality. In comparison, sharp Fourier restriction inequalities have a relatively short history, with the first works on the subject going back to Kunze [35], Foschi [23] and Hundertmark–Zharnitsky [30].

Works addressing the existence of extremizers for inequalities of Fourier restriction type tend to be a *tour de force* in classical analysis, using a variety of sophisticated techniques: bilinear estimates and refined estimates in $X^{s,b}$ -type spaces [16, 37, 43, 45], concentration compactness arguments tailored to the specific problem in question [26, 32, 33, 35, 47, 48, 49], variants and generalizations of the Brézis–Lieb lemma from functional analysis [21, 22, 26], Fourier integral operators [16, 26], symmetrization techniques [16], variational, perturbative and spectral analysis [16, 19, 26, 37], regularity theory for equations with critical scaling and additive combinatorics [17], among others. In contrast, a full characterization of extremizers has been given in a few selected cases using much more elementary methods. This is due to the presence of a large underlying group of symmetries which allows for several simplifications that ultimately reduce the problem to a simple geometric observation. We will try to illustrate this point in the upcoming sections.

Before doing so, let us briefly comment on some approaches that have been developed in the last decades in order to establish Tomas–Stein type Fourier restriction inequalities. For the sake of brevity, we specialize our discussion to inequalities (1.1) and (1.4), but a more general setting should be kept in mind. If T denotes the restriction operator $f \mapsto \hat{f}|_{\mathcal{M}}$, then its adjoint T^* , usually called the extension operator, is given by $g \mapsto \widehat{g\mu}$, where the Fourier transform of the measure $g\mu$ is given by

$$(1.6) \quad \widehat{g\mu}(x) = \int_{\mathcal{M}} g(\omega) e^{-ix \cdot \omega} d\mu_{\omega}. \quad (x \in \mathbb{R}^d)$$

The Tomas–Stein inequality at the endpoint $(p, q) = (\frac{2d+2}{d+3}, 2)$ is equivalent to the extension estimate

$$(1.7) \quad \|\widehat{g\mu}\|_{L^{\frac{2d+2}{d-1}}(\mathbb{R}^d)} \leq C \|g\|_{L^2(\mathcal{M}, \mu)}.$$

If $g \in L^2(\mathcal{M}, \mu)$, then the composition $T^*T(g)$ is well-defined, and a computation shows that

$$T^*T : g \mapsto g * \widehat{\mu}.$$

Since the operator norms satisfy $\|T\|^2 = \|T^*\|^2 = \|T^*T\|$, the study of these three operators is equivalent, even if the goal is to obtain sharp inequalities and determine optimal constants. So we focus on the operator T^*T . Boundedness of $T^*T(g)$ is only ensured if the Fourier transform $\widehat{\mu}(x)$ exhibits some sort of decay as $|x| \rightarrow \infty$. This in turn is a consequence of the principle of stationary phase, see [51, 59], since the nonvanishing curvature of $\mathcal{M}$ translates into a nondegenerate Hessian for the phase function of the oscillatory integral given by (1.6) with $g \equiv 1$. This is the starting point for the original argument of Tomas [56], which was then extended to the endpoint $\frac{2d+2}{d-1}$ by embedding T^*T into an analytic family of operators and invoking Stein’s complex interpolation theorem. It is hard not to notice the parallel between the operator T^*T and the averaging operator $f \mapsto f * \mu$, whose

L^p improving properties can be established via the same proof on the Fourier side, see for instance [51, pp. 370-371].

A second method to prove restriction estimates goes back to the work of Ginibre and Velo [27]. It consists of introducing a time parameter and treating the extension operator as an evolution operator. Two key ingredients for this approach are the Hausdorff–Young inequality and fractional integration in the form of the Hardy–Littlewood–Sobolev inequality.¹ These methods are more amenable to the needs of the partial differential equations community, and for instance allow to treat the case of mixed norm spaces.

In the special cases when the dual exponent p' is an even integer, one can devise yet another proof which comes from the world of bilinear estimates, see for instance [9, 25, 34]. One simply rewrites the left-hand side of inequality (1.7) as an L^2 norm, and appeals to Plancherel in order to reduce the problem to a multilinear convolution estimate. For instance, if $d = 3$, then $p' = \frac{2 \cdot 3 + 2}{3 - 1} = 4$, and

$$\|\widehat{g\mu}\|_{L^4(\mathbb{R}^3)}^2 = \|\widehat{g\mu} \widehat{g\mu}\|_{L^2(\mathbb{R}^3)} = \|g\mu * g\mu\|_{L^2(\mathbb{R}^3)}.$$

Similarly, if $d = 2$, then $p' = \frac{2 \cdot 2 + 2}{2 - 1} = 6$, and

$$\|\widehat{g\mu}\|_{L^6(\mathbb{R}^2)}^3 = \|g\mu * g\mu * g\mu\|_{L^2(\mathbb{R}^2)}.$$

In the bilinear case, the pointwise inequality

$$|g\mu * g\mu| \leq |g|\mu * |g|\mu$$

then reveals that one can restrict attention to nonnegative functions. This observation can greatly simplify matters, as it reduces an oscillatory problem to a question of geometric integration over a specific manifold. Furthermore, an application of the Cauchy–Schwarz inequality with respect to an appropriate measure implies the pointwise inequality

$$|g\mu * g\mu|^2 \leq (|g|^2\mu * |g|^2\mu)(\mu * \mu).$$

A good understanding of the convolution measure $\mu * \mu$ becomes a priority. Given integrable functions $g, h \in L^1(\mathcal{M}, \mu)$, the convolution $g\mu * h\mu$ is a finite measure defined on the Borel subsets $E \subset \mathbb{R}^d$ by

$$(g\mu * h\mu)(E) = \int_{\mathcal{M} \times \mathcal{M}} \chi_E(\eta + \zeta) g(\eta) h(\zeta) d\mu_\eta d\mu_\zeta.$$

It is clear that this measure is supported on the Minkowski sum $\mathcal{M} + \mathcal{M}$. In most situations of interest when some degree of curvature is present, one can check that $g\mu * h\mu$ is absolutely continuous with respect to Lebesgue measure on $\mathbb{R}^d$. In such cases, the measure $g\mu * h\mu$ can be identified with its Radon–Nikodym derivative with respect to Lebesgue measure, and for almost every $\xi \in \mathbb{R}^d$ we have that

$$(1.8) \quad (g\mu * h\mu)(\xi) = \int_{\mathcal{M} \times \mathcal{M}} \delta(\xi - \eta - \zeta) g(\eta) h(\zeta) d\mu_\eta d\mu_\zeta.$$

¹Interestingly, the full restriction conjecture on $\mathbb{R}^2$ can be proved with a combination of Hausdorff–Young and Hardy–Littlewood–Sobolev, see for instance [51, pp. 412-414].

Here $\delta(\cdot)$ denotes the d -dimensional Dirac delta distribution. As we shall exemplify in the course of the paper, expression (1.8) turns out to be very useful for computational purposes.

Overview. As we hope to have made apparent already, this note is meant as a short survey of a restricted part of the topic of sharp Strichartz and Fourier extension inequalities. In §2 we deal with noncompact surfaces, and discuss the cases of the paraboloid and the cone, where a full characterization of extremizers is known, and the cases of the hyperboloid and a quartic perturbation of the paraboloid, where extremizers fail to exist. Most of the material in this section is contained in the works [12, 23, 39, 43, 44]. In §3 we discuss the case of spheres. A full characterization of extremizers at the endpoint is only known in the case of $\mathbb{S}^2$. We mention some recent partial progress in the case of the circle $\mathbb{S}^1$, and observe how the methods in principle allow to refine some related inequalities. In particular, we obtain a new sharp extension inequality on $\mathbb{S}^1$ in the mixed norm space $L_{\text{rad}}^6 L_{\text{ang}}^2(\mathbb{R}^2)$. Most of the material in §3 is contained in the works [13, 14, 18, 24, 40]. We leave some final remarks to §4, where we hint at a possible unifying picture for the results that have been discussed. Finally, we include an Appendix with a brief introduction to integration on manifolds using delta calculus.

Remarks and further references. The style of this note is admittedly informal. In particular, some objects will not be rigorously defined, and most results will not be precisely formulated. None of the material is new, with the exception of the results in §3.2 and a few observations that we have not been able to find in the literature. The subject is becoming more popular, as shown by the increasing number of works that appeared in the last five years. We have attempted to give a rather complete set of references, which includes several interesting works [3, 4, 6, 7, 8, 11, 29, 31, 38, 41, 42, 46] that will not be discussed here. Given its young age, there are plenty of open problems in the area. Our contribution is to provide some more.

2. NONCOMPACT SURFACES

2.1. Paraboloids. Given $d \geq 1$, let us consider the d -dimensional paraboloid

$$\mathbb{P}^d = \{(\xi, \tau) \in \mathbb{R}^{d+1} : \tau = |\xi|^2\},$$

equipped with projection measure

$$\mu_d(\xi, \tau) = \delta(\tau - |\xi|^2) d\xi d\tau.$$

The validity of an $L^2(\mu_d) \rightarrow L^{2+\frac{4}{d}}$ extension estimate follows from Strichartz inequality (1.4) for the Schrödinger equation as discussed before. Extremizers for this inequality are known to exist in all dimensions [47], and to be Gaussians in low dimensions [2, 23, 30]. Extremizers are conjectured to be Gaussians in all dimensions [30], see also [15]. Let us specialize to the case $d = 2$ and follow mostly [23]. In view of identity (1.8), which itself is a consequence of formula (A.1) from the Appendix, the convolution of projection measure

on the two-dimensional paraboloid $\mathbb{P}^2$ is given by

$$(\mu_2 * \mu_2)(\xi, \tau) = \int_{\mathbb{R}^2 \times \mathbb{R}^2} \delta\left(\tau - |\eta|^2 - |\zeta|^2\right) d\eta d\zeta = \int_{\mathbb{R}^2} \delta(\tau - |\eta|^2 - |\xi - \eta|^2) d\eta.$$

Changing variables and computing in polar coordinates according to (A.5), we have that

$$(\mu_2 * \mu_2)(\xi, \tau) = \int_{\mathbb{R}^2} \delta\left(\tau - \frac{|\xi|^2}{2} - 2|\eta|^2\right) d\eta = \frac{\pi}{2} \chi\left(\tau \geq \frac{|\xi|^2}{2}\right).$$

We arrive at the crucial observation that the convolution measure $\mu_2 * \mu_2$ defines a function which is not only uniformly bounded, but also constant in the interior of its support $\{2\tau \geq |\xi|^2\}$. See [23, Lemma 3.2] for an alternative proof that uses the invariance of $\mu_2 * \mu_2$ under Galilean transformations and parabolic dilations. A successive application of Cauchy–Schwarz and Hölder’s inequality finishes the argument. Indeed, the pointwise bound

$$(2.1) \quad |(f\mu_2 * f\mu_2)(\xi, \tau)|^2 \leq (\mu_2 * \mu_2)(\xi, \tau) (|f|^2 \mu_2 * |f|^2 \mu_2)(\xi, \tau)$$

follows from an application of the Cauchy–Schwarz inequality with respect to the measure

$$\delta\left(\tau - |\eta|^2 - |\zeta|^2\right) d\eta d\zeta.$$

Integrating inequality (2.1) over $\mathbb{R}^{2+1}$, an application of Hölder’s inequality then reveals

$$(2.2) \quad \|f\mu_2 * f\mu_2\|_{L^2(\mathbb{R}^3)}^2 \leq \|\mu_2 * \mu_2\|_{L^\infty(\mathbb{R}^3)} \|f\|_{L^2(\mathbb{R}^2)}^4.$$

It is possible to turn both inequalities simultaneously into an equality. The conditions for equality in (2.1) translate into a functional equation

$$(2.3) \quad f(\eta)f(\zeta) = F(\eta + \zeta, |\eta|^2 + |\zeta|^2)$$

which should hold for some complex-valued function F defined on the support of the convolution $\mu_2 * \mu_2$, and almost every point (η, ζ) . An example of a solution to (2.3) is given by the Gaussian function $f(\eta) = \exp(-|\eta|^2)$ and the corresponding $F(\xi, \tau) = \exp(-\tau)$. All other solutions are obtained from this one by applying a symmetry of the Schrödinger equation, see [23, Proposition 7.15]. That they turn inequality (2.2) into an equality follows from the fact that the convolution $\mu_2 * \mu_2$ is constant inside its support.

The one-dimensional case $L^2(\mu_1) \rightarrow L^6$ admits a similar treatment. The threefold convolution $\mu_1 * \mu_1 * \mu_1$ is given by a constant function in the interior of its support $\{3\tau \geq \xi^2\}$, and the corresponding functional equation can be solved by similar methods. Gaussians are again seen to be the unique extremizers.

Alternative approaches are available: Hundertmark–Zharnitski [30] based their analysis on a novel representation of the Strichartz integral $\|\widehat{f\sigma}\|_{L^4}^4$, and Bennett et al. [2] identified a monotonicity property of such integrals under a certain quadratic heat-flow.

2.2. Cones. Given $d \geq 2$, consider the (one-sheeted) cone

$$\Gamma^d = \{(\xi, \tau) \in \mathbb{R}^{d+1} : \tau = |\xi| > 0\},$$

equipped with its Lorentz invariant measure

$$\nu_d(\xi, \tau) = \delta(\tau - |\xi|) \frac{d\xi d\tau}{|\xi|} = 2\delta(\tau^2 - |\xi|^2) \chi(\tau > 0) d\xi d\tau.$$

The second identity is a consequence of formula (A.4) from the Appendix. The validity of an $L^2(\nu_d) \rightarrow L^{2+\frac{4}{d-1}}$ extension estimate follows from Strichartz inequality (1.5) for the wave equation as discussed before. Extremizers for the cone are known to exist in all dimensions [45], and to be exponentials in low dimensions [12, 23]. Let us specialize to the case $d = 3$. The convolution of the Lorentz invariant measure on the three-dimensional cone Γ^3 is given by

$$(\nu_3 * \nu_3)(\xi, \tau) = \int_{\mathbb{R}^3} \frac{\delta(\tau - |\eta| - |\xi - \eta|)}{|\eta||\xi - \eta|} d\eta.$$

Given $\xi \in \mathbb{R}^3$, we write a generic vector $\eta \in \mathbb{R}^3$ in spherical coordinates, so that $d\eta = \rho^2 \sin \theta d\rho d\theta d\varphi$, where $\rho = |\eta| \geq 0$, $\theta \in [0, \pi]$ is the angle between ξ and η , and $\varphi \in [0, 2\pi]$ is an angular variable. Setting $\sigma = |\xi - \eta|$, the Jacobian of the change of variables $\eta \rightsquigarrow (\rho, \sigma, \varphi)$ into bipolar coordinates, see Figure 2 below, is given by

$$d\eta = \frac{\rho\sigma}{|\xi|} d\rho d\sigma d\varphi.$$

FIGURE 2. Bipolar coordinates

Letting $a = \rho - \sigma$ and $b = \rho + \sigma$, we invoke the change of variables formula (A.5) to further compute

$$\begin{aligned} (\nu_3 * \nu_3)(\xi, \tau) &= 2\pi \int_{\substack{|\rho - \sigma| \leq |\xi| \\ \rho + \sigma \geq |\xi|}} \frac{\delta(\tau - \rho - \sigma) \rho \sigma}{\rho \sigma |\xi|} d\rho d\sigma = \\ &= \frac{\pi}{|\xi|} \int_{\substack{|a| \leq |\xi| \\ b \geq |\xi|}} \delta(\tau - b) da db = 2\pi \chi(\tau \geq |\xi|). \end{aligned}$$

This again defines a constant function inside its support, and a combination of Cauchy–Schwarz and Hölder as before establishes the sharp inequality. The characterization of extremizers follows from the analysis of the functional equation

$$(2.4) \quad f(\eta)f(\zeta) = F(\eta + \zeta, |\eta| + |\zeta|),$$

which yields as a particular solution the exponential function $f(\eta) = \exp(-|\eta|)$ and the corresponding $F(\xi, \tau) = \exp(-\tau)$. All other solutions are obtained from this one by applying a symmetry of the wave equation, see [23, Proposition 7.23], and the lower dimension case $L^2(\nu_2) \rightarrow L^6$ admits a similar treatment.

2.3. Hyperboloids. We now switch to the (one-sheeted) hyperboloid

$$\mathbb{H}^d = \{(\xi, \tau) \in \mathbb{R}^{d+1} : \tau = \sqrt{1 + |\xi|^2}\},$$

equipped with the Lorentz invariant measure

$$\lambda_d(\xi, \tau) = \delta(\tau - \sqrt{1 + |\xi|^2}) \frac{d\xi d\tau}{\sqrt{1 + |\xi|^2}} = 2\delta(\tau^2 - |\xi|^2 - 1) \chi(\tau > 0) d\xi d\tau.$$

As established in [54], an $L^2(\lambda_d) \rightarrow L^{p'}$ extension estimate holds provided

$$(2.5) \quad \begin{aligned} 2 + \frac{4}{d} \leq p' \leq 2 + \frac{4}{d-1}, & \text{ if } d > 1, \\ 6 \leq p' < \infty, & \text{ if } d = 1. \end{aligned}$$

Note that the lower and upper bounds in the exponent range (2.5) correspond to the cases of the paraboloid and the cone, respectively. We focus on the case $d = 2$ and $p' = 4$, and take advantage of Lorentz symmetries to compute the convolution $\lambda_2 * \lambda_2$. Along the vertical axis of the hyperboloid $\mathbb{H}^2$,

$$(\lambda_2 * \lambda_2)(0, \tau) = \int_{\mathbb{R}^2} \delta(\tau - 2\sqrt{1 + |\eta|^2}) \frac{d\eta}{1 + |\eta|^2} = \frac{2\pi}{\tau} \chi(\tau \geq 2).$$

Lorentz invariance forces the convolution $\lambda_2 * \lambda_2$ to be constant along the level sets of the function $\tau^2 - |\xi|^2$. As a consequence,

$$(2.6) \quad (\lambda_2 * \lambda_2)(\xi, \tau) = \frac{2\pi}{\sqrt{\tau^2 - |\xi|^2}} \chi(\tau \geq \sqrt{4 + |\xi|^2}).$$

Contrary to the previous cases, this no longer defines a constant function inside its support. Since it is uniformly bounded (by π), the argument can still be salvaged to yield a sharp

extension inequality. Extremizers for this inequality, however, do not exist. We shall observe a similar phenomenon in the case of perturbed paraboloids, considered in the next subsection, and postpone a more detailed discussion until then.

2.4. Perturbations. Let us start with a brief discussion of a specific instance of a comparison principle from [39] that proved useful in establishing sharp inequalities for perturbed paraboloids. Let $(\mathbb{P}^2, \mu_2)$ denote the two-dimensional paraboloid considered in §2.1, and let $\tilde{\mu}_2$ denote the projection measures on the surface

$$\{(\xi, \tau) \in \mathbb{R}^{2+1} : \tau = |\xi|^2 + |\xi|^4\}.$$

Then the pointwise inequality

$$(2.7) \quad (\tilde{\mu}_2 * \tilde{\mu}_2)\left(\xi, \frac{|\xi|^2}{2} + \frac{|\xi|^4}{8} + \tau\right) \leq (\mu_2 * \mu_2)\left(\xi, \frac{|\xi|^2}{2} + \tau\right)$$

holds for every $\tau > 0$ and $\xi \in \mathbb{R}^d$, and is strict at almost every point of the support of the measure $\tilde{\mu}_2 * \tilde{\mu}_2$. The feature of the function $\xi \mapsto |\xi|^4$ that makes this possible is *convexity*. Any nonnegative, continuously differentiable, strictly convex function would do, see [39, Theorem 1.3] for a precise version of this comparison principle which holds in all dimensions. A sharp $L^2(\tilde{\mu}_2) \rightarrow L^4$ extension inequality can be obtained by concatenating Cauchy–Schwarz and Hölder as before, and extremizers do not exist because of the strict inequality in (2.7). Heuristically, extremizing sequences are forced to place their mass in arbitrarily small neighborhoods of the region where the convolution $\tilde{\mu}_2 * \tilde{\mu}_2$ attains its global maximum, for otherwise they would not come close to attaining the sharp constant. The analysis of the cases of equality in (2.7) reveals that this region has zero Lebesgue measure, and this forces any extremizing sequence to concentrate.

3. COMPACT SURFACES

3.1. Spheres. Consider the endpoint Tomas–Stein extension inequality on the sphere

$$(3.1) \quad \left(\int_{\mathbb{R}^d} |\widehat{f\sigma}(x)|^{p'} dx \right)^{\frac{1}{p'}} \leq C \|f\|_{L^2(\mathbb{S}^{d-1}, \sigma)},$$

where $\sigma = \sigma_{d-1}$ denotes surface measure on $\mathbb{S}^{d-1}$ and $p' = \frac{2d+2}{d-1}$. Extremizers for inequality (3.1) were first shown to exist when $d = 3$ in [16]. The precise form of nonnegative extremizers was later determined in [24], and they turn out to be constant functions. See also [26] for a conditional existence result in higher dimensions. It seems natural to conjecture that extremizers should be constants in all dimensions. Spheres are antipodally symmetric compact manifolds, and this brings in some additional difficulties which can already be observed at the level of convolution measures. Indeed, formula (A.4) from the Appendix implies

$$\sigma_{d-1}(\xi) = \delta(1 - |\xi|) d\xi = 2\delta(1 - |\xi|^2) d\xi.$$

Invoking (A.3), and then (A.4) once again, we have

$$(\sigma_{d-1} * \sigma_{d-1})(\xi) = 2 \int_{\mathbb{S}^{d-1}} \delta(1 - |\xi - \omega|^2) d\sigma_\omega = \frac{2}{|\xi|} \int_{\mathbb{S}^{d-1}} \delta\left(2\frac{\xi}{|\xi|} \cdot \omega - |\xi|\right) d\sigma_\omega.$$

Computing in polar coordinates according to (A.5), one concludes

$$(3.2) \quad (\sigma_{d-1} * \sigma_{d-1})(\xi) = \frac{V_{d-2}}{|\xi|} \left(1 - \frac{|\xi|^2}{4}\right)_+^{\frac{d-3}{2}},$$

where V_{d-2} denotes the surface measure of $\mathbb{S}^{d-2}$. When $d = 3$, and consequently $p' = 4$, the inequality in question is equivalent to a 4-linear estimate. Moreover, the last factor in (3.2) simplifies and one is left with

$$(3.3) \quad (\sigma_2 * \sigma_2)(\xi) = \frac{2\pi}{|\xi|} \chi(|\xi| \leq 2).$$

Expression (3.3) blows up at the origin $\xi = 0$, which prevents a straightforward adaptation of the methods from the previous section. In particular, the interaction between antipodal points causes difficulties that prevented the local analysis from [16] to identify the global extremizers of the problem. The work [24] resolves this issue in a simple manner, using the following geometric feature of the sphere: If the sum of three unit vectors $\omega_1, \omega_2, \omega_3 \in \mathbb{S}^d$ is again a unit vector, then necessarily

$$(3.4) \quad |\omega_1 + \omega_2|^2 + |\omega_2 + \omega_3|^2 + |\omega_3 + \omega_1|^2 = 4.$$

To see why this is true, one simply squares the assumption $|\omega_1 + \omega_2 + \omega_3| = 1$ and expands the left-hand side of (3.4). An application of the Cauchy–Schwarz inequality together with identity (3.4) reduces the analysis to antipodally symmetric functions, and at the same time neutralizes the singularity of the convolution measure at the origin. This reduces the 4-linear problem on f to a bilinear problem on its square f^2 . More precisely, one is left with establishing a monotonicity property for the quadratic form

$$H(g) = \int_{(\mathbb{S}^2)^2} \overline{g(\omega)} g(\nu) |\omega - \nu| d\sigma_\omega d\sigma_\nu,$$

where $g = f^2$ is now assumed to be merely integrable. This in turn is accomplished via spectral analysis. If c denotes the mean value of g over the sphere and $\mathbf{1}$ denotes the constant function equal to 1, one wants to show that $H(g) \leq H(c\mathbf{1})$. The crucial observation is that the quadratic form H is diagonal in a suitable basis. In fact, expanding $g = \sum_{k \geq 0} Y_k$ in spherical harmonics, we have

$$(3.5) \quad H(g) = 2\pi \sum_{k \geq 0} \Lambda_k \|Y_k\|_{L^2(\mathbb{S}^2)}^2 \leq 2\pi \Lambda_0 \|Y_0\|_{L^2(\mathbb{S}^2)}^2 = H(c\mathbf{1}),$$

where the eigenvalues Λ_k can be computed via the Funk–Hecke formula [20, p. 247]. It turns out that $\Lambda_k < 0$ when $k \geq 1$, we refer the reader to [24] for the full details. This approach was extended in [13] to establish sharp $L^2(\sigma_{d-1}) \rightarrow L^4$ extension estimates on $\mathbb{S}^{d-1}$ for $d = 4, 5, 6, 7$. Table 1 indicates the signs of the corresponding coefficients $\Lambda_k = \Lambda_k(\mathbb{S}^{d-1})$.

Note that the sum in (3.5) ranges over even values of k only since the function f , and therefore $g = f^2$, is assumed to be antipodally symmetric. In particular, the sign of the

	Λ_0	Λ_1	Λ_2	Λ_3	Λ_4	Λ_5	Λ_6	$\dots$
$d = 3$	+	−	−	−	−	−	−	$\dots$
$d = 4$	+	0	−	0	−	0	−	$\dots$
$d = 5, 6, 7$	+	+	−	−	−	−	−	$\dots$
$d \geq 8$	+	+	+	*	*	*	*	$\dots$

TABLE 1. Signs of the coefficients $\Lambda_k(\mathbb{S}^{d-1})$

coefficient Λ_1 is of no importance to the analysis. However, one starts to observe that $\Lambda_2(\mathbb{S}^{d-1}) > 0$ if $d \geq 8$, and this is the reason for the failure of this method of proof.

Threefold convolution measures can also be computed via delta calculus, at the expense of possibly more complicated expressions. For instance, the convolution $\sigma_2 * \sigma_2 * \sigma_2$ is a radial function supported on the ball of radius 3 centered at the origin, given by the expression

$$(\sigma_2 * \sigma_2 * \sigma_2)(\xi) = \begin{cases} 8\pi^2, & \text{if } 0 \leq |\xi| \leq 1, \\ 4\pi^2 \left(-1 + \frac{3}{|\xi|} \right), & \text{if } 1 \leq |\xi| \leq 3. \end{cases}$$

Notice that this defines a bounded, continuous function which is constant inside the unit ball, and decreases to zero on the annulus $\{1 \leq |\xi| \leq 3\}$. The lowest dimensional endpoint case $(d, p) = (2, 6)$ holds some hidden surprises. First of all, the estimate translates into an inequality involving a 6-linear form. The convolution $\sigma_1 * \sigma_1 * \sigma_1$ defines a radial function given by a complicated integral expression, see [14, Lemma 8], which is better illustrated in Figure 3 below.

FIGURE 3. Plot of the function $r \mapsto (\sigma_1 * \sigma_1 * \sigma_1)(r)$.

A difficulty is that the convolution $\sigma_1 * \sigma_1 * \sigma_1$ now blows up along a whole circle, and not just at one point. One would still like to reduce matters to a two-step analysis, and a possible program is as follows. One first reduces the 6-linear analysis to a trilinear analysis

on squares, and then performs a spectral analysis of the trilinear form

$$(3.6) \quad T(h_1, h_2, h_3) = \int_{(\mathbb{S}^1)^6} h_1(\omega_1) h_2(\omega_2) h_3(\omega_3) (|\omega_4 + \omega_5 + \omega_6|^2 - 1) \, d\Sigma_{\vec{\omega}},$$

where the measure Σ is given by

$$(3.7) \quad d\Sigma_{\vec{\omega}} = \delta(\omega_1 + \omega_2 + \omega_3 + \omega_4 + \omega_5 + \omega_6) \, d\sigma_{\omega_1} \, d\sigma_{\omega_2} \, d\sigma_{\omega_3} \, d\sigma_{\omega_4} \, d\sigma_{\omega_5} \, d\sigma_{\omega_6}.$$

This is later specialized to the case $h_1 = h_2 = h_3 = f^2$. Let us formulate these steps in a more precise manner:

Step 1. *Let $f \in L^2(\mathbb{S}^1)$ be nonnegative and antipodally symmetric. Then:*

$$\begin{aligned} \int_{(\mathbb{S}^1)^6} f(\omega_1) f(\omega_2) f(\omega_3) f(\omega_4) f(\omega_5) f(\omega_6) (|\omega_4 + \omega_5 + \omega_6|^2 - 1) \, d\Sigma_{\vec{\omega}} \\ \leq \int_{(\mathbb{S}^1)^6} f(\omega_1)^2 f(\omega_2)^2 f(\omega_3)^2 (|\omega_4 + \omega_5 + \omega_6|^2 - 1) \, d\Sigma_{\vec{\omega}}. \end{aligned}$$

Step 2. *Let $h \in L^1(\mathbb{S}^1)$ be a nonnegative and antipodally symmetric function. Let c denote the mean value of h over $\mathbb{S}^1$. Then $T(h, h, h) \leq c^3 T(\mathbf{1}, \mathbf{1}, \mathbf{1})$, with equality if and only if h is constant.*

Step 1 is an open problem, posed as a conjecture in [14]. Step 2 is the subject of papers [14, 40]. In short, one decomposes $h = c + g$ where g has mean zero, and analyzes each of the summands in

$$T(h, h, h) = T(c, c, c) + 3T(c, c, g) + 3T(c, g, g) + T(g, g, g)$$

separately. The linear term in g vanishes for symmetry considerations, the bilinear term is nonpositive and the trilinear term can be controlled in absolute value by the bilinear term. The proof of these statements relies on expanding the integrand in (3.6) in Fourier basis, and appealing to delicate estimates for integrals of sixfold products of Bessel functions. We refer the reader to the original papers for details, and proceed to discuss a related inequality which can be put in sharp form by invoking more elementary estimates for integrals of Bessel functions.

3.2. Estimates in mixed norm spaces. The connection between Bessel functions and spherical harmonics is easily seen via the formula

$$(3.8) \quad \widehat{Y_k \sigma}(x) = i^k J_{\frac{d}{2}-1+k}(|x|) |x|^{1-\frac{d}{2}} Y_k\left(\frac{x}{|x|}\right), \quad (x \in \mathbb{R}^d)$$

which follows from an application of the Funk–Hecke formula together with Rodrigues formula for Gegenbauer polynomials, see [53, 58]. Spherical harmonic expansions played an important role in the recent work of Córdoba [18] on certain Fourier extension inequalities in mixed norm spaces. The following inequality, which had already appeared in the Ph.D.

thesis of Vega [57], was reproved in [18] using different methods: For $d \geq 2$ and $q > \frac{2d}{d-1}$, there exists $C_{d,q} < \infty$ such that

$$(3.9) \quad \|\widehat{f\sigma}\|_{L_{\text{rad}}^q L_{\text{ang}}^2(\mathbb{R}^d)} \leq C_{d,q} \|f\|_{L^2(\mathbb{S}^{d-1})}.$$

Here the norm in $L_{\text{rad}}^q L_{\text{ang}}^2(\mathbb{R}^d)$ is given by the integral

$$\left(\int_0^\infty \left(\int_{\mathbb{S}^{d-1}} |\widehat{f\sigma}(r\omega)|^2 d\sigma_\omega \right)^{\frac{q}{2}} r^{d-1} dr \right)^{\frac{1}{q}}.$$

Note that inequality (3.9) follows in a simple way from Hölder's inequality for those exponents q for which the (adjoint) Tomas–Stein inequality is known to hold. We now indicate a possible path to obtain the sharp form of a number of instances of inequality (3.9). Given $f \in L^2(\mathbb{S}^{d-1})$, we expand it in normalized spherical harmonics

$$f = \sum_{k \geq 0} a_k Y_k,$$

with each $\|Y_k\|_{L^2} = 1$. Appealing to formula (3.8) and to orthogonality of the $\{Y_k\}$, we see that

$$\int_{\mathbb{S}^{d-1}} |\widehat{f\sigma}(r\omega)|^2 d\sigma_\omega = \sum_{k \geq 0} |a_k|^2 |J_{\frac{d}{2}-1+k}(r)|^2 r^{2-d}.$$

The q -th power of the left-hand side of inequality (3.9) can thus be rewritten as

$$(3.10) \quad \int_0^\infty \left(\sum_{k \geq 0} |a_k|^2 |J_{\frac{d}{2}-1+k}(r)|^2 \right)^{\frac{q}{2}} r^{(1-\frac{d}{2})q} r^{d-1} dr,$$

which is the starting point for the analysis in [18]. Whenever the exponent q is an even integer, we can take an alternative route and in principle obtain a sharp inequality. Let us illustrate this in the case $(d, q) = (2, 6)$, which corresponds to a weaker form of the $L^2(\sigma_1) \rightarrow L^6$ adjoint Tomas–Stein inequality discussed in the previous subsection. In this case, the integral (3.10) can be rewritten as

$$\sum_{k, \ell, m \geq 0} |a_k|^2 |a_\ell|^2 |a_m|^2 I(k, \ell, m),$$

where the integrals $I(k, \ell, m)$ are defined as

$$I(k, \ell, m) = \int_0^\infty J_k^2(r) J_\ell^2(r) J_m^2(r) r dr.$$

These integrals obey the following monotonicity property:

Proposition 3. *Let k, ℓ, m be nonnegative integers. Then:*

$$I(k, \ell, m) \leq I(0, 0, 0),$$

and equality holds if and only if $k = \ell = m = 0$.

As a consequence, we obtain a sharp form of this particular instance of inequality (3.9):

$$\begin{aligned} \|\widehat{f\sigma}\|_{L_{\text{rad}}^6 L_{\text{ang}}^2(\mathbb{R}^2)}^6 &= \sum_{k,\ell,m \geq 0} |a_k|^2 |a_\ell|^2 |a_m|^2 I(k, \ell, m) \\ &\leq I(0, 0, 0) \sum_{k,\ell,m \geq 0} |a_k|^2 |a_\ell|^2 |a_m|^2 = I(0, 0, 0) \|f\|_{L^2(\mathbb{S}^1)}^6, \end{aligned}$$

with constants being the unique extremizers.

Proof of Proposition 3. Reasoning as in [14, §3], we see that

$$I(k, \ell, m) = \int_{\mathbb{R}^2} \widehat{e_k \sigma} \widehat{e_k \sigma} \widehat{e_\ell \sigma} \widehat{e_\ell \sigma} \widehat{e_m \sigma} \widehat{e_m \sigma} dx = \int_{(\mathbb{S}^1)^6} (\omega_1 \overline{\omega_2})^k (\omega_3 \overline{\omega_4})^\ell (\omega_5 \overline{\omega_6})^m d\Sigma_{\vec{\omega}},$$

where the function $e_n : \mathbb{S}^1 \rightarrow \mathbb{C}$ is defined via $e_n(\omega) = \omega^n$ and the measure Σ is given by (3.7). The triangle inequality immediately implies

$$I(k, \ell, m) \leq I(0, 0, 0),$$

with equality if and only

$$(\omega_1 \overline{\omega_2})^k (\omega_3 \overline{\omega_4})^\ell (\omega_5 \overline{\omega_6})^m = 1,$$

for every $\omega_1, \dots, \omega_6 \in \mathbb{S}^1$ such that $\sum_{j=1}^6 \omega_j = 0$. One easily checks that this implies $k = \ell = m = 0$, and the proof is complete. $\square$

4. TOWARDS A UNIFYING PICTURE

Table 2 summarizes some of our discussion from the previous sections concerning the sharp form of a number of Fourier extension inequalities which can be recast as bilinear estimates for the appropriate convolution measures.

Manifold	Fourier extension inequality	Extremizers
Paraboloid	$L^2(\mathbb{P}^2, \mu_2) \rightarrow L^4(\mathbb{R}^3)$	Gaussians
Cone	$L^2(\Gamma^3, \nu_3) \rightarrow L^4(\mathbb{R}^4)$	Exponentials
Hyperboloid	$L^2(\mathbb{H}^2, \lambda_2) \rightarrow L^4(\mathbb{R}^3)$	Do not exist
Sphere	$L^2(\mathbb{S}^2, \sigma_2) \rightarrow L^4(\mathbb{R}^3)$	Constants

TABLE 2.

The two-dimensional surfaces in question (paraboloid, hyperboloid and sphere) can all be obtained as intersections of the three-dimensional cone with appropriately chosen hyperplanes. Figure 4 below illustrates this point. There, the ambient space $\mathbb{R}^4$ is endowed with coordinates $(\xi, \tau) \in \mathbb{R}^{3+1}$, where $\xi = (\xi_1, \xi') \in \mathbb{R}^{1+2}$. One cannot help noticing that the restriction of the exponential function $\exp(-|\xi|)$, which is an extremizer for the cone, to the different conic sections coincides with the corresponding extremizers given by Table 2. It is a constant function when $\tau = 1$ (sphere, red in Fig. 4) and a Gaussian when $\tau = -\xi_1 + 2$ (paraboloid, blue in Fig. 4). Furthermore, it yields the function

$\exp(-\sqrt{1+|\xi'|^2})$ when $\tau = \sqrt{1+|\xi'|^2}$ (hyperboloid, yellow in Fig. 4). Extremizers for the problem on the hyperboloid do not exist, but it was shown in [44, Lemma 5.4] that the function $f_a(\xi') = \exp(-a\sqrt{1+|\xi'|^2})$ produces an extremizing sequence $\{f_a/\|f_a\|_{L^2}\}$ as $a \rightarrow \infty$ for the $L^2(\lambda_2) \rightarrow L^4$ extension inequality on the hyperboloid.

FIGURE 4. Conic sections

APPENDIX A. INTEGRATION ON MANIFOLDS USING DELTA CALCULUS

Let $\mathcal{M}$ be a smooth k -dimensional submanifold of $\mathbb{R}^d$, with $0 < k < d$. On $\mathcal{M}$ one can define a canonical measure $\sigma = \sigma_{\mathcal{M}}$ which is naturally induced by the Euclidean metric structure of $\mathbb{R}^d$. Integration on the manifold $\mathcal{M}$ can be rigorously defined by means of differential forms [50], but the actual computation of integrals of the form

$$\int_{\mathcal{M}} \varphi(x) d\sigma$$

often appears to be a challenging task when $\mathcal{M}$ is a nontrivial manifold. The theory of distributions comes into help since these integrals can be viewed as pull-backs of Dirac delta distributions [28].

Suppose that $\mathcal{M}$ is (locally) implicitly described as the zero level set of a C^1 function f defined on an open subset Ω of $\mathbb{R}^d$ and taking values in $\mathbb{R}^{d-k}$,

$$\mathcal{M} = \mathcal{M}_f := \{x \in \Omega : f(x) = 0\},$$

and assume that the Jacobian matrix $Df(x)$ of the map f has maximal rank at every point $x \in \Omega$. Consider the usual Dirac delta distribution $\delta(\cdot)$ on $\mathbb{R}^{d-k}$ defined by

$$\langle \delta(\cdot), \varphi \rangle = \int_{\mathbb{R}^{d-k}} \delta(y) \varphi(y) dy = \varphi(0),$$

for any smooth test functions φ . We would like to make sense of the composition $\delta(f(x))$ as a distribution on Ω . There are several ways to define such a composition. One possibility is to approximate the delta distribution with a family of smooth bump functions,

$$\gamma_\varepsilon(y) = \varepsilon^{k-d} \gamma(\varepsilon^{-1}y), \quad \text{where } 0 \leq \gamma \in C_c^\infty(\mathbb{R}^{d-k}), \quad \int_{\mathbb{R}^{d-k}} \gamma(y) dy = 1.$$

One is led to define

$$\int_{\Omega} \delta(f(x)) \varphi(x) dx = \langle \delta(f), \varphi \rangle := \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \gamma_\varepsilon(f(x)) \varphi(x) dx,$$

for every $\varphi \in C_c^\infty(\Omega)$. This limit converges to the integral

$$(A.1) \quad \int_{\Omega} \delta(f(x)) \varphi(x) dx = \int_{\mathcal{M}} \frac{\varphi(x)}{J_f(x)} d\sigma,$$

where $J_f(x)$ is the Jacobian determinant of the function f at the point x ,

$$J_f(x) = \sqrt{\det(Df(x) \cdot Df(x)^t)} = |df_1(x) \wedge df_2(x) \wedge \cdots \wedge df_{d-k}(x)|,$$

and here $\wedge$ stands for the wedge product of differential forms. Therefore integrals over the manifold $\mathcal{M}$ can be expressed as integrals over $\mathbb{R}^d$ by means of delta distributions:

$$\int_{\mathcal{M}} \varphi(x) d\sigma = \int_{\mathbb{R}^d} \delta(f(x)) \varphi(x) J_f(x) dx.$$

Let us now have a look at some simple but useful algebraic rules which follow easily from these definitions, and allow for manipulation of integrals with delta distributions. These rules have been used in the previous sections to carry out explicit computations of convolution measures defined on the various manifolds considered there.

A.1. Hypersurfaces. In the codimension 1 case, $k = d - 1$, we have that $\mathcal{M}$ is a hypersurface defined by a scalar function. Identity (A.1) simplifies to

$$(A.2) \quad \int_{\Omega} \delta(f(x)) \varphi(x) dx = \int_{\mathcal{M}} \frac{\varphi(x)}{|\nabla f(x)|} d\sigma.$$

For example, in the case of the unit sphere $\mathbb{S}^{d-1}$ equipped with surface measure σ , we have that $f(x) = |x| - 1$, $|\nabla f(x)| = 1$, and

$$\int_{\mathbb{S}^{d-1}} \varphi(x) d\sigma = \int_{\mathbb{R}^d} \delta(|x| - 1) \varphi(x) dx.$$

For another example, consider the region $\Omega_- = \{x \in \Omega : f(x) < 0\}$, with boundary inside Ω given by $\mathcal{M}$, and with outgoing unit normal vector given by $\nu = \frac{\nabla f}{|\nabla f|}$. Combining the divergence theorem with formula (A.2), we have the following: For any smooth vector field $V(x)$ with compact support in Ω ,

$$\int_{\Omega_-} (\nabla \cdot V)(x) dx = \int_{\mathcal{M}} V(x) \cdot \nu(x) d\sigma = \int_{\Omega} \delta(f(x)) V(x) \cdot \nabla f(x) dx.$$

A.2. Products of delta distributions. We can make sense of the product of two delta distributions by simply setting

$$\delta(f(x)) \delta(g(x)) := \delta\left(\begin{smallmatrix} f(x) \\ g(x) \end{smallmatrix}\right),$$

whenever the right-hand side is well-defined as a distribution supported on the manifold $\mathcal{M}_{(f,g)} = \mathcal{M}_f \cap \mathcal{M}_g$. We can also make sense of the integration of the distribution $\delta(f(x))$ over the manifold $\mathcal{M}_g$ as follows:

$$\begin{aligned} \text{(A.3)} \quad \int_{\mathcal{M}_g} \delta(f(x)) \varphi(x) d\sigma_{\mathcal{M}_g} &= \int_{\Omega} \delta\left(\begin{smallmatrix} f(x) \\ g(x) \end{smallmatrix}\right) \varphi(x) J_g(x) dx \\ &= \int_{\mathcal{M}_f \cap \mathcal{M}_g} \varphi(x) \frac{J_g(x)}{J_{(f,g)}(x)} d\sigma_{\mathcal{M}_{(f,g)}}. \end{aligned}$$

In the case of codimension $d - k > 1$, the delta distribution can always be viewed (locally) as a product of $d - k$ delta distributions on hypersurfaces.

For example, let σ denote the surface measure on the $(d - 1)$ -dimensional unit sphere in $\mathbb{R}^d$, and let h be a smooth function on the unit sphere. The convolution $h\sigma * h\sigma$ is supported on the ball of radius 2 centered at the origin, and its value at a point x inside that ball can be written as an integral over the $(d - 2)$ -dimensional sphere Γ_x obtained as the intersection of the unit sphere with its translate by x . The sphere Γ_x has radius $\sqrt{1 - \frac{|x|^2}{4}}$. We can write $\Gamma_x = \mathcal{M}_f \cap \mathcal{M}_g$, with $f(y) = |y| - 1$ and $g(y) = |x - y| - 1$. If $y \in \Gamma_x$, then

$$J_{(f,g)}(y) = |df(y) \wedge dg(y)| = \left| \frac{x - y}{|x - y|} \wedge \frac{y}{|y|} \right| = |x \wedge y| = |x| \sqrt{1 - \frac{|x|^2}{4}}.$$

FIGURE 5. $\Gamma_x = \mathbb{S}^{d-1} \cap (x + \mathbb{S}^{d-1})$

Using formula (A.3), we obtain a generalization of formula (3.2): If $|x| \leq 2$, then

$$\begin{aligned}
 (h\sigma * h\sigma)(x) &= \int_{\mathbb{R}^d} \delta\left(\frac{|x-y|-1}{|y|-1}\right) h(x-y)h(y) \, dy \\
 &= \int_{\Gamma_x} \frac{h(x-y)h(y)}{J_{(f,g)}(y)} \, d\sigma_y = \frac{2}{|x| \sqrt{4-|x|^2}} \int_{\Gamma_x} h(x-y)h(y) \, d\sigma_y \\
 &= \frac{V_{d-2}}{|x|} \left(1 - \frac{|x|^2}{4}\right)^{\frac{d-3}{2}} \oint_{\Gamma_x} h(x-y)h(y) \, d\sigma_y,
 \end{aligned}$$

where V_{d-2} is the surface volume of the $(d-2)$ -dimensional unit sphere and $\oint$ denotes the averaged integral.

A.3. Multiplication by scalar functions. Let $\alpha : \Omega \rightarrow \mathbb{R}$ be a positive C^1 scalar function. Then, on the manifold $\mathcal{M}_{\alpha f} = \mathcal{M}_f$, we have that $J_{\alpha f} = \alpha^{d-k} J_f$, and consequently

$$(A.4) \quad \delta(\alpha(x)f(x)) = \alpha(x)^{-d+k} \delta(f(x)).$$

In particular, if α and β are smooth positive scalar functions which coincide on $\mathcal{M}_f$, then $\delta(\alpha f) = \delta(\beta f)$. For example, on the unit sphere, we have that

$$\delta(|x|^2 - 1) = \frac{1}{|x| + 1} \delta(|x| - 1) = \frac{1}{2} \delta(|x| - 1).$$

In a similar way, on the null cone, we have that

$$\delta(t^2 - |x|^2) = \frac{\delta(t - |x|)}{t + |x|} - \frac{\delta(t + |x|)}{t - |x|} = \frac{\delta(t - |x|)}{2|x|} + \frac{\delta(t + |x|)}{2|x|}.$$

A.4. Change of variables. Suppose that $x = \Psi(y)$ is a local diffeomorphism on $\mathbb{R}^d$. As for any distribution, we have the usual change of variables rule,

$$(A.5) \quad \int_{\Omega} \delta(f(x)) \varphi(x) dx = \int_{\Psi^{-1}(\Omega)} \delta(f(\Psi(y))) \varphi(\Psi(y)) |\det D\Psi(y)| dy.$$

If instead we consider the composition $g(x) = \Phi(f(x))$, with Φ a local diffeomorphism defined on a neighborhood of 0 in $\mathbb{R}^{d-k}$ and satisfying $\Phi(0) = 0$, then $\mathcal{M}_g = \mathcal{M}_f$, and

$$\delta(\Phi(f(x))) = |\det D\Phi(f(x))|^{-1} \delta(f(x)).$$

In particular, if L is a nonsingular $d \times d$ matrix and M is a nonsingular $(d-k) \times (d-k)$ matrix, then we have that

$$\int_{L^{-1}(\Omega)} \delta(Mf(Lx)) \varphi(x) dx = \frac{1}{|\det L \det M|} \int_{\Omega} \delta(f(y)) \varphi(L^{-1}y) dy.$$

ACKNOWLEDGEMENTS

The second author expresses his gratitude to Dmitriy Bilyk, Feng Dai, Vladimir Temlyakov and Sergey Tikhonov for organizing the workshop *Function spaces and high-dimensional approximation* at the Centre de Recerca Matemàtica-ICREA, May 2016, whose hospitality is greatly appreciated.

REFERENCES

- [1] W. Beckner, *Inequalities in Fourier analysis*, Ann. of Math. (2) **102** (1975), no. 1, 159–182.
- [2] J. Bennett, N. Bez, A. Carbery and D. Hundertmark, *Heat-flow monotonicity of Strichartz norms*, Anal. PDE **2** (2009), no. 2, 147–158.
- [3] J. Bennett, N. Bez and M. Iliopoulou, *Flow monotonicity and Strichartz inequalities*, Int. Math. Res. Not. IMRN (2015), no. 19, 9415–9437.
- [4] J. Bennett, N. Bez, C. Jeavons and N. Pattakos, *On sharp bilinear Strichartz estimates of Ozawa–Tsutsumi type*, Preprint, 2014. arXiv:1404.2466. To appear in the J. Math. Soc. Japan.
- [5] J. Bennett, A. Carbery, T. Tao, *On the multilinear restriction and Kakeya conjectures*, Acta Math. **196** (2006), no. 2, 261–302.
- [6] N. Bez and C. Jeavons, *A sharp Sobolev–Strichartz estimate for the wave equation*, Electron. Res. Announc. Math. Sci. **22** (2015), 46–54.
- [7] N. Bez and K. Rogers, *A sharp Strichartz estimate for the wave equation with data in the energy space*, J. Eur. Math. Soc. (JEMS) **15** (2013), no. 3, 805–823.
- [8] N. Bez and M. Sugimoto, *Optimal constants and extremisers for some smoothing estimates*, Preprint, 2012. arXiv:1206.5110.
- [9] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations*, Geom. Funct. Anal. **3** (1993), no. 2, 107–156.
- [10] J. Bourgain and C. Demeter, *The proof of the ℓ^2 decoupling conjecture*, Ann. of Math. (2) **182** (2015), no. 1, 351–389.
- [11] A. Bulut, *Maximizers for the Strichartz inequalities for the wave equation*, Differential Integral Equations **23** (2010), no. 11–12, 1035–1072.
- [12] E. Carneiro, *A sharp inequality for the Strichartz norm*, Int. Math. Res. Not. IMRN (2009), no. 16, 3127–3145.
- [13] E. Carneiro and D. Oliveira e Silva, *Some sharp restriction inequalities on the sphere*, Int. Math. Res. Not. IMRN (2015), no. 17, 8233–8267.

- [14] E. Carneiro, D. Foschi, D. Oliveira e Silva and C. Thiele, *A sharp trilinear inequality related to Fourier restriction on the circle*, Preprint, 2015. arXiv:1509.06674. To appear in Rev. Mat. Iberoam.
- [15] M. Christ and R. Quilodrn, *Gaussians rarely extremize adjoint Fourier restriction inequalities for paraboloids*, Proc. Amer. Math. Soc. **142** (2014), no. 3, 887–896.
- [16] M. Christ and S. Shao, *Existence of extremals for a Fourier restriction inequality*, Anal. PDE. **5** (2012), no. 2, 261–312.
- [17] M. Christ and S. Shao, *On the extremizers of an adjoint Fourier restriction inequality*, Adv. Math. **230** (2012), no. 3, 957–977.
- [18] A. Crdoba, *Singular integrals, maximal functions and Fourier restriction to spheres: the disk multiplier revisited*, Adv. Math. **290** (2016), 208–235.
- [19] T. Duyckaerts, F. Merle and S. Roudenko, *Maximizers for the Strichartz norm for small solutions of mass-critical NLS*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **10** (2011), no. 2, 427–476.
- [20] A. Erdlyi, W. Magnus, F. Oberhettinger and F. Tricomi, *Higher transcendental functions. Vol. II*. Robert E. Krieger Publishing Co. Inc., Melbourne, FL, 1981.
- [21] L. Fanelli, L. Vega and N. Visciglia, *On the existence of maximizers for a family of restriction theorems*, Bull. Lond. Math. Soc. **43** (2011), no. 4, 811–817.
- [22] L. Fanelli, L. Vega and N. Visciglia, *Existence of maximizers for Sobolev–Strichartz inequalities*, Adv. Math. **229** (2012), no. 3, 1912–1923.
- [23] D. Foschi, *Maximizers for the Strichartz inequality*, J. Eur. Math. Soc. (JEMS) **9** (2007), no. 4, 739–774.
- [24] D. Foschi, *Global maximizers for the sphere adjoint Fourier restriction inequality*, J. Funct. Anal. **268** (2015), no. 3, 690–702.
- [25] D. Foschi and S. Klainerman, *Bilinear space-time estimates for homogeneous wave equations*, Ann. Sci. cole Norm. Sup. (4) **33** (2000), no. 2, 211–274.
- [26] R. Frank, E. H. Lieb and J. Sabin, *Maximizers for the Stein–Tomas inequality*, Geom. Funct. Anal. **26** (2016), no. 4, 1095–1134.
- [27] J. Ginibre and G. Velo, *On a class of nonlinear Schrdinger equations. I. The Cauchy problem, general case*, J. Funct. Anal. **32** (1979), no. 1, 1–32.
- [28] L. Hrmander, *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 256. Springer-Verlag, Berlin, 1983.
- [29] D. Hundertmark and S. Shao, *Analyticity of extremizers to the Airy–Strichartz inequality*, Bull. Lond. Math. Soc. **44** (2012), no. 2, 336–352.
- [30] D. Hundertmark and V. Zharnitsky, *On sharp Strichartz inequalities in low dimensions*, Int. Math. Res. Not. IMRN (2006), Art. ID 34080, 1–18.
- [31] C. Jeavons, *A sharp bilinear estimate for the Klein–Gordon equation in arbitrary space-time dimensions*, Differential Integral Equations **27** (2014), no. 1–2, 137–156.
- [32] J.-C. Jiang, B. Pausader and S. Shao, *The linear profile decomposition for the fourth order Schrdinger equation*, J. Differential Equations **249** (2010), no. 10, 2521–2547.
- [33] J.-C. Jiang, S. Shao and B. Stovall, *Linear profile decompositions for a family of fourth order Schrdinger equations*, Preprint, 2014. arXiv:1410.7520.
- [34] S. Klainerman and M. Machedon, *Space-time estimates for null forms and the local existence theorem*, Comm. Pure Appl. Math. **46** (1993), 1221–1268.
- [35] M. Kunze, *On the existence of a maximizer for the Strichartz inequality*, Comm. Math. Phys. **243** (2003), no. 1, 137–162.
- [36] E. H. Lieb, *Sharp constants in the Hardy–Littlewood–Sobolev and related inequalities*, Ann. of Math. (2) **118** (1983), no. 2, 349–374.
- [37] D. Oliveira e Silva, *Extremizers for Fourier restriction inequalities: convex arcs*, J. Anal. Math. **124** (2014), 337–385.
- [38] D. Oliveira e Silva, *Nonexistence of extremizers for certain convex curves*, Preprint, 2012. arXiv:1210.0585. To appear in Mat. Res. Lett.

- [39] D. Oliveira e Silva and R. Quilodrn, *On extremizers for Strichartz estimates for higher order Schrdinger equations*, Preprint, 2016. arXiv:1606.02623.
- [40] D. Oliveira e Silva and C. Thiele, *Estimates for certain integrals of products of six Bessel functions*, Preprint, 2015. arXiv:1509.06309. To appear in Rev. Mat. Iberoam.
- [41] T. Ozawa and K. Rogers, *Sharp Morawetz estimates*, J. Anal. Math. **121** (2013), 163–175.
- [42] T. Ozawa and K. Rogers, *A sharp bilinear estimate for the Klein–Gordon equation in $\mathbb{R}^{1+1}$* , Int. Math. Res. Not. IMRN 2014, no. 5, 1367–1378.
- [43] R. Quilodrn, *On extremizing sequences for the adjoint restriction inequality on the cone*, J. Lond. Math. Soc. (2) **87** (2013), no. 1, 223–246.
- [44] R. Quilodrn, *Nonexistence of extremals for the adjoint restriction inequality on the hyperboloid*, J. Anal. Math. **125** (2015), 37–70.
- [45] J. Ramos, *A refinement of the Strichartz inequality for the wave equation with applications*, Adv. Math. **230** (2012), no. 2, 649–698.
- [46] A. Selvitella, *Remarks on the sharp constant for the Schrdinger Strichartz estimate and applications*, Electron. J. Differential Equations 2015, No. 270, 19 pp.
- [47] S. Shao, *Maximizers for the Strichartz and the Sobolev–Strichartz inequalities for the Schrdinger equation*, Electron. J. Differential Equations (2009), No. 3, 13 pp.
- [48] S. Shao, *The linear profile decomposition for the Airy equation and the existence of maximizers for the Airy–Strichartz inequality*, Anal. PDE **2** (2009), no. 1, 83–117.
- [49] S. Shao, *On existence of extremizers for the Tomas–Stein inequality for S^1* , J. Funct. Anal. **270** (2016), no. 10, 3996–4038.
- [50] M. Spivak, *A comprehensive introduction to differential geometry. Vol. I*. Second edition. Publish or Perish, Inc., Wilmington, Del., 1979.
- [51] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [52] E. M. Stein and R. Shakarchi, *Functional analysis. Introduction to further topics in analysis*. Princeton Lectures in Analysis, 4. Princeton University Press, Princeton, NJ, 2011.
- [53] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton Mathematical Series, 32. Princeton University Press, Princeton, NJ, 1971.
- [54] R. Strichartz, *Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations*, Duke Math. J. **44** (1977), no. 3, 705–714.
- [55] T. Tao, *Some recent progress on the restriction conjecture*, Fourier analysis and convexity, 217–243, Appl. Numer. Harmon. Anal., Birkhuser Boston, Boston, MA, 2004.
- [56] P. Tomas, *A restriction theorem for the Fourier transform*, Bull. Amer. Math. Soc. **81** (1975), 477–478.
- [57] L. Vega, *El multiplicador de Schrdinger, la funcion maximal y los operadores de restriccin*, Ph.D. thesis, Universidad Autnoma de Madrid, 1988.
- [58] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Second Edition. Cambridge University Press, Cambridge, 1966.
- [59] T. Wolff, *Lectures on harmonic analysis*, University Lecture Series, 29. American Mathematical Society, Providence, RI, 2003.

DAMIANO FOSCHI, DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSIT DI FERRARA, VIA MACHIAVELLI 30, 44121 FERRARA, ITALY

E-mail address: damiano.foschi@unife.it

DIOGO OLIVEIRA E SILVA, HAUSDORFF CENTER FOR MATHEMATICS, 53115 BONN, GERMANY

E-mail address: dosilva@math.uni-bonn.de