Changes in the ankle muscles co-activation pattern after 5 years following total ankle joint replacement

De la Fuente, Carlos; Martinez-Valdes, Eduardo; Cruz-Montecinos, Carlos; Guzman-Venegas, Rodrigo ; Arriagada, David; Peña y Lillo, Roberto ; Henriquez, Hugo ; Carpes, Felipe

DOI:

[10.1016/j.clinbiomech.2018.09.019](https://doi.org/10.1016/j.clinbiomech.2018.09.019)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

De la Fuente, C, Martinez-Valdes, E, Cruz-Montecinos, C, Guzman-Venegas, R, Arriagada, D, Peña y Lillo, R, Henriquez, H & Carpes, F 2018, 'Changes in the ankle muscles co-activation pattern after 5 years following total ankle joint replacement', *Clinical Biomechanics*, vol. 59, pp. 130-135.
<https://doi.org/10.1016/j.clinbiomech.2018.09.019>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Changes in the ankle muscles co-activation pattern after 5 years following total ankle joint replacement

Carlos De la Fuente^{a,f,k} (delafuente@gmail.com), Eduardo Martinez-Valdes^{b,c} (e.a.martinezvaldes@bham.ac.uk), Carlos Cruz-Montecinos^{d,e} (ccmkine@gmail.com), Rodrigo Guzman-Venegas^f (rguzman@uandes.cl), David Arriagada^g (david.arriagada.t@gmail.com), Roberto Peña y Lillo^{g,h} (robertopyl@gmail.com), Hugo Henríquez^{h,i} (hhenrquez@yahoo.com.ar), Felipe P Carpes^j ✉ (carpes@unipampa.edu.br).

^a Carrera de Kinesiología, Departamento de Cs. de la Salud, Facultad de Medicina, Pontificia Universidad Católica, #7820436 Santiago, Chile.

^b School of Sport, Exercise and Rehabilitation Sciences, Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), College of Life and Environmental Sciences, University of Birmingham, # Edgbaston B15 2TT Birmingham, United Kingdom.

^c Centro de Investigación en Fisiología del Ejercicio (CIFE), Universidad Mayor, Santiago, Chile

^d Laboratory of Biomechanics and Kinesiology, Hospital San José, #8380419 Santiago, Chile.

^e Department of Physical Therapy, Faculty of Medicine, Universidad de Chile, #8380453 Santiago, Chile.

^f Laboratorio LIBFE. Escuela de Kinesiología. Universidad de los Andes, #7620086 Santiago, Chile.

^g Kinesiología, Universidad de Santiago de Chile, #9170124 Santiago, Chile.

^h Servicio de Tobillo y Pie, Instituto Traumatológico, #8340220 Santiago, Chile.

ⁱ Facultad de Medicina, Universidad de Chile, #8380419 Santiago, Chile.

^j Laboratory of Neuromechanics, Universidade Federal do Pampa, Campus Uruguaiana, #97508000 Uruguaiana, Brazil.

^k Centro de Salud Deportivo, Clinica Santa Maria, #8340518, Santiago, Chile.

* Corresponding author (✉).

Felipe Carpes. Email address: carpes@unipampa.edu.br (F. Carpes). Address: #97508000 Uruguaiana, Brazil.

ABSTRACT

Background: The Hintegra® arthroplasty provides inversion-eversion stability, permits axial rotation, ankle flexion-extension, and improvements of the gait patterns are expected up to 12 months of rehabilitation. However, sensorimotor impairments are observed in ankle flexors/extensors muscles after rehabilitation, with potential negative effects on locomotion. Here we determined the timing and amplitude of co-activation of the tibialis anterior and medial gastrocnemius muscles during gait by assessing non-operated and operated legs of patients with total ankle replacement, 5 years after surgery.

Methods: Twenty-nine patients (age: 58 [5.5] years, height: 156.4 [6.5] cm, body mass: 72.9 [6.5] kg, 10 men, and 19 women) that underwent Hintegra® ankle arthroplasty were included. Inclusion criteria included 5 years prosthesis survivorship. The onset and offset of muscle activation (timing), as well as the amplitude of activation, were determined during barefoot walking at self-selected speed by surface electromyography. The timing, percentage, and index of co-activation between the tibialis anterior and medial gastrocnemius were quantified and compared between non-operated and operated legs.

Findings: The operated leg showed higher co-activation index and temporal overlapping between tibialis anterior and medial gastrocnemius during gait ($P < .001$).

Interpretation: The neuromuscular changes developed during the process of degeneration do not appear to be restored 5 years following arthroplasty. The insertion of an ankle implant may restore anatomy and alignment but neuromuscular adaptations to degeneration are not corrected by 5 years following joint replacement.

Abstract's word count: 229/250

Main text's word count: 3544/4000

Keywords: EMG; joint stiffness; total ankle replacement; walking; gastrocnemius; tibialis anterior.

1. Introduction

Lower limb osteoarthritis is associated with pain and impaired function with negative effects on locomotion. More specifically, ankle osteoarthritis impairs sagittal plane motion (dorsiflexion and plantarflexion) and torque (Al-Mohrougi et al., 2018). In many cases the joint degeneration will reach levels that require the total joint replacement. Ankle arthroplasty is a surgical procedure in which the tibiotalar joint is replaced (Giannini et al., 2000; Caravaggi et al., 2015). Eighty percent of candidates are patients who developed ankle osteoarthritis secondary to trauma (post-traumatic etiology) (Horisberger et al., 2009; Bloch et al., 2015), with far fewer outcomes reported compared to hip and knee arthroplasty (Al-Mohrougi et al., 2018). In the past 40 years more than 30 models of ankle prosthesis have been introduced, with few outcomes against hip and knee arthroplasty (Giannini et al., 2000; Henricson et al., 2011). In this regard, Hintegra® arthroplasty is a third generation prosthesis that provides inversion-eversion stability, permits axial rotation, ankle flexion-extension being one of the few ankle prosthesis showing good results concerning activities of daily living (Michael et al., 2008; Hintermann, 2005). Previous studies showed that the Hintegra prosthesis is built with stable components, with low rate of complications (Michael et al., 2008; Hintermann, 2005; Barg et al., 2011), allowing an adequate range of motion (Valderrabano et al., 2003; Michael et al., 2008; Hintermann, 2005). However, the lack of restoration of gait symmetry and joint function remains a problem after ankle arthroplasty (Caravaggi et al., 2015).

During the first year after Hintegra® arthroplasty, there are adaptations in the first three months of rehabilitation that negatively affects the gait biomechanics. These include lower maximal plantar-flexion moment, total adduction moment, medial ground reaction force, and higher anterior ground reaction force during gait by increasing mechanical loading on knee joint (Valderrabano et al., 2007). This adaptive period was previously reported as negative phase of rehabilitation (De la Fuente et al., 2014). These parameters are expected to improve towards the end of the first year after surgery (Valderrabano et al., 2007), with significant improvement of gait pattern (Valderrabano et al., 2007; Aidi et al., 2013) and quality of life (Esparragoza et al., 2011). It has been suggested that 5 years following total ankle joint replacement there is an increase in the American Orthopaedic Foot and Ankle Score (Zaidi et al., 2013). However, there is little information concerning the effect of ankle replacement on neuromuscular activity of ankle muscles (tibialis anterior and medial gastrocnemius) (Doets et al., 2007). Abnormal muscle co-activation could result in increased and abnormal kinetics on the prosthesis predisposing to loosening. Presumably the prosthesis is designed to replicate

normal anatomy of the ankle joint. If however the muscular co-activation does not change the forces will remain high.

An adequate control of tibialis anterior and medial gastrocnemius muscles depends both on the timing of agonist and antagonist activation (Hubley-Kozey et al., 2010; Rosa et al., 2010), and the intensity of their contractions, i.e. weight coefficients (Cappelini et al., 2006). These muscles sometimes present co-activation, an involuntary and concurrent activation of the antagonist, in opposition to the contraction of the agonist (Duchateau et al., 2014). Changes in co-activation occur during different physical activities e.g. running compared to walking (Cappelini et al., 2006), footwear selection (Alkjær et al., 2012) and in pathological processes e.g. cerebellar ataxia (Mari et al., 2014), foot and knee motor functions (Di Nardo et al., 2015), and ankle osteoarthritis (Von Tscharnier & Valderrabano, 2010). Furthermore, co-activation may affect gait pattern and the integrity of prostheses due to abnormal joint force development (De la Fuente et al., 2014).

After the Hintegra[®] arthroplasty placement, the gait pattern is expected to be restored, and co-activation between tibialis anterior and medial gastrocnemius should not occur at medium term after the first three months of rehabilitation (Michael et al., 2008; Hintermann, 2005). However, there is a lack of evidence concerning patterns of co-activation after ankle replacement when rehabilitation is completed. Therefore, here we determine the relative measures of amplitude and timing of co-activation of the tibialis anterior and medial gastrocnemius muscles during gait, by comparing the electromyographic activity of the non-operated and operated leg of patients with total ankle replacement, 5 years after surgery. Our null-hypotheses were: i) total ankle replacement after 5 years does not recover the timing of co-activation to the level of the non-operated leg during gait; ii) total ankle replacement after 5 years does not recover the temporal percentage of co-activation to the level of the non-operated leg during gait, and ii) total ankle replacement after 5 years does not recover the index of co-activation to the level of the non-operated leg during gait.

2. Methods

2.1. Study design

This is a cross-sectional, observational, analytical study design. The sample included twenty-nine patients (10 men and 19 women) that underwent unilateral total ankle replacement 5 years before the measurement session in which the operated and non-operated leg were compared during gait. The inclusion criteria were: i) unilateral

total ankle replacement with Hintegra[®] arthroplasty (Newdeal SA, Vienne, France); ii) at least 5 years following replacement; iii) rehabilitation treatment of one year; iv) posttraumatic arthritis; v) age between 40 and 70 years old; vi) passive dorsal range of motion greater than 5° at the sagittal plane (De la Fuente et al., 2014); vi) radiological stability (Horisberger et al., 2009; Bai et al., 2010; Guyer and Richardson, 2008); and v) have been rehabilitated at the *Instituto Traumatológico* (Santiago, Chile) by the same surgical and rehabilitation team. The exclusion criteria were: i) need for assisted locomotion; ii) major limitations in performance of daily life activities; iii) major periarticular tissue impairment; iv) ipsilateral or contralateral hip/knee osteoarthritis; v) inflammatory diseases; vi) neurological pathology; vii) active infection; and viii) cognitive impairment (von Tscharner and Valderrabano, 2010). This study was approved by the institutional review board of the *Instituto Traumatológico* (Santiago, Chile) according to the principles of the Declaration of Helsinki. All participants signed a consent term agreeing to participate in this study.

2.2. Sample size

The sample size was estimated “a priori” with a pilot experiment that included 6 patients [mean (standard deviation) 56.5 (2.1) years-old, 30.1 (1.8) kg/m², 3 men and 3 women] that fulfilled the inclusion and exclusion criteria. A sample size of 23 patients was estimated considering a difference between two dependent means (matched pairs), using two-tailed *t*-test with alpha error of 5% and statistical power of 80% for an estimated effect size of 0.62. Six additional patients were included to the sample to anticipate possible attrition (20% of estimation). The total patients assessed were 35 patients; 6 patients from the “a priori” determination of sample size and additional 29 patients. The statistical calculus was performed by G*Power software version 3.1.9.2. (Kiel University, Germany).

2.3. Surgery and physical therapy procedures

All participants had a Hintegra[®] prosthesis (Hintermann, 2005) (Newdeal SA, Vienne, France). In general terms, surgery involved an anterior longitudinal incision of 10 to 12 cm performed to dissect the retinaculum. Moreover, the soft tissue and periosteum from the bone were dissected. Resection of the talus and tibia were performed to insert arthroplasty components using Hintermann[®] distractor and oscillating saw. Osteophytes on the talar neck and anterior aspect of medial malleolus were also removed. After that, the tibial and talar

components were inserted. The last inserted component was the polyethylene component. The tissues were sutured, and the procedure was finalized by fitting a short leg cast. The patients were monitored for the next three weeks after surgery, once per week, and during the first 4 weeks after surgery patients were immobilized with a short leg cast and instructed to unload the operated leg and rest.

The physiotherapy intervention was performed from week 4 until week 52 (Ingrosso et al., 2009). From the week 4 to 12, the patients attended the rehabilitation service, where they used a walking boot. In this period, partial weight bearing was permitted using the assistance of canes. Furthermore, stretching to improve the dorsal flexion range of motion, strengthening of ankle, knee and hip muscles, and pain relief with physical agents were activities performed. The re-education of gait without assistance, bipedal heel rise, balance exercises, and pain relief management were performed until the end of week 52 (Martin et al., 2007). Afterwards, patients returned to the foot and ankle hospital unit every six months to be assessed in the follow-up period.

2.4. Data acquisition and processing

The data acquisition consisted of two stages: clinical assessment and surface electromyography recordings. The patients attended an interview, and the clinical assessment was performed at the foot and ankle service of the *Instituto Traumatológico* (Santiago, Chile). The age, body mass, height, AOFAS score (Kitaoka et al., 1994), passive dorsiflexion range of movement, difference in calf whilst standing (Saxena et al., 2011; Valdebarrano et al., 2006), and intensity of pain at rest and during walking assessed with a numerical verbal scale (0 no pain, 10 maximum possible pain) (Hintermann, 2005) and were part of the clinical assessment.

Surface electromyography recordings were performed one week after the clinical assessment at the *Centro de Investigaciones Medicas del Instituto Traumatológico* (Santiago, Chile). After a 5-min warm-up on a cycloergometer without external load and cadence of 60 rpm, patients walked barefoot at self-selected speed (von Tscharnier and Valdebarrano, 2010; De la Fuente et al., 2014) along a 5-meter flat surface on a straight-line. Patients performed five trials for familiarization within the walking space. Although we did not measure kinematics, a common heel strike pattern was observed among the patients. The electromyography signals were acquired during walking using a Myomonitor IV electromyography amplifier (Delsys, inc., Boston, USA). Two DE-2.3 single differential surface electromyography sensors (Delsys, inc., Boston, USA) with an inter-electrode

distance of 10 mm were used. The data collection employed a 16-bit analog-digital converter card (National Instrument Corp., Austin, TX, USA) operated by a Matlab software (Mathworks Inc., Massachusetts, USA) at a sampling rate of 1000 Hz, band-pass filtered (20–450 Hz), and hardware amplified with a gain of 1000 V/V. The muscles assessed were tibialis anterior and medial gastrocnemius based on their lower mean electromyography frequency and intensity previously found by Valderrabano et al. (2006) in unilateral ankle osteoarthritis, and the hypothesis of Doets et al. (2007), who suggested that after ankle joint replacement, higher co-activation between these muscles could exist during gait. The surface electromyography sensors were placed according to the European recommendations for surface electromyography (Hermens et al., 2000).

2.5. Signal treatment

The electromyography signals were filtered by a zero-lag 4th order finite impulse response Butterworth with a band pass of 20 to 450 Hz. Onset and offset times of muscle activation (Caravaggi et al., 2015) were identified using a continuous wavelet with the algorithm proposed by Merlo et al. (2003). The algorithm was performed using the Hermite-Rodriguez mother wavelet, 10% of noise power, 150 ms for the time of two detected activation intervals, and 5 ms for the spike rejection from non-rectified signals. The noise of electromyography signals was extracted when patients stood quietly before walking. Five consecutive electromyography bursts of the tibialis anterior and gastrocnemius medialis (corresponding to 5 strides) were used for the analysis.

All signals were full wave rectified (Figure 1). Due to the variability and non-accordance of normalization methods for analysis of electromyography signals (Rosa et al., 2010), a standardized treatment of signals by z-score method was performed. Each sample was subtracted from its expected value ($E[X]$) and divided by the standard deviation. This expresses the number of standard deviations by which the sample is above the $E[X]$. As the whole electromyography signals showed a normal distribution, the z-score was obtained using the arithmetic mean of the data.

2.6. Outcomes

The timing of co-activation: determined by the time overlap between the onsets and offsets activation (Rosa et al., 2010).

Percentage of co-activation: from the time activation onsets and offsets, the percentage of co-activation was quantified as the overlapped percentage of muscle activation (Rosa et al., 2010).

Co-activation index: determined by the overlapped amplitudes of muscle activation between the tibialis anterior and medial gastrocnemius. The co-activation index was implemented in discrete form by the trapezoidal method (Eq. 1) from the original continuous form previously defined in the literature (Rosa et al., 2010).

$$Coactivation\ index = 2 \times \frac{\sum_{CoAc_{onset}}^{CoAc_{offset}} \left[\frac{(X_{i+1} - X_i)}{2} \times \Delta t \right]}{\sum_{Antagonist}^{Antagonist} \left[\sum_{onset}^{offset} \left(\frac{(X_{i+1} - X_i)}{2} \times \Delta t \right) \right]_j} \times 100\% \quad (Eq. 1)$$

In the equation 1, $CoAc_{onset/offset}$ is the co-activation time of onset or offset, x_i is the electromyography sample, Δt is the interval time of data acquisition given the sampling frequency, and j represents the agonist or antagonist condition. To assess the intensity of concurrent contractions, the trapezoidal areas were obtained using the “cumtrapz function” from the full-rectified signals using the Matlab software (Mathworks Inc., Massachusetts, USA).

2.7. Statistical analysis

Data were reported as the median and interquartile range [IQ range] because the Shapiro-Wilk test revealed a non-parametric data distribution. Homoscedasticity was confirmed using the Levene’s test. To compare the timing, percentage, and index of co-activation of medial gastrocnemius and tibialis anterior between the operated and non-operated legs, a Wilcoxon Signed-Ranks test of two-tails was used with alpha error equal to 5%. To assess the possible existence of co-variable and interaction with outcomes, a multiple regression respect to analysis for age, body mass, height, AOFAS score, the passive dorsal range of movement, calf circumference difference between leg during a standing posture, the intensity of pain at rest and during gait were considered at $p < 0.05$. Data were analyzed using the Matlab software statistical toolbox (Mathworks Inc., Massachusetts, USA).

3. Results

The timing of co-activation between the tibialis anterior and medial gastrocnemius in the non-operated leg [median: 390 ms, IQ range: 80 ms] was lower than in the operated leg [median: 566 ms, IQ range: 104 ms, $p<0.001$]. The percentage of co-activation between the tibialis anterioris and gastrocnemius medialis in the non-operated leg [median: 0.00 %, IQ range: 1.84 %] was lower than in the operated leg [median: 100%, IQ range: 0.00 % $p<0.001$]. The index of co-activation between the tibialis anterior and medial gastrocnemius in the non-operated leg [median: 0.00 % IQ range: 1.94 %] was lower than in the operated leg [median: 30.96%, IQ range: 13.52%, $p<0.001$]. The changes in co-activation are depicted in the Figure 1.

The demographic characteristics of the sample regarding age, body mass, height, AOFAS score, passive dorsal range of movement, calf circumference difference between legs during standing posture, intensity of pain at rest, and during gait did not co-vary or interacted with any of the electromyography outcomes ($p>0.05$).

*** Table 1 around here ***

*** Figure 1 around here ***

4. Discussion

The most important finding of this work was that patients with the Hintegra® arthroplasty present abnormal plantar and dorsi-flexion co-activation patterns during gait after 5 years following total ankle joint replacement. Although the patients did not receive rehabilitation after 5 years following total ankle joint replacement due to the improvement observed in the AOFAS score, the present findings suggest the need for an appropriate sensorimotor rehabilitation program for patients in which problems at medium term of survival of the prosthesis still persist.

Although it can't be ensured (due to the lack of kinetics and kinematics assessment), the altered co-activation found in our study may suggest that both absorption and propulsion phases of walking could be committed 5 years after replacement surgery. Dorsal and plantar-flexor muscles need to be activated during different phases of gait in order to maintain the normal mechanics of the ankle joint during swing and load

response. This is important for the control of the anterior advance of the tibia or the generation of propulsion. Therefore, these mechanical actions need temporal muscle coordination (timing of activation). Based on the findings from Di Nardo et al. (2015) and Cappellini et al. (2006), the tibialis anterior and medial gastrocnemius muscles should present independent temporal activations during the different phases of normal gait (low speed). It contrasts with the results of patients treated with ankle prosthesis, since a temporal overlap between these muscles was found.

Based on the findings from Siegler et al. (2013), an altered kinetic condition can be present in patients with Hintegra[®] arthroplasty. Siegler et al. (2013) tri-dimensionally modeled the talus of 26 healthy adults and found that “the trochlear surface can be modeled as a skewed truncated conic saddle shape with its apex oriented laterally rather than medially”, which contrasts with the mechanical model of Inman, in which the Hintegra[®] arthroplasty is based (Hinterman 2005). Thus, the mechanical design of the Hintegra[®] arthroplasty could alter distribution or intensity of ankle joint forces during the stance phase, or both, such as the abnormal joint shear forces and delayed the first peak of vertical ground reaction force found after the negative rehabilitation phase in patients treated with an Hintegra[®] arthroplasty (De la Fuente et al., 2014). Our data suggests that these mechanical changes could result of sensorimotor adaptations, since an increased co-activation between dorsal and plantar flexors would reduce the velocity of the foot and tibia during stance phase, which reduces the vertical component of the ground reaction force at load response phase. However, an increased co-activation might also result of intrinsic instability of the prosthesis in the frontal plane, which could have an unstable effect over the foot based on mechanical findings from Siegler et al. (2013), further leading to the shifting of the medial gastrocnemius rather than a shift of the tibialis anterior activation, as shown here.

As the shifting in the medial gastrocnemius could occur, the propulsion phase might also be affected (Giannini et al., 2000; Henricson et al., 2011). This possible change is in accordance with de la Fuente et al. (2014), who proposed a gait pattern with external hip rotation that increases the activity of hip extensors (rather than plantar flexors) to generate the propulsion of the leg with Hintegra[®] arthroplasty, after the negative rehabilitation phase. Therefore, it is possible that patients with Hintegra[®] arthroplasty experience an over-stabilization of the ankle joint after at medium term, affecting the whole pattern of gait by an unstable load response and altered propulsion phase. This hypothesis still needs to be investigated.

On the other hand, Stubbs et al. (1998) suggested that the load over viscoelastic tissues creates a sensorimotor response called “muscle-ligament reflex” causing an altered activity of agonist and antagonist

muscles over time. Therefore, the surgery itself could cause the neuromuscular changes observed, since the tibialis anterior and ankle ligaments must be pulled to introduce the arthroplasty components into the ankle joint space. This is important as one wonders if neuromuscular changes are reversible with time. Our findings would suggest that at 5 years after ankle replacement they are not reversible. In making this statement perhaps prosthesis designers would be better to use an arthroplasty designed to tolerate the irreversible forces on the prosthesis. Finally, chronic pain could be another source of the neuromuscular changes observed in our study. Valderrabano et al., (2006) and von Tscharner et al. (2010) described that the time of exposure to pain in osteoarthritis is the worst stimulus to the nervous system changing the neuromuscular patterns of activations. This could induce changes in the ankle stiffness leading to pathological neuroplasticity changes, possibly present since before the arthroplasty surgery. However, in our study, the intensity of pain was not identified as a co-variable and did not show any interaction with the outcomes.

Due to the higher co-activation found in patients with Hintegra® arthroplasty at 5 years after ankle replacement, new rehabilitation approaches are needed to improve the locomotion in patient who shows the sensorimotor impairments described here. To our knowledge, this is the first pathological report at medium term in patients with Hintegra® arthroplasty. Future studies should identify *in vivo* whether the ankle axis of this arthroplasty is really positioned as Inman previously reported, since alterations in the prosthesis axis of rotation could lead to the sensorimotor impairments found. Also, the presence of other concomitant musculoskeletal problems (e.g., knee, hip or trunk disorders, metatarsal-cuneiform, cuneiform-navicular and talo-navicular osteoarthritis) requires further examination. Finally, it is important to investigate whether previous pathological sensorimotor conditions acquired before the arthroplasty are irreversible after the surgery (i.e., chronic pain). Regarding the limitations of our study, the sample is only representative of patients with lower AOFAS score, in contrast to the results with Hintegra® (Hinterman, 2005) and other arthroplasties (Aidi et al., 2013; Caravaggi et al., 2015). Also, the measurement of pain considered a verbal pain scale (Hinterman, 2005), which may not reflect the sensorimotor impairments showed by previous studies (Ervilha et al., 2004; Lindstrøm et al., 2011). Finally, we propose that a detailed mechanical analysis of gait, in addition to the analysis of co-activation alteration, is needed to understand the possible neuro-mechanical pathological new hypotheses discussed in our study.

301 **5. Conclusions**

302 Altered co-activation during gait is present after at 5 years after total ankle replacement with the Hintegra®
303 prosthesis. It may result of an attempt to compensate ankle instabilities following arthroplasty, which effects on
304 gait dynamics require attention during rehabilitation programs.

305

306 **Conflict of interests**

307 The authors declare no conflicts of interest.

308

309 **Funding**

310 This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-
311 profit sectors. FPC is supported by a CNPq research fellowship.

312

313 **Acknowledgements**

314 We acknowledge the assistance in data collection provided by the undergraduate students Guillermo Gonzalez,
315 Javiera Asecio and Roberto Peña from Universidad de Chile.

6. References

- Aidi R., Cro S., Gurusamy K., Siva N., Macgregor A., Henricson A., Goldberg A. 2013. The outcome of total ankle replacement: a systematic review and meta-analysis. *Bone Joint J.* 95-B(11),1500-1507.
- Alkjær T., Raffalt P., Petersen N., Simonsen E. 2012. Movement behavior of high-heeled walking: how does the nervous system control the ankle joint during an unstable walking condition?. *PLoS One.* 7(5),e37390.
- Al-Mahrouqi M.M., MacDonald D.A., Vicenzino B., Smith M.D. 2018. Physical impairments in adults with ankle osteoarthritis: a systematic review and meta-analysis. *J Orthop Sports Phys Ther.* 7:1-43.
- Bai L., Lee K., Song E., Yoon T., Seon J. 2011. Total ankle arthroplasty outcome comparison for post-traumatic and primary osteoarthritis. *Foot Ankle Int.* 31(12),1048-1058.
- Barg A., Knupp M., Anderson A., Hintermann B. 2011. Total ankle replacement in obese patients: component stability, weight change, and functional outcome in 118 consecutive patients. *Foot Ankle Int.* 32(10),925-932.
- Bloch B., Srinivasan S., Mangwani J. 2015. Current Concepts in the Management of Ankle Osteoarthritis: A Systematic Review. *J Foot Ankle Surg.* 54(5),932-939.
- Cappellini G., Ivanenko Y., Poppele R., Lacquaniti F. 2006. Motor patterns in human walking and running *J of Neurophysiol.* 95(6), 3426-3437.
- Caravaggi P., Lullini G., Leardini A., Berti L., Vannini F., Giannini S. 2015. Functional and clinical evaluation at 5-year follow-up of a three-component prosthesis and osteochondral allograft transplantation for total ankle replacement. *Clin Biomech (Bristol, Avon).* 30(1),59-65.
- De la Fuente C., Carcuro G., Ramírez-Campillo R., Campos C., Soza F. 2015. Prosthetic wear in subjects with Hintegra total ankle joint replacement secondary to trauma in functional phase. *Fisioterapia.* 37,165-174
- Di Nardo F., Mengarelli A., Maranesi E., Burattini L., Fioretti S. 2015. Assessment of the ankle muscle co-contraction during normal gait: a surface electromyography study. *J Electromyogr Kinesiol.* 25(2),347-354.
- Doets H.C., van Middelkoop M., Houdijk H., Nelissen R.G., Veeger H.E. 2007. Gait analysis after successful mobile bearing total ankle replacement. *Foot Ankle Int.* 28(3),313-322.
- Dyer J.O., Maupas E., de Andrade Melo S., Bourbonnais D., Nadeau S., Forget R. 2014. Changes in activation timing of knee and ankle extensors during gait are related to changes in heteronymous spinal pathways after stroke. *J Neuroeng Rehabil.* 11,148.
- Ervilha U.F., Arendt-Nielsen L., Duarte M., Graven-Nielsen T. 2004. The effect of muscle pain on elbow flexion and coactivation tasks. *Exp Brain Res.* 156(2),174-182.

345 Esparragoza L., Vidal C., Vaquero J. 2011. Comparative study of the quality of life between arthrodesis and
346 total arthroplasty substitution of the ankle. *J Foot Ankle Surg.* 50(4),383–387.

347 Giannini S., Leardini A., O’connor J. 2000. Total ankle replacement: Review of the designs and of the current
348 status. *Foot AnkleSurg.* 6,77-88.

349 Guyer A., Richardson G. 2008. Current concepts review: total ankle arthroplasty. *Foot Ankle Int.* 29(2),256-
350 264.

351 Henricson A., Nilsson J., Carlsson A. 2011. 10 year survival of total ankle arthroplasties: a report on 780 cases
352 from the Swedish Ankle Register. *Acta Orthop.* 82(6),655-659.

353 Hermens H.J., Freriks B., Disselhorst-Klug C., Rau G. 2000. Development of recommendations for SEMG
354 sensors and sensor placement procedures. *J Electromyogr Kinesiol.* 10(5), 361-374.

355 Hintermann B. 2005. Total ankle arthroplasty: historical overview, current concepts and future perspectives,
356 illustrated ed. Austria: Springer.

357 Horisberger M., Hintermann B., Valderrabano V. 2009. Alterations of plantar pressure distribution in
358 posttraumatic end-stage ankle osteoarthritis. *Clin Biomech (Bristol, Avon).* 24,303-307.

359 Hubley-Kozey C.L., Hatfield G.L., Davidson K.C. 2010. Temporal coactivation of abdominal muscles during
360 dynamic stability exercises. *J Strength Cond Res.*24(5),1246-1255.

361 Ingrosso S., Benedetti M., Leardini A., Casanelli S., Giannini S. 2009. Gait analysis in patients operated with a
362 novel total ankle prosthesis. *Gait Posture.* 30,132-137.

363 Kitaoka H., Alexander I., Adelar R. 1994. Clinical rating systems for the ankle, hindfoot, midfoot, hallux, and
364 lesser toes. *Foot Ankle Int.* 15(7),349–353.

365 Lindstrøm R., Schomacher J., Farina D., Rechter L., Falla D. 2011. Association between neck muscle
366 coactivation, pain, and strength in women with neck pain. *Man Ther.*16(1),80-86.

367 Mari S., Serrao M., Casali C., Conte C., Martino G., Ranavolo A., Coppola G., Draicchio F., Padua L., Sandrini
368 G, Pierelli F. 2014. Lower limb antagonist muscle co-activation and its relationship with gait parameters in
369 cerebellar ataxia. *Cerebellum.*13(2),226-236.

370 Martin R.L., Stewart G.W., Conti S.F. 2007. Posttraumatic ankle arthritis: an update on conservative and
371 surgical management. *J Orthop Sports Phys Ther.* 37(5),253–259.

372 Merlo A., Farina D., Merletti R. 2003. A fast and reliable technique for muscle activity detection from surface
373 EMG signals. *IEEE Trans Biomed Eng.* 50(3),316-323.

374 Michael J., Golshani A., Gargac S., Goswami T. 2008. Biomechanics of the joint clinical outcomes of total
 375 ankle replacement. *J Mech Behav Biomed Mater.* 1(4),276-294.
 376 Rosa M.C., Marques A., Demain S., Metcalf C.D., Rodrigues J. 2014. Methodologies to assess muscle co-
 377 contraction during gait in people with neurological impairment - a systematic literature review. *J Electromyogr*
 378 *Kinesiol.* 24(2),179-191.
 379 Saxena A., Ewen B., Mafulli N. 2011. Rehabilitation of the operated achilles tendon: parameters for predicting
 380 return to activity. *J Foot Ankle Surg.* 50(1),37-40.
 381 Siegler S., Toy J., Seale D., Pedowitz D. 2014. The Clinical Biomechanics Award 2013 -- presented by the
 382 International Society of Biomechanics: new observations on the morphology of the talar dome and its
 383 relationship to ankle kinematics. *Clin Biomech (Bristol, Avon).* 29(1),1-6.
 384 Stubbs M., Harris M., Solomonow M., Zhou B., Lu Y., Baratta R.V. 1998. Ligamento-muscular protective
 385 reflex in the lumbar spine of the feline. *J Electromyogr Kinesiol.* 8(4), 197-204.
 386 Valderrabano V., Hintermann B., Nigg B.M., Stefanyshyn D., Stergiou P. 2003. Kinematic changes after fusion
 387 and total replacement of the ankle: part 1:Range of motion. *Foot Ankle Int.* 24(12),881-887.
 388 Valderrabano V., Nigg B., von Tscharnner V., Stefanyshyn D., Goepfert B., Hintermann B. 2007. Gait analysis in
 389 Ankle Osteoarthritis and Total Ankle Replacement. *Clin Biomech (Bristol, Avon).* 22,894-904.
 390 Valderrabano V., von Tscharnner V., Nigg B., Hintermann B., Goepfert B., Fung T., Frank C.B., Herzog W.
 391 2006. Lower leg muscle atrophy in ankle osteoarthritis. *J Orthop Res.* 24(12),2159-2169.
 392 von Tscharnner V., Valderrabano V. 2010. Classification of multi muscle activation patterns of osteoarthritis
 393 patients during level walking. *J Electromyogr Kinesiol.* 20(4),676-683.
 394 Zaidi R., Cro S., Gurusamy K., Siva N., Macgregor A., Henricson A., Goldber A. 2013. The outcome of total
 395 ankle replacement: a systematic review and meta-analysis. *Bone Joint J.* 95-B(11):1500-1507.

396

397 **Figure caption**

398 **Figure 1. Temporal co-activation between medial gastrocnemius and tibialis anterior during gait in the**
399 **non-operated (top) and operated (bottom) legs from patient 1.** The EMG signals of the medial
400 gastrocnemius are shown in light gray and positive values. The tibialis anterior signals are shown in dark gray
401 and negative values. The line shows the identified time of muscle activity as proposed by Merlo et al (2003).
402 The upper box shows the color intensity scale of the temporal overlapping relative to the medial gastrocnemius.
403 The bottom boxes show the intensity of each overlap found. The signals were normalized in function of z-score.