Patient-reported outcome monitoring in a routine paediatric oncology setting

Kaur, Geetinder; Kyte, Derek; Reeve, Bryce B; Basch, Ethan; Calvert, Melanie

DOI:

[10.1016/S1470-2045\(18\)30785-X](https://doi.org/10.1016/S1470-2045(18)30785-X)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kaur, G, Kyte, D, Reeve, BB, Basch, E & Calvert, M 2019, 'Patient-reported outcome monitoring in a routine paediatric oncology setting: challenges and opportunities', *The Lancet Oncology*, vol. 20, no. 1, pp. 19-20.
[https://doi.org/10.1016/S1470-2045\(18\)30785-X](https://doi.org/10.1016/S1470-2045(18)30785-X)

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility 14/01/2019

[https://doi.org/10.1016/S1470-2045\(18\)30785-X](https://doi.org/10.1016/S1470-2045(18)30785-X)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Patient-reported outcome monitoring in a routine paediatric oncology setting: challenges and opportunities

Cancer affects one in every 500 children younger than 15 years in the UK, and around 1800 new cases are diagnosed in this population every year.¹ Despite advances in treatments and improved overall survival, children with cancer have a high symptom burden, related both to the underlying cancer and associated treatments that can substantially affect their physical, mental, social, and emotional wellbeing, and cognitive function.² Moreover, there are major psychosocial and financial implications for families of children with cancer, and substantial societal health-care costs.

Effective symptom management is the cornerstone of high-quality oncology care. Lack of recognition of the child's symptoms is likely to cause or worsen complications, delay alleviation of their symptoms, limit the ability to safely and effectively deliver treatments, increase hospital and emergency visits, and have a negative effect on clinical outcomes.³

Studies have shown that clinicians' and parents' perceptions of a child's symptoms do not accurately reflect the child's perspective, particularly for subjective attributes such as emotion, pain, cognition, and confidence for the future.⁴ This incongruence is especially important for common, unobservable symptoms such as nausea, which can only be detected through self-reporting. Eliciting the patient's voice through self-reporting of symptoms is crucial to optimising symptom management.

The importance of patient-centred care in improving the quality and effectiveness of health care is well recognised. One way to assess the effect of disease and treatment on symptoms is with the use of patient-reported outcome measures. As stated by the US Food and Drug Administration, patient-reported outcome measures show "the status of a patient's health condition that comes directly from the patient, without interpretation of the patient's response by a clinician or anyone else".⁵ A 2018 systematic review⁶ identified 38 patient-reported

outcome measures used in paediatric oncology research, measuring a total of 81 symptoms. Self-reporting of symptoms is feasible in the paediatric population because most patient-reported outcome instruments are specifically designed for ages ranging from 8 years to 18 years, with varying recall periods ranging from immediate recall, to up to 1 year. The feasibility of collecting symptom data at multiple timepoints in longitudinal studies supports the potential use of these instruments as clinical tools to monitor symptoms. Electronic reporting platforms have shown promising results for children with cancer, highlighting the need for further development.⁷ Communication of symptom data to clinical teams leads to improved parental understanding of the child's symptoms, better communication with clinicians, and improvement of the children's emotional health-related quality of life.⁸ There is also evidence from an adult oncology setting that symptom reporting with automated clinician alerts results in better health-related quality of life, fewer emergency room visits, fewer hospital admissions, longer duration of palliative chemotherapy, and improved survival compared with usual care.³

However, despite the availability of specific patient-reported outcome symptom tools in paediatric oncology and evidence of clinical effectiveness when used in adult patient populations, their use in routine clinical paediatric oncology practice is minimal. An international survey⁹ revealed that 94% of health-care professionals working in paediatric oncology practices across 52 countries valued routine collection and use of patient-reported outcomes. However, several logistical issues, such as time constraints, insufficient staff, and financial limitations, were identified as potential barriers. Other key barriers were technological, which included a lack of standardised patient-reported outcome platforms that can interface with electronic health records, an absence of financial incentives to develop such platforms, as well as a shortage of standardised processes for integrating patient-reported outcomes in routine clinical workflow.¹⁰

There is a need to develop the infrastructure for systematic collection of symptom data that are communicated, in real time, to clinical teams to enable better identification and timely, efficient clinical management. Patient-reported outcome systems that assess quality of life, adverse effects, resource needs, and practical issues associated with treatment (such as compliance to treatment) should be based on the needs of children, young people, and their families, be age appropriate, and easy for patients to use. They should also be clinically meaningful, actionable, and logistically practical from the clinical team's perspective.

The International Society for Quality of Life Research (ISOQoL) developed a user's guide for implementing patient-reported outcome assessment in routine clinical practice, describing methodological and practical considerations. Successful implementation of such systems in routine clinical practice would require the engagement of everyone involved, including the patients and their families, treating clinical teams, hospital management, and funders, and would also require formal evaluation to assess the effectiveness of the system. Designing and co-producing standardised systems for symptom self-reporting in partnership with key stakeholders could improve the quality and safety of care, outcomes, and quality of life of children with cancer.

**Geetinder Kaur, Derek Kyte, Bryce B Reeve, Ethan Basch, Melanie Calvert*

Centre for Patient Reported Outcome Research, Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK (GK, DK, MC); National Institute for Health Research Birmingham Biomedical Research Centre and National Institute for Health Research Surgical Reconstruction and Microbiology Research Centre, University Hospitals Birmingham NHS Foundation Trust and University of Birmingham, Birmingham, UK (DK, MC); Centre for Health Measurement, Department of Population Health Sciences and Department of Pediatrics, Duke University School of Medicine, Durham, NC, USA

(BBR); and UNC Lineberger Comprehensive Cancer Centre, School of Medicine, University of North Carolina, Chapel Hill, NC, USA (EB)

g.kaur.2@bham.ac.uk

DK reports grants from Innovate UK, National Institute for Health Research (NIHR) Birmingham Biomedical Research Centre, NIHR Surgical Reconstruction and Microbiology Research Centre (SRMRC) at the University Hospitals Birmingham NHS Foundation Trust, the University of Birmingham, Macmillan Cancer Support, and personal fees from Merck outside the submitted work. BBR is supported by a grant from the US National Cancer Institute to develop a system for children to self-report symptoms they experience while receiving cancer treatment. EB receives research funding from the US National Cancer Institute and Patient-Centered Outcomes Research Institute, is an Associate Editor for *JAMA*, and is a scientific advisor for Noona, Sivan, and the Centers for Medicare and Medicaid Services. MC reports personal fees from Astellas and Takeda, and grants from Innovate UK, NIHR, NIHR SRMRC at the University Hospitals Birmingham NHS Foundation Trust, the University of Birmingham, and Macmillan Cancer Support outside the submitted work. GK declares no competing interests. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health.

1 Cancer Research UK. Children's cancer incidence statistics

<http://www.cancerresearchuk.org/health-professional/cancer-statistics/childrenscancers/incidence#heading-Two> (accessed Oct 10, 2018).

2 Ullrich CK, Billett AL, Berde CB. Symptom management in children with cancer. In: Orkin SH, Nathan DG, Ginsburg D, et al, eds. *Nathan and Oski's hematology of infancy and childhood*. Philadelphia: Saunders, 2009: 2349–96.

3 Basch E, Deal AM, Kris MG, et al. Symptom monitoring with patient-reported outcomes during routine cancer treatment: a randomized controlled trial. *J Clin Oncol* 2016; **34**: 557–65.

4 Glaser AW, Davies K, Walker D, Brazier D. Influence of proxy respondents and mode of administration on health status assessment following central nervous system tumours in childhood. *Qual Life Res* 1997; **6**: 43–53.

5 US Food and Drug Administration. Guidance for industry: patient-reported outcome measures: use in medical product development to support labeling claims. 2009. <https://www.fda.gov/downloads/drugs/guidances/ucm193282.pdf> (accessed Oct 10, 2018).

6 Pinheiro LC, McFatrigh M, Lucas N, et al. Child and adolescent self-report symptom measurement in pediatric oncology research: a systematic literature review. *Qual Life Res* 2018; **27**: 291–319.

7 Leahy AB, Feudtner C, Basch E. Symptom monitoring in pediatric oncology using patient reported outcomes: why, how and where next. *Patient* 2018; **1**: 147–53.

8 Engelen V, Detmar S, Koopman H, et al. Reporting health-related quality of life scores to physicians during routine follow-up visits of pediatric oncology patients: is it effective? *Pediatr Blood Cancer* 2011; **58**: 766–74.

9 Schepers SA, Haverman L, Zadeh S, Grootenhuis MA, Wiener L. Healthcare professionals' preferences and perceived barriers for routine assessment of patient-reported outcomes in pediatric oncology practice: moving toward international processes of change. *Pediatr Blood Cancer* 2016; **63**: 2181–88.

10 Basch E. Patient reported outcomes-harnessing patients' voices to improve clinical care. *N Engl J Med* 2017; **376**: 105–08.