Improving outcomes in high-risk Myelodysplasia

Craddock, Charles

DOI:

[10.1200/JCO.2017.73.0812](https://doi.org/10.1200/JCO.2017.73.0812)

License:

Other (please specify with Rights Statement)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Craddock, C 2017, 'Improving outcomes in high-risk Myelodysplasia: Festina Lente', *Journal of Clinical Oncology*, pp. JCO2017730812. <https://doi.org/10.1200/JCO.2017.73.0812>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

© 2019 American Society of Clinical Oncology. All rights reserved.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Improving Outcomes in High-Risk Myelodysplasia: Festina Lente

Charles Craddock, *Queen Elizabeth Hospital, Birmingham, and University of Birmingham, Edgbaston, Birmingham, United Kingdom*

See accompanying article on page 2745

The myelodysplastic syndromes (MDS) represent a compendious group of clonal hematopoietic disorders characterized by ineffective hematopoiesis resulting in peripheral blood cytopenias and a significant risk of transformation to acute myeloid leukemia (AML). The degree of cytopenia, blast percentage, and presentation karyotype permit identification of patients with high-risk MDS whose outcome is particularly poor. Targeted mutational analysis has transformed our understanding of the pathogenesis of MDS and, at the same time, provided the tools for preliminary analyses of clonal structure. Acquired abnormalities in classes of genes determining chromatin structure, including mutations in *DNMT3A*, *TET2*, *IDH1*, and *IDH2* (which influence CpG island methylation status) and *EZH2* and *ASXL1* (which contribute to the control of histone acetylation) are commonly observed in MDS and likely contribute to disease pathogenesis.¹

Currently, the only curative treatment in MDS is allogeneic stem-cell transplantation. Over the last decade, pharmacologic reversal of acquired abnormalities in chromatin structure has emerged as a plausible therapeutic strategy in a range of hematologic malignancies, notably AML and MDS, where inhibitors of DNA methyltransferase (DNMT) and histone deacetylase enzymes have demonstrated activity in preclinical models.² Indeed, the most important therapeutic advance in the treatment of adult higher-risk MDS, particularly in older patients ineligible for intensive chemotherapy or allogeneic stem-cell transplantation, has been the demonstration that the DNMT inhibitors azacitidine (AZA) and decitabine can produce either complete remission (CR) or a hematologic improvement in a significant number of patients.^{3,4} Furthermore, in a prospective randomized trial, AZA was shown to improve overall survival (OS), thereby representing the first pharmacologic agent to alter the natural history of high-risk MDS.⁵ Importantly, recent prospective trials have also demonstrated improved OS after AZA therapy in older adults with AML who are not eligible for intensive chemotherapy.⁶ The current subcutaneous mode of administration of AZA is inconvenient and sometimes painful, but the recent development of a relatively well-tolerated formulation, CC486, with augmented pharmacokinetics promises to transform the ease of administration of an agent that has now become the standard of care in older patients with high-risk MDS.⁷ Of interest, CC486 also demonstrates clinical activity in patients who have experienced treatment failure with subcutaneous AZA, suggesting that the more prolonged exposure to AZA delivered by CC486 can result in clinical responses in patients resistant to AZA.

It remains the case, however, that only a minority of patients treated with AZA achieve a bone marrow CR or CRi (characterized by a reduction in bone marrow blast percentage to < 5% without complete peripheral count recovery), and disease progression remains inevitable. Consequently, the last decade has witnessed an active search for novel agents with the capacity to both improve response rates and prolong survival in AZA-treated patients. In preclinical models, the combination of AZA with histone deacetylase inhibitors, such as sodium valproate, etinostat, and vorinostat, markedly increases antitumor activity.² Indeed, two early-phase studies of combined AZA and vorinostat administration seemed encouraging, with CR rates reported between 50% and 70%.^{8,9} An alternative strategy that has shown promising results in early-phase trials is combination with the immunomodulatory drug lenalidomide.¹⁰ Consequently, the prospective randomized trial by Sekeres et al¹¹ reported in the article that accompanies this editorial, which demonstrated no increase in either overall response rate or survival in patients with high-risk MDS treated with either combination compared with AZA monotherapy, is important and consistent with recent studies showing no benefit of combining AZA with etinostat or pracinostat.^{12,13} Why might this meticulously conducted trial have failed to replicate the encouraging results of previous early-phase trials using both drug combinations? Such disparities are most commonly attributed to potential selection bias in early-phase trials, and it is therefore important to note that in this study, the demographic and molecular characteristics of the trial population appear representative of a typical population of patients with high-risk MDS. Alternatively, it is possible that the additional toxicity observed with the experimental combination arms resulted in either underdosing of AZA or premature cessation of combination therapy, and this possibility is supported by the trend toward improved survival in the vorinostat arm despite increased toxicity.

On the assumption that therapeutic decision making should be informed by Einstein's apocryphal observation that the definition of insanity is doing the same thing over and over again and expecting different results, it is perhaps important now to consider what alternative strategies might improve the clinical activity of AZA with regard to both increasing CR rate and prolonging OS. Rational construction of novel AZA-based combinations is hampered by our failure to fully understand both how AZA exerts its antitumor effect and the cellular basis of disease relapse in this clinical setting. Studies correlating upregulation of key cell cycle

regulators, such as *CDKN1A*, with AZA's activity in both in vitro and animal models of AML implicate induction of G1/S cell cycle arrest as one of the mechanisms of AZA-induced antitumor activity. These observations provide a rationale for combining AZA with inhibitors that target G2 checkpoint kinases, such as Wee1, Chk1, or ATR. Combination with other agents that exert a p53-independent mechanism of killing is also of interest, and it is noteworthy that in preliminary studies, coadministration of AZA with the *Bcl-2* inhibitor venetoclax has been reported to produce CR rates of up to 70% in newly diagnosed patients with high-risk MDS and AML.¹⁴

The major cause of treatment failure in AZA-treated patients is the almost-universal occurrence of disease resistance; consequently, strategies with the capacity to maintain bone marrow responses are required. The occurrence at relapse of subclones present at only a low frequency at diagnosis in AZA-treated patients supports the implementation of a sequential therapeutic strategy using targeted agents such as *FLT3ITD*, *IDH1*, or *IDH2* inhibitors in combination with AZA,¹⁵ potentially on the basis of sequential analysis of clonal structure. At the same time, immunophenotypic characterization of the stem/progenitor cell compartments containing leukemic stem cells permits serial quantitation of leukemic stem/progenitors during therapy.¹⁶ A current but largely untested hypothesis is that this population serves as a reservoir of disease resistance, which, if confirmed, raises the possibility of both using serial quantitation of this population as a biomarker of response to novel AZA-based combinations and highlights the importance of developing novel drug or antibody treatment strategies with the potential to target residual leukemic stem/progenitor cells.

A major factor limiting the effective use of AZA in clinical practice is the challenge of identifying the patients likely to obtain a major clinical benefit at the start of treatment, either in terms of CR acquisition or improved survival. A notable feature of the article by Sekeres et al¹¹ is the prospective correlation of clinical response with the results of next-generation sequencing of 40 candidate myeloid genes. In previous retrospective analyses, mutations in *DNMT3A* have been shown to correlate with an increased likelihood of achieving a clinical response to AZA, although this has not been consistently observed.¹⁷ It is therefore of considerable interest that this observation was replicated in this prospective study because just as retrospective phase I/II trials can yield unreproducible clinical observations, the same may also apply to the previous, mostly retrospective, analyses of the molecular predictors of response to AZA-based therapy. Just as importantly, this prospective evaluation provides a template for integrating mutational profiling into future trials of AZA-based combination therapy.

More generally, the increasing molecular stratification of diseases previously considered homogeneous clinical entities, such as MDS and AML, coupled with the tsunami of emergent novel therapies, poses profound questions about the scale and capacity of the translational infrastructure that will be required if we are to rapidly assess an exponentially increasing number of promising novel drug, antibody, and cellular therapies in ever-decreasing genetically defined disease populations. What is increasingly clear is that our current reliance on both trial designs and translational infrastructures that may have been fit for purpose in 2007 when the therapeutic cupboard was bare threatens to disadvantage

patients, academia, and the pharmaceutical sector in the next decade. Although the precise models that will be required to realize the therapeutic promise of the next decade will be nuanced according to factors such as disease incidence, molecular stratification, and therapeutic opportunities, certain common principles can be predicted to apply. At the heart of accelerated trial delivery is the establishment of resourced trial networks with a sufficient density of patients to allow the rapid assessment of a range of novel therapies as developed so effectively in stem-cell transplantation and planned for antibiotic trials.^{18,19} Such networks should be appropriately resourced and incentivised to reward rapid trial development with enhanced pharmacovigilance allowing trial delivery to regulatory standard, removing the current inefficient split between commercial and academic studies. Finally, integration of next-generation sequencing, studies of measurable residual disease and candidate biomarkers, where possible in real time, must be ensured so that both discovery science and trial delivery have the opportunity to impact clinical care as swiftly as possible.

AUTHOR'S DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at jco.org.

REFERENCES

- Kennedy JA, Ebert BL: Clinical implications of genetic mutations in myelodysplastic syndrome. *J Clin Oncol* 35:968-974, 2017
- Min C, Moore N, Shearstone JR, et al: Selective inhibitors of histone deacetylases 1 and 2 synergize with azacitidine in acute myeloid leukemia. *PLoS One* 12:e0169128, 2017
- Wijermans P, Lübbert M, Verhoef G, et al: Low-dose 5-aza-2'-deoxycytidine, a DNA hypomethylating agent, for the treatment of high-risk myelodysplastic syndrome: A multicenter phase II study in elderly patients. *J Clin Oncol* 18:956-962, 2000
- Silverman LR, Demakos EP, Peterson BL, et al: Randomized controlled trial of azacitidine in patients with the myelodysplastic syndrome: A study of the cancer and leukemia group B. *J Clin Oncol* 20:2429-2440, 2002
- Fenaux P, Mufti GJ, Hellstrom-Lindberg E, et al: Efficacy of azacitidine compared with that of conventional care regimens in the treatment of higher-risk myelodysplastic syndromes: A randomised, open-label, phase III study. *Lancet Oncol* 10:223-232, 2009
- Dombret H, Seymour JF, Butrym A, et al: International phase 3 study of azacitidine vs conventional care regimens in older patients with newly diagnosed AML with >30% blasts. *Blood* 126:291-299, 2015
- Garcia-Manero G, Gore SD, Kambhampati S, et al: Efficacy and safety of extended dosing schedules of CC-486 (oral azacitidine) in patients with lower-risk myelodysplastic syndromes. *Leukemia* 30:889-896, 2016
- Silverman LVA, Odchimar-Reissig R, LeBlanc A, et al: A phase I trial of the epigenetic modulators vorinostat, in combination with azacitidine (azaC) in patients with the myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML): A study of the New York Cancer Consortium. *Blood* 112:3656, 2008
- Garcia-Manero G, Yang H, Bueso-Ramos C, et al: Phase 1 study of the histone deacetylase inhibitor vorinostat (suberoylanilide hydroxamic acid [SAHA]) in patients with advanced leukemias and myelodysplastic syndromes. *Blood* 111:1060-1066, 2008
- Pollyea DA, Zehnder J, Coutre S, et al: Sequential azacitidine plus lenalidomide combination for elderly patients with untreated acute myeloid leukemia. *Haematologica* 98:591-596, 2013
- Sekeres MA, Othus M, List AF, et al: Randomized phase II study of azacitidine alone or in combination with lenalidomide or with vorinostat in higher-risk myelodysplastic syndromes and chronic myelomonocytic leukemia: North American Intergroup Study SWOG S1117. *J Clin Oncol* 35:2745-2753, 2017
- Prebet T, Sun Z, Figueroa ME, et al: Prolonged administration of azacitidine with or without entinostat for myelodysplastic syndrome and acute myeloid leukemia with myelodysplasia-related changes: Results of the US Leukemia Intergroup trial E1905. *J Clin Oncol* 32:1242-1248, 2014

13. Garcia-Manero G, Montalban-Bravo G, Berdeja JG, et al: Phase 2, randomized, double-blind study of pracinostat in combination with azacitidine in patients with untreated, higher-risk myelodysplastic syndromes. *Cancer* 123: 994-1002, 2017
14. Bogenberger JM, Kornblau SM, Pierceall WE, et al: BCL-2 family proteins as 5-azacytidine-sensitizing targets and determinants of response in myeloid malignancies. *Leukemia* 28:1657-1665, 2014
15. Mossner M, Jann JC, Wittig J, et al: Mutational hierarchies in myelodysplastic syndromes dynamically adapt and evolve upon therapy response and failure. *Blood* 128:1246-1259, 2016
16. Craddock C, Quek L, Goardon N, et al: Azacitidine fails to eradicate leukemic stem/progenitor cell populations in patients with acute myeloid leukemia and myelodysplasia. *Leukemia* 27:1028-1036, 2013

17. Traina F, Visconte V, Elson P, et al: Impact of molecular mutations on treatment response to DNMT inhibitors in myelodysplasia and related neoplasms. *Leukemia* 28:78-87, 2014
18. Steering Committee of the Blood and Marrow Transplant Clinical Trials Network: The Blood and Marrow Transplant Clinical Trials Network: An effective infrastructure for addressing important issues in hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 22:1747-57, 2016
19. McDonnell A, Rex JH, Goossens H, et al: Efficient delivery of investigational antibacterial agents via sustainable clinical trial networks. *Clin Infect Dis* 63: S57-S59, 2016 (suppl 2)

DOI: <https://doi.org/10.1200/JCO.2017.73.0812>; published at jco.org on May 26, 2017.

ASCO's *Journal of Global Oncology*

This online-only, open access journal fills a growing need for high-quality literature on the array of challenges that health care professionals in resource-limited settings face in caring for patients with cancer and in conducting research. Topics covered include:

- Cancer treatment and diagnosis
- Palliative and supportive care
- Prevention
- Barriers to care
- Epidemiology
- Health policy

Article types that will be considered include original reports, reviews, commentaries, correspondence, special articles, case reports, and editorials.

Submit your article today at submitjgo.ascopubs.org

American Society of Clinical Oncology

AUTHOR'S DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Improving Outcomes in High-Risk Myelodysplasia: Festina Lente

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/site/ifc.

Charles F Craddock

Honoraria: Celgene, Janssen Oncology, Novartis, Merck Sharp & Dohme

Consulting or Advisory Role: Celgene, Janssen Oncology, Merck Sharp & Dohme

Speakers' Bureau: Celgene, Janssen Oncology

Research Funding: Celgene

Acknowledgment

The author acknowledges valuable discussions with Tanja Stankovic, MD, PhD, and Paresh Vyas, MD, PhD, and programmatic support from the Bloodwise Trials Acceleration Programme, as well as research support from Celgene.