Biological nitrogen fixation in peatlands

Saiz, Ernesto; Sgouridis, Fotis; Drifjhout, Falko; Ullah, Sami

DOI:

[10.1016/j.soilbio.2019.01.011](https://doi.org/10.1016/j.soilbio.2019.01.011)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Saiz, E, Sgouridis, F, Drifjhout, F & Ullah, S 2019, 'Biological nitrogen fixation in peatlands: comparison between acetylene reduction assay and $^{15}\text{N}_2$ assimilation methods', *Soil Biology and Biochemistry*, vol. 131, pp. 157-165. <https://doi.org/10.1016/j.soilbio.2019.01.011>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Published in *Soil Biology and Biochemistry* on 16/01/2019

<https://doi.org/10.1016/j.soilbio.2019.01.011>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Note: This paper is accepted and published in Soil Biology and Biochemistry Journal and should be cited as:

■ Saiz, Ernesto, F Sgouridis, F Drijfhout, and S Ullah. 2019. Biological nitrogen fixation in peatlands: Comparison between acetylene reduction assay and $^{15}\text{N}_2$ assimilation methods. Soil Biology and Biochemistry Vol 131: 157-165
<https://doi.org/10.1016/j.soilbio.2019.01.011>

BIOLOGICAL NITROGEN FIXATION IN PEATLANDS: COMPARISON BETWEEN ACETYLENE REDUCTION ASSAY AND $^{15}\text{N}_2$ ASSIMILATION METHODS

Ernesto Saiz^a, Fotis Sgouridis^b, Falko P. Drijfhout^c, and Sami Ullah^{d*}

^a School of Geography, Geology, and the Environment, Keele University, Staffordshire, ST5 5BG, UK

^b School of Geographical Sciences, University of Bristol, BS8 1SS, UK

^c Chemical Ecology Group, School of Physical and Chemical Sciences, Keele University, Staffordshire, ST5 5BG, UK

^d School of Geography, Earth, and Environmental Sciences, and Birmingham Institute of Forest Research, University of Birmingham, B15 2TT, UK

*Corresponding author: Sami Ullah (s.ullah@bham.ac.uk).

21 Biological Nitrogen Fixation (BNF) is an essential microbial process supplying available
22 nitrogen (N) to *Sphagnum* mosses in ombrotrophic peatlands. Acetylene Reduction Assay
23 (ARA) and the $^{15}\text{N}_2$ assimilation are the main methods used for the measurement of BNF.
24 ARA is used as a proxy where the moles of ethylene (C_2H_4) produced from acetylene (C_2H_2)
25 during incubation of mosses and peat are used to estimate the moles of N being fixed using a
26 conversion factor (CF), thus relating the moles of C_2H_4 produced to the moles of N fixed. A
27 theoretical CF of 3:1 originally developed for agricultural soils using pure nitrogenase
28 enzymes is in use; in some cases a site specific CF is determined through parallel incubation
29 of mosses and peat with ARA and $^{15}\text{N}_2$ assimilation methods to enable the application of
30 ARA for subsequent BNF measurement at high resolution and low cost. However, in recent
31 literature, the reported site and/or species specific CF for peatlands varies by an order of
32 magnitude, thus raising the question if measured CFs are robust and consistent enough for the
33 estimation of BNF in peatlands. Thus, we measured BNF using the ARA and the direct $^{15}\text{N}_2$
34 assimilation methods in three different peatlands across the UK during the growing season
35 over two years. The incubations were carried out in parallel on the dominant *Sphagnum* spp.
36 (*S. cuspidatum*, *S. fallax*, *S. capillifolium*, and *S. papillosum*) and top bulk peat (0-15 cm).
37 Additional incubations were performed using the direct $^{15}\text{N}_2$ assimilation method with and
38 without C_2H_2 addition to evaluate if C_2H_2 was suppressing N assimilation through BNF all
39 together in peatlands. According to the results, the CF varied from 0.001 to 5.363, with a
40 median CF of 0.028 for both mosses and peat, which is far lower than the theoretical 3:1 CF.
41 The CF was also highly variable with differences up to 3 orders of magnitude across the
42 different *Sphagnum* species. The measured CF between years for the same species and across
43 the three peatland sites varied significantly suggesting an inconsistent performance of ARA
44 against the ^{15}N assimilation method. The generally low but highly varied CFs measured under

45 this study shows that C_2H_2 differentially interferes with the activity of diazotrophic microbes,
46 which results in an inconsistent CF at species, and site scales, and over time. In conclusion,
47 ARA is not suitable as a proxy method for estimating and/or modelling BNF in peatlands

48 Key words: Acetylene Reduction Assay, ARA, $^{15}N_2$ assimilation method, biological nitrogen
49 fixation, *Sphagnum*, diazotrophs, nitrogenase enzymes.

50

1. Introduction

Biological nitrogen fixation (BNF) is an essential microbial process for the provision of available nitrogen (N) to plants in nutrient-poor ombrotrophic peatlands that otherwise rely on atmospheric deposition of reactive N (Berg et al., 2013; Bragina et al., 2013). Ombrotrophic peatlands are usually dominated by *Sphagnum* spp. (mosses), which are adapted to acidic and nutrient-poor conditions (van den Elzen, et al., 2018). Moss-associated cyanobacteria and free-living diazotrophic bacteria fix atmospheric N₂ into the bioavailable NH₄⁺ form, thus enabling the plants to meet their N demands for capturing atmospheric carbon (Postgate, 1982). Nitrogenase enzyme in the diazotrophs is responsible for reducing N₂ into ammonium as below (Smith and Gallon, 1993):

There are two main methods for measuring BNF: direct ¹⁵N₂ assimilation (¹⁵N₂ method) and acetylene reduction assay (ARA; Sprent, 1979; Bellenger et al., 2014). The ¹⁵N₂ method allows for the direct quantification of BNF rates; however, the high cost of isotopic tracing is prohibitive for a widespread repeated use. It involves the incubation of samples with ¹⁵N₂ gas, for direct measurement of ¹⁵N incorporation into biomass by N fixers, followed by the determination of ¹⁵N signature in the incubated samples through mass spectrometry (Zehr and Montoya, 2007). Although it is a robust method, some practical problems have been reported when using this method. For example incubation chamber leakage (Chalk et al., 2017), oxygen depletion during long term incubation (over weeks) (Myrold et al., 1999), incomplete and slow equilibration between the ¹⁵N₂ gas and the water sample (Großkopf et al., 2012), and ¹⁵N₂ gas contamination (Dabundo et al., 2014) can result in under or overestimation of BNF.

The ARA is the most common method for measuring BNF in different ecosystem types. It is based on the nitrogenase enzyme preferential reduction of acetylene (C₂H₂) to ethylene (C₂H₄) that involves just two electron transfer, instead of reducing N₂ which involves eight, when C₂H₂ is present at relatively high concentrations (10% v/v; Koch and Evans, 1966; Schöllhorn and Burris, 1967; Hardy et al., 1968), according to the following equation (Bergersen, 1970; Staal et al., 2001):

Despite its simplicity ARA is an indirect method and a conversion factor (CF) is needed to estimate BNF rate equivalents based on the number of moles of C₂H₄ produced. The theoretical CF obtained from the formulas (1) and (2) relating the number of reducing equivalents is 4:1 (moles of C₂H₄ produced per mole of nitrogen fixed) (Zehr and Montoya, 2007). Empirical measurements in *in vitro* experiments of nitrogen fixing bacteria (*Azotobacter* and *Clostridium*) as well as *in situ* have found that the ratio of C₂H₄ produced to N fixed was between 3 and 4.5 (Hardy et al., 1968). However, some authors consider that a couple of electrons and protons are used to release one molecule of H₂ in equation 1, and therefore, the theoretical CF should be 3:1, which is what has been traditionally used in the soil literature (Postgate, 1982). Many authors have reported deviations from the theoretical CF when measuring BNF in peatlands (Chapman and Hemond, 1982; Schwintzer, 1983), in the laboratory with forest soil cores (Nohrstedt, 1983), or in different nitrogen-fixing systems (Bergersen, 1970). As a result, these authors strongly recommended site specific calibration of the ARA method using the ¹⁵N₂ method (Bergersen, 1970; Roskoski, 1981; Nohrstedt, 1983) for subsequent application of ARA at large scale over time.

Although there have been several studies relating ARA and the ¹⁵N₂ method in legumes and laboratory cultures (Bergersen, 1970), in forests (Roskoski, 1981; Nohrstedt, 1983), and in

arctic habitats (Liengen, 1999), the effect of C_2H_2 on diazotrophic microbial activity in peatlands and hence BNF has been overlooked. It is known that C_2H_2 interferes with different microbial processes typical of peatlands such as nitrification, blocking it; denitrification, inhibiting the respiratory reduction of N_2O to N_2 (Ryden, 1982); and methanotrophy, inhibiting the oxidation of methane (Kip et al., 2010). These metabolic processes provide energy to diazotrophs (Raghoebarsing et al., 2005) and substrate (for example, coupled and/or direct respiratory N_2O reduction and N fixation; Desloover et al., 2014; Farias et al., 2013), thus their inhibition or suppression in the presence of C_2H_2 might affect C_2H_4 production and hence estimation of BNF rates given that the presence of these microbes in the incubated media (mosses and peat) may vary over time (Raghoebarsing et al., 2005). In recent literature, it has been shown that methanotrophs play an important role in BNF (Larmola et al., 2014; Vile et al., 2014), and that they are present in association of *Sphagnum*-mosses other than cyanobacteria as a key N fixing microbe (Larmola et al., 2010) in peatlands all over the world (Kip et al., 2010).

Many studies have applied site specific CFs in peatlands where *Sphagnum* spp. were present such as 3.11 (Kox et al., 2016), 0.85 (Stewart et al., 2011) or 0.32 (Vile et al., 2014) for mosses, and 1.1 (Knorr et al., 2015) for peat, albeit all of them reported high variability, which raises the question if the site-specific CF is reliable and reproducible over time. In other cases, ARA is applied for *Sphagnum* mosses using the theoretical 3:1 CF (Rousk et al., 2018) or using a CF previously reported for the site (Rousk et al., 2015), whilst indicating that BNF rates for the sites would be underestimated because of the inhibitory effects of C_2H_2 on methanotrophs. The only study that focused on the effects of ARA on BNF rates estimation (Warren et al., 2017) was on peat soil, and therefore, no information exists on the effect of C_2H_2 on diazotrophic microbes associated with *Sphagnum* mosses including cyanobacteria and hence the usefulness of ARA as a proxy of BNF in peatlands.

Our main aim was to evaluate the usefulness of BNF method in peatlands. In this study, ARA was calibrated against the $^{15}\text{N}_2$ assimilation method with the objectives to assess: (1) if the conversion factor is consistent across *Sphagnum* species and peat at each site, across sites, and across wider geographic temperate peatland regions, (2) if the CF is consistent over time, and (3) if $^{15}\text{N}_2$ assimilation through BNF is completely suppressed in the presence of C_2H_2 . Knowing the reported interference of C_2H_2 with microbial activities that have a direct and/or indirect bearing on BNF activity such as methanotrophy, respiratory reduction of N_2O and nitrification (Larmola et al., 2014; Desloover et al., 2014; Farias et al., 2013; Ho and Bodelier, 2015; Sgouridis et al., 2016), we hypothesized that ARA as a proxy method will underestimate BNF rates in peatlands.

2. Material and methods

2.1. Study sites and sampling

Sphagnum mosses and peat samples were collected from three ombrotrophic peatlands in the UK: Migneint (52° 59' 20.8" N – 3° 48' 09.8" W) in Wales, Fenn's and Whixall (52° 55' 20.8" N – 2° 45' 58.6" W) in England, and Forsinard (58° 23' 42.2" N – 3° 56' 47.0" W) in Scotland (Fig. 1). These sites had different characteristics regarding mean annual temperature, rainfall, and atmospheric reactive nitrogen (Nr) deposition rates (Table 1) so that the comparative performance of the two methods could then be representative at large geographic scale.

Figure 1. Location of the study sites in the United Kingdom.

Moreover, the selected sites were exposed to variable atmospheric Nr deposition and the rationale for the selection was also to evaluate the comparative performance of the two techniques and to know if chronic Nr deposition might be affecting the CF given that a recent paper reported suppression of BNF in feather mosses exposed to high Nr deposition (Ackermann et al., 2012, Rousk and Michelsen, 2016). Two sampling campaigns were undertaken at Migneint and Fenn's and Whixall sites during the growing season, in 2016 and 2017, respectively; and one campaign at Forsinard in 2017 for *in situ* method performance incubations. Additionally, at Migneint site, samples were collected in spring 2016 for laboratory-based CF determination.

The vegetation in these sites consisted of mosses and ericoid shrubs. For moss associated BNF quantification, samples of four dominant *Sphagnum* species were collected at each site: *Sphagnum cuspidatum* and *Sphagnum fallax* (most common in hollows); and *Sphagnum papillosum* and *Sphagnum capillifolium* (in hummocks). During the 2016 campaigns, bulk

peat (0-15 cm) was also collected from hollows and hummocks, while in 2017 peat was collect only from hollows. *Sphagnum* and peat samples were collected from five random locations within each site to capture the wider inherent spatial variability of each site.

Table 1. Mean annual temperature, precipitation and reactive nitrogen (Nr) deposition in the study sites.

Site	Mean annual temperature (°C)	Mean annual precipitation (mm)	Atmospheric Nr deposition (kg N ha ⁻¹ yr ⁻¹)
Forsinard (Scotland)	6.9	1104	6
Migneint (Wales)	7.3	2236	17
Fenn's and Whixall (England)	9.5	747	26
Source: Met Office, Air Pollution Information System (APIS).			

2.2. ¹⁵N₂ assimilation method

For each species and peat four out of five replicates were incubated with ¹⁵N₂ (98 atom% Cambridge Isotope Laboratories Inc., USA), with the fifth being the control (incubated using ambient air). Each replicate consisted of 20 live moss shoots (~ upper 5 cm) of the selected moss species or 10 g of peat after passing it through a 2 mm sieve. Shoots and peat were placed in 50 ml serum vials which were capped with air tight rubber septa. Following the closure, 5 ml headspace air was drawn using gas tight syringe and replaced with 5 ml of the ¹⁵N₂ (98 atom%) gas (10% headspace concentration) and the bottles were then placed 'upside

down' (avoiding cap shade) in the same area from where the samples were collected and incubated for 24 hours to avoid issues of oxygen depletion during long-term incubation (Myrold et al., 1999). In case of peat, samples were placed under the moss carpet (dark conditions). Parallel incubations were run for ARA (details below). Additionally, in order to control some of the main factors affecting BNF (temperature, light), one set of moss and peat samples from Migneint were incubated under laboratory conditions with the temperature set at 20 °C (± 2) and the light/dark cycle of 12 hour, while maintaining light intensity through artificial light (photosynthetically active radiation of $\sim 2000 \mu\text{mol m}^{-2} \text{s}^{-1}$) to determine CF under optimal laboratory conditions for comparison with field-based incubations.

After the 24 hour incubation, the vials were opened and aerated to flush out any remaining $^{15}\text{N}_2$ gas and bring it to ambient conditions. Immediately after aeration, the samples were transported in a cool box to the laboratory and then weighed and dried at 70 °C for 72 hours. Dried samples were manually pulverised ($< 2 \text{ mm}$) and subsamples ($\sim 1 \text{ g}$) were sent to the Life Sciences Mass Spectrometry Facility at the Centre for Ecology and Hydrology, Lancaster for ^{15}N content analysis by Isotope Ratio Mass Spectrometry (IRMS) using a Carlo Erba NA1500 (Italy) elemental analyser coupled to a Dennis Leigh Technologies (UK) isotope ratio mass spectrometer. In-house working standards of natural abundance (flour and soil) were analysed every twelfth sample giving an analytical precision of 0.36 ‰. They were calibrated against the certified reference material Internal Atomic Energy Agency standard AEA-N1 (nitrogen isotope in ammonium sulphate). All the control and enriched moss and peat samples were analysed in duplicate (Jardine and Cunjak, 2005) and their variability was within the limits of the analytical precision of the IRMS. We also analysed duplicates of three non-enriched tissues samples on the IRMS and the resulting reproducibility of the analysis including a cross-laboratory check (details in section 2.3 below) was within the analytical precision of the IRMS. Subsequently, we only ran one control and four enriched incubations

during BNF measurements while ensuring that the duplicate runs of each sample on the IRMS was within the analytical precision limits. This check was critical given that the experiment relied on the incubation of one non-enriched sample in the field for calculating BNF rates using the following formula (Liengen, 1999):

$$Y = \left(\frac{\text{atom}\% \text{ } ^{15}\text{N}_{\text{excess}}}{100} \right) \times \left(\frac{\text{total N}_{\text{sample}} \times 10^9}{t \times 28} \right) \times \left(\frac{100}{\% ^{15}\text{N}_{\text{headspace}}} \right)$$

where Y (nmol N g⁻¹ dw h⁻¹) is the amount of N fixed during the experiment, atom% ¹⁵N_{excess} is the difference between atom%¹⁵N_{sample} and atom%¹⁵N_{control}, total N is the total amount of nitrogen in the sample (g N 100 g⁻¹ dw), t is the incubation time, 28 is the molecular weight of N₂ (g mol⁻¹), and %¹⁵N is the percentage of ¹⁵N out of the total amount of N gas in each incubation bottle.

2.3. ¹⁵N₂ Gas quality control

Contamination of commercial ¹⁵N₂ gas with ¹⁵N-labelled nitrate and ammonium can interfere with the detection of BNF (Dabundo et al., 2014). We evaluated the potential contamination of the ¹⁵N₂ gas as well as the possibility of abiotic uptake of ¹⁵N₂ gas by incubating six samples of dried (105 °C) mosses for 24 hours, three with ¹⁵N₂ enriched gas as above and three without. After the incubation, the samples were processed as above and were sent to two different laboratories (CEH Lancaster and Bristol University) for ¹⁵N analysis using IRMS to ensure cross laboratory checks (Bahlmann et al., 2010). The results obtained (average δ ¹⁵N in sample of enriched ones: -0.360; and of non-enriched ones: -0.387) showed a difference of -0.03 δ ¹⁵N between the treatments. Therefore, we used this averaged

difference ($-0.03 \delta^{15}\text{N}$) as a threshold below which any difference between the control and enriched samples incubated for direct BNF measurement was not considered.

2.4. Acetylene reduction assay (ARA)

Following the placement of 20 shoots of mosses or 10 g of field moist peat in serum bottles ($n = 5$) and capping with septa, 10 % of the headspace was replaced with (10 % v/v) pure and fresh C_2H_2 obtained by adding deionised water to calcium carbide. Immediately after doing so, a 3 ml gas sample was obtained (T_0), and replaced by the same gas mixture to maintain atmospheric pressure within the vials. Gas samples were subsequently collected following the same procedure at 6 and 24 hours.

To check for possible contamination of the acetylene gas with ethylene and endogenous production of ethylene by mosses and peat, we carried out quality control incubations. Mosses and peat were incubated with and without the addition of C_2H_2 (each with three replicates), whereas three bottles received C_2H_2 but no sample and three bottles were incubated under ambient air without sample and C_2H_2 . The results showed no endogenous C_2H_4 production or gas contamination, and negligible level of C_2H_4 in air was detected which was later used for background correction while calculating ethylene production rates.

The gas samples were analysed for C_2H_4 concentration using a gas chromatograph (Varian 39000) equipped with a Restek-Alumina BOND/MAPD column (30 mm x 0.53 mm x 10 μm) and a flame ionization detector (FID) using He as a carrier gas. The temperatures of the injector and detector were 200 $^\circ\text{C}$, and for the column was 135 $^\circ\text{C}$. The head pressure was 3.4 psi and the carrier flow 3.2 ml min^{-1} . The injection was manual. C_2H_4 production rates were calculated by linear regression between time intervals T_0 - T_{24} using a standard calibration

curve for each of the daily batch samples. Using standards injection after 10 samples each, the quality of the runs were checked and where needed, corrected for any drift in the signal.

2.5. ARA – $^{15}\text{N}_2$ direct assimilation conversion factor (CF ratio)

The ARA conversion factor was calculated by dividing moles of C_2H_4 produced (ARA method) by the moles of N fixed ($^{15}\text{N}_2$ direct assimilation) for each parallel incubation for different species and peat collected from the Migneint site and incubated under laboratory conditions (Vile et al. 2014) in 2016. Following the CF determination under laboratory conditions, CFs were then estimated for the field-based incubations for the Fenn's and Whixall (2016-17), Migneint (2016-17), and Forsinard (2017) sites. CFs were calculated per site as well as per species and peat type within each site.

2.6. BNF determination with $^{15}\text{N}_2$ assimilation method with and without C_2H_2 addition

In 2017, during the growing season, samples (*Sphagnum* mosses and peat) collected from Fenn's and Whixall, Migneint, and Forsinard sites, were incubated for BNF measurement using the $^{15}\text{N}_2$ assimilation method as described above, where 3 replicates were further amended with C_2H_2 (10 % v/v) and 3 without to evaluate if the presence of C_2H_2 will completely inhibit N_2 reduction to NH_4^+ by diazotrophs given that under high C_2H_2 concentration, diazotrophs have been shown to preferentially reduce C_2H_2 than reducing N_2 (Koch and Evans, 1966; Schöllhorn and Burris, 1967).

2.7. Statistical analysis

Data were tested for normality and homogeneity of variance and since they were found non-normal and non-homogeneous, only non-parametric statistical tests were applied. To test differences in paired samples, we used bootstrapped t-test. Differences in C_2H_4 production (ARA) and BNF ($^{15}\text{N}_2$ method) and the effect of C_2H_2 on BNF rates were tested using the

Wilcoxon ranked sum test. To evaluate the effect of the site and the species we used the Kruskal-Wallis test. All the analysis was performed on the IBM SPSS Statistics program, version 24.

3. Results

3.1. Conversion factors

The CFs obtained showed great variability across species, sites and time ranging from 0.001 to 5.363 moles of C₂H₄ produced per mol of N fixed (Table 2). The mean ($\pm$ standard deviation) obtained for all the species and sites was 0.45 ± 2.373 , while the median ($\pm$ median absolute deviation) is 0.028 ± 0.022 (see supplementary Table S1) which is far lower than the theoretical ratio 3:1. Across *Sphagnum* species within sites and between years, the CF varied orders of magnitude (Table 2). CF values in peat also differed substantially between laboratory-based, and *in situ* incubations in 2016 and 2017 at the Migneint site, while in case of Fenn's and Whixall and Forsinard the data were <LOD except in once instance (Table 2). Overall, the available data suggest high variability in CF values of peat between sites and years (Table S1). In case of *S. cuspidatum* we observed extreme CF values during the laboratory incubations in 2016 from the Migneint site, *in situ* incubation at the Migneint site in 2016, and *in situ* incubations at the Forsinard site in 2017, which resulted in variations up to three orders of magnitude. Exclusion of these extreme CF values still resulted in significant differences where the median CFs by species and year and between lab-based and *in situ* incubations exhibited variations of up to two orders of magnitude (see supplementary Table S2).

287 Table 2. Conversion factors obtained as the ratio between C₂H₄ production measured by the ARA method divided by the N fixed using the ¹⁵N₂
288 method. Data shown for species (rows) is the median (± median absolute deviation - MAD) of the three replicates (four for Migneint and Fenn's
289 and Whixall 2016). Mean conversion factors (± standard deviation) for the *Sphagnum* species and peat are also shown. LOD: under the limit of
290 detection. ND: no data.

Site	Lab incubations Migneint	Migneint		Fenn's & Whixall		Forsinard	Median of the three sites for the different species (±MAD)
Name / Year	2016	2016	2017	2016	2017	2017	2017
<i>S. cuspidatum</i>	0.457 (±0.071)	5.363 (±1.740)	0.056 (±0.014)	0.002 (±0.001)	0.002 (±0.001)	0.114 (±0.003)	0.056 (±0.054)
<i>S. capillifolium</i>	0.18 (±0.087) ⁴	0.025 (±0.011)	0.010 (±0.0002)	0.035 (±0.005)	0.002 (±0.001)	0.010 (±)	0.010 (±0.0002)
<i>S. fallax</i>	0.053 (±0.035)	0.047 (±0.020)	0.036 (±0.004)	0.033 (±)	0.005 (±0.002)	0.008 (±0.008)	0.008 (±0.003)
<i>S. papillosum</i>	0.015 (±0.015)	0.014 (±0.008)	0.043 (±0.003)	0.010 (±0.003)	0.031 (±0.020)	0.094 (±0.026)	0.043 (±0.012)
Median (±MAD)	0.095 (±0.084)	0.045 (±0.036)	0.039 (±0.010)	0.012 (±0.011)	0.005 (±0.003)	0.081 (±0.060)	0.026 (±0.018)

	Mean (±SD)	0.205 (±0.252)	1.989 (±5.334)	0.091 (±0.191)	0.018 (±0.016)	0.010 (±0.015)	0.203 (±0.323)	0.089 (±0.089)		
	Peat hollows	0.001 (±0.0001)	0.011 (±0.007)	LOD	LOD	LOD	LOD			
	Peat hummocks	0.005 (±0.004)	0.027 (±0.022)	ND	0.020 (±0.013)	ND	ND			
	Median (±MAD)	0.001 (±0.0002)	0.018 (±0.010)		0.020 (±0.013)					
	Mean (±SD)	0.004 (±0.005)	0.019 (±0.019)		0.020 (±0.018)					

291

292 BNF rates measured by the $^{15}\text{N}_2$ method and the ARA method showed significant differences.
293 The median ARA rates of mosses and peat per sites in 2017 estimated by applying the
294 theoretical CF of 3:1 for the C_2H_4 produced during parallel incubations, were significantly
295 lower (Fig. 2; C_2H_4 produced also shown) than the direct BNF rates measured based on the
296 $^{15}\text{N}_2$ assimilation method. The difference between the median BNF rates based on ARA and
297 ^{15}N assimilation measurements was more prominent for the Fenn's and Whixall site (ARA
298 being four hundred times lower than $^{15}\text{N}_2$ method). Moreover, by applying the median field-
299 based CF obtained for each of the sites, the BNF rates obtained by the ARA method were
300 also relatively lower than the ones obtained using the $^{15}\text{N}_2$ assimilation method (Fig. 2),
301 which suggests an important underestimation of BNF by even applying the species, peat and
302 site specific CFs values that we measured.

303 BNF measured by the $^{15}\text{N}_2$ assimilation method was higher than the ARA (C_2H_4 produced, no
304 CF applied) for each *Sphagnum* species in each site, albeit the difference varied widely
305 among species (Fig.3). At Migneint the ^{15}N assimilation rate was 20 (*S. cuspidatum*) to 98 (*S.*
306 *capillifolium*) times larger than the C_2H_4 production; at Fenn's and Whixall it was 54 (*S.*
307 *papillosum*) to 271 (*S. cuspidatum*) times larger; and at Forsinard it was 7 (*S. capillifolium*) to
308 50 (*S. fallax*) times. Note that no detectable fixation was found in peat. This large range in the
309 ratio of ^{15}N fixed to C_2H_4 produced signified lack of consistency among species and sites.

310 3.2. BNF by $^{15}\text{N}_2$ method with and without C_2H_2

311 C_2H_2 failed to completely inhibit N_2 reduction to NH_4^+ using the ^{15}N uptake as a direct
312 evidence (Fig. 4). The partial suppression of BNF by the C_2H_2 was also inconsistent across
313 the three sites compared to BNF rates in the absence of C_2H_2 . The relative percentage of BNF
314 suppression calculated at each site (Suppression % = $[(\text{BNF with } \text{C}_2\text{H}_2 - \text{BNF without } \text{C}_2\text{H}_2) / \text{BNF without } \text{C}_2\text{H}_2] \times 100$)

/ BNF without C_2H_2] * 100) based on the mean was 74% at Migneint, 87% at Fenn's and
 Whixall, and 99% at Forsinard. However, based on the median, we found that in Migneint
 there was no suppression but a relative enhancement of 6%, whilst in Fenn's and Whixall, the
 relative percentage of BNF suppression was 64% and in Forsinard 99%, showing a marked
 variability. The differences in the percentage of BNF suppression among species within each
 site also varied substantially (ranging from none to complete suppression and even
 enhancements) showing an inconsistent suppression pattern (Fig. S1).

Figure 2. Rates of C_2H_4 produced ($C_2H_4\ nmol\ g^{-1}\ DW\ d^{-1}$), BNF rates estimated ($nmol\ N\ g^{-1}\ DW\ d^{-1}$) using the theoretical 3:1 CF of ARA method, BNF rates estimated ($nmol\ N\ g^{-1}\ DW\ d^{-1}$) using site specific CFs (0.039 for Migneint; 0.005 for Fenn's & Whixall; and 0.081 for Forsinard), and direct BNF rates measured using $^{15}N_2$ assimilation method ($nmol\ N\ g^{-1}\ DW\ d^{-1}$) at each site in 2017. The box shows 25th, 50th (central line) and 75th percentile with whiskers showing the min and maximum values. The x sign show mean value and the dot an outlier (<1.5 IQR). Note the log-scale y-axis ($n=15$).

330

331 Figure 3. Estimated C₂H₄ produced (C₂H₄ nmol g⁻¹ DW d⁻¹) using ARA method and direct
 332 BNF measurements (nmol N g⁻¹ DW d⁻¹) using ¹⁵N₂ assimilation method, for each of the
 333 Sphagnum species and peat within each site in 2017. Shown are the median values, ± median
 334 absolute deviation (n=3). No bars mean no N₂ fixation or no C₂H₂ reduction detected due to
 335 values being <LOD. Note the different log-scale y-axis.

Figure 4. Estimated BNF rates ($\text{nmol N g}^{-1} \text{DW d}^{-1}$) in *Sphagnum* mosses and peat, by site, in 2017, using the $^{15}\text{N}_2$ assimilation method with and without C_2H_2 addition. The box shows 25th, 50th (central line) and 75th percentile with whiskers showing the min and maximum values. The x sign show the mean value and the dot an outlier (<1.5 IQR). Note the log-scale y-axis ($n=15$).

4. Discussion

Both for *Sphagnum* spp. and peat, the rates of C_2H_4 produced were lower than the direct assimilation of N determined through the $^{15}\text{N}_2$ assimilation method. This resulted in extremely low species-specific CFs (Table 2) signifying potentially serious underestimation of BNF rates by the ARA method, particularly in *Sphagnum* dominated peatlands. Hardy et al. (1968) reported CF ratios between 3 and 4.5 after empirical measurements of BNF activity in bacterial cultures and nitrogenase enzyme preparations as well as in free-living bacteria using parallel ARA and $^{15}\text{N}_2$ direct assimilation methods. Following this publication, the estimated theoretical CF of 3:1 has been applied in various ecosystems including peatlands (Basilier, 1979; Markham, 2009; Rousk et al., 2018). However, many studies have reported greater CFs than the theoretical one ranging from 3.11 to 4.5 for peat and *Sphagnum* spp. together (Basilier, 1980; Chapman and Hemond, 1982; Urban and Eisenreich, 1988; Kox et

al., 2016; Warren et al., 2017). One potential plausible explanation for the discrepancies over the theoretical CF was the possibility of significant endogenous C₂H₄ production in subsurface peat (Schwintzer, 1983; Kox et al., 2016). Conversely, lower CFs than the theoretical CF have also been reported for *Sphagnum* spp. such as 2.48 (Sorensen et al., 2006), 0.85 (Stewart et al., 2011), 0.32 (Vile et al., 2014), or *Sphagnum* peat 1.1 (Knorr et al., 2015), as well as for bryophytes 0.25 (Menge and Hedin, 2009). Therefore the reported site specific CF of peatlands show marked deviations from the theoretical CF and this is consistent with the range of CF measured under this study, albeit we have gone further in estimating CFs per species and in different peatlands across an Nr deposition gradient. High rates of Nr deposition did not shut down BNF and we did not detect any effect on the comparative performance of the methods and the resulting calculation of the CFs. The high variability of these measured CFs suggests that the *Sphagnum* microbiome and its species-specific distribution (Kostka et al., 2016), as well as the inhibitory effects of C₂H₂ on microbial processes such as methanotrophy, nitrification and nitrous oxide reduction may be at play, thus leading to highly inconsistent CFs across species, sites and time. We speculate that such a differential effect could be responsible for the extreme CF values estimated in case of *S. cuspidatum* (Lab incubations Migneint 2016, Migneint 2016) and hence as a *hot spot* of biogeochemical processes (McClaine et al., 2003).

The variability in the measured CFs in this study is further substantiated by the fact that the presence of C₂H₂ differentially affected the suppression of N₂ fixation, but did not completely suppress it across the sites as demonstrated under pure in vitro incubations of nitrogenase enzymes in the presence of C₂H₂ (Koch and Evans, 1966; Schöllhorn and Burris, 1967). This differential suppression response under the ARA must have led to the variable CF ratios we estimated in this study. It appears that the diversity of diazotrophic communities from autotrophic cyanobacteria to chemolithotrophic methanotrophs associated with *Sphagnum*

379 mosses in peatlands may have been affected differentially by C_2H_2 with varied C_2H_4
380 production across species, sites and time. The CF is estimated to save resources for
381 widespread application of ARA in peatlands; however, the difference in CFs at species and
382 site level and over time is not consistent and thus we recommend the use of ^{15}N assimilation
383 method for measuring BNF in peatlands.

384 The very low rates of C_2H_4 production by the ARA in our study could be explained by the
385 presence of methanotrophs and the inhibitory effects of C_2H_2 on the methane monooxygenase
386 enzyme in these bacteria; thus depriving the methanotrophs of a significant bacterial energy
387 to fuel BNF (Flett et al., 1975; De Bont and Mulder, 1976). Widespread presence of
388 methanotrophs associated with *Sphagnum* species has been established (Basiliko et al., 2004;
389 Raghoebarsing et al., 2005; Kip et al., 2010; Larmola et al., 2010). Furthermore, Larmola et
390 al. (2014) and Vile et al. (2014) have demonstrated that methanotrophs can contribute up to
391 40% to total BNF in peatlands, therefore the use of the ARA method could underestimate the
392 BNF rates up to a similar percentage and even more, as shown by our study (~53% on
393 average), which is in agreement with a peat BNF study (~55%) in Minnesota, USA (Warren
394 et al., 2017). C_2H_2 also shuts down nitrification, and stops the reduction of N_2O to N_2 (Ryden,
395 1982). The inhibition of these processes, particularly of nitrification, leads to an increase in
396 plant available nitrogen (NH_4^+) that may in turn limit biological nitrogen fixation (Stewart et
397 al. 2013) particularly in ecosystems such as peatlands with a tightly coupled N cycle such as
398 peatlands. Moreover, inhibition of N_2O reduction which could potentially provide energy
399 and substrate for BNF, might also be a factor in affecting C_2H_4 production rates and hence
400 low and inconsistent CF in the end as observed in this study. For example, Farias et al. (2013)
401 and Desloover et al. (2014) reported respiratory reduction of N_2O to N_2 and its subsequent
402 fixation by diazotrophs in pure bacterial cultures and thus inhibition of N_2O reduction to N_2
403 by C_2H_2 in peatlands might be affecting N fixation rates.

High microbial diversity has been found in *Sphagnum* species between different habitats (e.g. hummocks vs hollows) within peatlands (Opelt et al., 2007). Two main functional groups of bacteria have been studied in *Sphagnum* mosses, nitrogen-fixers and methane-oxidizers (some of which are able to fix nitrogen as well; Auman et al., 2001), and important differences in the community diversity of these two types of bacteria between *Sphagnum* species have been reported (Bragina et al., 2013). This microbial community diversity could potentially explain the high variability of the CF between *Sphagnum* spp. observed in our study, due to the differential presence of these kinds of bacteria, and the different level of interference in their associated microbiological processes in the presence of C₂H₂. Further microbiological studies are recommended to verify the net inhibitory impact against the abundance and expression of N fixers, methanotrophs, nitrifiers and N₂O reducers.

4.1. Conclusions

The conversion factors measured under this study using the direct ¹⁵N assimilation and ARA methods were inconsistent across species, site, and peat and over time. This lack of reproducibility and deviation from the theoretical CF of 3:1 show that ARA as a proxy method cannot fully reflect the BNF activity and hence fixation rates in peatlands. This lack of consistency and partial, yet differential suppression of N₂ fixation in the presence of C₂H₂ led to lower CF values and hence underestimation of BNF. Direct interference and/or inhibition of microbes, particularly methanotrophs seem to result in the differential suppression of N₂ fixation. Therefore, caution is needed when estimating and modelling BNF rates based on the ARA method in peatlands. We conclude that the ARA method is not suitable for BNF measurement in *Sphagnum*-dominated peatlands.

Acknowledgements

We acknowledge the financial and in-kind support of the Keele University ACORN Program, Dr Andrew Stott at the NERC's Life Sciences Mass Spectrometry Facility at Lancaster and the British Bryological Society. We thank Jesus Fincias for assistance during the field sampling campaigns. We thank the National Trust and Natural England for allowing us access to their sites for research in the UK. Thanks to the management team at the Fenns-Whixal, Migneint and Forsinard sites for helping us in the field with relevant information and logistics.

References

- Ackermann, K., Zackrisson, O., Rousk, J., Jones, D.L., DeLuca, T.H. 2012. N₂ fixation in feather mosses is a sensitive indicator of N deposition in boreal forests. *Ecosystems* 15, 986-998.
- Auman, A., Speake, C., Lidstrom, M., 2001. nifH sequences and nitrogen fixation in type I and type II methanotrophs. *Applied and Environmental Microbiology* 67, 4009-4016.
- Bahlmann, E., Bernasconi, S.M., Bouillon, S., Houtekamer, M., Korntheuer, M., Langenberg, F., Mayr, C., Metzke, M., Middelburg, J.J., Nagel, B., Struck, U., Voß, M., Emeis, K.C., 2010. Performance evaluation of nitrogen isotope ratio determination in marine and lacustrine sediments: an inter-laboratory comparison. *Organic Geochemistry* 41, 3–12.
- Basilier, K., 1979. Moss-associated nitrogen fixation in some mire and coniferous forest environments around Uppsala, Sweden. *Lindbergia* 5, 84-88.
- Basilier, K. 1980. Fixation and Uptake of Nitrogen in *Sphagnum* Blue-Green-Algal Associations. *Oikos* 34, 239-242.

449 Basiliko, N., Knowles, R., Moore, T., 2004. Roles of moss species and habitat in methane
450 consumption potential in a northern peatland. *Wetlands* 24, 178-185.

451 Bellenger, J.P., Xu, Y., Zhang, X., Morel, F.M.M., Kraepiel, A.M.L., 2014. Possible
452 contribution of alternative nitrogenases to nitrogen fixation by asymbiotic N₂-fixing
453 bacteria in soils. *Soil Biology & Biochemistry* 69, 413-420.

454 Berg, A., Danielsson, Å., Svensson, B., 2013. Transfer of fixed-N from N-fixing
455 cyanobacteria associated with the moss *Sphagnum riparium* results in enhanced growth
456 of the moss. *Plant and Soil* 362, 271-278.

457 Bergersen F. J., 1970. The quantitative relationship between nitrogen fixation and the
458 acetylene-reduction assay. *Australian Journal of Biological Sciences* 23, 1015-1025.

459 Bragina, A., Berg, C., Mueller, H., Moser, D., Berg, G., 2013. Insights into functional
460 bacterial diversity and its effects on Alpine bog ecosystem functioning. *Scientific*
461 *Reports* 3, 1955.

462 Chalk, P.M., He, J.Z., Peoples, M.B., Chen, D., 2017. ¹⁵N₂ as a tracer of biological N₂
463 fixation: A 75-year retrospective. *Soil Biology & Biogeochemistry* 106, 36-50.

464 Chapman, R.R., Hemond, H.F., 1982. Dinitrogen Fixation by Surface Peat and *Sphagnum* in
465 an Ombrotrophic Bog. *Canadian Journal of Botany-Revue Canadienne De Botanique* 60,
466 538-543.

467 Dabundo, R., Lehmann, M.F., Treibergs, L., Tobias, C.R., Altabet, M.A., Moisander, P.H.,
468 Granger, J., 2014. The Contamination of Commercial N-15(2) Gas Stocks with N-15-
469 Labeled Nitrate and Ammonium and Consequences for Nitrogen Fixation
470 Measurements. *PLoS ONE* 9, e110335.

471 De Bont, J.A., Mulder, E.G., 1976. Invalidity of the acetylene reduction assay in alkane-
 472 utilizing, nitrogen-fixing bacteria. *Applied and Environmental Microbiology* 31, 640-
 473 647.

474 Desloover, J., Roobroeck, D., Heylen, K., Puig, S., Boeckx, P., Verstraete, W., Boon, N.,
 475 2014. Pathway of nitrous oxide consumption in isolated *Pseudomonas stutzeri* strains
 476 under anoxic and oxic conditions. *Environmental Microbiology* 16, 3143-3152.

477 Farias, L., Faundez, J., Fernandez, C., Cornejo, M., Sanhueza, S., Carrasco, C. (2013).
 478 Biological N₂O fixation in the eastern south Pacific Ocean and marine cyanobacterial
 479 cultures. *PLoS ONE* 8, UNSP e63956.

480 Flett, R., Rudd, J., Hamilton, R., 1975. Acetylene-Reduction Assays for Nitrogen-Fixation in
 481 Freshwaters - Note of Caution. *Applied Microbiology* 29, 580-583.

482 Grosskopf, T., Mohr, W., Baustian, T., Schunck, H., Gill, D., Kuypers, M.M.M., Lavik, G.,
 483 Schmitz, R.A., Wallace, D.W.R., LaRoche, J., 2012. Doubling of marine dinitrogen-
 484 fixation rates based on direct measurements. *Nature* 488, 361–4.

485 Hardy, R.W.F., Holsten, R.D., Jackson, E.K., Burns, R.C., 1968. The acetylene–ethylene
 486 assay for N₂ fixation: laboratory and field evaluation. *Plant Physiology* 43, 1185–1207.

487 Ho, A., Bodelier, P.L.E., 2015. Diazotrophic methanotrophs in peatlands: the missing link?
 488 *Plant and Soil* 389, 419-423.

489 Jardine, T., Cunjak, R., 2005. Analytical error in stable isotope ecology. *Oecologia* 144, 528–
 490 533.

491 Kip, N., van Winden, J.F., Pan, Y., Bodrossy, L., Reichart, G., Smolders, A.J.P., Jetten,
 492 M.S.M., Damste, J.S.S., Op den Camp, H.J.M., 2010. Global prevalence of methane
 493 oxidation by symbiotic bacteria in peat-moss ecosystems. *Nature Geoscience* 3, 617-621.

494 Knorr, K., Horn, M.A., Borken, W., 2015. Significant nonsymbiotic nitrogen fixation in
 495 Patagonian ombrotrophic bogs. *Global Change Biology* 21, 2357-2365.

496 Koch, B., Evans, H.J., 1966. Reduction of acetylene to ethylene by soybean root nodules.
 497 *Plant Physiology* 41, 1748-49.

498 Kostka, J.E., Weston, D.J., Glass, J.B., Lilleskov, E.A., Shaw, A.J., Turetsky, M.R., 2016.
 499 The *Sphagnum* microbiome: new insights from an ancient plant lineage. *New*
 500 *Phytologist* 211, 57-64.

501 Kox, M.A.R., Luke, C., Fritz, C., van den Elzen, E., van Alen, T., Op den Camp, H.J.M.,
 502 Lamers, L.P.M., Jetten, M.S.M., Ettwig, K.F., 2016. Effects of nitrogen fertilization on
 503 diazotrophic activity of microorganisms associated with *Sphagnum magellanicum*. *Plant*
 504 *and Soil* 406, 83-100.

505 Larmola, T., Leppanen, S.M., Tuittila, E., Aarva, M., Merila, P., Fritze, H., Tirola, M., 2014.
 506 Methanotrophy induces nitrogen fixation during peatland development. *Proceedings of*
 507 *the National Academy of Sciences of the United States of America* 111, 734-739.

508 Larmola, T., Tuittila, E., Tirola, M., Nykänen, H., Martikainen, P.J., Yrjälä, K., Tuomivirta,
 509 T., Fritze, H., 2010. The role of *Sphagnum* mosses in the methane cycling of a boreal
 510 mire. *Ecology* 91, 2356-2365.

511 Liengen, T., 1999. Conversion factor between acetylene reduction and nitrogen fixation in
512 free-living cyanobacteria from high arctic habitats. *Canadian Journal of Microbiology*
513 45, 223-229.

514 Markham, J.H., 2009. Variation in moss-associated nitrogen fixation in boreal stands.
515 *Oecologia* 161, 353–359.

516 McClain, M. E., Boyer, E. W., Dent, C. L., Gergel, S. E., Grimm, N. B., Groffman, P. M.,
517 Hart, S. C., Harvey, J. W., Johnston, C. A., Mayorga, E., McDowell, W. H., Pinay, G.,
518 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial and
519 aquatic ecosystems. *Ecosystems* 6, 301-312.

520 Menge, D.N.L., Hedin, L.O., 2009. Nitrogen fixation in different biogeochemical niches
521 along a 120 000-year chronosequence in New Zealand. *Ecology* 90, 2190-2201.

522 Myrold, D.D., Ruess, W.R., Klug, M.J., 1999. Dinitrogen fixation. In: Robertson, G.P.,
523 Coleman, D.C., Bledsoe, C.S., Sollins, P. (Eds.), *Standard soil methods for long-term*
524 *ecological research*. Oxford University Press, Oxford, pp. 241-257.

525 Nohrstedt, H. 1983. Conversion factor between acetylene reduction and nitrogen-fixation in
526 soil: effect of water content and nitrogenase activity. *Soil Biology & Biochemistry* 15,
527 275-279.

528 Opelt, K., Berg, C., Schoenmann, S., Eberl, L., Berg, G., 2007. High specificity but
529 contrasting biodiversity of *Sphagnum*-associated bacterial and plant communities in bog
530 ecosystems independent of the geographical region. *ISME Journal* 1, 502-516.

531 Postgate, J.R., 1982. *The fundamentals of nitrogen fixation*, Cambridge University Press,
532 Cambridge.

533 Raghoebarsing, A., Smolders, A., Schmid, M., Rijpstra, W., Wolters-Arts, M., Derksen, J.,
 534 Jetten, M., Schouten, S., Damste, J., Lamers, L., Roelofs, J., den Camp, H., Strous, M.,
 535 2005. Methanotrophic symbionts provide carbon for photosynthesis in peat bogs. *Nature*
 536 436, 1153-1156.

537 Roskoski, J., 1981. Comparative C₂H₂ reduction and ¹⁵N₂ fixation in deciduous wood litter.
 538 *Soil Biology & Biochemistry* 13, 83-85.

539 Rousk, K., Michelsen, A., 2016. The sensitivity of moss-associated nitrogen fixation towards
 540 repeated nitrogen input. *PLoS ONE* 11, e0146655.

541 Rousk, K., Sorensen, P.L., Lett, S., Michelsen, A., 2015. Across-Habitat Comparison of
 542 Diazotroph Activity in the Subarctic. *Microbial Ecology* 69, 778-787.

543 Rousk, K., Vestergaard, M., Christensen, S., 2018. Are nitrous oxide emissions and nitrogen
 544 fixation linked in temperate bogs? *Soil Biology & Biochemistry* 123, 74-79.

545 Ryden, J. 1982. Effects of Acetylene on Nitrification and Denitrification in 2 Soils during
 546 Incubation with Ammonium-Nitrate. *Journal of Soil Science* 33, 263-270.

547 Schöllhorn, R., Burris, R.H., 1967. Acetylene as a competitive inhibitor of N₂ fixation.
 548 *Proceedings of the National Academy of Sciences of the United States of America* 58,
 549 213-16.

550 Schwintzer, C.R., 1983. Non-Symbiotic and Symbiotic Nitrogen-Fixation in a Weakly
 551 Minerotrophic Peatland. *American Journal of Botany* 70, 1071-1078.

552 Sgouridis, F., Stott, A., Ullah, S., 2016. Application of the ¹⁵N gas-flux method for measuring
 553 in situ N₂ and N₂O fluxes due to denitrification in natural and semi-natural terrestrial

554 ecosystems and comparison with the acetylene inhibition technique. *Biogeosciences* 13,
555 1821-1835.

556 Smith, R.J., Gallon, J.R., 1993. Nitrogen fixation. In Lea, P.J. & Leegood, R.C.(Eds), *Plant*
557 *biochemistry and molecular biology*, John Wiley & Sons, Chichester, pp. 129-153.

558 Sorensen, P.L., Jonasson S., Michelsen A., 2006. Nitrogen fixation, denitrification, and
559 ecosystem nitrogen pools in relation to vegetation development in the subarctic. *Arctic*
560 *Antarctic and Alpine Research* 38, 263–272.

561 Sprent, J.I., 1979. *The biology of nitrogen-fixing organisms*, McGraw-Hill, London.

562 Staal, M., Lintel-Hekkert, S., Harren, F., Stal, L., 2001. Nitrogenase activity in cyanobacteria
563 measured by the acetylene reduction assay: a comparison between batch incubation and
564 on-line monitoring. *Environmental Microbiology* 3, 343-351.

565 Stewart, K.J., Coxson, D., Grogan, P., 2011. Nitrogen Inputs by Associative Cyanobacteria
566 across a Low Arctic Tundra Landscape. *Arctic Antarctic and Alpine Research* 43, 267-
567 278.

568 Stewart, K.J., Brummell, M.E., Coxson, D.S., Siciliano, S.D., 2013. How is nitrogen fixation
569 in the high arctic linked to greenhouse gas emissions? *Plant and Soil* 362, 215–229.

570 Urban N., Eisenreich, S., 1988. Nitrogen cycling in a forested Minnesota bog. *Canadian*
571 *Journal of Botany* 66, 435– 449.

572 van den Elzen, E., van den Berg, L.J.L., van der Weijden, B., Fritz, C., Sheppard, L.J.,
573 Lamers, L.P.M., 2018. Effects of airborne ammonium and nitrate pollution strongly

574 differ in peat bogs, but symbiotic nitrogen fixation remains unaffected. Science of the
575 Total Environment 610, 732-740.

576 Vile, M., Kelman Wieder, R., Živkovic, T., Scott, K., Vitt, D., Hartsock, J., Iosue, C., Quinn,
577 J., Petix, M., Fillingim, H., Popma, J., Dynarski, K., Jackman, T., Albright, C., Wykoff,
578 D., 2014. N-fixation by methanotrophs sustains carbon and nitrogen accumulation in
579 pristine peatlands. Biogeochemistry 121, 317-328.

580 Warren, M.J., Lin, X., Gaby, J.C., Kretz, C.B., Kolton, M., Morton, P.L., Pett-Ridge, J.,
581 Weston, D.J., Schadt, C.W., Kostka, J.E., Glass, J.B., 2017. Molybdenum-Based
582 Diazotrophy in a *Sphagnum* Peatland in Northern Minnesota. Applied and
583 Environmental Microbiology 83, UNSP e01174.

584 Zehr, J.P., Montoya, J.P., 2007. Measuring N₂ fixation in the field. In Bothe, H., Ferguson,
585 S.J., Newton, W.E. (Eds), Biology of the nitrogen cycle. Amsterdam, Netherlands:
586 Elsevier, pp. 193-205.

587