Which violence against women educational strategies are effective for pre-qualifying healthcare students? A systematic review

Sammut, Dana; Bradbury-Jones, Caroline

DOI:

[10.1177/1524838019843198](https://doi.org/10.1177/1524838019843198)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Sammut, D & Bradbury-Jones, C 2019, 'Which violence against women educational strategies are effective for pre-qualifying healthcare students? A systematic review', *Trauma, Violence and Abuse*.
<https://doi.org/10.1177/1524838019843198>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 13/03/2019

Sammut, D., Kuruppu, J., Hegarty, K., & Bradbury-Jones, C., Which Violence Against Women Educational Strategies Are Effective for Prequalifying Health-Care Students?: A Systematic Review, Trauma, Violence, and Abuse © The Author(s) 2019. DOI: 10.1177/1524838019843198.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Trauma, Violence, & Abuse

Which violence against women educational strategies are effective for pre-qualifying healthcare students? A systematic review

Journal:	<i>Trauma, Violence, & Abuse</i>
Manuscript ID	TVA-18-101.R1
Manuscript Type:	Review Manuscripts
Keywords:	Assessment < Domestic Violence, Battered women < Domestic Violence, Disclosure of domestic violence < Domestic Violence, Intervention/treatment < Domestic Violence, Adult victims < Sexual Assault

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Which violence against women educational strategies are effective for pre-qualifying healthcare students? A systematic review

Background

Gender-based violence (GBV) is a broad term describing behaviors that cause harm to individuals for reasons associated with gender (Bloom, 2008). While both men and women can be victims of GBV, women are disproportionately affected (World Health Organization (WHO), 2013). Internationally, the forms of GBV most widely reported and studied are intimate partner violence (IPV) and sexual violence, with estimates suggesting that 35% of women worldwide have experienced either or both during their lifetime (WHO, 2013). Women suffer both short- and long-term “exacerbated consequences” when compared with male GBV victims, including reduced ability to work and raise a family (United Nations Statistics Division (UNSD), 2015), complications with pregnancy and childbirth, and increased risk of alcohol misuse, depression and suicide (WHO, 2013). For these reasons, this review concentrates exclusively on violence against women (VAW).

While there is debate about the efficacy of routine screening for IPV, it remains essential for clinicians to have the skills and confidence to ask and address the issue competently in practice, regardless of whether or not abuse has been disclosed. In a meta-analysis of qualitative studies, Feder, Hutson, Ramsay, & Taket (2006) found that women largely felt inquiry about IPV to be appropriate when the issue was approached sensitively by healthcare professionals. Moreover, Leung, Phillips, Bryant, and Hegarty (2018) found that physicians’ perceived ‘readiness’ and ‘preparedness’ were multifaceted concepts with an important bearing on their perceived role to intervene in IPV. Through early detection of abuse (as well as identifying those at greater risk), healthcare professionals are well placed to make referrals and/or facilitate the implementation of evidence-based interventions appropriate to the context (O’Doherty, Taket, Valpied, and Hegarty, 2016).

Regional differences are significant, with some parts of the world experiencing a higher concentration of specific violence forms. For example, the practice of female genital mutilation/cutting (FGM/C) is known to be more concentrated in 29 countries across Africa and the Middle East, with more than 125 million females believed to be currently affected (United Nations Children's Fund, 2013). Similarly, attitudes towards IPV vary considerably across countries and cultures, with levels of women's acceptance varying from 3% in parts of Europe to 92% in Guinea (UNSD, 2015). Within Europe, it is noteworthy that legal definitions of GBV vary between countries, with some forms of VAW yet to be criminalized in certain European Union member states (European Commission, 2016). Discrepancies in approaches to GBV also exist among U.S. states, with a variety of intervention programs continuing in some states despite minimal evidence to support their effectiveness (Langhinrichsen-Rohling, 2010). These regional differences highlight the importance of conducting research with an international perspective to ensure that educational efforts are comprehensive.

A number of studies have identified lack of GBV educational strategies as being a problem, for both pre-qualifying (Doran & Hutchinson, 2017; Valpied, Aprico, Clewett, & Hegarty, 2017) and post-qualifying health professionals (Crombie, Hooker, & Reisenhofer, 2016; Sprague, Madden, Simunovic, Godin, Pham, & Bhandari, 2012). The term 'qualifying' here refers to ability to practice professionally in line with legal requirements (i.e. education, training, registration with professional body). Other studies report that despite having received some formal education and training, qualified practitioners continue to demonstrate poor GBV knowledge and identification rates (Haist, Wilson, Lineberry, & Griffith, 2007; Hinderliter, Doughty, Delaney, Pitula, & Campbell, 2003). Notably, in the study by Hinderliter *et al.* (2003), this was despite self-reported improved confidence among surveyed participants following their IPV education programs, suggesting that feelings of competence do not always

align with objective outcome improvements. On the other hand, practitioners' perceptions of their own knowledge and abilities are undoubtedly an important factor for response to GBV. Authors have noted that lack of confidence in one's own ability to recognize and address GBV is a common concern among students (Bradbury-Jones & Broadhurst, 2015) and qualified practitioners (Taylor, Bradbury-Jones, Kroll, & Duncan, 2013), likely linking to poorer rates of GBV recognition.

Previous literature reviews have been conducted with a similar focus of interest to our study, although none were equal in scope (i.e. including multiple forms of GBV, various pre-professional disciplines and an international focus). However, their findings are relevant and complement many conclusions drawn here. For example, Crombie, Hooker and Reisenhofer's (2016) scoping review of nursing and midwifery IPV education provided insights into gaps in the literature which needed exploring, including the need for rigorous evaluation of existing education programs. Hamberger's (2007) review of IPV curricula for medical students in the U.S. made similar recommendations for research. Sawyer, Coles, Williams, and Williams (2016) produced a systematic review examining the effects of several IPV education programs for both pre- and post-qualifying allied health professionals. The authors concluded that there was an overall positive association between IPV education and improvements in knowledge, attitudes, skills and behaviors, although it was noted that findings were limited by the overall quality of included studies. The latter review differs from ours in that it asked *whether* IPV educational interventions are effective, with less attention given to *how*. Additionally, its sole focus on IPV, inclusion of qualified populations and exclusion of medical students distinguishes it from the present review, with only three of its 18 included studies matching those shortlisted for inclusion here.

To our knowledge, there have been no comparative studies of pre-qualifying GBV educational strategies across multiple healthcare disciplines. It is important, however, that educational interventions are developed in line with the best available evidence to maximize their impact and efficacy. It is essential that students are competent to recognize and address GBV before qualifying, given the significant health impacts of VAW (WHO, 2013).

Review Method

Aim and Design

This review aimed to identify, collate and critique existing evidence about educational strategies on GBV for pre-qualifying healthcare students, with a view to ascertaining best practice in this subject area and informing future pre-qualifying training programs. A systematic literature review was conducted to synthesize findings from quantitative, qualitative and mixed methods primary research studies, to ensure that all relevant data relating to the research question could be captured (Grant & Booth, 2009). **Inclusion of quantitative and qualitative studies allowed for consideration of a wider variety of outcome measures, including both objectively measured and self-reported results, and resulted in a wider pool of data from which to draw conclusions. We therefore use the word 'effectiveness' to encompass a variety of criteria (summarized under 'measured outcomes' column in Table 3).**

Search Strategy

A computerized search of six databases was undertaken: Medline, PsychINFO, Embase, CINAHL, Applied Social Sciences Index and Abstracts (ASSIA), and Nursing & Allied Health Database. Searches were conducted between September 2017 and July 2018. PICO criteria (Population, Intervention, Comparison and Outcomes) were determined in advance to facilitate a systematic approach and to ensure articles were selected appropriately (Table 1). Searches were conducted using a combination of keywords and Medical Subject Headings (MeSH) to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

allow greater sensitivity in the search strategy. Boolean operators and truncation were used to broaden and/or narrow the search appropriately. A summary of the search terms is included in Table 2. Finally, a review of the reference lists of all key articles identified was undertaken, in order to find any further potential papers for inclusion.

[Insert Tables 1 and 2 about here]

Inclusion and Exclusion Criteria

To meet this review’s aims, only studies discussing and/or evaluating a GBV-related education intervention were included. Sample populations had to be students on a pre-qualifying healthcare course or degree. No date restriction was applied to searches. Exclusion criteria included studies with only a partial focus on the education intervention; studies focusing on violence against men and/or children (including adolescents), as well as non-gender-specific elder abuse; unpublished and non-peer-reviewed research; studies not published in English; and non-empirical or secondary research (including reviews, editorials, opinions and discussion papers).

Quality Assessment Method

Following selection, the quality of all included studies was evaluated. Appraisal tools were selected according to study design and purpose so that methodologies would be assessed appropriately (Whittemore & Knafl, 2005). A BestBETs (2012) tool for appraising educational interventions was used for all quantitative and mixed-methods studies, as it allowed for consideration of both quantitative and qualitative criteria. However, as this tool was considerably more quantitative-focused, a Critical Appraisal Skills Programme (CASP, 2017) checklist (recommended by Ciliska, Thomas, & Buffett; 2008) was used to appraise the single qualitative study shortlisted for inclusion.

Meade and Richardson (1997) describe the process of assigning numerical quality scores to individual studies as “arbitrary and unscientific” (p. 535). Instead, they suggest that the merits and limitations of each study should be presented clearly in the review write-up, so that readers can judge for themselves the usefulness of each. This approach was adopted here, with an additional summary of methodological rigor and data relevance for each paper, as proposed by Tranter, Irvine and Collins (2011) (adapted from Whitemore & Knafl, 2005). Whitemore and Knafl (2005) also suggest that each study’s quality should be considered in a “meaningful way” (p. 550): for example, if discrepancies in findings may be reasonably attributed to poor methodology. As this can only be determined later on in the review process, no studies were excluded from data analysis on the basis of quality alone.

Data Extraction and Synthesis

A data extraction table was created using subheadings which helped to summarize the studies in a meaningful way, capturing information as it related to the research question (Table 3). Statistical meta-analysis was initially attempted for the quantitative studies, but due to the heterogeneity of study designs and outcome measures, a meaningful statistical summation would not have been possible. Therefore, a narrative summary and thematic synthesis of the studies’ findings was undertaken. This was an inductive process, with themes emerging from the data. Themes and coding strategies were cross-checked between two authors to maximize rigor. The Cochrane Consumers and Communication Review Group (2013) guidelines for narrative synthesis were followed to ensure robustness within this process.

Results

Search Outcome and Characteristics of Included Studies

A total of 488 studies were retrieved from searches after duplicates were removed. These were screened against the PICO criteria and excluded first by title ($n = 320$) and then by abstract (n

1
2
3 = 113), leaving 55 for full-text review. Of these, 38 were excluded for not meeting the inclusion
4
5 criteria, leaving 17 studies in total. This process is outlined in the Preferred Reporting Items
6
7 for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram (Figure 1) (Moher,
8
9 Liberati, Tetzlaff, Altman, & The PRISMA Group, 2009). [Insert Figure 1 here]
10
11
12
13
14 Of the 17 studies, 14 were conducted in the USA, and one each in Canada, Australia and
15
16 Ireland. All 17 studies were focused on IPV/domestic violence (DV), general interpersonal or
17
18 sexual violence. There was no discussion of FGM/C, forced marriage, honor violence or human
19
20 trafficking within any of the studies. This is discussed further in the present review's
21
22 limitations. Thirteen used a quantitative design, three used a mixed methods approach and one
23
24 used a qualitative phenomenological design.
25
26
27
28
29 Of the four studies that collected qualitative data, two explored results thematically (Helton &
30
31 Evans, 2001; Pomeroy, Parrish, Bost, Cowlagi, Cook, & Stepura, 2011) and two used open-
32
33 ended questions to collect students' feedback and opinions regarding the intervention (Cerulli,
34
35 Nichols-Hadeed, Raimondi, Stone, & Cerulli, 2015; Smith, Wight, & Homer, 2017). The
36
37 studies each describe the effects of a GBV-related education intervention, and the majority
38
39 reported a significant improvement (where applicable) in the measured outcomes post-
40
41 intervention. Four themes were identified following synthesis: (1) Learning Mechanism, (2)
42
43 Focus of Learning, (3) Duration of Course and (4) Gender of the Audience.
44
45
46
47 [Insert Table 3 about here]
48
49

50
51 **Quality of Included Studies**

52
53 The included studies varied in design and quality (see Table 4 and Table 5). All except one
54
55 presented their aims and outcomes clearly; the quality of the study by Heron, Hassani, Houry,
56
57 Quest, and Ander (2010) was compromised due to lack of clarity. Only two papers used a RCT
58
59 study design (Danley, Gansky, Chow, & Gerbert, 2004; Edwardsen, Morse, & Frankel, 2006).
60

The majority of included studies used quasi-experimental pretest-posttest designs, which can pose threats to internal and external validity. For example, if both pretest and posttest utilize the same assessment instrument, posttest results may be skewed by the participants' familiarity with the content being assessed (Knapp, 2016).

Participant and assessor blinding was only discussed in three studies (Edwardsen, Morse, & Frankel, 2006; Elman, Hooks, Tabak, Regehr, & Freeman, 2004; Pomeroy *et al.*, 2013). If not discussed, it was assumed that blinding did not take place. One study (Heron *et al.*, 2010) mentioned blinding within its abstract, but failed to discuss further within the write-up, and so was rated 'unclear' for this criterion. Moreover, only one study (Milone, Burg, Duerson, Hagen, & Pauly, 2009) provided evidence of sample size estimation.

Seven studies (Elman *et al.*, 2004; Ernst, Houry, Nick, & Weiss, 1998; Ernst, Houry, Weiss, & Szerlip, 2000; Haase, Short, Chapman, & Dersch, 1999; Heron *et al.*, 2010; Jonassen, Pugnaire, Mazor, Regan, Jacobson, Gammon, Doepel, & Cohen, 1999; Pomeroy *et al.*, 2013) included some level of delay between intervention and posttest. There is mixed evidence regarding the significance of delaying posttest, with some authors suggesting that delays provide greater evidence of learning retention (Schmitt, 2010) and others finding no significant difference between immediate and delayed posttest results (Loyens, Jones, Mikkers, & van Gog, 2015). Nevertheless, four included papers (Danley *et al.*, 2004; Edwardsen, Morse, & Frankel, 2006; McAndrew, Pierre, & Kojanis, 2014; Milone *et al.*, 2010) made reference to their short follow-up time when discussing the limitations of their studies.

[Insert Tables 4 and 5 about here]

Theme 1: Learning Mechanism

This theme considers the ways in which learning is delivered to students, primarily exploring the outcomes of didactic and interactive approaches. For the purposes of this discussion,

didactic refers to passive learning, while *interactive* refers to any educational methods requiring active engagement from the students. The latter includes strategies involving standardized patients (SPs), role play, group discussion or even interactive online tutorials.

Buranosky, Hess, McNeil, Aiken, and Chang's (2012) study of 279 medical students, which featured various teaching forms, found that interactive activities had a greater impact than didactic methods for improving measured outcomes. Specifically, two of the post-intervention outcomes (IPV-related awareness and support for universal screening) were significantly improved ($p < 0.001$ for both) for students who elected to undertake additional experiential training, while results showed no significant improvement for those who participated in core (didactic) activities alone. However, it is worth noting that as the experiential activities were optional, this group's improvement may be partly attributable to greater overall interest in the subject. Nevertheless, other studies drew similar conclusions. Milone *et al.* (2010) found that there was no statistically significant difference between medical students ($n = 102$) who had and had not attended a voluntary didactic lecture (all participants had subsequently attended a mandatory SP encounter), again suggesting a limited effect from the didactic intervention. Similarly, Pomeroy *et al.*'s (2011) study on 63 social work students featured two intervention groups (one interactive and one didactic) and one control group. This study found that while the didactic group demonstrated increased knowledge post-intervention, the interactive group showed greater understanding of how knowledge could be integrated in practice.

Helton and Evans (2001), who collected qualitative data from 18 nursing students, emphasized the importance of practical experiences for the students. One student commented: "Going to court really brought to life the amazingly large number of women out there being abused," while another stated: "[The practical experiences] got us out of the textbook comfort zone and into the domestic violence war zone the clients face." In Cerulli *et al.*'s (2015) mixed methods

study, pharmacy students (n = 237) reported that they would prefer a more interactive session – offering suggestions such as small-group discussion and opportunities to interact with mock patients – to the didactic session they attended.

One study which did not conform to this theme was that of Smith, Wight, and Homer (2017), which saw midwifery students give feedback following an all-day workshop. Despite commenting that both didactic and interactive activities had been useful, the aspect of the workshop cited as ‘most useful’ was an authentic practice video (44% of 102 students) which had no interactive component. However, a continuing theme was that students wanted their learning to be practice-focused rather than simply theoretical (discussed in theme 2).

Overall, interactive approaches to teaching students about GBV/VAW seemed more effective.

Theme 2: Focus of Learning

This theme discusses students’ preferences and outcomes relating to the focus of their learning – specifically, whether learning was practice-focused or mainly theoretical in content. While this idea links closely with the discussion of *learning mechanism*, there are subtle differences requiring separate attention. For example, as highlighted by Smith, Wight, and Homer’s (2017) study, didactic activities (such as a video screening) can still be practice-focused, demonstrating real-life scenarios and focusing on how knowledge and skills can be *applied*. This idea was echoed in Cerulli *et al.*’s (2015) study, where the majority of lecture content was focused on communicating factual evidence (e.g. IPV prevalence rates), with less attention given to how this information related to practice. In their qualitative feedback, students indicated that they would have liked more profession-specific content, including discussion of pharmacists’ professional liability and legal obligations when intervening with IPV in practice.

In the study by Edwardsen, Morse, and Frankel (2006), the educational intervention (delivered to both the intervention and control group) featured two hours of didactic and interactive IPV education, including introduction to a mnemonic for facilitating clinical interviews. The intervention group was then provided with a pocket-sized card describing the mnemonic and additional guidance for use in practice, which was later used in a facilitated discussion before the posttest. While there was no difference in overall content covered by the intervention and control group, this modest practice-focused intervention produced consistently improved results in the intervention group compared with the control. However, the improvement was only statistically significant in two of the seven assessed criteria. Hence, because both groups received a similar education program, the difference in preparedness between the intervention and control group was not as considerable as those studies whose control groups received no GBV-related education at all.

Elman *et al.* (2004) raised the issue of transfer of learning – that is, whether improved knowledge and testing scores will translate into better GBV identification in practice. In their objective structured clinical examination (OSCE), medical students ($n = 110$) were more likely to enquire about GBV in the scenario with a more obvious transfer context (i.e. the prenatal patient, as pregnancy had been highlighted as a risk factor for abuse during the theoretical learning session). The educational intervention tested in this study could certainly be called practice-focused, as all students were given the opportunity to interact with a SP following the seminar. However, the authors argue that the lower rates of enquiry about GBV for the elderly musculoskeletal patient could partly be attributed to the fact that the case presented no obvious transfer from the seminar.

Contrary to the above, the study by Heron *et al.* (2010) found that participation in a SP OSCE did not improve medical students' ($n = 41$) self-reported comfort levels to care for IPV victims,

despite this being a clear practice-focused learning opportunity. The authors offered no explanation for this negative finding. Notably, this study was of questionable quality (see Table 4) and its findings should be interpreted with caution.

This theme primarily highlights that practice-focused interventions with a clear transfer context were received more favorably by students.

Theme 3: Duration of Course

Courses ranged in duration from a 15-minute multimedia tutorial (Danley *et al.*, 2004) to a 40-hour experiential program (Buranosky *et al.*, 2012). Two studies (Danley *et al.*, 2004; McAndrew, Pierre & Kojanis, 2014) evaluating brief (≤ 1 hour) computer-based tutorials found that while knowledge scores were significantly improved post-intervention, attitudes and opinion scales were much less so ($p < 0.05$ in only two out of eight items in both studies). Within Danley *et al.*'s (2004) study, two attitudinal questions were consistently non-significant in terms of differences between the control and intervention groups ("There are specific things I can do to help" and "[I] believe I can recognize and help"), suggesting that self-efficacy in particular was difficult to alter. In contrast, the two significantly improved opinions in the study by McAndrew, Pierre and Kojanis (2014) were perceived self-efficacy and perceived impact of constraints (reverse scored), both suggesting an improvement in the students' belief in their ability to effect change. Nevertheless, overall opinions and attitudes were much less significantly improved than knowledge within both studies.

Jonassen *et al.* (1999), who compared two medical student cohorts receiving the same intervention over different lengths of time, found that the longer intervention (lasting 3.5 days) ($n = 67$ students) produced sustained improvements in all three assessed domains (knowledge, attitudes and skills) at 6-month follow-up. In contrast, the cohort receiving the 2-day course ($n = 77$ students) maintained only their skills after 6 months. Notably, the students on the shorter

course rated its duration more favorably. However, in Cerulli *et al.*'s study pharmacy students' feedback on a 1.5-hour didactic lecture overwhelmingly advised that the session was not long enough to thoroughly cover the content. Many students suggested that while the content was informative, the lecture did not adequately prepare them to address IPV in practice, with one individual commenting: "I'm not sure how much this lecture could help us 'spot' one [IPV victim]". This refers back to theme 2: *focus of learning*.

Overall, while all studies discussed within this theme reported positive outcomes to some extent, it seems that students' attitudes were more effectively altered following longer courses.

Theme 4: Gender of the Audience

Four studies (Buranosky *et al.*, 2012; Ernst *et al.*, 1998; Ernst, *et al.*, 2000; Milone *et al.*, 2009) identified a gendered difference in knowledge scores and attitudes of participants, with females consistently outperforming males. Further, in the qualitative study by Helton and Evans (2001) nursing students (n = 18) were more likely to identify with victims of the same gender as themselves. Kennedy, Vellinga, Bonner, Stewart, and McGrath (2013) discussed the literature evidence for the gendered difference in male and female rape myth acceptance, although their own study did not analyze demographic data so as to maintain participant anonymity.

The study on dental students (n = 65) by Everett, Kingsley, Demopoulos, Herschaft, Lamun, Moonie, Bungum, and Chino (2012) found that female participants were statistically more likely ($p = 0.008$) to have received previous IPV education in comparison with male participants, although the study provided no discussion of demographic data in the results (i.e. relationship between gender and changes in students' awareness and professional beliefs). However, the reason for this imbalance may be reflected in Haase *et al.*'s (1999) study, where women were disproportionately interested in undertaking the elective on DV (67% of students on the elective were female, despite comprising only 44% of the cohort (n = 115 medical

students); $p = 0.002$). The elective was found to significantly impact knowledge of resources ($p = 0.000$) and improve identification of DV victims ($p = 0.04$). However, after using two-way ANOVA (analysis of variance) the authors concluded that the difference in scores between the DV-educated and non-educated groups was not attributable to gender. The authors provided no basic data in their report, and it is unclear whether they used ANOVA testing within the intervention group itself to factor out the gender effect. It is therefore impossible to deduce whether females outperformed males in this study. Similarly, the study by Danley *et al.* (2004) found there was no statistically significant difference in the scores of male and female participants ($n = 174$). There is no explanation for this finding and it may simply reflect the baseline knowledge and attitudes of the study's sample.

These findings make it difficult to draw conclusions about the overall effect of audience gender across the studies. However, the fact that gendered differences were noted or addressed within so many papers meant we could not reasonably exclude these points from thematic analysis.

Discussion

This review was conducted to identify best educational practices in GBV for pre-qualifying healthcare students. The reviewed educational strategies varied in structure and content, and assessed numerous outcomes to identify post-learning changes and improvements. Narrative synthesis of the 17 studies generated four themes: *learning mechanism*; *focus of learning*; *duration of course*; and *gender of the audience*. Figure 2 shows a visual representation of these themes. This depiction groups the first three themes together for being 'intrinsic' to the educational program (i.e. inherent aspects which can be manipulated by educators), while the final 'extrinsic' theme is not related to the programs' structure or content. [Insert Figure 2 here]

The first theme, *learning mechanism*, discussed the relative effect sizes of interactive and didactic intervention strategies. Overall, interactive approaches to learning yielded better

1
2
3 results than did didactic. This theme echoes other literature findings on the subject of
4
5 continuing medical education in qualified physicians (Bloom, 2005; Mansouri & Lockyer,
6
7 2007) and other health professionals (O'Brien, Freemantle, Oxman, Wolf, Davis, & Herrin,
8
9 2001). However, much of this previous research has measured 'effectiveness' in relation to
10
11 improved performance in clinical settings and improved patient outcomes (Forsetlund,
12
13 Bjørndal, Rashidian, Jamtvedt, O'Brien, Wolf, Davis, Odgaard-Jensen, & Oxman, 2009).
14
15 These outcomes are not possible to measure in student populations, and as discussed within the
16
17 second theme, *focus of learning*, improved test scores might not always translate into better
18
19 practice. Supporting this idea, a qualitative study by Bradbury-Jones and Broadhurst (2015)
20
21 reports that students themselves often foresee difficulties in linking theory with practice, which
22
23 again highlights the importance of a more practice-focused approach. Similarly, a quasi-
24
25 experimental study by Kripke, Steele, O'Brien, and Novack (1998) assessed the knowledge,
26
27 skills and attitudes of 55 medical residents before, immediately after and again at 6-months
28
29 following a DV workshop. Despite sustained improvements in all three domains at 6-month
30
31 follow-up, no improvement was observed in the residents' actual screening behaviors or DV
32
33 identification rates. The authors argued that creating long-term behavior change should be the
34
35 next goal for GBV educators (Kripke *et al.*, 1998).
36
37
38
39
40
41
42

43 Natan, Khater, Ighbariyea, and Herbet (2016) suggested that behavioral change theories such
44
45 as the Theory of Planned Behaviour (TPB) (developed by Ajzen, 1991) can be used to predict
46
47 students' intention to screen for DV. According to the TPB, a number of variables underlie
48
49 individuals' intentions to perform an action, including behavioral attitudes, normative beliefs
50
51 and perceived behavioral control (Ajzen, 1991). Extrapolating from this, it could be argued
52
53 that these variables ought to be measured at baseline in future study samples, so that
54
55 educational interventions can be developed to target and alter those variables which will predict
56
57 intention to act (Nelson, Cook & Ingram, 2013). However, even if successful, improvements
58
59
60

in intended behaviors may still fail to amount to improvements in practice. These arguments suggest that the outcomes which are measurable in student populations (objective knowledge, self-reported preparedness, perceived ability to effect change, etc.) are less valid than outcomes measurable in qualified populations (increased rates of identification, referral to outside agencies, etc.). That said, the former group of outcomes undoubtedly has a demonstrable effect on *likelihood* of competently performing the latter tasks in practice, and their value must not be discounted. Future researchers may wish to explore the effects of incorporating principles of behavior change into GBV education to help ensure learning evolves into effective practice.

No theoretical consistencies were noted among the papers contributing to this review.

Another theme, *duration of course*, suggested that longer interventions are more effective at altering students' opinions and attitudes. This theme mirrors the findings of a 69-study meta-analysis by Anderson and Whiston (2005), which looked at the effectiveness of various university-level sexual assault education programs. Anderson and Whiston (2005) argue that multiple in-depth sessions produce better outcomes than single-session programs, consistent with findings from later studies (Mansouri & Lockyer, 2007; Marinopoulos, Dorman, Ratanawongsa, Wilson, Ashar, Magaziner, Miller, Thomas, Prokopowicz, Qayyum, & Bass, 2007). This argument also echoes Buranosky *et al.*'s (2012) findings, where outcome improvements correlated with participation in additional training activities, leading the authors to conclude that "once is not enough" (p. 1192).

This is not to say that brief single-session interventions are ineffective. Only four studies included in this review (Buranosky *et al.*, 2012; Haase *et al.*, 1999; Helton & Evans, 2001; Jonassen *et al.*, 1999) featured interventions with multiple sessions taking place over the course of more than one day, yet all except one study (Heron *et al.*, 2010) reported positive findings to some degree. However, based on the trends found in this review and other larger studies, it

seems that longer multifaceted interventions are more effective for producing sustained improvements, particularly in attitudinal outcomes.

The final theme, *gender of the audience*, discussed the trend of gendered differences in students' baseline and post-intervention knowledge and attitudes toward GBV, with females frequently outperforming males. There is a wealth of literature supporting the idea that female healthcare students are less accepting of common rape myths and more positive in their attitudes toward victims of sexual violence (Anderson & Quinn, 2009; Sivagnanam, Bairy, & D'Souza, 2005). However, there is debate surrounding the applicability of this information in educational settings. For example, Brecklin and Forde's (2001) 45-study meta-analysis concluded that rape education programs are more effective (here referring to improved attitudinal outcomes) when delivered in single-gender settings. Conversely, Anderson and Whiston's (2005) 69-study meta-analysis reported very mixed findings on this same topic, with women and men statistically performing better in mixed-gender groups for the majority of measures. One exception to this was the finding that females in single-gender groups showed more positive behavioral intentions, although this conclusion was drawn from the results of only four studies (Anderson & Whiston, 2005). This was also a key difference between these two meta-analyses, with Brecklin and Forde (2001) looking only at attitudinal changes, while Anderson and Whiston (2005) further addressed knowledge and behavioral indices. Behavioral indices are arguably more important in this discussion, as improved performance in practice (both intended and actual) is the main goal of these educational strategies.

These mixed findings suggest that more research is needed before conclusions can be drawn about the effectiveness of single- versus mixed-gender audiences in GBV education. That said, educators should still consider the possibility that male practitioners may encounter specific barriers in practice to which females will be less susceptible (for example, female victims may

1
2
3 feel more reluctant to disclose information to male professionals) (Hester, Williamson, Regan,
4 Coulter, Chantler, Gangoli, Davenport, & Green, 2012). Therefore, while the subject of
5 audience gender itself cannot be conclusively settled, the gendered issues practitioners may
6 encounter should not be overlooked.
7
8
9
10
11
12

13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Limitations

There are some methodological limitations to this study. First, limited human resources and time constraints meant that certain content and study types were systematically excluded during screening. This included non-peer-reviewed and unpublished research. It is generally accepted that published studies are more likely to report significant findings (Cook & Therrien, 2017), and by excluding unpublished research the present review may have drawn biased conclusions. It was also beyond the scope of this review to draw on every aspect of learning theory, meaning many pertinent academic perspectives were omitted. This is discussed further under recommendations for future research. Second, statistical meta-analysis was not possible in this review, nor moderation analysis. However, narrative analysis allowed trends to be identified and offered the additional advantage of inclusion of qualitative data.

The studies contributing to this review's analysis were themselves of varying quality, which could potentially compromise the soundness of conclusions drawn here. Further, the generalizability of this review's results is impacted due to its failure to include studies discussing many wider forms of GBV (honor violence, FGM/C, etc.), which may be partly attributable to the exclusion of non-English language studies. As discussed earlier, some forms of GBV are more culturally driven than others and may therefore be discussed to a greater extent in non-English language journals. That said, a recent European-focused mapping review and synthesis by Bradbury-Jones, Appleton, Clark, and Paavilainen (2017) found that IPV and sexual abuse are the GBV forms that receive the majority of literature attention, consistent with

findings here. While these two forms likely represent the largest global GBV burden, it was disappointing that no other forms featured in this review. Inclusion of studies from non-English-speaking parts of the world could have provided a useful basis for discussion of cultural variation in GBV teaching and learning.

Recommendations for Educational Practice and Future Research

Educators should take heed of the present review’s findings when designing and implementing GBV-focused learning programs. The visual summary of themes (Figure 2) may be useful in the development of future curricula as a summary of aspects of which to be mindful. A key recommendation for future research is to incorporate a focus on wider learning theory – for example, Kolb’s (1984) cycle of experiential learning; Honey and Mumford’s (1986) learning styles; and McGill and Beaty’s (1995) action learning approach. These theoretical approaches could offer a valuable framework upon which to base future educational strategies, incorporating a focus on internal cognitive processes in addition to the external educational factors considered here. Research and education programs should also strive to give greater attention to the wider forms of GBV.

The U.S. is taking steps locally and federally to recognize freedom from domestic violence as a human right (Cornell University, 2018). However, government policy can only go so far in bringing about actual change to healthcare practice. Since the introduction of the Affordable Care Act (ACA) in 2012, which mandates insurance coverage of IPV screening and counselling, U.S. clinicians have continued to demonstrate inadequate screening practices, with many reporting lack of confidence as a significant barrier (Tavrow, Bloom, and Withers, 2017). Chapin, Coleman, and Varner (2011) note that hospitals and other healthcare institutions are often left to design and implement their own policies governing specific practices, yet this too can result in inconsistencies from one organization to another. By introducing effective

GBV educational strategies at pre-qualifying level, these problems can be addressed at the earliest opportunity in healthcare practitioners' careers.

Conclusion

To date, this is the first internationally-focused literature review to combine evidence on the subject of GBV educational strategies for pre-qualifying healthcare students. Findings suggest that interactive and practice-focused learning interventions produce the best results (i.e. knowledge scores, self-reported comfort and confidence, etc.) and that courses of longer duration are more effective in instilling attitudinal change. Future research should consider the gendered differences in GBV learning and whether these differences have an impact on practice. Global efforts to improve outcomes for GBV victims must begin by ensuring healthcare providers receive adequate training *before* they enter professional practice. Despite its limitations, findings from this review make a valuable contribution to knowledge on the subject of GBV learning and offer practical guidance to educators developing future curricula.

REFERENCES *(References included in the present review are marked with an asterisk.)*

Ajzen, I. (1991). The theory of planned behaviour. *Organizational Behavior and Human Decision Processes*, 50(2), 179-211.

Anderson, I. & Quinn, A. (2009). Gender differences in medical students' attitudes towards male and female rape victims. *Psychology, Health & Medicine*, 14(1), 105-110.

Anderson, L. A. & Whiston, S. C. (2005). Sexual assault education programs: A meta-analytic examination of their effectiveness. *Psychology of Women Quarterly*, 29, 374-388.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BestBETs (2012) *BETs CA worksheets*. Retrieved from <http://bestbets.org/links/BET-CA-worksheets.php>, on 19 February, 2018.

Bloom, B. S. (2005). Effects of continuing medical education on improving physician clinical care and patient health: A review of systematic reviews. *International Journal of Technology Assessment in Health Care*, 21(3), 380-385.

Bloom, S. S. (2008). *Violence against women and girls. A compendium of monitoring and evaluation indicators*. Retrieved from <https://bit.ly/2L8BDRe>, on 20 November, 2017.

Bradbury-Jones, C., Appleton, J. V., Clark, M. & Paavilainen, E. (2017). A profile of gender-based violence research in Europe: Findings from a focused mapping review and synthesis. *Trauma, Violence & Abuse*. In press.

Bradbury Jones, C. & Broadhurst, K. (2015). Are we failing to prepare nursing and midwifery students to deal with domestic abuse? Findings from a qualitative study. *Journal of Advanced Nursing*, 71(9), 2062-2072.

Brecklin, L. R. & Forde, D. R. (2001). A meta-analysis of rape education programs. *Violence and Victims*, 16(3), 303-321.

*Buranosky, R., Hess, R., McNeil, M. A., Aiken, A. M. & Chang, J. C. (2012). Once is not enough: Effective strategies for medical student education on intimate partner violence. *Violence Against Women*, 18(10), 1192-1212.

*Cerulli, C., Nichols-Hadeed, C., Raimondi, C., Stone, J. T. & Cerulli, J. (2015). Facilitating intimate partner violence education among pharmacy students: What do future pharmacists want to know? *Currents in Pharmacy Teaching and Learning*, 7(3), 283-291.

Chapin, J. R., Coleman, G. & Varner, E. (2011). Improving medical screening for intimate partner violence through self-efficacy. *Journal of Injury and Violence Research*, 3(1), 19-23.

Ciliska, D., Thomas, H. & Buffett, C. (2008). *An introduction to evidence-informed public health and a compendium of critical appraisal tools for public health practice*. Ontario: National Collaborating Centre for Methods and Tools (NCCMT). Retrieved from <https://bit.ly/2LnuAPV>, on 19 February, 2018.

Cochrane Consumers and Communication Review Group (2013). *Cochrane Consumers and Communication Review Group: Data synthesis and analysis*. Retrieved from <http://cccr.org.cochrane.org>, on 20 February, 2018.

Cook, B. G. & Therrien, W. J. (2017). Null effects and publication bias in special education research. *Behavioral Disorders*, 42(4), 149-158.

Cornell University (2018). *Freedom from Domestic Violence as a Fundamental Human Right Resolutions, Presidential Proclamations, and Other Statements of Principle*. Retrieved from <https://bit.ly/2wbtEZX>, on 8 August, 2018.

Critical Appraisal Skills Programme (CASP) (2017). *CASP qualitative research checklist*. Retrieved from <http://www.casp-uk.net/casp-tools-checklists>, on 19 February, 2018.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Crombie, N., Hooker, L. & Reisenhofer, S. (2016). Nurse and midwifery education and intimate partner violence: A scoping review. *Journal of Clinical Nursing*, 26, 2100-2125.

*Danley, D., Gansky, S. A., Chow, D. & Gerbert, B. (2004). Preparing dental students to recognize and respond to domestic violence: The impact of a brief tutorial. *Journal of the American Dental Association*, 135(1), 67-73.

Doran, F. & Hutchinson, M. (2017). Student nurses' knowledge and attitudes towards domestic violence: Results of survey highlight need for continued attention to undergraduate curriculum. *Journal of Clinical Nursing*, 26(15-16), 2286-2296.

*Edwardsen, E. A., Morse, D. S. & Frankel, R. M. (2006). Structured practice opportunities with a mnemonic affect medical student interviewing skills for intimate partner violence. *Teaching and Learning in Medicine*, 18(1), 62-68.

*Elman, D., Hooks, R., Tabak, D., Regehr, G. & Freeman, R. (2004). The effectiveness of unannounced standardised patients in the clinical setting as a teaching intervention. *Medical Education*, 38(9), 969-973.

*Ernst, A., Houry, D., Nick, T. G. & Weiss, S. J. (1998). Domestic Violence Awareness and Prevalence in a First-year Medical School Class. *Academic Emergency Medicine*, 5, 64-68.

*Ernst, A., Houry, D., Weiss, S. J. & Szerlip, H. (2000). Domestic violence awareness in a medical school class: 2-year follow-up. *Southern Medical Journal*, 93(8), 772-776.

European Commission (2016). *Special Eurobarometer 449: Gender-based violence*.

European Union Open Data Portal. Retrieved from http://data.europa.eu/euodp/en/data/dataset/S2115_85_3_449_ENG, on 8 February, 2018.

*Everett, R. J., Kingsley, K., Demopoulos, C. A., Herschaft, E. E., Lamun, C., Moonie, S., Bungum, T. J. & Chino, M. (2012). Awareness and beliefs regarding intimate partner violence among first-year dental students. *Journal of Dental Education*, 77(3), 316-322.

Feder, G., Hutson, M., Ramsay, J. & Taket, A. (2006). Women Exposed to Intimate Partner Violence: Expectations and Experiences When They Encounter Health Care Professionals: A Meta-analysis of Qualitative Studies. *Archives of Internal Medicine*, 166, 22-37.

Forsetlund, L., Bjørndal, A., Rashidian, A., Jamtvedt, G., O'Brien, M. A., Wolf, F. M., Davis, D., Odgaard-Jensen, J. & Oxman, A. D. (2009). Continuing education meetings and workshops: Effects on professional practice and health care outcomes. *Cochrane Database of Systematic Reviews*, Issue 2. CD003030.

Grant, M. J. & Booth, A. (2009). A typology of reviews: An analysis of 14 review types and associated methodologies. *Health Information and Libraries Journal*, 26(2), 91-108.

*Haase, C. E., Short, P. D., Chapman, D. M. & Dersch, S. A. (1999). Domestic Violence Education in Medical School: Does It Make a Difference? *Academic Emergency Medicine*, 6(8), 855-857.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Haist, S. A., Wilson, J., Lineberry, M. J. & Griffith, C. H. (2007). A Randomized Controlled Trial Using Insinuated Standardized Patients to Assess Residents' Domestic Violence Skills Following a Two-Hour Workshop. *Teaching and Learning in Medicine*, 19(4), 336-342.

Hamberger, L. K. (2007). Preparing the next generation of physicians: Medical school and residency-based intimate partner violence curriculum and evaluation. *Trauma, Violence and Abuse*, 8(2), 214-225.

*Helton, S. M. & Evans, G. W. (2001). “She looked just like me”: A domestic violence learning module. *Issues in Mental Health Nursing*, 22(5), 503-516.

*Heron, S. L., Hassani, D. M., Houry, D., Quest, T. & Ander, D. S. (2010). Standardized patients to teach medical students about intimate partner violence. *Western Journal of Emergency Medicine*, 11(5), 500-505.

Hester, M., Williamson, E., Regan, L., Coulter, M., Chantler, K., Gangoli, G., Davenport, R. & Green, L. (2012). *Exploring the service and support needs of male, lesbian, gay, bi-sexual and transgendered and black and other minority ethnic victims of domestic and sexual violence*. Retrieved from <https://bit.ly/2Ln5q3C>, on 19 April, 2018.

Hinderliter, D., Doughty, A. S., Delaney, K., Pitula, C. R. & Campbell, J. (2003). The effect of intimate partner violence education on nurse practitioners' feelings of competence and ability to screen patients. *Journal of Nursing Education*, 42(10), 449-454.

Honey, P. & Mumford, A. (1986). *Using your learning styles*. Maidenhead: Honey.

*Jonassen, J. A., Pugnaire, M. P., Mazor, K., Regan, M., Jacobson, E., Gammon, W., Doepel, D. & Cohen, A. J. (1999). The effect of a domestic violence interclerkship on the knowledge, attitudes, and skills of third-year medical students. *Academic Medicine*, 74(7), 821-828.

*Kennedy, K. M., Vellinga, A., Bonner, N., Stewart, B. & McGrath, D. (2013). How teaching on the care of the victim of sexual violence alters undergraduate medical students' awareness of the key issues involved in patient care and their attitudes to such patients. *Journal of Forensic and Legal Medicine*, 20(6), 582-587.

Kolb, D. (1984). *Experiential learning: Experience as the source of learning and development*. Englewood Cliffs: Prentice-Hall.

Knapp, T. R. (2016). Why is the one-group pretest–posttest design still used? *Clinical Nursing Research*, 25(5), 467-472.

Kripke, E. N., Steele, G., O'Brien, M. K. & Novack, D. H. (1998). Domestic violence training program for residents. *Journal of General Internal Medicine*, 13(12), 839-841.

Langhinrichsen-Rohling, J. (2010). Controversies Involving Gender and Intimate Partner Violence in the United States. *Sex Roles*, 62, 179-193.

Leung, T. P., Phillips, L., Bryant, C. & Hegarty, K. (2018). How family doctors perceived their 'readiness' and 'preparedness' to identify and respond to intimate partner abuse: A qualitative study. *Family Practice*, 35(4), 517-523.

Loyens, S. M. M., Jones, S. H., Mikkers, J. & van Gog, T. (2015). Problem-based learning as a facilitator of conceptual change. *Learning and Instruction*, 38, 34-42.

Mansouri, M. & Lockyer, J. (2007). A meta-analysis of continuing medical education effectiveness. *Journal of Continuing Education in the Health Professions*, 27(1), 6-15.

Marinopoulos, S. S., Dorman, T., Ratanawongsa, N., Wilson, L. M., Ashar, B. H., Magaziner, J. L., Miller, R. G., Thomas, P. A., Prokopowicz, G. P., Qayyum, R. & Bass, E. B. (2007). Effectiveness of continuing medical education. *Evidence Report/Technology Assessment (Full Report)*, 149, 1-69.

*McAndrew, M., Pierre, G. C. & Kojanis, L. C. (2014). Effectiveness of an online tutorial on intimate partner violence for dental students: A pilot study. *Journal of Dental Education*, 78(8), 1176-1181.

McGill, I. & Beaty, L. (1995). *Action learning: A guide for professional, management and educational development*. 2nd ed. London: Kogan Page.

Meade, M. O. & Richardson, W. S. (1997). Selecting and appraising studies for a systematic review. *Annals of Internal Medicine*, 127(7), 531-537.

*Milone, J. M., Burg, M. A., Duerson, M. C., Hagen, M. G. & Pauly, R. R. (2010). The effect of lecture and a standardized patient encounter on medical student rape myth acceptance and

attitudes toward screening patients for a history of sexual assault. *Teaching and Learning in Medicine*, 22(1), 37-44.

Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G. & The PRISMA Group (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine*, 6(7), e1000097.

Natan, M. B., Khater, M., Ighbariyea, R. & Herbet, H. (2016). Readiness of nursing students to screen women for domestic violence. *Nurse Education Today*, 44, 98-102.

Nelson, J., Cook, P. & Ingram, J. (2013). Utility of the theory of planned behavior to predict nursing staff blood pressure monitoring behaviors. *Journal of Clinical Nursing*, 23, 461-470.

O'Brien, M. A., Freemantle, N., Oxman, A. D., Wolf, F., Davis, D. & Herrin, J. (2001). Continuing education meetings and workshops: Effects on professional practice and health care outcomes. *Cochrane Database of Systematic Reviews*, Issue 1. CD003030.

O'Doherty, L., Taket, A., Valpied, J. & Hegarty, K. (2016). Receiving care for intimate partner violence in primary care: Barriers and enablers for women participating in the weave randomised controlled trial. *Social Science & Medicine*, 160, 35-42.

*Pomeroy, E., Parrish, D. E., Bost, J., Cowlagi, G., Cook, P. & Stepura, K. (2011). Educating students about interpersonal violence: Comparing two methods. *Journal of Social Work Education*, 47(3), 525-544.

Sawyer, S., Coles, J., Williams, A. & Williams, B. (2016). A systematic review of intimate partner violence educational interventions delivered to allied health care practitioners. *Medical Education*, 50(11), 1107-1121.

Schmitt, N. (2010). *Researching vocabulary: A vocabulary research manual*. Basingstoke: Palgrave Macmillan.

Sivagnanam, G., Bairy, K. L. & D'Souza, U. (2005). Attitude towards rape: A comparative study among prospective physicians of Malaysia. *Medical Journal of Malaysia*, 60, 286-293.

*Smith, R., Wight, R. & Homer, C. S. E. (2017). 'Asking the hard questions': Improving midwifery students' confidence with domestic violence screening in pregnancy. *Midwifery Education in Practice*, 28, 27-33.

Sprague, S., Madden, K., Simunovic, N., Godin, K., Pham, N. K. & Bhandari, M. (2012). Barriers to screening for intimate partner violence. *Women & Health*, 52(6), 587-605.

Tavrow, P., Bloom, B. E. & Withers, M. H. (2017). Intimate partner violence screening practices in California after passage of the Affordable Care Act. *Violence Against Women*, 23(7), 871-886.

Taylor, J., Bradbury-Jones, C., Kroll, T. & Duncan, F. (2013). Health professionals' beliefs about domestic abuse and the issue of disclosure: A critical incident technique study. *Health and Social Care in the Community*, 21(5), 489-499.

Tranter, S., Irvine, F. & Collins, E. (2012). Innovations aimed at improving the physical health of the seriously mentally ill: An integrative review. *Journal of Clinical Nursing*, 21, 1199-1214.

United Nations Children's Fund (UNICEF) (2013). *Female genital mutilation/cutting: A statistical overview and exploration of the dynamics of change*. Retrieved from https://www.unicef.org/publications/index_69875.html, on 7 February, 2018.

United Nations Statistics Division (2015). Violence against women. In *The world's women 2015: Trends and statistics* [online], 139-161. Retrieved from <https://unstats.un.org/unsd/gender/chapter6/chapter6.html>, on 7 February, 2018.

Valpied, J., Aprico, K., Clewett, J. & Hegarty, K. (2017). Are future doctors taught to respond to intimate partner violence? A study of Australian medical schools. *Journal of Interpersonal Violence*, 32(16), 2419-2432.

Whittemore, R. & Knafl, K. (2005). The integrative review: Updated methodology. *Journal of Advanced Nursing*, 52(5), 546-554.

World Health Organization (2013). *Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence*. Retrieved from <https://bit.ly/2PAijL9>, on 20 November, 2017.

POPULATION	Pre-qualifying healthcare students (nurses, midwives, doctors, dentists, physiotherapists, occupational therapists, radiographers, dieticians, podiatrists, paramedics, orthoptists, prosthetists/orthotists, social workers, speech and language therapists, pharmacists, and any other allied health professionals).
INTERVENTION	Any female-focused GBV education intervention or training program for identified population (including intimate partner violence or abuse, domestic violence or abuse, spouse violence or abuse, battered women, family violence or abuse, rape, human trafficking, sexual violence or abuse, female genital mutilation/cutting or female circumcision, forced marriage, honor crimes and honor killings).
COMPARISON	Any or none.
OUTCOMES	Self-reported or objectively measured improvement relating to the subject of GBV (including knowledge, attitudes, beliefs and confidence in practice).

Table 1: PICO Criteria

POPULATION	PHENOMENON OF INTEREST	INTERVENTION
Medic* student*	Speech and language therap*	Gender-based violence
Nurs* student*	student*	Gender violence
Dentist* student*	SALT student*	Gender abuse
Dental student*	Speech therap*	Intimate partner violence
Pharmac* student*	student*	Domestic violence
Midwifery student*	Paramedic*	Spous* abuse
Student midwives	student*	Battered women
Occupational therap*	Orthopti* student*	Family violence
student*	Prostheti* student*	Family abuse
Physiotherap*	Orthoti* student*	Rape
student*	Social work*	Human trafficking
student*	student*	Sexual abuse
Radiograph* student*	Podiatr* student*	Sexual violence
Dietetic* student*	Allied health student*	Female circumcision
Dietitian* student*	Health care student*	Female genital mutilation
	Healthcare student*	Female genital cutting
		FGM
		Forced marriage*
		Honor killing*
		Honor crime*

Table 2: Summary of Search Terms

Table 3: Summary of Included Studies

Author (year) and country	Aim	Study design and data collection	Sample	Educational intervention	Measured outcomes	Summary of findings
Buranosky <i>et al.</i> (2012) USA	To assess how various types and amounts of IPV education for medical students affected knowledge and attitudes.	Correlational study design. Survey instrument consisting of demographic questions plus 25 items about education, knowledge and attitudes concerning IPV.	279 medical students from first- to fourth-year of study. Gender ratio: 45% male, 52% female.	Various teaching forms: core didactic training delivered in lectures (first year) and small group sessions during clerkships (third year). Additional elective experiential programs included 40-hour training in local emergency department, and volunteer work at a local IPV shelter clinic.	<ul style="list-style-type: none"> • IPV-related knowledge (objectively measured). • IPV-related awareness (self-reported). • Comfort in screening for IPV (self-reported). • Support of universal screening for IPV (self-reported). • Attitude regarding importance of interventions for IPV (self-reported). 	<ul style="list-style-type: none"> • Higher knowledge scores were obtained by female students and those with a history of IPV. • Knowledge scores increased with each additional year of study. • Knowledge scores increased with participation in increased number of core or experiential activities. • Participation in experiential activities was more significant a factor in improving students' self-reported attitudes than was participation in core (didactic) activities.
Cerulli <i>et al.</i> (2015) USA	To evaluate an IPV didactic session adapted for pharmacy students and describe student quantitative and qualitative feedback on the session.	Mixed methods; posttest-only design, with quantitative and qualitative data measurements. Questionnaire used to collect data.	237 pharmacy students, year of study not specified. Gender ratio: Not specified.	1.5-hour evidence-based IPV lecture.	<ul style="list-style-type: none"> • Students' opinions and feedback regarding the educational intervention were collected. 	<ul style="list-style-type: none"> • Students expressed the belief that their ability to recognize IPV had improved as a result of the lecture. • In qualitative feedback, students advised that course content should relate more specifically to pharmacy. • Students felt that 1.5 hours was not enough time to cover the material thoroughly, and indicated a wish for more interactive content (e.g. role-play with mock patients).
Danley <i>et al.</i> (2004) USA	To evaluate the impact of a brief, interactive multimedia tutorial designed to prepare dentists to recognize and respond to DV.	RCT using three study groups in a modified Solomon four-group design: a pretest and posttest experimental group; a posttest only group; and a two-test control group. Data was collected through Likert-style questions via computer.	174 subjects, of which 161 were dental students† (the remaining 13 were faculty members). Students from second- to fourth-year of study were recruited. Gender ratio: 53% male, 47% female.	One-time interactive multimedia tutorial featuring actors, where subjects are asked to interact with a virtual patient. Participation required between 15 and 25 minutes.	<ul style="list-style-type: none"> • Knowledge and attitudes about DV were assessed on the basis of four main criteria (Asking, Validating, Documenting, Referring; AVDR) (self-reported). • Beliefs about own knowledge of DV-related concepts (self-reported). • Attitudes about other related aspects of DV (self-reported). 	<ul style="list-style-type: none"> • After the tutorial, both experimental groups demonstrated significant improvements in knowledge and attitudes in all four AVDR criteria as compared with the control group. • The two experimental groups did not differ significantly from each other, suggesting that improvements could not be attributed to a testing effect. • Regarding attitudes about 'other related aspects of DV', only 2 out of 8 posttest scores for both intervention groups were significantly different from control group scores.

1	Edwardsen,	To determine if	Prospective	43 first-year	1-hour didactic IPV	• Perceived usefulness of	• The intervention group perceived the
2	Morse and	a teaching	randomized trial with	medical students	presentation, followed by	the mnemonic (self-	mnemonic to be more useful than did
3	Frankel (2006)	module with a	intervention and	(43% of cohort).	1-hour faculty-facilitated	reported).	the control group.
4	USA	mnemonic	control group. A	Gender ratio: Not	small-group discussion	• Ability to discern a	• The intervention group consistently
5		improves IPV	video-taped	specified.	featuring SPs for practice	history of abuse during	outperformed the control group during
6		interviewing	competency-based		of interviewing skills.	interaction with SP	evaluation. However, of the seven
7		skills among	evaluation using SPs		Intervention group	(objectively measured).	criteria assessed, only two offered a
8		medical	was used to assess		attended additional		statistically significant group difference.
9		students.	the intervention		training session prior to		
10			results. A posttest		small-group discussion		
11			questionnaire was		which involved		
12			also given to assess		structured discussion on		
13			perceived usefulness		use of the mnemonic.		
14	Elman <i>et al.</i>	To examine the	Quasi-experimental	110 third-year	Interactive 2-hour	• Frequency with which	• There was a statistically significant
15	(2004)	effect of	study design with	medical students	seminar, including	students inquired about	increase in frequency of inquiring about
16	Canada	unannounced	intervention and	(61% of cohort).	opportunity to practice	family violence during	family violence by students in the
17		SPs in the	control group.	Gender ratio: Not	management of a SP.	end-of-rotation OSCE	intervention group.
18		clinical setting	Intervention	specified.	The intervention group	(objectively measured).	• The improvement was more pronounced
19		as a teaching	outcomes were		was assigned to receive a		in the group of students interacting with
20		strategy for	assessed during end-		similar interaction during		a prenatal SP case, as compared with the
21		medical students	of-rotation OSCE.		clinical rotation; they		musculoskeletal SP case group, possibly
22		on the subject of			were unaware the		due to the more obvious transfer context
23		family violence.			‘patient’ was an actor		of the former.
24					until after the encounter.		
25	Ernst <i>et al.</i>	To determine	One-group pretest-	141 first-year	Series of lectures (3	• Actual knowledge	• Students’ knowledge was significantly
26	(1998)	knowledge	posttest study design.	medical students	hours in total) about DV	(objectively measured).	increased for specific facts (3 out of 14)
27	USA	about DV, the	Confidential and	participated in	including demonstration	• Actual personal history of	following the intervention.
28		effectiveness of	anonymous written	both pre- and	of mock patient. Lectures	DV (objectively	• Statistically significant differences in
29		formal	survey used to collect	posttest.	covered historical	measured by ISA).	responses to specific questions (3 out of
30		instruction	data; this was	Gender ratio: 49%	perspective of DV,	• Perceived personal	14 in posttest) were observed between
31		about DV, and	completed once	male, 50% female	current laws, risk factors,	history of DV (self-	male and female students, with females
32		the prevalence	before and one month	(NB. One student	victimization processes,	reported).	outperforming males in all 3 questions.
33		of DV in a first-	after the intervention.	not accounted	screening tools		• After the intervention, more students
34		year medical	The Index of Spouse	for).	(including ISA), and		identified themselves as having
35		school class.	Abuse (ISA) was		barriers and facilitators		experienced DV. However, there were
36			administered once		to identification in		discrepancies between the students’
37			only (pretest) to		practice.		actual and perceived levels of personal
38			determine baseline				experience of DV.
39			levels of abuse				
40			among the group.				

Ernst <i>et al.</i> (2000) USA	To evaluate the long-term effectiveness (LTE) and long-term retention (LTR) of formal instruction about DV.	Longitudinal one-group pretest-posttest study design. General knowledge survey on DV used to collect data; this was given before, one month after, and 2 years after the formal instruction.	104 medical students (70% of those who received the intervention participated in this 2-year follow-up). The students were first-year at the time of the intervention. Gender ratio: 51% male, 49% female.	A series of lectures about DV, including a demonstration with a mock patient being seen for DV-related injuries. This educational session was approximately 3 hours in length.	<ul style="list-style-type: none"> • Information learned and retained (LTE and LTR) (objectively measured). • Information learned but forgotten (neither LTE nor LTR) (objectively measured). • Information already part of the students' knowledge base (objectively measured). 	<ul style="list-style-type: none"> • Responses to the general knowledge survey on DV largely showed improvement between pretest and 3-year follow-up. • Between the first and second survey (administered 2 months following formal instruction)[‡], there was more significant improvement in DV-related knowledge than between the first and third survey, suggesting reduced LTE and LTR over time.
Everett <i>et al.</i> (2012) USA	To assess first-year dental students' awareness and beliefs regarding IPV before and after a one-hour educational seminar.	One-group pretest-posttest study design. Voluntary survey pre- and post-intervention; eight-item questionnaire.	65 first-year dental students. Gender ratio: 70.8% male, 29.2% female.	One-hour educational seminar facilitated by an experienced IPV outreach coordinator, involving a presentation, supplemental resources and contact information, and question-and-answer session.	<ul style="list-style-type: none"> • Awareness of IPV as a health or dental profession issue and awareness of IPV resources (self-reported). • Professional beliefs regarding the dental profession and IPV issues (self-reported). • Personal beliefs regarding IPV education and intervention (self-reported). 	<ul style="list-style-type: none"> • Respondents' awareness of IPV as a health or dental profession issue and awareness of IPV resources increased significantly following the educational session. • Professional beliefs were largely improved following the educational session. • Willingness to participate in further IPV education and self-reported comfort to participate in IPV intervention both increased significantly following the educational session.
Haase <i>et al.</i> (1999) USA	To investigate whether an elective DV course delivered to medical students resulted in improved attitudes and abilities during clinical years.	Posttest-only study design. Data collected through 11-item survey instrument which was administered 2 years after the DV course.	115 medical students. Of these, one fourth reported having participated in the elective DV course. Students were first-year at the time of the intervention. Gender ratio: 56% male, 44% female.	Nine hours of didactic sessions over a six-week period. These sessions covered strategies for identification of victims, intervention methods, and reading of landmark DV literature. Participation in a class project was also required to complete the course (no further information given).	<ul style="list-style-type: none"> • Perceived preparedness to recognize and address DV (self-reported). • Comfort (self-reported). • Frequency of screening for DV during clinical rotations (self-reported). • History of successfully identifying DV during clinical rotations (self-reported). • Awareness of local DV advocacy program (objectively measured). 	<ul style="list-style-type: none"> • Women were disproportionately interested in undertaking the elective DV course. However, there was not a significant gender effect on students' scoring (measured by a Domestic Violence Sensitivity Scale; DVSS). • The DV-educated group scored significantly higher on the DVSS. • DV-educated students reported higher levels of comfort, frequency of screening and history of successfully identifying DV during clinical rotations.

1	Helton and	To examine	Qualitative	The journals of 18	22-hour mandatory	• No measured outcomes	• Five common themes emerged from the
2	Evans (2001)	senior	phenomenological	baccalaureate	course involving	due to qualitative	students' journals: (1) clinical
3	USA	baccalaureate	study design. Data	nursing students	orientation, pretesting	phenomenological study	encounters were frightening and
4		nursing	collected from	(out of 60 in total)	and posttesting, review	design.	emotionally difficult to attend; (2)
5		students'	participants' journals	were randomly	of didactic material,		surprise at the incidence of violence; (3)
6		experiences of	(voluntary and	selected for	learning module		identifying with the victims; (4)
7		completing an	anonymous); this was	analysis.	notebook and video, 8-		dispelling of stereotypes about
8		experiential DV	the only written		hour court experience,		perpetrators and victims; and (5)
9		learning module	requirement for the	Gender ratio: 10%	three 60-90-minute group		consideration of the moral issues of
10		(DVLM).	DVLM.	male, 90% female	therapy sessions (one		'good' and 'bad' in relation to nursing
11				(total cohort).	with shelter victims and		practice.
12					two with perpetrators in		
13					court-ordered group),		
14					and three-hour debriefing		
15	Heron <i>et al.</i>	To implement	Prospective	41 fourth-year	One-hour interactive	• Competency in IPV (self-	• There was no significant improvement
16	(2010)	and assess an	observational study	medical students.	didactic training in small	reported and objectively	in students' self-reported comfort levels
17	USA	IPV OCSE	design. 360-degree		groups. Additionally, the	measured).	between pre- and posttest.
18		module using	evaluation form was	Gender ratio: 49%	OSCE featuring SPs was	• Comfort (self-reported	• Correlation between self-reported
19		SPs, with the	used to assess	female, 51% male	conducted prior to the	and objectively	comfort level and OSCE-assessed
20		goal of	competency and	(NB. Ratio	posttest, and although it	measured).	competencies was positive.
21		improving	comfort levels during	reflects the overall	served as a means of	• Communication,	
22		students'	OSCE. Pre- and	cohort, whose size	assessment, it could be	professionalism and	
23		competency,	posttests were also	was not	considered as an	patient care (objectively	
24		self-efficacy,	used to assess	specified).	interactive experience	measured).	
25		communication	knowledge, attitudes,		and/or intervention in its		
26		skills and	beliefs and comfort		own right, as it would		
27		professionalism.	levels surrounding		likely have impacted the		
28			IPV, but these were		students' posttest		
29			not graded, and no		responses.		
30			basic data are				
31	Jonassen <i>et al.</i>	To determine	Two-group pretest-	67 medical	First cohort: 3.5-day	• Knowledge, attitudes and	• Students' knowledge, attitudes and
32	(1999)	whether	posttest study design.	students from first	interclerkship on	skills pertaining to DV	skills were all significantly improved
33	USA	participation in	Posttest was	cohort and 77	domestic violence,	(objectively measured	following DVI participation (both
34		a domestic	implemented once	students from	including a series of	through written	cohorts).
35		violence	immediately after	second cohort	didactic and interactive	assessment).	• Students from the first cohort
36		interclerkship	intervention and	participated in the	content (lectures, films,	• Feedback regarding	maintained improvements in all three
37		(DVI) improved	again at 6 months. A	pretest, immediate	panels, standardized	course format, content	domains (knowledge, attitudes and
38		the knowledge,	third cohort which	posttest and 6-	patient interviews, role	and effectiveness (self-	skills) at 6-month follow-up, whereas
39		attitudes and	had not received the	month posttest	plays and small-group	reported).	students from the second cohort
40		skills of two	DVI was used as a	(written	discussions). Students		maintained their improved skills only.
41		successive	control (certain	measurements			

	cohorts of medical students.	posttest criteria only). Data was collected via written assessments, Likert scale feedback, questionnaires regarding clinical experiences, and OSCE assessments.	only). The OSCE assessment scores of both full cohorts (98 and 107 students respectively) were compared with a third cohort of 93 students.	also interacted with real DV survivors. Second cohort: Same intervention as first cohort but lasting 2 days.	<ul style="list-style-type: none"> • Reports of students' clinical experiences with DV (self-reported). • Preparedness for DV-based OSCE (self-reported). • Performance in addressing DV (objectively measured though OSCE). 	<ul style="list-style-type: none"> • Students who participated in the shorter DVI rated the course more favorably. • Both intervention cohorts performed significantly better in their DV-based OSCE assessments than did the third (control) cohort. All three cohorts performed equally in non-DV-based OSCE stations. • Students from both intervention cohorts reported a significantly higher sense of preparedness for the DV-based OSCE than the control cohort.
Kennedy <i>et al.</i> (2013) <i>Ireland</i>	To evaluate the effectiveness of a teaching session on the care of the victim of sexual violence for undergraduate medical students.	Quasi-experimental study design. Pre- and post-intervention survey using Likert scale.	88 third-year medical students. Gender ratio: 44% male, 56% female (NB. Ratio reflects the 105 students who originally consented to take part in the study).	Two-hour interactive lecture including a training DVD to help maintain student engagement.	<ul style="list-style-type: none"> • Awareness of key issues relating to care of patients who have experienced sexual violence (self-reported). • Insight into common rape myths (self-reported). 	<ul style="list-style-type: none"> • Students improved their knowledge of patient care and insight into common rape myths following the educational intervention.
McAndrew, Pierre and Kojanis (2014) <i>USA</i>	To determine the effectiveness of an online DV tutorial for changing dental students' knowledge, attitudes, beliefs and behaviors.	Quasi-experimental pretest-posttest study design with control and intervention group. Pre- and post-intervention survey (Physician Readiness to Manage Intimate Partner Violence Survey; PREMIS) used to collect data.	25 forth-year dental students. Gender ratio: Not specified.	One-hour web-based tutorial divided into 10 modules. The intervention group took the survey before and after the online training, while the control group took the same survey twice <i>before</i> completing the online training.	<ul style="list-style-type: none"> • Perceived preparation to manage IPV (self-reported). • Perceived knowledge (self-reported). • Actual knowledge (objectively measured). • Opinion scales to assess perceived readiness to manage IPV based on 8 criteria (self-reported). 	<ul style="list-style-type: none"> • The intervention group's composite scale scores were significantly increased post-intervention for perceived preparation, perceived knowledge and actual knowledge. • There was a significant improvement in 2 of the 8 opinion scale scores for the intervention group; the remaining 6 were improved but not significant. • The intervention group's posttest scores were greater than those of the control group, but not significant.
Milone <i>et al.</i> (2009) <i>USA</i>	To determine if medical students' attitudes toward rape and sexual assault can be changed through	One-group pretest-posttest design. Anonymous 21-item questionnaire completed before interventions, again after first intervention, and	102 second-year medical students. Gender ratio: 46% male, 54% female. (NB. Ratio reflects the 127 students in	Intervention 1: 45-minute lecture on the subject of sexual assault, including instructions on screening techniques and treatment of victims. Lecture attendance was voluntary.	<ul style="list-style-type: none"> • Rape myth acceptance (self-reported). • Attitudes toward screening (self-reported). 	<ul style="list-style-type: none"> • Significant differences in both rape myth acceptance and attitudes toward screening were observed between pretest and first posttest, but not between first and second posttest. • No significant difference was observed between the responses of students who had and had not attended the lecture.

	educational interventions.	again after second intervention.	the cohort, and not the 102 who completed the posttest).	Intervention 2: SP encounter involving history-taking and a physical exam.		<ul style="list-style-type: none">• Significant differences were observed between male and female responses (females were less accepting of rape myths and showed more positive attitudes toward screening).
Pomeroy <i>et al.</i> (2013) <i>USA</i>	To compare the effects of two educational methods on social work students' knowledge and attitudes towards interpersonal violence among college students.	Mixed methods; three-group pretest-posttest study design with qualitative measurement procedures. Data was collected through focus groups.	63 social work students, year of study not specified (48 participated in the pre-intervention focus groups, and 40 in the post-intervention focus groups). Gender ratio: 19% male, 81% female.	Three groups were compared (two intervention and one control); the interventions were peer theatre (interactive; 90-minute presentation) and peer education (didactic; 90-minute lecture). The comparison group watched a family orientated video that contained no violence (90 minutes in length).	<ul style="list-style-type: none">• Focus group questions were centered on beliefs about relationship violence, sexual assault and stalking.	<ul style="list-style-type: none">• Themes from pre-intervention focus groups: (1) relationship violence: naïve awareness; (2) sexual assault: concerned but uninformed; (3) stalking: serious versus familiar, funny and flattering.• Post-intervention, the comparison group statements remained similar to pre-intervention; the peer education group results showed increased knowledge acquisition; and the peer theatre group results suggested integration of knowledge, awareness and practical application.
Smith, Wight and Homer (2017) <i>Australia</i>	To evaluate the effectiveness of an educational intervention to increase student midwives' confidence to screen for and respond to DV disclosure during pregnancy.	Mixed methods; one-group pretest-posttest design. Three surveys used to collect data: initial online knowledge survey, pre-workshop confidence survey (immediately prior) and post-workshop confidence survey (immediately after). Quantitative and qualitative questions were included.	72 midwifery students completed the pretest survey and 102 completed the posttest survey. 40% of the sample were in their third-year of study§. Gender ratio: Not specified.	Interdisciplinary and interactive one-day workshop covering theory and practice.	<ul style="list-style-type: none">• Confidence levels in 8 DV-related topic areas (self-reported).• Open-ended survey questions, pre- and post-workshop, asked the students to identify activities which may increase their confidence to screen for and respond to disclosures of DV.	<ul style="list-style-type: none">• A statistically significant increase in students' confidence levels was reported across all 8 topic areas following the workshop.• In the pretest survey, students suggested that activities centered on the <i>process</i> of addressing DV in practice would be most useful in helping to increase their confidence.• In the posttest survey, students reported that the most useful activities in the workshop were those focused on the <i>process</i> of addressing DV, consistent with their pretest views.• Both didactic and interactive activities were cited as being useful.

†Although only 93% of the sample in Danley *et al.*'s (2004) study were students, the paper was deemed to be useful in answering the research question.

‡2-month follow-up survey was part of a previous study by the authors.

§Approximately 60% of the students in Smith *et al.*'s (2017) study were registered nurses completing a 12-month preregistration midwifery program. Although this group was already registered in a healthcare profession, they were yet to qualify in the profession being studied by this research (midwifery), and so the paper was not excluded. It would not have been possible for this review to exclude all papers featuring already-registered health professionals seeking to qualify in a second health-related area, as oftentimes this is not an explored characteristic.

Table 4: Summary of Critical Appraisal (Quantitative and Mixed Methods)

	Aims clearly stated	Study design suitable to meet objectives	Outcomes	Quantitative and qualitative methods used appropriately	Teachers and learners blinded to study purpose	Assessors and examiners blinded to learning method	Evidence of sample size estimates	Ethical approval obtained if appropriate†	Data collection instrument (validity)	Data collection instrument (reliability)	Time elapsed between intervention and posttest‡	Data analysis	Presentation of results	Discussion	Summary score for methodological rigor§	Summary score for data relevance§
Buranosky <i>et al.</i>	G	F	G	G	n/a	n/a	P	n/a	F	P	n/a	G	G	G	2	3
Cerulli <i>et al.</i>	G	F	F	G	P	P	P	n/a	P	P	n/a	F	G	G	2	2
Danley <i>et al.</i>	G	G	G	G	P	P	P	G	F	P	P	G	G	G	2	3
Edwardsen <i>et al.</i>	G	G	G	G	G	G	P	G	G	G	P	G	G	G	3	3
Elman <i>et al.</i>	G	F	P	G	G	P	P	G	P	P	G	G	G	G	2	1
Ernst <i>et al.</i> (1998)	G	F	F	G	U	P	P	G	F	P	G	G	G	G	2	2
Ernst <i>et al.</i> (2000)	G	F	F	G	P	P	P	G	P	P	G	G	G	G	2	2
Everett <i>et al.</i>	G	F	G	G	P	P	P	n/a	P	P	P	G	G	G	2	3
Haase <i>et al.</i>	G	F	P	G	P	P	P	n/a	P	F	G	F	F	F	2	1
Heron <i>et al.</i>	F	U	U	G	U	U	P	G	F	P	G	U	F	G	1	1
Jonassen <i>et al.</i>	G	G	G	G	P	P	P	U	F	F	G	G	G	F	2	3
Kennedy <i>et al.</i>	G	F	G	G	P	P	P	G	G	G	P	G	G	G	2	2
McAndrew <i>et al.</i>	G	G	G	G	P	P	P	G	G	G	P	G	G	G	2	3
Milone <i>et al.</i>	G	F	G	G	P	P	G	G	P	G	P	F	G	G	2	3
Pomeroy <i>et al.</i>	G	G	G	G	G	U	P	G	n/a	n/a	G	G	G	G	3	3
Smith <i>et al.</i>	G	F	G	G	P	P	P	G	G	P	P	G	G	G	2	3

G = Good; F = Fair; P = Poor; U = Unclear; n/a = Not applicable

†Some studies were given exempt status by the university in which the research took place.

‡Not applicable given to studies where time elapsed between intervention and posttest was not relevant (e.g. if opinions and feedback were the only assessed outcomes).

§As adapted by Tranter, Irvine and Collins (2011) after Whitemore and Knafl (2005).

Table 5: Summary of Critical Appraisal (Qualitative)

	Clear statement of research aims	Appropriateness of qualitative methodology	Research design appropriate to address aims	Appropriateness of recruitment strategy	Data collection	Researcher-participant relationship considered	Consideration of ethical issues	Data analysis sufficiently rigorous	Clear statement of findings	Value of the research	Summary score for methodological rigor†	Summary score for data relevance†
Helton and Evans	G	G	G	F	G	P	G	G	G	F	2	3

G = Good; F = Fair; P = Poor; U = Unclear; n/a = Not applicable
†As adapted by Tranter, Irvine and Collins (2011) after Whittemore and Knafl (2005).

Table 6: Critical Findings

Critical Findings
<ol style="list-style-type: none">1. Interactive approaches to learning yield better results than didactic approaches.2. Students prefer their learning to be practice-focused rather than strictly theoretical.3. Courses of longer duration seem to be more effective in altering students' attitudes and opinions.4. Gendered differences might be significant, although more research is needed before conclusions can be drawn. Females tended to outperform males in assessments.

Table 7: Implications for Policy, Practice and Research

Implications for Policy, Practice and Research
<ol style="list-style-type: none">1. GBV learning opportunities should have a practical focus and should aim to incorporate an interactive element for improved results.2. More research is needed on the subject of single- versus mixed-gender audiences in GBV education for pre-qualifying healthcare students.3. Existing and future education programs should give greater attention to the wider forms of GBV (including female genital mutilation/cutting, forced marriage, honor violence and human trafficking). Future research may also strive to incorporate a focus on wider learning theory and consider its application in the development of GBV curricula.

Figure 1: PRISMA Flow Chart

Figure 2: Visual Summary of Themes