The impact of cannabinoid receptor 2 deficiency on neutrophil recruitment and inflammation

Hussain, Mohammed Tayab; Greaves, David R.; Iqbal, Asif Jilani

DOI:

[10.1089/dna.2019.5024](https://doi.org/10.1089/dna.2019.5024)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Hussain, MT, Greaves, DR & Iqbal, AJ 2019, 'The impact of cannabinoid receptor 2 deficiency on neutrophil recruitment and inflammation', *DNA and Cell Biology*, vol. 38, no. 10, pp. 1025-1029.
<https://doi.org/10.1089/dna.2019.5024>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Final publication is available from Mary Ann Liebert, Inc., publishers <http://dx.doi.org/10.1089/dna.2019.5024>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The impact of cannabinoid receptor 2 (CB₂) deficiency on neutrophil recruitment and inflammation

M.T. Hussain¹, D.R. Greaves² and A.J. Iqbal¹.

¹Institute of Cardiovascular Sciences (ICVS), College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom.

² Sir William Dunn School of Pathology, University of Oxford, Oxford, United Kingdom.

Correspondence:

Asif J Iqbal, Institute of Cardiovascular Sciences, College of Medical and Dental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT.

Phone: +441214144043

Email: a.j.iqbal@bham.ac.uk

Keywords:

Neutrophils, inflammation, cannabinoids, cannabinoid receptors

Invited for DNACB Bit

Abstract

Neutrophil trafficking into damaged or infected tissues is essential for the initiation of inflammation, clearance of pathogens and damaged cells, and ultimately tissue repair. Neutrophil recruitment is highly dependent on the step-wise induction of adhesion molecules and pro-migratory chemokines and cytokines. A number of studies in animal models have shown the efficacy of CB₂ agonists in limiting inflammation in a range of pre-clinical models of inflammation, including colitis, atherosclerosis, multiple sclerosis and ischemia-reperfusion injury. Recent work in pre-clinical models of inflammation raises two questions; by what mechanisms do CB₂ agonists provide anti-inflammatory effects during acute inflammation and what challenges exist in the translation of CB₂ modulating therapeutics into the clinic.

Neutrophil recruitment in acute inflammation

Neutrophil migration into inflamed tissues is driven by soluble factors released by activated endothelial and tissue resident immune cells. As inflammation reduces vascular flow rates, neutrophils in the blood undergo margination, selectin-mediated weak adhesion and rolling (Zarbock and Ley, 2009). The process of neutrophil firm adhesion and migration is mediated by G-protein coupled receptors (GPCRs) and integrin ligation (Lefort and Ley, 2012). Interactions between lymphocyte function-associated antigen 1 (LFA-1), an $\alpha\beta$ -integrin, and its ligand intracellular adhesion molecule 1 (ICAM-1) facilitates neutrophil recruitment by inducing pro-migratory changes including; further integrin clustering, cytoskeletal changes, increases in intracellular calcium levels and weakening of endothelial cell: endothelial cell junctions (Ley et al., 2007).

Following migration through the endothelium, neutrophils will cross the pericyte sheath and vascular basement membrane into inflamed tissues (Nourshargh et al., 2010). Remodelling of the tissue is essential for neutrophil recruitment under physiological and pathophysiological conditions; enzymatic degradation of the basement membrane by matrix metalloproteinase-9 (MMP-9) being an example (Yabluchanskiy et al., 2013). Clinically, stroke patients have been found to have increased circulating levels of MMP-9, this is accompanied by remodelling of the blood brain barrier and neutrophil infiltration (Turner and Sharp, 2016). In murine models, MMP-9 deficient mice had been observed to have reduced counts of airway infiltrating neutrophils following influenza infection; decreased neutrophil recruitment was accompanied by increased viral titres in these mice (Bradley et al., 2012).

Recently, there has been increasing interest in the effect of neurotransmitters/neurochemicals on leukocyte migration. The cannabinoid system has received continuous attention, now more so than ever, against the backdrop of increased recreational and medicinal use of cannabis, cannabinoids and cannabinoid-like compounds (Kerage et al., 2019).

Cannabinoid receptor regulation of inflammation

Cannabinoid receptors 1 and 2 (CB₁ and CB₂) are class A rhodopsin-like GPCRs with endogenous lipid ligands. CB₁ is highly expressed throughout the central nervous system (CNS) accompanied by restricted expression of CB₂ in the brain (Laprairie et al., 2012; Chen et al., 2017). CB₁ expression has also been reported in a number of peripheral tissues, such as the liver, pancreas and skeletal muscle. CNS expression of CB₁ mediates the psychoactive effects of cannabinoids such as delta-9-tetrahydrocannabinol (Δ^9 -THC) (Mackie, 2007). CB₁ expression outside the CNS may play an important role in appetite regulation and tissue repair (Dodd et al., 2010; Cooper and Regnell, 2014; Krott et al., 2016). CB₂ is most widely expressed in cells of the immune system, including but not restricted to, B-cells, monocytes, neutrophils and in particular, cells derived from macrophages (Galiègue et al., 1995; Van Sickle et al., 2005; Ofek et al., 2006). CB₁ and CB₂ activation cause the downstream activation of a number of signalling pathways (Figure.1) (Mackie, 2008).

Pharmacological activation of cannabinoid receptors has been reported to improve the clinical outcomes of a number of chronic inflammatory conditions in mice. For example, by increasing the half-life of 2-arachidonoylglycerol (2-AG), an endogenous CB₁ and CB₂ ligand, Alhouayek *et al.* reported a marked reduction in colonic and

peripheral inflammation in mice subjected to chemically induced colitis. The authors demonstrated that increased circulating levels of 2-AG reduced weight loss, decreased pro-inflammatory cytokine levels, improved colonic histological endpoints and intestinal barrier integrity (Alhouayek et al., 2011). The synthetic CB₂ ligand JWH-133 displayed similar efficacy in attenuating chemically induced colitis in IL-10^{-/-} mice. Singh *et al.* have shown that JWH-133 treated colitis mice presented with reduced activated leukocytes infiltrating the intestinal lamina propria and mesenteric lymph nodes (Singh et al., 2011). Synthetic activation of CB₂ by using low oral doses of Δ⁹-THC in atherosclerotic mice has also been shown to be effective at ameliorating disease progression; immune cells recovered from Δ⁹-THC treated mice had diminished proliferative capacity and reduced interferon-gamma secretion. Increases in interferon-gamma secretion was of particular significance as it correlates to vascular plaque progression from stable to unstable state, thereby enhancing risk of mortality (Van der Wal and Becker, 1999; Steffens et al., 2005). More recently, Zhao *et al.* have shown that CB₂ activation with the synthetic ligand, WIN55212-2, also reduced plaque burden and immune cell infiltration in atherosclerotic mice. Interestingly, the authors demonstrated that this reduction in vascular disease was due to decreased adhesion molecule expression (Zhao et al., 2010). Alterations in the recruitment of immune cells, including neutrophils, T-cells and macrophages by CB₂ agonists has also been reported in a number of pre-clinical models of human inflammatory conditions, including, sepsis (Csóka et al., 2009; Sardinha et al., 2014), multiple sclerosis (Maresz et al., 2007; Wen et al., 2015) and ischemia-reperfusion injury (Di Filippo et al., 2004; Bátkai et al., 2012).

For the most part, the vast majority of pre-clinical studies focus entirely on the use of CB₂ deficient mice or CB₂ specific pharmaceuticals. Importantly, Wen *et al.*

demonstrated that the anti-inflammatory effects of the CB₂ agonist WWL70 in wild-type mice undergoing experimental autoimmune encephalomyelitis were lost in CB₂ deficient mice verifying the efficacy of CB₂ agonists in targeting immune cell recruitment (Wen et al., 2015).

Effects of CB₂ deficiency on neutrophil recruitment

Kapellos *et al.* have built further on previous pre-clinical studies by studying the phenotype of immune cell trafficking in CB₂ deficient animals. The majority of the pre-clinical studies on the effects of cannabinoid receptors on myeloid cell biology *in vitro* and *in vivo* have used pharmacological tools rather than global or cell type specific knockout mice. A clear example of the dangers of relying solely on cannabinoid receptor agonists and antagonists is provided by the studies of Taylor *et al.* who demonstrated that macrophage chemotaxis in response to well characterised CB₂ agonists occurred with equal potency in CB₂ deficient mice (Taylor et al., 2015).

Under steady state, mice with global deficiencies in CB₂ were shown to have elevated neutrophil counts in the bone marrow (Kapellos et al., 2017). The observed changes in neutrophil counts may result from endogenous CB₂ agonists contributing to either myelopoiesis and/or the egress of myeloid cells from the bone marrow under normal physiological conditions (McHugh et al., 2008; Pasquevich et al., 2015). Despite not observing any significant changes in neutrophil recruitment to the lung following the induction of endotoxemia, Kapellos and colleagues reported a greater retention of neutrophils in the spleen of CB₂ deficient mice 2 hours following LPS administration. Levels of pro-inflammatory cytokines and soluble factors were observed to be significantly elevated in the spleens of CB₂ deficient mice, namely, CXCL10, CCL3,

IL-6 and MMP-9. Additionally, MMP-9 was found to be significantly elevated in the serum of CB₂ deficient animals. The authors concluded that CB₂ deficiency delayed the host response to LPS by decreasing neutrophil migration into and within lymphoid organs, reducing immune cell cross-talk with neutrophils, and pro-longing host exposure to LPS (Kapellos et al., 2017).

In a subsequent study using the dorsal air pouch model of inflammation, Kapellos and colleagues observed enhanced neutrophil recruitment in CB₂^{-/-} air pouches following zymosan injection. Enhanced neutrophil recruitment was independent of any changes in circulating blood cell counts. Importantly, the authors confirmed the effect of CB₂ deficiency was directly affecting neutrophils as bone marrow chimera experiments demonstrated that CB₂ deficiency on haematopoietic cells, but not stromal cells, led to increased immune cell recruitment in response to inflammation. Following zymosan administration, CB₂ deficient neutrophils overexpressed pro-migratory genes and their air pouch exudates were observed to have increased levels of pro-inflammatory cytokines and factors, like MMP-9, CCL22 and CXCL10. Interestingly, purified CB₂ deficient neutrophils were reported to have greater adherence to immobilised ICAM. In the same study the authors explored the effects of pharmacological modulation of CB₂ signalling on human neutrophils. Human neutrophils treated with the CB₂ agonist, JWH133, were shown to have reduced adhesion and transmigration across cytokine stimulated endothelial cell monolayers (Kapellos et al., 2019).

While CB₂ is largely seen as a GPCR that mediates anti-inflammatory effects, there are some studies which propose a pro-inflammatory role for CB₂ agonists both *in vivo* and *in vitro* (Turcotte et al., 2015). A potential explanation for these differences in various pre-clinical models of inflammation are variations in endocannabinoid levels

between different tissue microenvironments in combination with which cell types are responsible for driving immune cell recruitment and the progression of inflammation.

Outlook and future perspectives

Currently, the recreational and medicinal use of cannabinoids, whether in the form of cannabis, synthetic cannabinoids or cannabis extracts such as cannabidiol (CBD), THC or cannabis oils, is at an all-time high with ever increasing societal acceptance and changes in local and national legislation. Cannabis oils and cannabis itself, require special attention as the both contain a mixture of CBD, THC or Δ^9 -tetrahydrocannabivarin (Δ^9 -THCV), all three are able to interact with both CB₁ and CB₂. Irrespective of the ability of cannabinoids to modulate neutrophil recruitment via interactions with CB₂, there are some important considerations before translating basic science findings to the clinic.

A challenge in producing compounds to modulate CB₁ or CB₂ signalling in vivo is the specificity of any one compound for either receptor. For example, Gp-1a was reported as a selective agonist for CB₂ but Soethoudt *et al.* demonstrated its promiscuity in being able to bind both cannabinoid receptors. With regards to Δ^9 -THC and 2-AG, they have been reported to have a >100-fold binding bias for CB₂, however, interactions with CB₁ were still observed (Mussinu et al., 2003; Soethoudt et al., 2017). In addition to issues with specificity, CBD, a CB₁ and CB₂ ligand, behaves very differently dependent on its concentration. It has been reported that CBD behaves as an antagonist of CB₁ at μ M concentrations but an inverse agonist at nM concentrations (Thomas et al., 2007). Current data also indicates that cannabinoids may interact at sites other than the orthosteric sites on cannabinoid receptors (Pertwee, 2008). What

impact concentration and off target effects can have on CB₂ interactions across a range of different CB₂ ligands still requires further investigation. However, the use of pre-clinical models lacking global expression of CB₂ is one way to better understand the potential immune modulating effects of CB₂ activation in humans without the biochemical limitations currently associated with the use of pharmacological tool compounds alone for research.

Aside from biochemical considerations, genetic studies in mice have highlighted a potential pitfall of CB₂ agonists as therapeutics. CB₂ deficiency seemingly prolonged host exposure to LPS, decreased viral clearance and broke down immune cell cross talk, such as neutrophil migration to the marginal zone of the spleen for involvement in antigen presentation to T-cells (Kesteman et al., 2008; Bradley et al., 2012; Kapellos et al., 2017). CB₂ deficiency also reduced levels of the factor MMP-9, similar to the effects seen in animals treated with the CB₂ antagonist, SR144528 (Netherland et al., 2010). In line with this, CB₂ agonists have been reported to inhibit MMP-9 release in pre-clinical models of inflammation (Netherland et al., 2010; Kapellos et al., 2019). This raises questions, as with all anti-inflammatory therapeutics; will CB₂ agonists leave individuals more susceptible to infections?

With regards to the clinical efficacy of a CB₂ agonist, lessons can be drawn from the CB₁ antagonist, rimonabant. Rimonabant was developed as a first in class drug intended for treating obesity by aiding weight-loss. After gaining FDA approval in 2006, the drug was withdrawn worldwide after two years. Detrimentally, rimonabant induced psychiatric alterations inducing depression, anxiety and suicidal ideation (Leite et al., 2009). With the restricted expression of CB₂ in the brain, it remains unclear how a CB₂ specific agonist would impact the psychiatric health of recipients (Rogers, 2015).

Despite this potential issue a number of CB₂ agonists, such as GW842166X and LY2828360, have been taken to clinical trials in the context of pain management (Dhopeshwarkar and Mackie, 2014). Interestingly, these compounds were found not to provide meaningful analgesia despite the efficacy observed in pre-clinical models of pain management (Ostenfeld et al., 2011;Pereira et al., 2013). However, The CB₂ agonist, lenabasum, successfully passed phase 2 trials in the context of dermatomyositis where the compound was observed to provide greater improvement in a dermatological damage index after four weeks of administration (Werth et al., 2018). Lenabasum is currently being used in a phase 3 clinical trial for patients with dermatomyositis with a completion date of September 2021 (Werth et al., 2019). For now, the administration of candidate CB₂ agonists during clinical trials has been short term in nature, whether long-term use of CB₂ agonists recreates the psychiatric side-effects remains to be seen. Overall, whether a CB₂ specific agonist will fare any better than rimonabant, or if the efficacy observed in animals is comparable to that in humans, is something that requires greater investigation in acute inflammation in humans.

Various challenges still remain when considering cannabinoids as efficacious and specific therapeutics but the use of pre-clinical disease models in CB₂ deficient animals highlight that CB₂ agonists merit serious consideration as new class of anti-inflammatory drug candidates.

Acknowledgements

This work was supported by funding from the British Heart Foundation (BHF) grant (RG/15/10/31485), Academy of Medical Sciences (SBF003\1156) and Birmingham Fellowship.

Conflict of interest disclosures

The authors declare no conflicts of interest regarding the publication of this paper.

References

- Alhouayek, M., Lambert, D.M., Delzenne, N.M., Cani, P.D., and Muccioli, G.G. (2011). Increasing endogenous 2-arachidonoylglycerol levels counteracts colitis and related systemic inflammation. *The FASEB Journal* 25, 2711-2721.
- Bátkai, S., Mukhopadhyay, P., Horváth, B., Rajesh, M., Gao, R.Y., Mahadevan, A., Amere, M., Battista, N., Lichtman, A.H., Gauson, L.A., Maccarrone, M., Pertwee, R.G., and Pacher, P. (2012). Δ 8-Tetrahydrocannabivarin prevents hepatic ischaemia/reperfusion injury by decreasing oxidative stress and inflammatory responses through cannabinoid CB2 receptors. *British journal of pharmacology* 165, 2450-2461.
- Bradley, L.M., Douglass, M.F., Chatterjee, D., Akira, S., and Baaten, B.J. (2012). Matrix metalloprotease 9 mediates neutrophil migration into the airways in response to influenza virus-induced toll-like receptor signaling. *PLoS Pathog* 8, e1002641.
- Chen, D.-J., Gao, M., Gao, F.-F., Su, Q.-X., and Wu, J. (2017). Brain cannabinoid receptor 2: expression, function and modulation. *Acta pharmacologica Sinica* 38, 312-316.
- Cooper, M.E., and Regnell, S.E. (2014). The hepatic cannabinoid 1 receptor as a modulator of hepatic energy state and food intake. *British journal of clinical pharmacology* 77, 21-30.
- Csóka, B., Németh, Z.H., Mukhopadhyay, P., Spolarics, Z., Rajesh, M., Federici, S., Deitch, E.A., Bátкаи, S., Pacher, P., and Haskó, G. (2009). CB2 Cannabinoid Receptors Contribute to Bacterial Invasion and Mortality in Polymicrobial Sepsis. *PLOS ONE* 4, e6409.

- Dhopeshwarkar, A., and Mackie, K. (2014). CB2 Cannabinoid Receptors as a Therapeutic Target—What Does the Future Hold? *Molecular Pharmacology* 86, 430.
- Di Filippo, C., Rossi, F., Rossi, S., and D'amico, M. (2004). Cannabinoid CB2 receptor activation reduces mouse myocardial ischemia-reperfusion injury: involvement of cytokine/chemokines and PMN. *Journal of Leukocyte Biology* 75, 453-459.
- Dodd, G.T., Mancini, G., Lutz, B., and Luckman, S.M. (2010). The peptide hemopressin acts through CB1 cannabinoid receptors to reduce food intake in rats and mice. *J Neurosci* 30, 7369-7376.
- Galiègue, S., Mary, S., Marchand, J., Dussossoy, D., Carrière, D., Carayon, P., Bouaboula, M., Shire, D., Le Fur, G., and Casellas, P. (1995). Expression of Central and Peripheral Cannabinoid Receptors in Human Immune Tissues and Leukocyte Subpopulations. *European Journal of Biochemistry* 232, 54-61.
- Kapellos, T.S., Recio, C., Greaves, D.R., and Iqbal, A.J. (2017). Cannabinoid Receptor 2 Modulates Neutrophil Recruitment in a Murine Model of Endotoxemia. *Mediators of Inflammation* 2017, 15.
- Kapellos, T.S., Taylor, L., Feuerborn, A., Valaris, S., Hussain, M.T., Rainger, G.E., Greaves, D.R., and Iqbal, A.J. (2019). Cannabinoid receptor 2 deficiency exacerbates inflammation and neutrophil recruitment. *The FASEB Journal* 33, 6154-6167.
- Kerage, D., Sloan, E.K., Mattarollo, S.R., and McCombe, P.A. (2019). Interaction of neurotransmitters and neurochemicals with lymphocytes. *Journal of Neuroimmunology* 332, 99-111.
- Kesteman, N., Vansanten, G., Pajak, B., Goyert, S.M., and Moser, M. (2008). Injection of lipopolysaccharide induces the migration of splenic neutrophils to the T cell

- area of the white pulp: role of CD14 and CXC chemokines. *J Leukoc Biol* 83, 640-647.
- Krott, L.M., Piscitelli, F., Heine, M., Borrino, S., Scheja, L., Silvestri, C., Heeren, J., and Di Marzo, V. (2016). Endocannabinoid regulation in white and brown adipose tissue following thermogenic activation. *Journal of lipid research* 57, 464-473.
- Laprairie, R.B., Kelly, M.E.M., and Denovan-Wright, E.M. (2012). The dynamic nature of type 1 cannabinoid receptor (CB(1)) gene transcription. *British journal of pharmacology* 167, 1583-1595.
- Lefort, C.T., and Ley, K. (2012). Neutrophil arrest by LFA-1 activation. *Frontiers in immunology* 3, 157-157.
- Leite, C.E., Mocelin, C.A., Petersen, G.O., Leal, M.B., and Thiesen, F.V. (2009). Rimonabant: an antagonist drug of the endocannabinoid system for the treatment of obesity. *Pharmacol Rep* 61, 217-224.
- Ley, K., Laudanna, C., Cybulsky, M.I., and Nourshargh, S. (2007). Getting to the site of inflammation: the leukocyte adhesion cascade updated. *Nature Reviews Immunology* 7, 678.
- Mackie, K. (2007). Understanding Cannabinoid Psychoactivity with Mouse Genetic Models. *PLOS Biology* 5, e280.
- Mackie, K. (2008). Cannabinoid Receptors: Where They are and What They do. *Journal of Neuroendocrinology* 20, 10-14.
- Maresz, K., Pryce, G., Ponomarev, E.D., Marsicano, G., Croxford, J.L., Shriver, L.P., Ledent, C., Cheng, X., Carrier, E.J., Mann, M.K., Giovannoni, G., Pertwee, R.G., Yamamura, T., Buckley, N.E., Hillard, C.J., Lutz, B., Baker, D., and Dittel, B.N. (2007). Direct suppression of CNS autoimmune inflammation via the

- cannabinoid receptor CB1 on neurons and CB2 on autoreactive T cells. *Nature Medicine* 13, 492.
- Mchugh, D., Tanner, C., Mechoulam, R., Pertwee, R.G., and Ross, R.A. (2008). Inhibition of human neutrophil chemotaxis by endogenous cannabinoids and phytocannabinoids: evidence for a site distinct from CB1 and CB2. *Molecular Pharmacology* 73, 441.
- Mussinu, J.M., Ruiu, S., Mule, A.C., Pau, A., Carai, M.A., Loriga, G., Murineddu, G., and Pinna, G.A. (2003). Tricyclic pyrazoles. Part 1: synthesis and biological evaluation of novel 1,4-dihydroindeno[1,2-c]pyrazol-based ligands for CB1 and CB2 cannabinoid receptors. *Bioorg Med Chem* 11, 251-263.
- Netherland, C.D., Pickle, T.G., Bales, A., and Thewke, D.P. (2010). Cannabinoid receptor type 2 (CB2) deficiency alters atherosclerotic lesion formation in hyperlipidemic Ldlr-null mice. *Atherosclerosis* 213, 102-108.
- Nourshargh, S., Hordijk, P.L., and Sixt, M. (2010). Breaching multiple barriers: leukocyte motility through venular walls and the interstitium. *Nature Reviews Molecular Cell Biology* 11, 366.
- Ofek, O., Karsak, M., Leclerc, N., Fogel, M., Frenkel, B., Wright, K., Tam, J., Attar-Namdar, M., Kram, V., Shohami, E., Mechoulam, R., Zimmer, A., and Bab, I. (2006). Peripheral cannabinoid receptor, CB2, regulates bone mass. *Proceedings of the National Academy of Sciences of the United States of America* 103, 696-701.
- Ostenfeld, T., Price, J., Albanese, M., Bullman, J., Guillard, F., Meyer, I., Leeson, R., Costantin, C., Ziviani, L., Nocini, P.F., and Milleri, S. (2011). A randomized, controlled study to investigate the analgesic efficacy of single doses of the

- cannabinoid receptor-2 agonist GW842166, ibuprofen or placebo in patients with acute pain following third molar tooth extraction. *Clin J Pain* 27, 668-676.
- Pasquevich, K.A., Bieber, K., Günter, M., Grauer, M., Pötz, O., Schleicher, U., Biedermann, T., Beer-Hammer, S., Bühring, H.-J., Rammensee, H.-G., Zender, L., Autenrieth, I.B., Lengerke, C., and Autenrieth, S.E. (2015). Innate immune system favors emergency monopoiesis at the expense of DC-differentiation to control systemic bacterial infection in mice. *European Journal of Immunology* 45, 2821-2833.
- Pereira, A., Chappell, A., Dethy, J., Hoeck, H., Arendt-Nielsen, L., Verfaillie, S., Boulanger, B., Jullion, A., Johnson, M., and Mcnearney, T. (2013). A proof-of-concept (POC) study including experimental pain models (EPMs) to assess the effects of a CB2 agonist (LY2828360) in the treatment of patients with osteoarthritic (OA) knee pain. *Clinical Pharmacology and Therapeutics* 93.
- Pertwee, R.G. (2008). The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *British journal of pharmacology* 153, 199-215.
- Rogers, N. (2015). Cannabinoid receptor with an 'identity crisis' gets a second look. *Nat Med* 21, 966-967.
- Sardinha, J., Kelly, M.E.M., Zhou, J., and Lehmann, C. (2014). Experimental Cannabinoid 2 Receptor-Mediated Immune Modulation in Sepsis. *Mediators of Inflammation* 2014, 7.
- Singh, U., Singh, N., Singh, B., Price, R., Nagarkatti, M., and Nagarkatti, P. (2011). Cannabinoids Receptor-2 (CB2) agonist ameliorates spontaneous colitis by attenuating neutrophils, mast cells, natural killer cells and inducing endogenous cannabinoids (162.3). *The Journal of Immunology* 186, 162.163.

- Soethoudt, M., Grether, U., Fingerle, J., Grim, T.W., Fezza, F., De Petrocellis, L., Ullmer, C., Rothenhausler, B., Perret, C., Van Gils, N., Finlay, D., Macdonald, C., Chicca, A., Gens, M.D., Stuart, J., De Vries, H., Mastrangelo, N., Xia, L., Alachouzos, G., Baggelaar, M.P., Martella, A., Mock, E.D., Deng, H., Heitman, L.H., Connor, M., Di Marzo, V., Gertsch, J., Lichtman, A.H., Maccarrone, M., Pacher, P., Glass, M., and Van Der Stelt, M. (2017). Cannabinoid CB2 receptor ligand profiling reveals biased signalling and off-target activity. *Nat Commun* 8, 13958.
- Steffens, S., Veillard, N.R., Arnaud, C., Pelli, G., Burger, F., Staub, C., Zimmer, A., Frossard, J.-L., and Mach, F. (2005). Low dose oral cannabinoid therapy reduces progression of atherosclerosis in mice. *Nature* 434, 782-786.
- Taylor, L., Christou, I., Kapellos, T.S., Buchan, A., Brodermann, M.H., Gianella-Borradori, M., Russell, A., Iqbal, A.J., and Greaves, D.R. (2015). Primary Macrophage Chemotaxis Induced by Cannabinoid Receptor 2 Agonists Occurs Independently of the CB2 Receptor. *Sci Rep* 5, 10682.
- Thomas, A., Baillie, G.L., Phillips, A.M., Razdan, R.K., Ross, R.A., and Pertwee, R.G. (2007). Cannabidiol displays unexpectedly high potency as an antagonist of CB1 and CB2 receptor agonists in vitro. *Br J Pharmacol* 150, 613-623.
- Turcotte, C., Chouinard, F., Lefebvre, J.S., and Flamand, N. (2015). Regulation of inflammation by cannabinoids, the endocannabinoids 2-arachidonoyl-glycerol and arachidonoyl-ethanolamide, and their metabolites. *Journal of Leukocyte Biology* 97, 1049-1070.
- Turner, R.J., and Sharp, F.R. (2016). Implications of MMP9 for Blood Brain Barrier Disruption and Hemorrhagic Transformation Following Ischemic Stroke. *Frontiers in cellular neuroscience* 10, 56-56.

- Van Der Wal, A.C., and Becker, A.E. (1999). Atherosclerotic plaque rupture – pathologic basis of plaque stability and instability. *Cardiovascular Research* 41, 334-344.
- Van Sickle, M.D., Duncan, M., Kingsley, P.J., Mouihate, A., Urbani, P., Mackie, K., Stella, N., Makriyannis, A., Piomelli, D., Davison, J.S., Marnett, L.J., Di Marzo, V., Pittman, Q.J., Patel, K.D., and Sharkey, K.A. (2005). Identification and functional characterization of brainstem cannabinoid CB2 receptors. *Science* 310, 329.
- Wen, J., Ribeiro, R., Tanaka, M., and Zhang, Y. (2015). Activation of CB2 receptor is required for the therapeutic effect of ABHD6 inhibition in experimental autoimmune encephalomyelitis. *Neuropharmacology* 99, 196-209.
- Werth, V., Oddis, C.V., Lundberg, I.E., Fiorentino, D., Cornwall, C., Dgetluck, N., Constantine, S., and White, B. (2019). SAT0303 DESIGN OF PHASE 3 STUDY OF LENABASUM FOR THE TREATMENT OF DERMATOMYOSITIS. *Annals of the Rheumatic Diseases* 78, 1228-1228.
- Werth, V.P., Hejazi, E., Pena, S.M., Haber, J.S., Okawa, J., Feng, R., Gabre, K., Concha, J.S., Cornwall, C., Dgetluck, N., Constantine, S., and White, B. (2018). FRI0470 A phase 2 study of safety and efficacy of lenabasum (JBT-101), a cannabinoid receptor type 2 agonist, in refractory skin-predominant dermatomyositis. *Annals of the Rheumatic Diseases* 77, 763-764.
- Yabluchanskiy, A., Ma, Y., Iyer, R.P., Hall, M.E., and Lindsey, M.L. (2013). Matrix metalloproteinase-9: Many shades of function in cardiovascular disease. *Physiology (Bethesda, Md.)* 28, 391-403.
- Zarbock, A., and Ley, K. (2009). Neutrophil adhesion and activation under flow. *Microcirculation (New York, N.Y. : 1994)* 16, 31-42.

Zhao, Y., Yuan, Z., Liu, Y., Xue, J., Tian, Y., Liu, W., Zhang, W., Shen, Y., Xu, W., Liang, X., and Chen, T. (2010). Activation of Cannabinoid CB2 Receptor Ameliorates Atherosclerosis Associated With Suppression of Adhesion Molecules. *Journal of Cardiovascular Pharmacology* 55, 292-298.