Enrichment of organic nitrogen in primary biological particles during advection over the North Atlantic

Dall'Osto, Manuel; Santl-Temkiv, Tina; O'Dowd, Colin; Harrison, Roy M.

DOI:

[10.1016/j.atmosenv.2019.117160](https://doi.org/10.1016/j.atmosenv.2019.117160)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Dall'Osto, M, Santl-Temkiv, T, O'Dowd, C & Harrison, RM 2020, 'Enrichment of organic nitrogen in primary biological particles during advection over the North Atlantic', *Atmospheric Environment*, vol. 222, 117160, pp. 1-9. <https://doi.org/10.1016/j.atmosenv.2019.117160>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Enrichment of Organic Nitrogen in Primary Biological Particles during advection over the North Atlantic

Manuel Dall'Osto^{1*}, Tina Santl-Temkiv², Colin O'Dowd³ and Roy M. Harrison⁴

¹ Institut de Ciències del Mar, CSIC Pg Marítim de la Barceloneta 37-49, 08003 Barcelona, Spain Email: dallosto@icm.csic.es

²Aarhus University, Department of Environmental Science, Roskilde, Denmark

³Department of Experimental Physics and Environmental Change Institute, National University of Ireland Galway, Ireland

⁴National Centre for Atmospheric Science Division of Environmental Health & Risk Management School of Geography, Earth & Environmental Sciences University of Birmingham, Edgbaston, Birmingham B15 2TT United Kingdom

Key points:

- Single particle data collected over Saharan desert-Tropical-North Atlantic Ocean.
- Data used to study Saharan dust (SD) and Primary Biogenic Aerosol Particles (PABP).
- Minor presence of organic nitrogen and alkylamines in SD-PBAB at their origin.
- PBAP are found to be enriched in alkylamines in North Atlantic Ocean air.
- Atmospheric biogeochemical processes may be responsible for detected alkylamines.

Keywords: Organic nitrogen, biological particles, ATOFMS, single particle, aerosol aging.

Abstract

Using a number of datasets from single particle Aerosol Time-Of-Flight Mass Spectrometry (ATOFMS) measurements, we show only a minor presence of Organic Nitrogen (ON) species in Saharan dust particles (aerosol size range 0.2-3 μ m) collected at their origin. ON enhancement is not observed on Saharan dust particles after atmospheric transport (48-96 hours) either over the Tropical Ocean in the vicinity of Cape Verde, over the North Atlantic Ocean. In the negative ATOFMS mass spectra, signals due to ON species were found weak in biological particles (rich in potassium and phosphate) in their source area, and signals due to alkylamines in the positive ATOFMS mass spectra were not found. In striking contrast, biological particles travelling within aerosol dust plumes are found to be enriched in ON species - including alkylamines - in North Atlantic Ocean air (Mace Head, Ireland), as seen in both positive and negative ATOFMS mass spectra. Contrary to filter based aerosol techniques which report ON enrichment within Saharan dust, our single particle mass spectrometry data - allowing study of the aerosol mixing state - suggests that the aging biological particles and not the associated transported aging Saharan dust may be a source of ON species. We suggest biogeochemical processes occurring in the atmosphere in which biological particles are responsible for ON production. This may be an important source of nutrients to the ocean via atmospheric deposition.

1. Introduction

Organic nitrogen (ON) represents a consistent fraction of the total atmospheric nitrogen and is an ubiquitous component of the atmospheric aerosols (Galloway et al., 2008; DeLong, E., 2009). Aerosol deposition and in particular the deposition of ON, to marine environments, where auto- and heterotrophic microbial communities are often nutrient-limited, may affect community composition, productivity and consequently the marine carbon cycle, including carbon sequestration (Cornell et al., 2003; Cape et al., 2011). Organic nitrogen is found as a complex mixture of materials from multiple natural and anthropogenic sources. Kanakidou et al. (2012) estimated that 45% of the total ON atmospheric flux to the ocean comes from combustion sources, 32% from primary biological aerosol particles (PBAP), 1% from soil dust, 20% from ocean primary emissions and 3% from ocean secondary emissions. However, large uncertainties are associated with N emissions and the magnitude and impacts of anthropogenic atmospheric nitrogen inputs on the ocean (Kanakidou et al., 2016; Jickells et al., 2017, Kanakidou et al., 2018). Altieri et al. (2016) suggested that anthropogenic activity is not a significant source of organic N to the marine atmosphere over the North Atlantic, despite large pollution sources present in North America. Other bio-aerosol studies also suggest that PBAP may account for almost a third of ON in the marine atmosphere (Cote et al. 2008, Tittensor et al., 2010; Vařtilingom et al. 2013, Krumnis et al. 2014, Miriokefalitakis et al., 2017; Amato et al. 2017, Šantl-Temkiv et al. 2018). A fraction of PBAP may be able to sustain microbial activity in the atmosphere and could also therefore be involved in atmospheric ON processing (Hill et al., 2007). For example, Šantl-Temkiv et al (2013) suggested that bacterial cells present in cloud droplets may have been responsible for the process of ON mineralization to inorganic nitrogen that the studies observed in the clouds.

The correct apportionment of ON sources is particularly important in the marine atmosphere because of the role that depositing ON plays in enhancing ocean productivity (Duce et al., 2008). Anthropogenic, agricultural and biomass burning sources of ON are mainly important in cases when terrestrial air masses travel over marine areas (Jickells et al., 2013). Less is known of the importance of ON marine emissions. Primary marine emissions of ON in the form of amino acids and their derivatives have been reported

138 (Graber and Rudich, 2006; Altieri et al. 2009; Geddes et al. 2009). Gas-to-particle
139 secondary organic aerosol reactions of gaseous amines with sulphuric acid have also
140 been found in marine aerosol over the North Atlantic (Facchini et al., 2008) and Pacific
141 Ocean (Sorooshian et al. 2008; Miyazaki et al. 2010; 2011). The Tropical NE Atlantic
142 marine atmosphere presents a peculiarly intricate scenario with a diverse, variable and
143 complex mixture of ON aerosol sources (Baker et al., 2010). Lesworth et al. (2010)
144 reported a wide range of ON concentrations associated with both coarse ($>1\mu\text{m}$) and fine
145 ($<1\mu\text{m}$) particles, apportioning them to African dust sources and to anthropogenic
146 emissions from Europe. Other studies carried out in the Eastern Mediterranean (Violaky et
147 al. 2010; Violaki and Mihalopoulos 2010) also stress that combustion processes and
148 African dust are an important source of ON. By contrast, ON of marine origin seems to
149 play a small role in these sub-tropical marine regions (Muller et al., 2009). Although these
150 studies stress the importance of dust for the marine ON budgets in air, the low sampling
151 time resolution (1-4 days) and off-line analysis that the authors used, do not allow them to
152 conclude whether the ON is associated directly with dust particles or with the large
153 quantities of PBAP that are ejected into the atmosphere with the dust events (Kellogg and
154 Griffin 2006).

155

156 Aside from the complexity and diverse sources of ON, there are also two main issues
157 associated with sampling and analysing ambient ON aerosols. First, the concentration of
158 ON is often estimated by measuring concentrations of total nitrogen and subtracting the
159 concentrations of NO_3^- and NH_4^+ , which propagates errors from the three individual
160 species and leads to uncertainties in the ON analytical concentrations. Second, most of
161 the current measurements are carried out with off-line filter measurements (Sullivan and
162 Prather, 2005), which do not allow information on the ON mixing state and hence
163 challenge the aerosol source apportionment. An aerosol population is broadly "internally
164 mixed" when all single particles have the same chemical composition (equal to the bulk
165 composition). By contrast, an aerosol population is "externally mixed" when single particles
166 have a different chemical composition. In order to reduce uncertainty on ON-containing
167 aerosols, Jickells et al. (2013) stressed the importance of deploying new technologies.
168 Detection of PBAPs by real-time (RT) techniques for autonomous, online detection and
169 characterization of PBAP properties has become increasingly important for research

170 purposes (Hoffman et al., 2019). Single particle mass spectrometry (SPMS) - techniques
171 that ablate and ionize single aerosol particles - is capable of PBAP detection, although the
172 interpretation of SPMS mass spectra is challenging and requires extensive human
173 interaction. For example, amino acids are ubiquitous compounds with an integral role in
174 atmospheric bioaerosol compositions (Ge et al., 2011). Previous SPMS identified amino
175 acids in PBAP particles (Ferguson et al., 2004; Russell et al., 2004; Czerwieniec et al,
176 2005, Schneider et al., 2011). However, difference in ionization laser wavelengths may
177 produce different single-particle mass spectra, and therefore, specific markers for amino
178 acids do not exist.

179 This study uses state-of-the-art Single Particle Aerosol Time-Of-Flight Mass Spectrometer
180 (ATOFMS; Su et al., 2004) to shed some light on natural aerosols containing ON species.
181 The ATOFMS data herein presented allows us to determine if the aerosol population is
182 internally or externally mixed, a peculiar feature absent in most filtration-based aerosol
183 sampling techniques (Sullivan and Prather, 2005). It is also worth reminding that the
184 ATOFMS is a particularly good aerosol instrument for studying amines and organic
185 nitrogen in general because the LDI laser wavelength (266 nm) ionizes them very
186 efficiently (Angelino et al., 2001; Dall'Osto et al., 2016; Healy et al., 2015).

187
188 The large ATOFMS datasets used in this work were previously used to study the variation
189 of the mixing state of Saharan dust particles with atmospheric transport (Dall'Osto et al.,
190 2010a) during the DODO project; and to study the aerosol properties associated with air
191 masses arriving into the North East Atlantic during the 2008 Mace Head EUCAARI
192 intensive observing period (Dall'Osto et al., 2010b). The current study aims - for the first
193 time - to specifically presents an ATOFMS analysis looking at the properties of specific
194 clusters - including biological particles - and comparing the temporal trends with black
195 carbon and dimethyl sulphide concentrations across different locations. We focus our
196 attention on Saharan dust and biological particles found at their origins, as well as on
197 those sampled during ambient field measurements over both Tropical and North Atlantic
198 Oceans. The aims of this study include: (1) to elucidate if Saharan dust or the associated
199 terrestrial PBAP contains ON, (2) to understand if ON in natural aerosols has mainly a
200 primary or a secondary origin and (3) to discuss whether aerosol aging results in a
201 degradation or production of ON.

202 **2. Methodology**

203

204 **2.1 Atmospheric measurements**

205

206 **2.1.1 ATOFMS**

207

208 We deployed an ATOFMS (Model 3800-100, TSI, Inc.), which collects bipolar mass
209 spectra of individual aerosol particles. Briefly, ambient aerosol is focused into a narrow
210 particle beam for sizes between 100 nm and 3 μm (vacuum aerodynamic diameter),
211 following which laser desorption/ionization take place and a positive and a negative single
212 particle mass spectra are obtained. The mass spectrum is only qualitative because the
213 peaks intensities depend strongly on the particle matrix, the coupling between the laser
214 and the particle and the shot to shot variability of the laser (Dall'Osto and Harrison, 2006).
215 However, recent studies (Jeong et al., 2011) report excellent correlations for inorganic
216 species (sulfate, nitrate, and ammonium) but weaker ones between total organic and
217 elemental carbon (EC) detected with ATOFMS and other instruments (Jeong et al., 2011).
218 Furthermore, using a different approach, Healy et al. (2012, 2013) demonstrated that the
219 ATOFMS can deliver good quantitative results for Elemental Carbon (EC) and Organic
220 Carbon (OC) containing particles, when compared with other measurement techniques.
221 Overall, the ATOFMS gives a measure of particle number as function of size and
222 chemical composition, allowing the determination of the mixing state.

223

224 **2.1.2 Dimethyl sulphide (DMS) measurements**

225

226 DMS measurements were carried during the North Atlantic Marine Boundary Layer
227 Experiment (NAMBLEX) took place at Mace Head, Ireland during summer 2002 (Lewis et
228 al.; 2005), further information can be found elsewhere (Heard et al., 2006).

229

230 **2.1.3 Black Carbon (BC) measurements**

231

Aerosol absorption (and Black Carbon mass) was measured using both a McGee Scientific Aethalometer AE-16 and a Multi-Angle Absorption Photometer (MAAP) (Dall'Osto et al., 2010b).

2.2 Laboratory and field studies

In this study, we utilized previously published Saharan dust ATOFMS data from our laboratories (Dall'Osto et al., 2010a) for the presence of ON. The Sahara Sahel Dust Corridor (SSDC) is a huge zone lying between latitudes 12° N and 28° N and running 4000 km east-west from Chad to Mauritania (Moreno et al., 2006). During low wind conditions, four different samples of SSDC desert soil, road, and aeolian dusts were collected. A summary of the SSDC locations is discussed elsewhere (Moreno et al., 2006; Dall'Osto et al., 2010a). Briefly, three dust samples were taken from the Saharan region, and one from Algeria. Each dust sample was put into a flask in a sonicator, a suspension of dust particles under filtered lab air flow was created, and single particles mass spectra were obtained with the ATOFMS. About 1,000 single particle mass spectra were collected for each of the four dust samples.

For this study, we also utilized ATOFMS datasets collected in two previous field studies. This gives the possibility to compare single particle mass spectra from airborne particles with those obtained from the Saharan dust particles analyzed in the laboratory. During August and September 2002 The North Atlantic Marine Boundary Layer Experiment (NAMBLEX) took place at Mace Head, Ireland (Heard et al., 2006). The Atmospheric Research Station at Mace Head allows westerly exposure to the North Atlantic ocean (clean sector, 180 degrees through west to 300 degrees) and the opportunity to study atmospheric composition under Northern Hemispheric background conditions. Also, within the Dust Outflow and Deposition to the Ocean (SOLAS-DODO) project (McConnell et al. 2008), an ATOFMS was deployed during a research cruise (P332) in the vicinity of the Cape Verde Islands (January-February 2006) on board the research vessel FS Poseidon as part of the UK-SOLAS programme. The study areas are summarized in Fig. 1:

- Fig 1, Sites A, B: Four samples of desert soil were collected from several regions of the Sahara-Sahel Dust Corridor during low wind conditions and analysed in a laboratory study (4,205 ATOFMS mass spectra, Dall'Osto et al. 2010a).
- Fig 1, Site C: A cruise (research vessel FS Poseidon; P332, Winter 2006) within the Dust Outflow and Deposition to the Ocean (SOLAS-DODO), where an ATOFMS collected 187,205 ATOFMS single particle mass spectra (Rijkenberg et al. 2008).
- Fig. 1, Site D: An ATOFMS operated at the Mace Head Atmospheric Research Station during the North Atlantic Marine Boundary Layer Experiment (NAMBLEX, Summer 2002) collecting 191,504 ATOFMS single particle mass spectra (Dall'Osto et al., 2004).

2.3 Data analysis

The ATOFMS data presented cover a large aerosol size range (100-3000 nm) at a high time resolution (1 hour) which were collected over a large area (Figure 1). ATOFMS single particle positive and negative mass spectra were imported into YAADA (Yet Another ATOFMS Data Analyzer, Song et al., 1999) and single-particle mass spectra were grouped with adaptive resonance theory neural network, ART-2a (learning rate 0.05, vigilance factor 0.85, and 20 iterations). Further details of the parameters can be found elsewhere (Dall'Osto and Harrison, 2006). Briefly, the ART-2a area matrix (AM) of a single particle mass spectra represents the average intensity for each m/z for all particles within a group, reflecting the typical mass spectra of specific aerosol groups.

Finally, air mass back trajectories of the air masses arriving at the different study areas were calculated at four arrival times for each day of the campaign (00:00, 06:00, 12:00 and 18:00 UTC), depicting the path taken by the air mass reaching the sampling site over the previous five days. The back trajectories were run using the on-line HYSPLIT model developed by the National Oceanic and Atmospheric Administration (NOAA) (Draxler and Rolph, 2003).

3. Results

295 The ART-2a algorithm generated a number of clusters used to describe the ATOFMS
 296 mass spectra. Most of the obtained particle types were associated with anthropogenic
 297 sources (fresh and aged combustion related aerosols), sea salt, ship emissions,
 298 secondary aerosol of inorganic nature, some of which were already discussed elsewhere
 299 (Dall'Osto et al., 2004; 2006, 2010a,b). Table 1 shows a summary of the previous results,
 300 and put it in context with the current specific ON analysis. Table 1 shows that Saharan
 301 dust particles and biological particles were detected in all the ATOFMS datasets
 302 presented. ATOFMS has already proved to be a good tool able to separate dust (mainly
 303 Ca-rich or Al-Si rich) and biological particles (Ferguson et al., 2004; Pratt et al., 2009).
 304 Briefly, biological mass spectral signatures can be differentiated from dust on the basis of
 305 abundant organic and phosphorus ions, as well as a lack of key dust markers, such as
 306 aluminium and silicates. Within the main objective of this work, the ATOFMS is a very
 307 sensitive instrument towards ON (m/z - 26 and m/z -42, due to $[CN]^-$ and $[CNO]^-$,
 308 respectively), and many alkylamines (Angelino et al. 2001; Moffet et al. 2008; Pratt et al.
 309 2009; Healy et al., 2015; Dall'Osto et al., 2016).
 310 Figure 2 shows the positive and negative ART-2a ATOFMS mass spectra of dust particles
 311 and PBAP detected at their source (Saharan dust particles collected in Africa; Fig. 2 a, b,
 312 respectively); during the SOLAS-DODO cruise around Cape Verde (Fig. 2 c, d,
 313 respectively) and during the NAMBLEX field study at Mace Head, Ireland (Fig. 2e, f,
 314 respectively). It is clear that ATOFMS mass spectra of Saharan dust particles (Fig. 2 a, c,
 315 e) shows peaks due to most common ion species characteristic of crustal materials
 316 including aluminum (m/z 27 $[Al]^+$), calcium (m/z 40 $[Ca]^+$ and 56 $[CaO]^+$), titanium (m/z 48
 317 $[Ti]^+$ and 64 $[TiO]^+$), lithium (m/z 7 $[Li]^+$) and iron (54 $^{54}[Fe]^+$ and 56 $^{54}[Fe]^+$). Peaks for other
 318 metals such as sodium (m/z 23 $[Na]^+$), magnesium (m/z 24 $^{24}[Mg]^+$, 25 $^{25}[Mg]^+$) and
 319 potassium (m/z 39 $[K]^+$) can also be seen. In addition, other peaks for calcium (57
 320 $[CaOH]^+$, 96 $[Ca_2O]^+$ and 113 $[(CaO)_2H]^+$) were detected but at low intensity. The negative
 321 particle mass spectra in Figure 2 (a, c, e) show ions from oxygen (m/z -16 $^{16}[O]^-$, -17 $[OH]^-$),
 322 chloride (m/z -35 $^{35}[Cl]^-$, -37 $^{37}[Cl]^-$), phosphate (m/z -63 $[PO_2]^-$ and m/z -79 $[PO_3]^-$), a
 323 small abundance of -46 $[NO_2]^-$ and -62 $[NO_3]^-$. It is important to note that in these Al/Si-
 324 rich Saharan dust particles, peaks of silicate at m/z -60 $[SiO_2]^-$ and -76 $[SiO_3]^-$ dominate
 325 the negative spectra. This is further discussed in Dall'Osto et al. (2004, 2010a), where

Al/Si-rich dust particles consistent with Saharan dusts which are dominated by SiO_2 and Al_2O_3 (Goudie and Middleton, 2001) were observed.

Few particles found in Saharan soil dust samples (<4%) were internally mixed with ON species. We have previously reported that both organic and organic nitrogen species were more commonly associated with Saharan dust particles collected in the source region relative to the atmospheric Saharan dust particles sampled during the two field studies (SOLAS-DODO and NAMBLEX, Dall'Osto et al. 2010a). It is possible that some of these ON components may be mineralized (a process by which organic N is converted to plant-available inorganic forms) during atmospheric transport. This is supported by the observation that Saharan dust particles with shorter atmospheric residence, i.e. collected near the Cape Verde Islands (Fig 1, 2c, DODO experiment), showing increased internally mixed nitrate (possibly mineralized ON). In addition, this can be seen also in the Saharan dust particles with longer atmospheric residence times, i.e. collected on the coast of Ireland (Fig. 1, 2e, NAMBLEX experiment) showing also a very high degree of internally mixed secondary species including nitrate (m/z -46, see also Dall'Osto et al., 2010a). Overall, our studies suggest that fresh Saharan dust is not a major source of ON. It is important to stress that no clear peaks associated to alkylamines (Healy et al., 2015; Dall'Osto et al., 2016) were detected in any of the Saharan dust particle types (Fig. 2 a, c, e).

With regard to biological particles, those were detected in all the datasets: the Saharan soil dust samples and the two ATOFMS ambient field studies (Table 1, Fig. 2). A strong peak at m/z 39 is usually seen in the positive mass spectra, whereas the negative spectrum is usually dominated by phosphate $[\text{PO}_2]^-$ m/z -63 and $[\text{PO}_3]^-$ m/z -79) and often also internally mixed with ON species (Pratt et al. 2009). Biological particles detected in Saharan dust samples and over the Tropical Ocean had a weak signal associated with ON and amines. Our results - showed in Fig. 2 - suggest that fresh PBAP are lacking in ON signal, both at their origin (dust samples collected in Africa) and after a few days of transport (Cape Verde region). By striking contrast, 19% of PBAP detected at Mace Head were internally mixed with ON compounds. By querying the ATOFMS NAMBLEX dataset, it was found that the majority (>99%) of the signal due to ON and amines species was associated with a specific PBAP type (Fig. 2 f), indicating that this is a major source of ON sampled over the North Atlantic. Additionally, strong peaks of alkylamines are seen in the

358 positive mass spectra, including m/z 58 [$C_2H_5NH=CH_2$]⁺, m/z 59 [$N(CH_3)_3$]⁺ and m/z 86
 359 [$(C_2H_5)_2N=CH_2$]⁺ (Dall'Osto et al., 2016). The peak of m/z 74 and m/z 104 (potentially
 360 [$C_2H_4NO_2$]⁺ and [$C_3H_6NO_3$]⁺, respectively) were previously attributed to organic nitrogen
 361 peaks including alkylamines (Angelino et al., 2001). A previous study in the city of
 362 Barcelona (Spain) found that m/z 104 was found to be to be linked alkylamines of primary
 363 aerosol origin, whereas m/z 74 alkylamines of secondary aerosol origin (Dall'Osto et al.,
 364 2016). This particle type (Fig. 2 f) presented an aerosol size distribution centred at about
 365 2 μ m, shifted towards an even larger modes than Al-Si Saharan particles (Dall'Osto et al.
 366 2004). An overview the differences in spectra for the PBAP NAMBLEX (aged transported
 367 PBAP) and its origin (PBAP Saharan region) or in the Tropical Ocean (PBAP DODO) can
 368 be seen in Figure 3, which show the subtraction spectrum for PBAP NAMBLEX minus
 369 PBAP DODO (Fig. 4a) and for PBAP NAMBLEX minus PBAP Saharan dust origin (Fig.
 370 4b). The positive ion spectrum for PBAP NAMBLEX is enhanced in organic nitrogen and
 371 alkylamines (m/z 59, 74, 86, 104), whilst the negative is enriched in nitrate peaks (m/z -46,
 372 -62) and organic nitrogen (m/z - 42). A representative ATOFMS single particle mass
 373 spectrum of the PBAP particle carrying ON (Fig. 2 f) is presented in Figure 4, showing a
 374 strong peak at m/z 39 likely to be due to potassium [K], and minor peaks at m/z 23 [Na]
 375 and m/z 56 [Fe] in the positive mass spectra. In the negative mass spectra, ON peaks are
 376 found at m/z -26 [CN] and m/z -42 [CNO], along with phosphate (m/z -63 and m/z -79) and
 377 methanesulfonic acid MSA (m/z -95). The mass spectrum has features that are generally
 378 attributed to vegetative debris (potassium-phosphate) except that it is internally mixed with
 379 ON species. Major ion peaks in the positive mass spectra at m/z 58, 59, 74, 86, and 104,
 380 118 and 154 are all consistent with ON ATOFMS mass spectra peaks detected in previous
 381 field and laboratory studies (Angelino et al. 2001, Moffet et al. 2008, Healy et al., 2015:
 382 Dall'Osto et al., 2016).

383 In order to investigate the origin of the dominant PBAP associated with ON during
 384 NAMBLEX, we separated the field study sampling period accordingly to three main air
 385 mass categories (marine Tropical, marine Polar-Arctic and European continental Polar –
 386 mT, mP-mA and cP respectively, Dall'Osto et al. 2010b). We found - as expected - higher
 387 values of BC for cP air masses (209 ± 197 ng m^{-3}) relative to mT (51 ± 23 ng m^{-3}) and the
 388 very clean mP-mA ones (33 ± 21 ng m^{-3}) (Figure 4a). DMS concentrations - by contrast -
 389 were much higher for mT air masses (507 ± 120 pptV) than for cP (105 ± 110 pptV) and mP-

mA (140 ± 120 pptV) (Figure 4b). Other studies (Dall'Osto et al. 2010b, Bassford et al. 1999) have also indicated that during mT air masses the vast majority of DMS and MSA observed at the Mace Head site had been transported significant distances from the mid tropical Atlantic. Figure 4c shows that biological particles were detected at about double the concentration (t-test, 95%, 1.31 ± 1 and 2.67 ± 2 respectively) for mT relative to cP air masses. This PBAP type was found in association with Al-Si particles detected in the same study and attributed to dust particles, mainly detected during mT air masses and to a lesser extent during cP air masses (Dall'Osto et al 2004, 2010a, Figure 3d). An increase in microbial concentrations during dust storms has previously been reported and can be two orders of magnitude higher than under non-dusty conditions (Hara and Zhang 2012, Yamaguchi et al 2014, Mazar et al. 2016). In conclusion, the PBAP associated with ON were likely transported to the North Atlantic together with dust events, which may have originated in the Sahara-Sahel region and travelled over the Tropical Ocean. During their atmospheric travel time, the PBAP injected with dust events, seem to first be low in ON as we observed during the SOLAS-DODO measurements and then get enriched with ON with atmospheric transport as we observed during the NAMBLEX measurements.

4. Discussion and Conclusion

The first conclusion of this study is that (within the $0.2\text{-}3\mu\text{m}$ size) the ON aerosol concentrations reported over the Tropical Ocean - at least during the study period - are likely to be contained within the terrestrial PBAP travelling in association with Saharan dust and not with the dust particle themselves. Saharan dust is also not found to be enhanced in ON species when detected over the North Atlantic. Overall, our single particle mass spectrometry measurements collected at high time resolution over a large geographical area, suggest that Saharan dust particles are not rich in ON, either at their origin, or in the Cape Verde region. The second conclusion of this study is that the PBAP that are injected into the atmosphere together with dust events are the aerosol particles that contain ON compounds. Furthermore, it appears that these biological particles get enriched in ON species during their atmospheric residence. The low associated BC concentrations make an anthropogenic influence unlikely.

421 Previous ATOFMS studies discovered existence of different ON groups (alkylamines,
422 proteins, amino acids etc.) in aerosols and reported their involvement in atmospheric
423 processes, including indoor tobacco smoke (Dall'Osto et al., 2007), cloud/fog processing
424 (Rehbein et al., 2011), primary traffic pollution (Angelino et al., 2001; Dall'Osto et al.,
425 2016), non-salt organic aerosol products through reaction with oxidizing agents (Murphy et
426 al., 2007, Pratt et al., 2009) and marine aerosol reactions (Dall'Osto et al., 2012; Healy et
427 al., 2015). However, it is important to note that - in this study - ON was found internally
428 mixed with biological particles and externally mixed with other aerosols, therefore we can
429 conclude that ON is not formed by condensation of secondary semi-volatile organic
430 nitrogen species as these compounds would condense on the whole aerosol population.

431
432 An explanation for the ON enrichment of the aging PBAP can only be speculative at this
433 stage. Previous studies have demonstrated that viable airborne micro-organisms present
434 in the atmosphere can contribute to atmospheric chemistry through degradation
435 processes, as well as chemical change due to the release or desorption of molecules from
436 microbiological entities (Deguillaume et al. 2008; Cote et al., 2008). Airborne metabolically
437 active microorganisms, for example fungal and bacterial cells, can transform chemical
438 constituents of the atmosphere by metabolic activity and have been implicated in the
439 nitrogen and carbon cycling in clouds (Hill et al. 2007, Cote et al 2008, Väitilingom et al
440 2013, Krumnis et al. 2014, Amato et al 2017, Šantl-Temkiv et al 2018). Nitrogen can be
441 microbially incorporated into organic compounds through the processes of fixation and
442 assimilation. Different prokaryotes - including Cyanobacteria - which are commonly found
443 in the atmosphere (reviewed in Genitsaris et al. 2011), are capable of nitrogen fixation and
444 can convert atmospheric nitrogen into biologically accessible ammonia. Assimilation of
445 inorganic nitrogen is performed by diverse bacteria and fungi, who can convert nitrate and
446 ammonia into different forms of ON. It is not known at this stage if the association of ON
447 with aging biological particles is due to the metabolic activity of microorganisms externally
448 mixed with Saharan dust plumes, which have travelled to the North Atlantic region. The
449 long distance transport of microorganisms with dust particles has been shown to
450 negatively affect the viability of microbial cells as <40% of all bacterial cells were found
451 viable compared to >70% of non-dust-associated cells (Hara and Zhang 2012). Despite
452 this, the absolute number of viable microorganisms travelling with dust storms remained

high. It may be that these viable microbial cells may maintain their metabolic activity and affect the atmospheric nitrogen cycle. Further multidisciplinary studies on the impact of airborne micro-organisms and possible chemical mechanisms are recommended to underpin these findings.

Another possible explanation is that the biological particles enriched in ON have a marine origin. However, we detected such biological particles mainly during dust events. Usually, concentration of bacteria over the sea are much lower than over land, and higher concentrations of aerosolized microorganisms during dust events relative to clean background marine conditions are normally observed (Kellog and Griffin 2006, Prospero et al., 2005; Griffin et al., 2006). Furthermore, the ATOFMS mass spectra of ocean-derived particle types are enriched in elements (mainly Na and Mg, Gaston et al., 2011) not found in the biological particles reported in this study.

In conclusion, using on-line single particle aerosol mass spectrometry, we show that fresh and aged Saharan dust aerosols are not a major source of ON. Furthermore, we find that biological particles – likely of terrestrial origin – are enriched in ON during atmospheric transport. We hypothesize that microbial processes may generate ON through nitrogen fixation and assimilation during the atmospheric residence of microbial cells, which results in an enrichment of ON compounds within the cell. Coarser particles possess higher dry deposition rates and such microorganisms internally mixed with ON may be an important source of nitrogen, which is an essential nutrient in North Atlantic Ocean marine ecosystems. González Benítez et al. (2009) found that dry deposition of particles was the main contributor of N-containing species in bulk rain samples, which were not strongly associated with either combustion or agricultural sources alone (González Benítez et al. 2010). Our data do not rule out a suggestion that SOA production of amines is important for the marine atmosphere when considering the fine particle mode (Facchini et al. 2008, Muller et al. 2009), but we propose an additional source for the total ON concentrations in

the coarse aerosols mode that has a natural origin. Further studies are needed in order to validate the event reported in this study.

Acknowledgements

We express our gratitude to the Master and crew of FS Poseidon for support during the cruise. Both the Mace Head sampling and DODO cruise were supported financially by the U.K. Natural Environment Research Council. This work was also funded by the Ireland Higher Education Authority Program for Research in Third Level Institutes – Cycle 5. We thank Dr. A. Lewis for DMS data, as discussed in Heard et al. (2015). The study was also supported by the Spanish Ministry of Economy through project PI-ICE (CTM2017–89117-R) and the Ramon y Cajal fellowship (RYC-2012-11922).

References

- Amato, P., Joly, M., Besaury, L., Oudart, A., Taib, N., Mone, A.I., Deguillaume, L., Delort, A.M., Debroas, D., 2017. Active microorganisms thrive among extremely diverse communities in cloud water. PLoS One 12 e0182869.
- Angelino, S.; Suess, D. T.; Prather, K. A., 2001 Formation of aerosol particles from reactions of secondary and tertiary alkylamines: Characterization by aerosol time-of-flight mass spectrometry. Environ. Sci. Technol., 35 (15), 3130–3138.
- Altieri, K. E., B. J. Turpin and S. P. Seitzinger, 2009. "Composition of Dissolved Organic Nitrogen in Continental Precipitation Investigated by Ultra-High Resolution FT-ICR Mass Spectrometry." Environmental Science & Technology 43(18): 6950-6955.

- Altieri, K. E., Fawcett, S. E., Peters, A. J., Sigman, D. M., and Hastings, M. G.: Marine biogenic source of atmospheric organic nitrogen in the subtropical North Atlantic, 2016, *P. Natl. Acad. Sci. USA*, 113, 925–930, <https://doi.org/10.1073/pnas.1516847113>.
- Baker, A. R., T. Lesworth, C. Adams, T. D. Jickells, and L. Ganzeveld, 2010, Estimation of atmospheric nutrient inputs to the Atlantic Ocean from 50°N to 50°S based on large scale field sampling: Fixed nitrogen and dry deposition of phosphorus, *Global Biogeochem. Cycles*, 24, GB3006, doi:10.1029/2009GB003634.
- Bassford, M. R., Nickless, G., Simmonds, P. G., Lewis, A. C., Pilling, M. J., and Evans, M. J., 1999. The concurrent observation of methyl iodide and dimethyl sulphide in marine air; implications for sources of atmospheric methyl iodide, *Atmos. Environ.*, 33, 2373–2383.
- Cape JN, Cornell SE, Jickells TD, Nemitz E. 2011 Organic nitrogen in the atmosphere—where does it come from? A review of sources and methods. *Atmos. Res.* 102, 30–48. (doi:10.1016/j.atmosres.2011.07.009)
- Cornell S.E., Jickells T.D., Cape J.N., Rowland A.P., Duce R.A., 2003. Organic nitrogen deposition on land and coastal environments: a review of methods and data, *Atmospheric Environment*, 37 2173-2191.
- Cote, G., Kos, R., Mortazavi, P., and Ariya, P. A., 2008 Microbial and “de novo” Transformation of Dicarboxylic Acids by Three Airborne Fungi, *Sci. Total Environ.*, 340, 530–532.
- Czerwieniec, G. A., S. C. Russell, H. J. Tobias, M. E. Pitesky, D. P. Fergenson, P. Steele, A. Srivastava, J. M. Horn, M. Frank, E. E. Gard, et al. 2005. Stable isotope labeling of entire *Bacillus atrophaeus* spores and vegetative cells using bioaerosol mass spectrometry. *Anal. Chem.* 77 (4):1081–1087. doi:10.1021/ac0488098
- Dall'Osto, M., Beddows, D.C.S., Kinnersley, R.P., Harrison, R.M., Donovan, R.J. and Heal M.R., 2004. Characterization of individual airborne particles by using Aerosol Time-of-Flight Mass Spectrometry (ATOFMS) at Mace Head, Ireland. *Journal of Geophysical Research* 109, D21302, doi: 10.1029/2004/JD004747.
- Dall'Osto, M. and Harrison R.M., 2006. Chemical characterisation of single airborne particles in Athens (Greece) by ATOFMS. *Atmospheric Environment* 40(39): 7614.
- Dall'Osto, M., Harrison, R. M., Charpantidou, E., Loupa, G., and Rapsomanikis, S., 2007: Characterisation of indoor airborne particles by using real-time aerosol mass spectrometry, *Sci. Total Environ.*, 384, 120–133.
- Dall'Osto, M.; R. M. Harrison, E. J. Highwood, C. O'Dowd, D. Ceburnis, X. Querol, E. P. Achterberg, 2010a. Variation of the mixing state of Saharan dust particles with atmospheric transport, *Atmospheric Environment*, Volume 44, Issue 26 Pages 3135-3146, ISSN 1352-2310, DOI: 10.1016/j.atmosenv.2010.05.030.

571 Dall'Osto, M., Ceburnis, D., Martucci, G., Bialek, J., Dupuy, R., Jennings, S. G.,
 572 Berresheim, H., Wenger, J., Healy, R., Facchini, M. C., Rinaldi, M., Giulianelli, L., Finessi,
 573 E., Worsnop, D., Ehn, M., Mikkilä, J., Kulmala, M., and O'Dowd, C. D., 2010b. Aerosol
 574 properties associated with air masses arriving into the North East Atlantic during the 2008
 575 Mace Head EUCAARI intensive observing period: an overview, *Atmos. Chem. Phys.*, 10,
 576 8413-8435, doi:10.5194/acp-10-8413-2010.
 577
 578 Dall'Osto, M., Ceburnis, D., Monahan, C., Worsnop, D. R., Bialek, J., Kulmala, M., Kurtén,
 579 T., Ehn, M., Wenger, J., Sodeau, J., Healy, R., and O'Dowd, C., 2012: Nitrogenated and
 580 aliphatic organic vapors as possible drivers for marine secondary organic aerosol growth,
 581 *J. Geophys. Res.-Atmos.*, 117, D12311, doi:10.1029/2012JD017522
 582
 583 Dall'Osto, M., Beddows, D. C. S., McGillicuddy, E. J., Esser-Gietl, J. K., Harrison, R. M.,
 584 and Wenger, J. C., 2016. On the simultaneous deployment of two single-particle mass
 585 spectrometers at an urban background and a roadside site during SAPUSS, *Atmos.*
 586 *Chem. Phys.*, 2016, 16, 9693-9710, <https://doi.org/10.5194/acp-16-9693>.
 587
 588 Deguillaume, L., Leriche, M., Amato, P., Ariya, P. A., Delort, A.-M., Pöschl, U.,
 589 Chaumerliac, N., Bauer, H., Flossmann, A. I., and Morris, C. E., 2008. Microbiology and
 590 atmospheric processes: chemical interactions of primary biological aerosols,
 591 *Biogeosciences*, 5, 1073-1084, doi:10.5194/bg-5-1073-2008.
 592
 593 DeLong, E. 2009 The microbial ocean from genomes to biomes. *Nature* 459, 200–206.
 594 (doi:10.1038/nature08059)
 595
 596 Draxler, R. R. and Rolph, G. D., 2003. HYSPLIT (Hybrid Single-Particle Lagrangian
 597 Integrated Trajectory) model v 4.9, NOAA Air Resource Laboratory, Silver Spring MD,
 598 available at: <http://www.arl.noaa.gov/ready/hysplit4.html>.
 599
 600 Duce, R.A., La Roche, J., Altieri, K., Arrigo, K.R., Baker, A.R., Capone, D.G., Cornell, S.,
 601 Dentener, F., Galloway, J., Ganeshram, R.S., Geider, R.J., Jickells, T., Kuypers, M.M.,
 602 Langlois, R., Liss, P.S., Liu, S.M., Middleburg, J.J., Moore, C.M., Nickovic, S., Oschlies,
 603 A., Pedersen, T., Prospero, J., Schlitzer, R., Seitzinger, S., Sorensen, L.L., Uematsu, M.,
 604 Ulloa, O., Voss, M., Ward, B., Zamora, L., 2008. Impacts of atmospheric anthropogenic
 605 nitrogen on the open ocean. *Science* 320, 893e897
 606
 607 Facchini, M. C., S. Decesari, M. Rinaldi, C. Carbone, E. Finessi, M. Mircea, S. Fuzzi, F.
 608 Moretti, E. Tagliavini, D. Ceburnis and C. D. O'Dowd, 2008. "Important Source of Marine
 609 Secondary Organic Aerosol from Biogenic Amines." *Environmental Science & Technology*
 610 42(24): 9116-9121.
 611
 612 Fergenson, D. P., Pitesky, M. E., Tobias, H. J., Steele, P. T., Czerwieniec, G. A., Russell,
 613 D. H., Lebrilla, C. B., Horn, J. M., Coffee, K. R., Srivastava, A., Pillai, S. P., Shih, M.-T.
 614 P., Hall, H. L., Ramponi, A. J., Chang, J. T., Langlois, R. G., Estacio, P. L., Hadley, R. T.,
 615 Frank, M. and Gard, E. E., 2004, Reagentless Detection and Classification of Individual
 616 Bioaerosol Particles in Seconds, *Anal. Chem.*, 76, 373-378.
 617

Huffman, J. A., A. E. Perring, N. J. Savage, BernaB.rd Clot, B. Crouzy, F. Tummon, Ofir
 Shoshanim, B. Damit, J. Schneider, V. Sivaprakasam, M. A. Zawadowicz, I. Crawford, M.
 Gallagher, D. Topping, D. C. Doughty, S. C. Hill & Y. Pan (2019): Real-time sensing of
 bioaerosols: Review and current perspectives, *Aerosol Science and Technology*, DOI:
 10.1080/02786826.2019.1664724
 Galloway, J. N., A. R. Townsend, J. W. Erisman, M. Bekunda, Z. Cai, J. R. Freney, L. A.
 Martinelli, S. P. Seitzinger, and M. A. Sutton, 2008, Transformations of the nitrogen cycle:
 Recent trends, questions, and potential solutions, *Science*, 320, 889–892,
 doi:10.1126/science.1136674.
 Gaston, C. J., H. Furutani, S. A. Guazzotti, K. R. Coffee, T. S. Bates, P. K. Quinn, L. I.
 Aluwihare, B. G. Mitchell, and K. A. Prather, 2011. Unique ocean-derived particles serve
 as a proxy for changes in ocean chemistry, *J. Geophys. Res.*, 116, D18310,
 doi:10.1029/2010JD015289.
 Ge, X., Wexler, A. S., and Clegg, S. L.: Atmospheric amines – Part I: A review, *Atmos.*
Environ., 45, 524–546, 2011.
 Geddes, S; Zahardis, J; Petrucci, GA., 2009. Chemical transformations of peptide
 containing fine particles: oxidative processing, accretion reactions and implications to the
 atmospheric fate of cell-derived materials in organic aerosol. *J Atmos. Chem.* 63:187–202
 DOI 10.1007/s10874-010-9161-2
 Genitsaris, S., Kormas, K. & Moustaka-Gouni, 2011. M. Airborne Algae and
 Cyanobacteria: Occurrence and Related Health Effects Savvas. *Front Biosci* 1, 772–787.
 González Benítez, J.M., Cape, J.N., Heal, M.R., van Dijk, N., Díez, A.V., 2009.
 Atmospheric nitrogen deposition in south-east Scotland: quantification of the organic
 nitrogen fraction in wet, dry and bulk deposition. *Atmospheric Environment* 43,
 4087e4094.
 Gonzalez Benitez, J.M., Cape, J.N., Heal, M.R., 2010. Gaseous and particulate water-
 soluble organic and inorganic nitrogen in rural air in southern Scotland, *Atmos. Environ.*
 44, 1506-1514.
 Goudie, A. S., and N. J. Middleton (2001), Saharan dust storms: Nature and
 consequences, *Earth Sci. Rev.*, 56(1– 4), 179–204.
 Graber, E. R. and Rudich, Y., 2006. Atmospheric HULIS: How humic-like are they? A
 comprehensive and critical review, *Atmos. Chem. Phys.*, 6, 729-753, doi:10.5194/acp-6-
 729-2006.
 Griffin, D., Westphal, D., and Gray, M., 2006. Airborne microorganisms in the African
 desert dust corridor over the mid-Atlantic ridge, *Ocean Drilling Program, Leg 209*,
Aerobiologia, 22, 211–226.

662 Hara, K. & Zhang, D., 2012. Bacterial abundance and viability in long-range transported
663 dust. *Atmos. Environ.* 47, 20–25.

664 Healy, R. M., Sciare, J., Poulain, L., Kamili, K., Merkel, M., Müller, T., Wiedensohler, A.,
665 Eckhardt, S., Stohl, A., Sarda-Estève, R., McGillicuddy, E., O'Connor, I. P., Sodeau, J. R.,
666 and Wenger, J. C, 2012.: Sources and mixing state of size-resolved elemental carbon
667 particles in a European megacity: Paris, *Atmos. Chem. Phys.*, 12, 1681-1700,
668 <https://doi.org/10.5194/acp-12-1681-2012>.

669
670 Healy, R. M., Sciare, J., Poulain, L., Crippa, M., Wiedensohler, A., Prévôt, A. S. H.,
671 Baltensperger, U., Sarda-Estève, R., McGuire, M. L., Jeong, C.-H., McGillicuddy, E.,
672 O'Connor, I. P., Sodeau, J. R., Evans, G. J., and Wenger, J. C, 2013: Quantitative
673 determination of carbonaceous particle mixing state in Paris using single-particle mass
674 spectrometer and aerosol mass spectrometer measurements, *Atmos. Chem. Phys.*, 13,
675 9479-9496, <https://doi.org/10.5194/acp-13-9479-2013>.

676
677 Healy, R.M., Evans, G.J., Murphy, M., Sierau, B., Arndt, J., McGillicuddy, E., O'Connor,
678 I.P., Sodeau, J.R., Wenger, J.C, 2015 Single-particle speciation of alkylamines in ambient
679 aerosol at five European sites. *Anal. Bioanal. Chem.*, 407, 5899–5909

680
681 Heard, D.E., Read, K.A., Methven, J., Al-Haider, S., Bloss, W.J., Johnson, G.P., Pilling,
682 M.J., Seakins, P.W., Smith, S.C., Sommariva, R., Stanton, J.C., Still, T.J., Ingham, T., B
683 rooks, B., De Leeuw, G., Jackson, A.V., McQuaid, J.B., Morgan, R., Smith, M.H.,
684 Carpenter, L.J., Carslaw, N., Hamilton, J., Hopkins, J.R., Lee, J.D., Lewis, A.C., Purvis,
685 R.M., Wevill, D.J., Brough, N., Green, T., Mills, G., Penkett, S.A., Plane, J.M.C., Saiz-
686 Lopez, A., Worton, D., Monks, P.S., Fleming, Z., Rickard, A.R., Alfarra, M.R., Allan, J.D.,
687 Bower, K., Coe, H., Cubison, M., Flynn, M., McFiggans, G., Gallagher, M., Norton, E.G.,
688 O'Dowd, C.D., Shillito, J., Topping, D., Vaughan, G., Williams, P., Bitter, M., Ball, S.M.,
689 Jones, R.L., Povey, I.M., O'Doherty, S., Simmonds, P.G., Allen, A., Kinnersley, R.P.,
690 Beddows, D.C.S., Dall'Osto, M., Harrison, R.M., Donovan, R.J., Heal, M.R., Jennings,
691 S.G., Noone, C., Spain, G., 2006. The North Atlantic Marine Boundary Layer Experiment
692 (NAMBLEX). Overview of the campaign held at Mace Head, Ireland, in summer 2002.
693 *Atmospheric Chemistry and Physics* 6, 241e2272.

694
695 Hill, K. A., Shepson, P. B., Galbavy, E. S., Anastasio, C., Kourtev, P. S., Konopka, A. &
696 Stirm, B. H. 2007 Processing of atmospheric nitrogen by clouds above a forest
697 environment. *J. Geophys. Res.* 112, 1–16. (doi:10. 1029/2006JD008002)

698
699 Jeong, C.-H., McGuire, M. L., Godri, K. J., Slowik, J. G., Rehbein, P. J. G., and Evans, G.
700 J., 2011. Quantification of aerosol chemical composition using continuous single particle
701 measurements, *Atmos. Chem. Phys.*, 11, 7027–7044, doi:10.5194/acp-11-7027-2011.

702
703 Kanakidou, M., Duce, R.A., Prospero, J.M., Baker, A.R., Benitez-Nelson, C., Dentener,
704 F.J., Hunter, K.A., Liss, P.S., Mahowald, N., Okin, G.S., Sarin, M., Tsigaridis, K., Uematsu,
705 M., Zamora, L.M., Zhu, T., 2012. Atmospheric fluxes of organic N and P to the global
706 ocean. *Glob. Biogeochem. Cycles* 26, 1–12. <https://doi.org/10.1029/2011GB004277>.

707
708 Kanakidou, M., Myriokefalitakis, S., Daskalakis, N., Fanourgakis, G., Nenes, A., Baker,

709 A.R., Tsigaridis, K., Mihalopoulos, N., 2016. Past, present, and future atmospheric
710 nitrogen deposition. *J. Atmos. Sci.* 73, 2039–2047. [https://doi.org/10.1175/JAS-D-](https://doi.org/10.1175/JAS-D-15-0278.1)
711 15-0278.1.

712
713 Kanakidou, M., Myriokefalitakis, S., Tsigaridis, K., 2018. Aerosols in atmospheric
714 chemistry and biogeochemical cycles of nutrients. *Environ. Res. Lett.* 13, 1–22. [https://doi.](https://doi.org/10.1088/1748-9326/aabcbd)
715 [org/10.1088/1748-9326/aabcbd](https://doi.org/10.1088/1748-9326/aabcbd)

716
717 Kellogg, C. A. and Griffin, D.W., 2006: Aerobiology and the global transport of desert dust,
718 *Trends Ecol Evol*, 21, 638–644, doi: 10.1016/j.tree.2006.07.004.

719
720 Krumins, V., Mainelis, G., Kerkhof, L. J. & Fennell, D. E., 2014. Substrate-Dependent
721 rRNA Production in an Airborne Bacterium. *Environ. Sci. Technol. Lett.* 376–381.
[doi:dx.doi.org/10.1021/ez500245y](https://doi.org/10.1021/ez500245y)

722
723 Jickells T, Baker AR, Cape JN, Cornell SE, Nemitz E., 2013. The cycling of organic
724 nitrogen through the atmosphere. *Philos Trans R Soc Lond B Biol Sci.* May
725 27;368(1621):20130115. doi: 10.1098/rstb.2013.0115.

726
727 Jickells, T. D., Buitenhuis, E. T., Altieri, K., Baker, A. R., Capone, D., Duce, R. A.,
728 Dentener, F., Fennel, K., Kanakidou, M., LaRoche, J., Lee, K., Liss, P. S., Middelburg, J.
729 J., Moore, J.K., Okin, G., Oschlies, A., Sarin, M., Seitzinger, S., Sharples, J., Singh, A.,
730 Suntharalingam, P., Uematsu, M., and Zamora, L. M. 2017: A re-evaluation of the
731 magnitude and impacts of anthropogenic atmospheric nitrogen inputs on the ocean, *Global*
732 *Biogeochem. Cy.*, 31, 289–305, <https://doi.org/10.1002/2016GB005586>

733
734 Lesworth, T., A. R. Baker, and T. Jickells, 2010. Aerosol organic nitrogen over the remote
735 Atlantic Ocean, *Atmos. Environ.*, 44, 1887–1893.

736
737 Lewis, A. C., Hopkins, J. R., Carpenter, L. J., Stanton, J., Read, K. A., and Pilling, M. J.,
738 2005. Sources and sinks of acetone, methanol, and acetaldehyde in North Atlantic marine
739 air, *Atmos. Chem. Phys.*, 5, 1963–1974 <http://www.atmos-chemphys.net/5/1963/2005/>

740
741 Mazar, Y., Cytryn, E., Erel, Y. & Rudich, Y., 2016. Effect of Dust Storms on the
742 Atmospheric Microbiome in the Eastern Mediterranean. *Environ. Sci. Technol.* **50**, 4194–
4202.

743
744 McConnell, C.L., Highwood, E.J., Coe, H., Formenti, P., Anderson, B., Osborne, S., Nava,
745 S., Desboeufs, K., Chen, G., Harrison, M.A.J., 2008. Seasonal variations of the physical
746 and optical characteristics of Saharan dust: Results from the Dust Outflow and Deposition
747 to the Ocean (SOLAS-DODO) experiment. *Journal of Geophysical Research-Atmospheres*
748 113, D14.

749
750 Myriokefalitakis, S., Fanourgakis, G., Kanakidou, M., 2017. The contribution of bioaerosols
751 to the organic carbon budget of the atmosphere. In: Karakostas, T.S. (Ed.),
752 *Perspectives on Atmospheric Sciences* pp. 845–851. Switzerland. [https://doi.org/10.](https://doi.org/10.1007/978-3-319-35095-0)
[1007/978-3-319-35095-0](https://doi.org/10.1007/978-3-319-35095-0).

- Miyazaki, Y., K. Kawamura, and M. Sawano, 2010. Size distributions of organic nitrogen and carbon in remote marine aerosols: Evidence of marine biological origin based on their isotopic ratios, *Geophys. Res. Lett.*, 37, L06803, doi:10.1029/2010GL042483.
- Miyazaki, Y., Kawamura, K., Jung, J., Furutani, H., and Uematsu, M., 2011 Latitudinal distributions of organic nitrogen and organic carbon in marine aerosols over the western North Pacific, *Atmos. Chem. Phys.*, 11, 3037-3049, doi:10.5194/acp-11-3037-2011.
- Moffet, R. C., de Foy, B., Molina, L. T., Molina, M. J., and Prather, K. A, 2008.: Measurement of ambient aerosols in northern Mexico City by single particle mass spectrometry, *Atmos. Chem. Phys.*, 8, 4499–4516, doi:10.5194/acp-8-4499-2008.
- Moreno, T., Querol, X., Castillo, S., Alastuey, A., Cuevas, E., Herrmann, L., Mounkaila, M., Elvira, J., Gibbons, W., 2006. Geochemical variations in aeolian mineral particles from the Sahara-Sahel Dust Corridor. *Chemosphere* 65 (2), 261e270
- Muller, C., Y. Iinuma, J. Karstensen, D. van Pinxteren, S. Lehmann, T. Gnauk and H. Herrmann, 2009. "Seasonal variation of aliphatic amines in marine sub-micrometer particles at the Cape Verde islands." *Atmospheric Chemistry And Physics* 9(24): 9587-9597.
- Murphy, S. M., Sorooshian, A., Kroll, J. H., Ng, N. L., Chhabra, P., Tong, C., Surratt, J. D., Knipping, E., Flagan, R. C., and Seinfeld, J. H., 2007: Secondary aerosol formation from atmospheric reactions of aliphatic amines, *Atmos. Chem. Phys.*, 7, 2313–2337, doi:10.5194/acp-7-2313-2007
- Pratt, K. A., Hatch, L. E., and Prather, K. A., 2009: Seasonal volatility dependence of ambient particle phase amines, *Environ. Sci. Technol.*, 43, 5276–5281,.
- Pratt, K., P. J. DeMott, J. R. French, Z. Wang, D. L. Westphal, A. J. Heymsfield, C. H. Twohy, A. J. Prenni & K. A. Prather, 2009 In situ detection of biological particles in cloud ice-crystals. *Nature Geoscience*, DOI: 10.1038/ngeo521
- Prospero, J., Blades, E., Mathison, G., and Naidu, R., 2005. Interhemispheric transport of viable fungi and bacteria from Africa to the Caribbean with soil dust, *Aerobiologia*, 21, 1–19.
- Rehbein, P. J. G., Jeong, C.-H., McGuire, M. L., Yao, X., Corbin, J.C., and Evans, G. J.. 2011: Cloud and fog processing enhanced gas-to-particle partitioning of trimethylamine, *Environ. Sci. Technol.*, 45, 4346, doi:10.1021/ES1042113
- Rijkenberg, M.J.A., Powell, C.F., Dall'Osto, M., Nielsdottir, M.C., Patey, M.D., Hill, P.G., Baker, A.R., Jickells, T.D., Harrison, R.M. and Achterberg, E.P., 2008. Changes in iron speciation following a Saharan dust event in the tropical North Atlantic Ocean. *Marine Chemistry* 110(1-2), 56-67.

- Russell, S. C., G. Czerwieniec, C. Lebrilla, H. Tobias, D. P. Fergenson, P. Steele, M. Pitesky, J. Horn, A. Srivastava, M. Frank, et al. 2004. Toward understanding the ionization of biomarkers from micrometer particles by bioaerosol mass spectrometry. *J. Amer. Soc. Mass Spectr.* 15 (6):900–909. doi:10.1016/j.jasms.2004.02.013
- Šantl-Temkiv, T., Finster, K., Dittmar, T., Hansen, B. M., Thyrhaug, R., Nielsen, N. W., and Karlson, U. G, 2013.: Hailstones: a window into the microbial and chemical inventory of a storm cloud, *PLoS ONE*, 8, e53550, doi:10.1371/journal.pone.0053550, 2013
- Šantl-Temkiv, T., Gosewinkel, U., Starnawski, P., Lever, M. & Finster, K., 2018. Aeolian dispersal of bacteria in southwest Greenland: their sources, abundance, diversity and physiological states. *Fems Microbiology Ecol.* doi:10.1093/femsec/fiy031
- Schneider, J., Freutel, F., Zorn, S. R., Chen, Q., Farmer, D. K., Jimenez, J. L., Martin, S. T., Artaxo, P., Wiedensohler, A., and Borrmann, S, 2011: Mass-spectrometric identification of primary biological particle markers and application to pristine submicron aerosol measurements in Amazonia, *Atmos. Chem. Phys.*, 11, 11415–11429, <https://doi.org/10.5194/acp-11-11415-2011>.
- Song, X. H., Hopke, P. K., Fergenson, D. P. and Prather, K. A., 1999 Classification of single particles analyzed by ATOFMS using an artificial neural network, ART-2A, *Anal. Chem.*, 71(4), 860–865.
- Sorooshian, A., Padró, L. T., Nenes, A., Feingold, G., McComiskey, A., Hersey, S. P., Gates, H., Jonsson, H. H., Miller, S. D., Stephens, G. L., Flagan, R. C., and Seinfeld, J. H., 2009. On the link between ocean biota emissions, aerosol, and maritime clouds: Airborne, ground, and satellite measurements off the coast of California, *Global Biogeochem. Cy.*, 23, GB4007, doi:10.1029/2009GB003464.
- Sullivan, R.C., Prather, K.A., 2005. Recent Advances in our understanding of atmospheric chemistry and climate made possible by on-line aerosol analysis instrumentation. *Analytical Chemistry* 77 (12), 3861e3885.
- Su, Y. X., Sipin, M. F., Furutani, H., and Prather, K. A., 2004. Development and Characterization of an Aerosol Time-of-Flight Mass Spectrometer with Increased Detection Efficiency, *Anal. Chem.*, 76, 712–719.
- Tittensor, D. P., Mora, C., Jetz, W., Lotze, H. K., Ricard, D., Vanden Berghe, E. & Worm, B. 2010 Global patterns and predictors of marine biodiversity across taxa. *Nature* 466, 1098–1101. (doi:10.1038/nature09329)
- Vařtillingom, M., Attard, E., Gaiani, N., Sancelme, M., Deguillaume, L., Flossmann, A. I., Amato, P., Delort, A.-M., 2012. Long-term features of cloud microbiology at the Puy de Dôme (France). *Atmos. Environ.* 56, 88–100. Vařtillingom, M., Deguillaume, L., Vinatier, V., Sancelme, M., Amato, P., Chaumerliac, N., Delort, A.-M., 2013. Potential impact of microbial activity on the oxidant capacity and organic carbon budget in clouds. *Proc. Natl. Acad. Sci. U. S. A.* 110, 559–564.

848
849 Violaki, K. and N. Mihalopoulos, 2010. "Water-soluble organic nitrogen (WSON) in size-
850 segregated atmospheric particles over the Eastern Mediterranean." *Atmospheric*
851 *Environment* 44(35): 4339-4345.
852
853 Violaki, K., Zarbas, P., Mihalopoulos, N., 2010. Long-term measurements of dissolved
854 organic nitrogen (DON) in atmospheric deposition in the Eastern Mediterranean: fluxes,
855 origin and biogeochemical implications. *Marine Chemistry* 120, 181–188

856 Yamaguchi, N., Ichijo, T., Baba, T. & Nasu, M. 2014. Long-range Transportation of
857 Bacterial Cells by Asian Dust. **36**, 145–151.

858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893

LIST OF TABLES AND FIGURES

Table 1. ATOFMS particle clusters identified from the Saharan dust soil samples and from the NAMBLEX and DODO ambient field studies.

Figure 1 Map of sampling locations: (A-B) shows the Saharan dust samples locations; Air mass arriving at Cape Verde (C) are for of a 5-day air mass back trajectory at mid-day at 500 m above the position of the ship during the dust event on 4 Feb 2006, (D) an marine Tropical (mT) air mass arriving at Mace Head (Ireland) at 12:00 LT on 8 December 2002 (Dall'Osto et al. 2004, Rijkenberg et al. 2008, Dall'Osto et al. 2010b). Aerosol transport time between point C and D was estimated to be 3-4 days from air mass trajectories analysis. Green and orange circles shows Saharan dust particles (Dust, orange) and primary biological aerosol particles (Bio, green). In point (D) these are enriched - with transport - in secondary organic aerosols (SOA, Dall'Osto et al., 2010a) and in Organic Nitrogen (ON, this study).

Figure 2 Positive (+) and negative (-) ART-2a ATOFMS mass spectra of Saharan dust particles and PBAP detected during Saharan origin (a, b, respectively); SOLAS-DODO cruise around Cape Verde (c, d, respectively) and NAMBLEX at Mace Head (e, f, respectively).

Figure 3 Positive (+) and negative (-) ART-2a ATOFMS mass spectra for PBAP NAMBLEX minus PBAP DODO (a) and PBAP NAMBLEX minus PBAP Saharan origin (b). Peaks below the horizontal line represents components less abundant in the NAMBLEX than the DODO/Origin; above the line are those with greater abundance.

Figure 4. Single particle positive and negative mass spectra of a single particle (2.3 μm) sampled during the NAMBLEX field study

Figure 5: (a) Black Carbon (Aethalometer AE16) concentrations (ng m^{-3}) (b) DMS concentrations (pptV), (c-d) ATOFMS counts for K-P-ON (biological particles) and Dust (Saharan Dust) for cP (European continental Polar), mT (marine Tropical) and mP-mA (marine Polar-Arctic) air masses (respectively); the three main air mass categories are further described in Dall'Osto et al., (2010b). ATOFMS counts are average counts per hour under mT and cP air masses (47 and 176 hours, respectively; total K-P-ON ATOFMS spectra 130 and 230, respectively).

942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959

960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978

	Number of ATOFMS mass spectra	Saharan dust particles (% by number)	Biological particles (% by number)	Others (% by number)
Saharan soil samples	4,205	55	10	>35
DODO (Tropical Ocean)	187,205	1.1	0.01	>98
NAMBLEX (North Atlantic Ocean)	191,504	1.0	0.01	>98

Table 1. ATOFMS particle clusters identified from the Saharan dust soil samples and from the NAMBLEX and DODO ambient field studies.

979
980
981
982
983
984
985

986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004

Figure 1. Map of sampling locations: (A-B) shows the Saharan dust samples locations; Air mass arriving at Cape Verde (C) are for of a 5-day air mass back trajectory at mid-day at 500 m above the position of the ship during the dust event on 4 Feb 2006, (D) an marine Tropical (mT) air mass arriving at Mace Head (Ireland) at 12:00 LT on 8 December 2002 (Dall'Osto et al. 2004, Rijkensberg el al. 2008, Dall'Osto et al. 2010b). Aerosol transport time between point C and D was estimated to be 3-4 days from air mass trajectories analysis. Green and orange circles shows Saharan dust particles (Dust, orange) and primary biological aerosol particles (Bio, green). In point (D) these are enriched - with transport - in secondary organic aerosols (SOA, Dall'Osto et al., 2010a) and in Organic Nitrogen (ON, this study).

(a)

(b)

(c)

(d)

(e)

(f)

Figure 2 Positive (+) and negative (-) ART-2a ATOFMS mass spectra of Saharan dust particles and PBAP detected during Saharan origin (a, b, respectively); SOLAS-DODO cruise around Cape Verde (c, d, respectively) and NAMBLEX at Mace Head (e, f, respectively).

1034
1035
1036

1037
1038
1039

1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050

Figure 3 Positive (+) and negative (-) ART-2a ATOFMS mass spectra for PBAP NAMBLEX minus PBAP DODO (a) and PBAP NAMBLEX minus PBAP Saharan origin (b). Peaks below the horizontal line represents components less abundant in the NAMBLEX than the DODO/Origin; above the line are those with greater abundance.

1051
1052
1053

1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077

Figure 4. Single particle positive and negative mass spectra of a single particle (2.3 μm) sampled during the NAMBLEX field study

1078
1079
1080
1081

1082
1083
1084
1085
1086
1087
1088
1089
1090

Figure 5: (a) Black Carbon (Aethalometer AE16) concentrations (ng m^{-3}) (b) DMS concentrations (pptV), (c-d) ATOFMS counts for K-P-ON (biological particles) and Dust (Saharan Dust) for cP (European continental Polar), mT (marine Tropical) and mP-mA (marine Polar-Arctic) air masses (respectively); the three main air mass categories are further described in Dall'Osto et al., (2010b). ATOFMS counts are average counts per hour under mT and cP air masses (47 and 176 hours, respectively; total K-P-ON ATOFMS spectra 130 and 230, respectively).