'If I die, I die, I don't care about my health'

Paudyal, Vibhu; MacLure, Katie; Forbes-McKay, Katrina; McKenzie, Myra; MacLeod, Joan; Smith, Ann; Stewart, Derek

DOI:

[10.1111/hsc.12850](https://doi.org/10.1111/hsc.12850)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Paudyal, V, MacLure, K, Forbes-McKay, K, McKenzie, M, MacLeod, J, Smith, A & Stewart, D 2020, 'If I die, I die, I don't care about my health': perspectives on self-care of people experiencing homelessness', *Health and Social Care in the Community*, vol. 28, no. 1, pp. 160-172. <https://doi.org/10.1111/hsc.12850>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: audyal V, MacLure K, Forbes-McKay K, et al. 'If I die, I die, I don't care about my health': Perspectives on self-care of people experiencing homelessness. *Health Soc Care Community*. 2020;28:160–172. <https://doi.org/10.1111/hsc.12850>, which has been published in final form at <https://doi.org/10.1111/hsc.12850>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

'If I die, I die, I don't care about my health': perspectives on self-care of people experiencing homelessness.

Vibhu Paudyal PhD^{1*} Institute of Clinical Sciences, University of Birmingham, UK

Katie MacLure PhD² Faculty of Health and Social Care, Robert Gordon University, Aberdeen, UK k.m.maclure@rgu.ac.uk

Katrina Forbes-McKay³ Applied Social Studies, Robert Gordon University, Aberdeen, UK k.e.forbes-mckay@rgu.ac.uk

Myra McKenzie, PhD, Faculty of Health and Social Care, Robert Gordon University, Aberdeen, UK m.mackenzie3@rgu.ac.uk

Joan MacLeod, Lead Pharmacist, NHS Grampian Lead Pharmacist (South Cluster) Aberdeen City Community Health Partnership, Aberdeen joan.macleod@nhs.net

Ann Smith, MRPharmS, NHS Grampian Lead Pharmacist (North Cluster) Aberdeen City Community Health Partnership, Aberdeen annsmith@nhs.net

Derek Stewart, PhD, Professor of Clinical Pharmacy and Practice, College of Pharmacy, Qatar University. d.stewart@qu.edu.qa

*Correspondence: Vibhu Paudyal PhD, Institute of Clinical Sciences, University of Birmingham, Birmingham, B15 2TT, UK. Phone: +44 (0)121 4142538; Email: v.paudyal@bham.ac.uk

'If I die, I die, I don't care about my health': perspectives on self-care of people experiencing homelessness.

Abstract:

Self-care, which refers to what people do to prevent disease and maintain good health, can alleviate negative health consequences of people experiencing homelessness. The aim of the study was to apply a theoretically informed approach in exploring engagement of people experiencing homelessness in self-care and to identify factors that can be targeted in future health and social care interventions. Qualitative semi-structured interviews were conducted with 28 participants opportunistically recruited from a specialist homelessness healthcare centre (SHHC) of North East Scotland, United Kingdom (UK). An interview schedule was developed based on the theoretical domains framework (TDF). Interviews were audio-recorded and transcribed verbatim. Six aspects of self-care were explored including (i) self-awareness of physical and mental health, (ii) health literacy including health seeking behaviour, (iii) healthy eating, (iv) risk avoidance or mitigation, (v) physical activity and sleep, and (vi) maintaining personal hygiene. Thematic analysis was conducted by two independent researchers following the Framework Approach. Participants described low engagement in self-care. Most barriers to engagement in self-care related to TDF domain 'environmental context and resources'. Participants often resorted to stealing or begging for food. Many perceived having low health literacy to interpret health related information. Visits to churches and charities to get a shower or to obtain free meals were commonplace. Participants expressed pessimism that there was 'nothing' they could do to improve their health and described perceived barriers often too big for them to overcome. Alienation, lack of social support, and the perception that they had done irreversible damage to their health prevented their involvement in self-care. The theme of 'social circle' held examples of both enabler and barriers in participants' uptake of risky

behaviours. Health and social services should work with persons experiencing homelessness in designing and delivering targeted interventions that address contextual barriers, multi-morbidity, health literacy and self-efficacy.

Keywords: Self-care, Homelessness, Health Behaviours

What is known about this topic?

- Ill health is a potential cause and consequence of homelessness but self-care can prevent and mitigate ill health
- A need to better understand self-care needs of people experiencing homelessness has recently been emphasised in health and social care policies across the UK
- There is a dearth of research exploring wider aspects of self-care amongst people experiencing homelessness as previous research has considered nutrition and diet, and risky behaviours in isolation.

What this paper adds?

- Study participants experiencing homelessness indicated low engagement in self-care across various domains such as diet, physical and mental health
- Low engagement in self-care was linked to a lack of resources, multi-morbidity, low health literacy and social influences
- Targeted interventions that address contextual barriers, multi-morbidity, health literacy and self-efficacy can improve participation in self-care

91 **Introduction**

92 In the United Kingdom (UK), people are considered homeless if they no longer
93 have a legal right to occupy their accommodation or if it would no longer be
94 reasonable (e.g. due to safety concerns) to continue living there (GOV.UK, 2015).
95 As such, homelessness takes many forms including sleeping rough, living in
96 derelict buildings, residing in temporary shelters, living in squats or sofa surfing
97 (Homeless link, 2016). Homelessness is on the rise across urban areas of the
98 Western World including the home countries of the UK (Scotland, England, Wales
99 and Northern Ireland) and has been linked to economic austerity. In 2018, nearly
100 twice as many people slept rough on any given night in England compared to
101 2010 (Homeless link, 2017; GOV.UK 2018). In Scotland, over 34,000 people
102 made homelessness applications to their local authority in 2016-17 requesting
103 accommodation (Scottish Public Health Observatory, 2018).

104

105 People experiencing homelessness face significant disadvantages in attaining and
106 maintaining a healthy lifestyle (Baggett *et al.* 2013; Aldridge *et al.* 2017;
107 University of Sheffield, 2012; Fazel *et al.* 2014). They do, therefore, experience
108 poor health outcomes with a prevalence of mental health illness, alcohol and drug
109 misuse, and communicable diseases higher than in the general population.
110 Opioid poisoning, heart failure, infectious diseases, and external causes such as
111 accidents, often contribute to the higher rate of mortality amongst street dwellers
112 (Hwang *et al.* 2005).¹⁰ Those occupying homeless shelters are also known to die
113 at an earlier age than the general population, with the average age of death
114 being 47 years (Hassanally *et al.* 2018).

115

116 Amongst multiple forms of homelessness, rough sleeping pre-disposes individuals
117 to much vulnerability. Government policies in the UK aim to tackle rough sleeping
118 through devolved administrations allowing England, Scotland, Wales and Northern
119 Ireland to develop their own legislations and strategies in preventing and

managing rough sleeping. In England, rough sleeping strategy was published in 2018 (Gov.UK, 2018) which aims to eliminate homelessness by 2027 by increasing bed spaces in city council accommodations, increasing access to substance misuse and mental health treatment and promoting joined-up care across sectors. 'Housing first' is one of the key interventions to supporting this strategy. Housing First aims to provide 'a stable, independent home and intensive personalised support and case management to homeless people with multiple and complex needs'. It aims to recognise housing as a matter of right than a reward (Homelessness Link, 2016). Further funding to tackle rough sleeping has been allocated by targeting areas with high proportion of rough sleeping in England. Such funding is allocated to offer dedicated support teams and securing additional bed spaces for people experiencing homelessness (Gov.UK, 2018). Other policy interventions to prevent rough sleeping includes Scottish Government's abolition of the priority needs assessment when offering accommodation to persons experiencing homelessness, entitling anyone finding themselves homelessness to settled accommodation and not just to families with children as was the case prior to the Act (The Scottish Government, 2012). 'Ending homelessness and rough sleeping: action plan' published by the Scottish government in 2018 aims to also tackle homelessness by tackling the root causes including additional support to people with adverse childhood experiences, and developing adversity and trauma informed workforce (The Scottish Government, 2018).

Addressing health inequalities requires a specific focus on the disadvantaged population. In particular, preventative services are known to be effective in alleviating the health impact of homelessness. Self-care, as defined by the World Health Organisation, is the ability of individuals, families and communities to promote health, prevent disease, maintain health, and to cope with illness and disability with or without the support of a healthcare provider (WHO, 2013), has been shown to prevent and mitigate ill health including long term illnesses. The

principles of self-care which can be applied to prevention and management of ill health are known to have arisen from a number of theoretical models such as the theory of self-regulation. Self-regulation models emphasise the importance of self-efficacy (Bandura, 2005), which relates to an individual's belief in their ability to learn and perform specific behaviours; and self-management (Lorig and Holman, 2003) which relates to adoption into practice of such behaviours. Self-efficacy often reinforces self-management. Self-management strategies, including patient-led self-care support groups, have also been shown to improve clinical outcomes amongst patients in a variety of long term illnesses (Minet *et al.* 2010), including effects on mortality, hospitalisation and quality of life (Ditewig *et al.* 2010). Supporting self-care can increase patient satisfaction of health and social care services, and enables greater integration of health and social care. In the UK, self-care features in the National Health Services plan as one of the key building blocks for a patient-centred health service (Department of Health, 2018).

It is important however to understand that within the spectrum of patient care, most care is shared care involving primary, secondary or tertiary health care and social care, and can involve a small or large components of self-care (Department of Health, 2005). In chronic and debilitating health conditions, people's participation in self-care is often minimal, whereas self-care occupies greater share in management of acute and non-debilitating conditions. Self-care practice is also dependent on context-specific factors including available resources and individuals hence should not be blamed for non-participation in self-care.

The seven pillars of self-care provide a framework to consider a wide range of activities relevant from the self-carer perspectives (International Self Care Foundation, 2018) (table 1). These include awareness of physical and mental health, health literacy and health seeking behaviour, healthy eating, hygiene, physical activity and sleep, and risk avoidance. The seven pillars of self-care framework, proposed by the International Self Care Foundations postulates that

unhealthy behaviours such as smoking, excess consumption of alcohol, poor diet and insufficient exercise often tend to cluster together (International Self Care Foundations, 2018). Similarly, healthy behaviours in the seven pillars also cluster together. **Therefore, promoting one healthy behaviour may motivate individuals to uptake other healthy behaviours.** We have previously used the seven pillars framework to identify appropriate interventions to promote self-care in offshore workers (Smith *et al.* 2018, Gibson Smith *et al.* 2018a).

Table 1 to appear here

Ill health is a potential cause and consequence of homelessness. A need to better understand supporting self-care and self-management for people experiencing homelessness, have recently been emphasised (The Queen's Nursing Institute, 2016). There is a dearth of research exploring wider aspects of self-care amongst people experiencing homelessness as previous research has looked at aspects such as nutrition and diet (Seale *et al.* 2016), risky behaviours (Roerecke *et al.* 2013), and health information seeking (McInnes *et al.* 2013) in isolation.

The aim of this study was to apply a theoretically informed approach in exploring engagement of people experiencing homelessness in undertaking self-care and to identify associated barriers that can be targeted in future health and social care interventions to promote self-care.

Method

Qualitative semi-structured, face-to-face, interviews were conducted with patients registered with an SHHC in North East of Scotland, UK between October 2015 and January 2016. This facility provides services to a patient population of approximately 400, of whom approximately 50% are on methadone therapy. Patients aged 18 years and over, presenting for the consultation during the data

collection days and those referred by the SHHC staff, were invited to participate. An effort was made to achieve variation in age and sex of the study participants. Researchers on site, who operated in pairs, provided further information about the research. Signed, informed consent was obtained by the researchers prior to interview commencement.

An interview schedule was developed based on the research aim, experience of the research team, available literature, and the Theoretical Domains Framework (TDF) (Cane *et al.* 2012; Francis *et al.* 2012). TDF is a framework consisting of 33 behavioural theories incorporated into 14 domains which allows researchers to identify barriers, facilitators or determinants of a particular behaviour. These include environmental context and resources, knowledge, skills, intentions, goals and behavioural regulations (table 2). TDF has been used extensively in qualitative studies to identify target behaviours for future interventions and to characterise implementation problems (Cane *et al.* 2012; Atkins *et al.* 2017). The researchers have previously used TDF in qualitative studies in identifying barriers of: access to primary healthcare by persons experiencing homeless (Gunner *et al.* 2019) and effective transition of care of such persons across services (Gibson-Smith *et al.* 2018b). When using TDF, it is imperative that the framework is used from the outset, including the development of an interview schedule, as the use of TDF at later stages of the research provides challenges in mapping the data against TDF domains (Cane *et al.* 2012; Atkins *et al.* 2017).

The interview schedule was reviewed for credibility by the research team, including a general practitioner (GP) and a nurse practitioner based at SHHC, a GP practice support pharmacist, a community pharmacist and academic health services researchers. Six pillars of self-care were explored (table 1). The seventh pillar of self-care 'rational and responsible use of medicines and products' was explored in another study (Paudyal *et al.* 2017).

Table 2 to appear here

The interview schedule was piloted amongst four participants. No change in the interview schedule was needed hence the pilot transcripts were analysed together with the main study transcripts. Interviews lasted a maximum of 30 minutes, with trained researchers, were audio-recorded and transcribed verbatim. Participants were recruited until data saturation was achieved, when no new themes emerge, as realised by the researchers during transcription and preliminary analysis of the data. Saturation was assumed based on the repetition of the themes from the subsequent interviews in the context of available data (Saunders *et al.* 2018).

The Framework Analysis technique (Ritchie *et al.* 2003) was used to guide the analytical process. The data pertaining to each pillar of self-care were coded into a matrix design based on the TDF (table 2). A framework was developed for each of the six pillars of self-care behaviours. Data relevant to these behaviours were mapped to the TDF domains and relevant themes under each domain were listed.

Researchers (VP, KM and DS) met to discuss initial coding after analysing the first four transcripts. Duplicate, independent checking of the transcripts and analysis was undertaken. Six undergraduate pharmacy students, including two visiting students, conducted duplicate independent analysis of the transcripts based on the coding.

Ethical and governance (R&D) approval for the study was granted by NHS East Midlands Committee (15/EM/0404) and NHS Grampian (2015RG005) respectively.

Results

Twenty-eight patients were interviewed, the majority of whom were male (n=21) with drug misuse being the key reason leading to homelessness (n=17) (table 3). The mean age was 42 years (range: 25-67 years). Most participants had faced homelessness for between six months and four years (n=17) (table 3).

Table 3 to appear here

Results from the thematic analysis are described below under each pillar of self-care. Narratives are presented alongside illustrative quotes in this section. The results are then mapped against TDF domains to relate the factors and barriers in relation to participant engagement with each pillar of self-care (table 4).

Self-awareness of physical and mental health

Most participants demonstrated knowledge and awareness of their health conditions and the impact of homelessness had on the onset and severity of their illnesses. Health conditions such as mental illness including drug and alcohol misuse, infections, ulcers, asthma, back pain and fatigue were commonly experienced as expressed by participants during the interviews. Participants described their capabilities and motivation to adopt better physical and mental health were compromised due to a lack of stable accommodation. Participants described feeling 'useless' and having suicidal ideation.

'I tried to kill myself about 5 times. It [homelessness] kicked your self-esteem to death.' 40 year old male

Most participants mentioned that they didn't attempt to change anything about their health while facing homelessness as health was not high amongst the list of priorities given the adversities they were facing.

293 'You care about your drugs, and at the time you think if you (I) die, you (I) die,...
294 you (I) don't care about your health...it doesn't matter, that's what you thought.
295 It's a dark place to be.' 34 year old male

296

297 Participants also mentioned having experienced a lot of stigma and discrimination
298 in society which negatively impacted their physical and mental well-being.

299

300 '...it (homelessness) affects you. People think you are a flaming drug addict,
301 scumbag, all they think ken [sic *know*], look at this mink, ken, sitting begging, get
302 a job, ken. It's nae good, you feel like snapping, and punching the *** out of folk,
303 but you cannae can you. You have got to hold yourself back. Especially on a
304 Saturday night...: '...I've been asked, like by a couple of guys, gay men for sex,
305 ken. Its nae good, they think you're homeless, they think you will do anything for
306 money, ken cause you are begging, ken.' 36 year old male

307

308 **Healthy eating**

309 Most participants described having adequate knowledge on the importance of
310 healthy meals to maintain good health. However, most reported poor access to
311 healthy meals due to lack of resources. One participant described experiences of
312 surviving on chocolates for several weeks. For a few participants, drugs or alcohol
313 would take precedence over food. Lack of appropriate space to prepare and cook
314 meals was commonly mentioned as a barrier. Visits to churches and charities for
315 free meals and accessing cheaper food sources, such as fast food chains, were
316 commonplace behaviours. Participants often had to rely on food given by those
317 passing by when sleeping rough.

318

319 'When you're sitting on the street folk would give you a coke and a sandwich or
320 something, sometimes I would have four, or five or six sandwiches that I would go
321 through but the nutritional (value) is low, so you, you lose a lot, your weight just

322 falls off you... Just nae eating right and taking drugs and alcohol it's just, the
323 weight just falls off you.' 34 year old male

324

325 One participant described the extreme experience of hunger lasting several days
326 where he had no other option than to steal food from a retailer.

327

328 'Ehm, basically I never ate for days, and then it would get to the point that I would
329 get so hungry that I would need to steal a sandwich or something out a shop.'39
330 year old male

331

332 Table 4 to appear here

333

334 **Health literacy and seeking health information**

335 Some participants described experiences of actively seeking health information
336 from their health and social care professionals for a diverse range of health issues
337 including substance misuse. Participants demonstrated awareness of where to
338 seek health information, with the preferred source of information being GPs and
339 nurses at the SHHC and social service counsellors. Participants who had very
340 recently moved to temporary accommodation also mentioned use of the internet
341 to seek health information. However, most participants identified themselves as
342 having low literacy skills and often not being able to interpret health information.

343

344 'I've looked up the internet [about health condition] a couple of times but I don't
345 understand it.' 43 year old Male

346

347 Some participants expressed feeling emotional in relation to discussing their
348 health with their healthcare professionals. This was due to their health being
349 closely linked to the life circumstances they were facing and being uncomfortable
350 discussing such issues with other people.

351

352 'I just don't like new people [healthcare professionals]. I just don't like having to
353 kinda having to repeat everything. I get myself in a muddle and I get all stressed
354 out.' 33 year old female

355 **Personal Hygiene**

356 Maintaining good personal hygiene was a priority for some participants. Those
357 who demonstrated motivation and intentions to remain free from substance
358 misuse mentioned that there was no excuse for not maintaining personal hygiene
359 even when sleeping rough. Some mentioned being advised by charities regarding
360 where to go on a daily basis for a shower. Others would pop in to friends' houses,
361 railway stations, and fast food restaurants to get a 'wash'. Some participants
362 described experiencing insecurity in public clean up facilities which prevented
363 them from using them on a regular basis.

364

365 'We've had to go to McDonalds to have a wash and stuff like that, there is, you will
366 find places, ken fit [sic *know what*] I mean. There is no excuse to be sitting in
367 some state some people are in. OK, your clothes are getting ripped cause you are
368 sitting on pavements all day and stuff like that, ken, you are going to look a mess,
369 doesn't mean that you have got to be stinking, a stinking mess, you know. But it is
370 hard.' 47 year old female

371

372 Other participants described that maintaining good hygiene was challenging due
373 to other life priorities and getting housed in stable accommodation was the only
374 way to maintain good personal hygiene. Therefore some participants were not
375 being personally motivated to wash or dress themselves properly even when they
376 had options and facilities available. Participants expressed emotions when
377 mentioning accounts of being stigmatised because of their poor personal hygiene.

378

379 '...you don't care about long hair, if it's greasy, you don't care if you walk onto a
380 bus and everybody walks off the bus 'cause you smell.' 34 year old male

381

Risk avoidance or mitigation

Most participants admitted to their current or past use of illicit drugs, hazardous drinking of alcohol, and smoking habits. Most participants who admitted substance misuse also mentioned being on opioid replacement therapy (ORT). Social influences were described as key to participants choosing to adopt or give up risky behaviours. However, some participants described lack of self-regulation and behavioural control in helping them to come off the substances. Some also described alcohol and substance misuse as a coping mechanism.

'Never succeeded, I've tried to give it [alcohol and drug misuse] up but think it's the only thing that keeps my nerves tied together just now so, but just, I think I've done too much damage to repair it anyway so If I'm going to die now, I'm gonna die. I've made my choices, so I've made my bed I'll have to lie in it, sorta thing. That's about it.' 39 year old male

Participants described going to extremes to obtain money for illicit substances including robbery and prostitution. Some participants described accounts of successfully giving up risky behaviours such as substance and alcohol misuse.

'When I was on drugs and [I felt] that...if I died one day, I died and then well my life was... when you're on drugs, you see [drugs] making these people die around you. I've been lying in a bed and the boy next to me was dead. It's just, oh well, I'd go into his pockets and take his money and drugs and walk out the house. Aye, that's the way you are. It's a weird, it's a horrible thing drugs. It does it to you, ken [sic *know*], heroin.' 54 year old male

'I've been totally clean off everything for just over a year in April.. I ended up being in mental health hospital for almost three months. That was like an extended rehab sorta thing, so I like stayed away from everything for 3 months which gave

411 me a fighting chance and I've been clean ever since then. Think I've had one drink
412 in last New Year since then.' 39 year old male

413

414 **Physical activity and sleep**

415 Most participants mentioned engagement in physical activity was beyond their list
416 of priorities. Morbidity, disability or lack of accommodation preventing them
417 taking up physical activities. A few participants expressed disinterest and lack of
418 motivation in engaging in the discussion about aspects of physical activity during
419 the interviews and such lack of self-efficacy was linked to adverse life
420 circumstances.

421

422 'No I've no done nothing (physical exercise) – just nothing at all. Just can't get
423 motivated. That's how I'm waiting to see the psychiatrist and to get on my anxiety.
424 'Cause I couldn't even come down here. Couldn't leave the house or and that's
425 what made us depressed 'cause I like just going right out, ken getting up and
426 going out. I couldn't get out on my own. 34 year old female

427

428 Some participants living in temporary, council offered accommodation described
429 the use of a gym or walking to maintain health. Many described having very little
430 or no sleep while facing homelessness. Lack of stable accommodation was a key
431 barrier to attaining quality sleep. Some mentioned using illicit substances to
432 enable better sleep. Participants described stigma, theft and violence while
433 sleeping rough.

434

435 '... you're sleeping in car parks and everything, freezing cold, ...so you don't get to
436 sleep ken [sic know] and folk say, "oh you should come to mines if you're ever
437 stuck", but you never bother, ken, because you knock on somebody's door in the
438 middle of the night they're hardly happy to see you but, eh aye, it was an absolute
439 nightmare because there was no churches letting people in or anything.' 49 year
440 old female

Discussion and conclusion

This study has explored homeless from the participants' perspectives on wider aspects of self-care through the use of theory. Engagement in self-care was perceived to be low across several pillars of self-care theory including healthy eating, health information seeking, maintenance of personal hygiene, risk avoidance and mitigation, and maintenance of personal hygiene mainly due to context, resource specific barriers and lack of self-efficacy due to poor perceived health and adverse life circumstances. The use of TDF allowed the barriers and facilitators of participant engagement in self-care to be mapped across domains that could be targeted in future interventions. Most of the barriers related to non-engagement in self-care identified in this study centred on the 'environmental context and resources' domain of TDF and this included lack of stable accommodation. Participants often expressed lack of motivation to uptake healthy behaviours, often compromised by other life priorities.

In this study, participants alluded to the role of charities and social support in enabling them to undertake self-care such as in enabling a healthy diet or maintaining personal hygiene. There is scope for health and social care professionals to offer such provision at the health or social care centres or to make referrals to services. Emphasis has been placed on healthcare professionals to recognise and screen for nutritional need of people experiencing homelessness and their families (The Queen's Nursing Institute, 2016). Participants in this study expressed low health literacy and hence health and social care professionals need to be aware of these barriers when referring people experiencing homelessness to sources of information. Participants described being emotionally vulnerable when discussing their health and self-care issues with the healthcare professionals because issues were closely linked to their life circumstances. Our previous study identified that rapport with health and social care workers was a key factor in

homeless people's preference to use SHHC facilities, even when they had relocated to permanent accommodation (Paudyal *et al.* 2018).

Most of the barriers to the uptake of self-care including healthy eating and physical activity were linked by participants of this study to their lack of accommodation. In England, the Homeless Reduction Act is coming into effect in 2018 (Paudyal and Saunders, 2018) following a similar homelessness legislation in Wales in 2014 (The Welsh government, 2014). While the effectiveness of this Act is yet to be evaluated, the Act mandates health and social care services to refer people who are at risk of or facing homelessness to local authorities for the provision of accommodation. Policy interventions such as **The Housing First initiative (Homeless Link, 2016) are likely to address context and resource related barriers. Research evidence demonstrates that Housing First initiative decreases homelessness and increases housing retention rates and decreases the use of emergency health services, and emergency shelters, particularly in people with severe mental health and substance misuse problems (Woodhall-Melnik and Dunn, 2018).** Screening people experiencing homelessness for mental and physical health conditions during their housing needs assessment provides an effective strategy for early intervention (Weinstein *et al.* 2013).

There is a lack of previous literature exploring wider aspects of self-care within a population of people experiencing homelessness as the literature often tends to focus on a single behaviour at a time. Therefore, only a limited comparison to previous literature could be undertaken. A recent study in the United States showed that poorer self-rated health was associated with the desire to reduce hazardous drinking and increase fruit and vegetable consumption in this population (Taylor *et al.* 2016). A recent review of the literature demonstrated evidence of malnutrition including saturated fat, low fruit and vegetable intake and numerous micro-nutrient deficiencies, amongst people experiencing

homelessness, often leading to physical and mental health consequences (Sprake *et al.* 2014). Furthermore, that review noted the search for food often takes priority over healthcare and access to medicines (Paudyal *et al.* 2017).

Strengths and limitations

This study is, to the best of our knowledge, the first to explore people experiencing homelessness' perspectives on wider self-care aspects. Duplicate and independent analysis of the data enabled trustworthiness of the findings. Use of theory enabled mapping of the key barriers and facilitators of engagement in self-care across domains of the TDF, the pillars of self-care and provides specific targets for future interventions.

This study has some limitations. Some participants of this study had recently been temporarily or permanently housed despite the use of the SHHC and were waiting to relocate to mainstream general practices. Such participants provided their retrospective accounts. The study participants were predominantly male, however, this reflects the data trend of persons experiencing homelessness. Only the patients with good rapport with the healthcare staff were included. This approach was used to ensure the safety of both the researchers and research participants. Therefore, views may not be representative of all participants from the study setting. In addition, the researchers used the transcripts and initial analysis when assuming data saturation. As reported in the literature, this is a common barrier to ascertaining saturation in qualitative studies (Saunders *et al.* 2018).

Practice and research implications

The results of this study suggest that promotion of self-care amongst people experiencing homelessness requires addressing the resource-related barriers such as provision of stable accommodation and their co-morbidities. Such barriers

collectively compromises their self-efficacy and motivation to uptake self-care. In addition, the results provide recommendations for the development, implementation and evaluation of health and social care interventions that can positively impact on their self-confidence, belief about capabilities, intentions and behavioural regulations. The Medical Research Council, UK provides a framework (Craig *et al.* 2013) for development of complex interventions. This study provides targeted areas for multi-faceted interventions and the data provides a valuable foundation on which to base development of interventions. It has been postulated that unhealthy behaviours, such as poor diet, drugs and alcohol misuse, tend to “cluster” together in individuals (International Self Care Foundations, 2018), so as the healthy behaviours cluster amongst certain sectors of the population. Such multi-faceted targeted interventions can be delivered at temporary accommodations, charities, outreach services, or health and social care settings that can enable homeless population to develop their self-confidence, improve health seeking behaviour and their intentions to lead a healthy lifestyle. People sleeping rough will also benefit from provision of healthy diet, tailored health related information, facilities for personal hygiene under one roof.

Poor mental health including the experience of stigma and discrimination was a recurrent theme in the data. Poor mental health can often be the cause and consequences of homelessness (Bowen *et al.* 2019). Various barriers to people’s access to mental health services have been described in the literature with concurrent substance misuse and history of self-harm often excluding patients access to mental health services (Gunner *et al.* 2019). Hence, people experiencing homelessness may benefit from multi-morbidity models of case management, and these are best embedded as part of housing-related interventions such as the Housing First initiative (Aubry *et al.* 2015). The Assertive Community Treatment (ACT) is one example where multi-morbidity including mental health and substance misuse is managed by a multidisciplinary

team with home based treatment and out of hours availability by also integrating peer support (Nugter *et al.* 2016). Stigma and discrimination were also commonly cited in relation to societal attitude towards homelessness and people experiencing homelessness. However, previous research showed that people experiencing homelessness also face stigma and discrimination when accessing health services (Paudyal *et al.* 2018; Gunner *et al.* 2019). Anti-stigma interventions for healthcare professionals such as the 'targeting the roots of healthcare provider stigma' which involves improving the ability of healthcare professionals to cope with the feelings and emotions when working with vulnerable patients; improving their competence and the confidence of staff; and addressing the lack of awareness of one's own prejudices have been shown to minimise perceived stigma and discrimination (Knaak and Patten, 2016). In addition, health and social care workers are able to better support people experiencing homelessness when they have the knowledge of patients' backgrounds and life circumstances (Padget and Henwood, 2012).

While health professionals based in specialist homelessness healthcare facilities may be more aware of the factors associated with non-engagement of people experiencing homelessness in self-care, as identified in this study, many homeless patients use mainstream services or may not come in contact with healthcare staff. Wider awareness will enable health promotion and self-care improvement in this population. Health and social services should avoid blaming individuals for their behaviours and low perceived engagement in self-care as often many of these barriers including context and societal factors need system based approach for change.

Conclusion

Low engagement in self-care was noted amongst the study participants. There is scope for targeted interventions focused on specific determinants to promote

each pillar of self-care by addressing contextual barriers, physical and mental co-morbidities, health literacy and people's self-efficacy. Health and social services should work with persons experiencing homelessness in designing and delivering targeted interventions.

Acknowledgements: We would like to thank Lynsey Cumming, Emma Petrie, Caitlin Stewart and Heather Wright for data collection and analysis, and Carl Mopho and Kathleen MacMillan from University of Dalhousie, Canada for their part in the data analysis.

Funding: This study was funded by Robert Gordon University

References

- Aldridge, R. W., Story, A., Hwang, S. W., Nordentoft, M., Luchenski, S. A., Hartwell, G & Hayward, A. C. (2017). Morbidity and mortality in homeless individuals, prisoners, sex workers, and individuals with substance use disorders in high-income countries: a systematic review and meta-analysis. *The Lancet* 391(10117):241-50. [https://doi.org/10.1016/S0140-6736\(17\)31869-X](https://doi.org/10.1016/S0140-6736(17)31869-X)
- Atkins L, Francis J, Islam R, O'Connor D, Patey A, Ivers N, Foy R, Duncan EM, Colquhoun H, Grimshaw JM, Lawton R. (2017). A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implementation Science*: 12(1):77. <https://doi.org/10.1186/s13012-017-0605-9>
- Atkins, L., Francis. J., Islam. R., O'Connor, D., Patey. A., Ivers. N., Foy, R., Duncan, E.M., Colquhoun, H., Grimshaw, J.M., Lawton, R. (2017). A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implementation Science* 12(1):77. <https://doi.org/10.1186/s13012-017-0605-9>
- Aubry, T., Goering, P., Veldhuizen, S., Adair, C.E., Bourque, J., Distasio, J., Latimer, E., Stergiopoulos, V., Somers, J., Streiner, D.L. and Tsemberis, S. (2015). A multiple-city RCT of housing first with assertive community treatment for homeless Canadians with serious mental illness. *Psychiatric Services*, 67(3), pp.275-281. doi: 10.1176/appi.ps.201400587.
- Baggett, T. P., Hwang, S. W., O'Connell, J. J., Porneala, B. C., Stringfellow, E. J., Orav, E. J., Rigotti, N. A. (2013). Mortality among homeless adults in Boston: shifts in causes of death over a 15-year period. *JAMA internal medicine*, 173(3), 189-195. Doi:10.1001/jamainternmed.2013.1604
- Bandura A. (2005). The primacy of self-regulation in health promotion. *Applied Psychology-an International Review*. 54[2]:245-254. <http://dx.doi.org/10.1111/j.1464-0597.2005.00208.x>
- Bowen, M., Marwick, S., Marshall, T., Saunders, K., Burwood, S., Yahyouche, A., Stewart, D. and Paudyal, V. (2019). Multimorbidity and emergency department visits by a homeless population: a database study in specialist general practice. *British Journal of General Practice*, p.bjgp19X704609. DOI: <https://doi.org/10.3399/bjgp19X704609>
- Cane, J., O'Connor, D., & Michie, S. (2012). Validation of the theoretical domains framework for use in behaviour change and implementation research. *Implementation science*, 7(1), 37. <https://doi.org/10.1186/1748-5908-7-37>
- Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2013).
- Department of communities and Local Government (2015). Statutory Homelessness in England. Available: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/437273/201503_Statutory_Homelessness.pdf. Accessed 20 March 2016.
- Department of Health (2018). Self care. Available: <http://webarchive.nationalarchives.gov.uk/20090216230833/http://www.dh.gov.uk/en/Healthcare/Selfcare/index.htm>. Accessed 23 April 2018.
- Department of Health. (2005). Self care: A real choice; Self care: A practical option. Department of Health. London.

640 Developing and evaluating complex interventions: the new Medical Research Council
641 guidance. *International journal of nursing studies*, 50(5), 587-592.
642 <https://doi.org/10.1136/bmj.a1655>

- 643 • Ditlew, J. B., Blok, H., Havers, J., & van Veenendaal, H. (2010). Effectiveness of self-
644 management interventions on mortality, hospital readmissions, chronic heart failure
645 hospitalization rate and quality of life in patients with chronic heart failure: a systematic
646 review. *Patient education and Counseling*, 78(3), 297-315. doi: 10.1016/j.pec.2010.01.016.
- 647 • Fazel, S., Geddes, J. R., & Kushel, M. (2014). The health of homeless people in high-income
648 countries: descriptive epidemiology, health consequences, and clinical and policy
649 recommendations. *The Lancet*, 384(9953), 1529-1540. 10.1016/S0140-6736(14)61132-6
- 650 • Francis, J. J., O'Connor, D., & Curran, J. (2012). Theories of behaviour change synthesised
651 into a set of theoretical groupings: introducing a thematic series on the theoretical domains
652 framework. *Implementation Science*, 7(1), 35. DOI: 10.1186/1748-5908-7-35
- 653 • Gibson Smith, K.L., Paudyal, V., Klein, S. and Stewart, D.C. (2018). Health, self-care and the
654 offshore workforce: opportunities for behaviour change interventions, an epidemiological
655 survey. *Rural and Remote Health* [online], 18(2). doi: 10.22605/RRH4319
- 656 • Gov. UK. (2018). The rough sleeping strategy. Available
657 <https://www.gov.uk/government/publications/the-rough-sleeping-strategy>. Accessed 14 May
658 2019.
- 659 • Gov.UK. (2018). Funding allocated for councils to help rough
660 sleepers [https://www.gov.uk/government/news/funding-allocated-for-councils-to-help-rough-](https://www.gov.uk/government/news/funding-allocated-for-councils-to-help-rough-sleepers)
661 [sleepers](https://www.gov.uk/government/news/funding-allocated-for-councils-to-help-rough-sleepers) . Accessed 20 July 2019.
- 662 • GOV.UK. (2019) Homelessness statistics: Statutory homelessness in England. Available at:
663 <https://www.gov.uk/government/collections/homelessness-statistics>. Accessed 16 March 2019.
- 664 • Gunner, E., Chandan, S.K., Marwick, S., Saunders, K., Burwood, S., Yahyouche, A. and
665 Paudyal, V. (2019). Provision and accessibility of primary healthcare services for people who
666 are homeless: a qualitative study of patient perspectives in the UK. *British Journal of General*
667 *Practice*. p.bjgp19X704633. DOI: <https://doi.org/10.3399/bjgp19X704633>
- 668 • Hassanally, K., Asaria, M. (2018). Homeless mortality data from East London. *London Journal*
669 *of Primary Care*, 4:1-4. <https://doi.org/10.1080/17571472.2018.1458443>
- 670 • Homeless link (2016). About homelessness. Available: [http://homeless.org.uk/about-](http://homeless.org.uk/about-homelessness#UjxuEj_mDBY)
671 [homelessness#UjxuEj_mDBY](http://homeless.org.uk/about-homelessness#UjxuEj_mDBY). Accessed 05 March 2016.
- 672 • Homeless Link. 2016. Housing First in England: The principles.
673 [https://www.homeless.org.uk/sites/default/files/site-](https://www.homeless.org.uk/sites/default/files/site-attachments/Housing%20First%20in%20England%20The%20Principles.pdf)
674 [attachments/Housing%20First%20in%20England%20The%20Principles.pdf](https://www.homeless.org.uk/sites/default/files/site-attachments/Housing%20First%20in%20England%20The%20Principles.pdf). Accessed 14 May
675 2019.
- 676 • Homelessness link (2017). Rough sleeping — our analysis. Available at:
677 [http://www.homeless.org.uk/facts/homelessness-in-numbers/rough-sleeping/rough-sleeping-](http://www.homeless.org.uk/facts/homelessness-in-numbers/rough-sleeping/rough-sleeping-our-analysis)
678 [our-analysis](http://www.homeless.org.uk/facts/homelessness-in-numbers/rough-sleeping/rough-sleeping-our-analysis). (accessed 20 April 2017).
- 679 • Hwang, S. W., Tolomiczenko, G., Kouyoumdjian, F. G., & Garner, R. E. (2005). Interventions to
680 improve the health of the homeless. *American journal of preventive medicine*, 29(4), 311-311.
681 DOI: <https://doi.org/10.1016/j.amepre.2005.06.017>
- 682 • International Self Care Foundations (2018). The seven pillars of self care. Available:
683 <http://isfglobal.org/practise-self-care/the-seven-pillars-of-self-care/>. Accessed Accessed 23
684 April 2018.
- 685 • Knaak, S., Patten, S. (2016). A grounded theory model for reducing stigma in health
686 professionals in Canada. *Acta Psych Scan*, 134:53-62. doi: 10.1111/acps.12612.
- 687 • Lorig, K.R., Holman, H.R. (2003). Self-management education: History, definition, outcomes,
688 and mechanisms. *Annals of Behavioral Medicine*. 26[1]:1-7. DOI:
689 10.1207/S15324796ABM2601_01
- 690 • McInnes, D. K., Li, A. E., & Hogan, T. P. (2013). Opportunities for engaging low-income,
691 vulnerable populations in health care: a systematic review of homeless persons' access to and
692 use of information technologies. *American journal of public health*, 103(S2), e11-e24. DOI:
693 10.2105/AJPH.2013.301623.
- 694 • Minet, L., Møller, S., Vach, W., Wagner, L., & Henriksen, J. E. (2010). Mediating the effect of
695 self-care management intervention in type 2 diabetes: a meta-analysis of 47 randomised
696 controlled trials. *Patient Education and Counseling*, 80(1), 29-41. doi:
697 10.1016/j.pec.2009.09.033.
- 698 • Nugter, M.A., Engelsbel, F., Bähler, M., Keet, R. and van Veldhuizen, R., 2016. Outcomes of
699 FLEXIBLE assertive community treatment (FACT) implementation: a prospective real life
700 study. *Community Mental Health Journal*, 52(8), pp.898-907. [https://doi.org/10.1007/s10597-](https://doi.org/10.1007/s10597-015-9831-2)
701 [015-9831-2](https://doi.org/10.1007/s10597-015-9831-2)
- 702 • Padgett, D.K. & Henwood, B.F. (2012). Qualitative research for and in practice: Findings from
703 studies with homeless adults who have serious mental illness and co-occurring substance
704 abuse. *Clinical Social Work Journal*, 40(2), 187-193. 10.1007/s10615-011-0354-1

- Paudyal, V., MacLure, K., Buchanan, C., Wilson, L., Macleod, J., & Stewart, D. (2017). 'When you are homeless, you are not thinking about your medication, but your food, shelter or heat for the night': behavioural determinants of homeless patients' adherence to prescribed medicines. *Public health*, 148, 1-8. doi: 10.1016/j.puhe.2017.03.002.
- Paudyal, V., & Saunders, K. (2018). Introduction of homeless reduction act in England: impact on health services. *Lancet*, 392 (10143), 195-197.
- Ritchie, J., Spencer, L., & O'Connor, W. (2003). Carrying out qualitative analysis. *Qualitative research practice: A guide for social science students and researchers*. SAGE Publications. London.
- Roerecke, M., & Rehm, J. (2013). Alcohol use disorders and mortality: a systematic review and meta-analysis. *Addiction*, 108(9), 1562-1578. doi: 10.1111/add.12231.
- Saunders, B., Sim, J., Kingstone, T., Baker, S., Waterfield, J., Bartlam, B., Burroughs, H., Jinks C. (2018). Saturation in qualitative research: exploring its conceptualization and operationalization. *Quality & quantity*, 1;52(4):1893-907. <https://doi.org/10.1007/s11135-017-0574-8>
- Seale, J. V., Fallaize, R., & Lovegrove, J. A. (2016). Nutrition and the homeless: the underestimated challenge. *Nutrition research reviews*, 29(2), 143-151. DOI: 10.1017/S0954422416000068
- Smith KG, Paudyal V, Quinn F, Klein S, Stewart D. (2018). Offshore workers and health behaviour change: an exploration using the Theoretical Domains Framework. *International maritime Health*, 69(4):248-56. doi: 10.5603/IMH.2018.0040
- Smith, K. G., Paudyal, V., MacLure, K., Forbes-McKay, K., Buchanan, C., Wilson, L., Stewart, D. (2018). Relocating patients from a specialist homeless healthcare centre to general practices: a multi-perspective study. *Br J Gen Pract*, 68 (667), e105-e113. doi: 10.3399/bjgp18X694577
- Sprake, E.F., Russell, J.M., Barker, M.E. (2014). Food choice and nutrient intake amongst homeless people. *Journal of Human Nutrition and Dietetics*, 27(3):242-50. doi: 10.1111/jhn.12130
- Taylor EM, Kendzor DE, Reitzel LR, Businelle MS. (2016). Health risk factors and desire to change among homeless adults. *American journal of health behaviour*, 1;40(4):455-60. doi: 10.5993/AJHB.40.4.7
- The Queen's Nursing Institute (2016). Food, Nutrition and Homelessness. Available: https://www.qni.org.uk/wp-content/uploads/2016/09/nutrition_guidance.pdf. Accessed 23 April 2018.
- The Scottish Government. (2012). The Homelessness (Abolition of Priority Need Test) (Scotland) Order 2012. Available: <http://www.legislation.gov.uk/ssi/2012/330/contents/made>. Accessed 14 May 2018.
- The Scottish Government. (2018). Ending homelessness and rough sleeping: action plan. <https://www.gov.scot/publications/ending-homelessness-together-high-level-action-plan/>. Accessed 20 July 2019.
- The Scottish Public Health Observatory (2018). Homelessness: demographics. Available: <http://www.scotpho.org.uk/life-circumstances/homelessness/data/demographics/>. Accessed 23 April 2018.
- The University of Sheffield and Crisis (UK) (2012). An analysis of the mortality of homeless people in early twenty-first century England. Available: https://www.crisis.org.uk/media/236799/crisis_homelessness_kills_es2012.pdf. Accessed 23 April 2018.
- The Welsh Government. (2014). The Housing (Wales) Act 2014. Available: <http://www.legislation.gov.uk/anaw/2014/7/contents>. Accessed 14 May 2018.
- Weinstein, L. C., LaNoue, M., Collins, E., Henwood, B. F., & Drake, R. E. (2013). Health care integration for formerly homeless people with serious mental illness. *Journal of Dual Diagnosis*, 9(1), 72-77. <https://doi.org/10.1080/15504263.2012.750089>
- Woodhall-Melnik, J.R. and Dunn, J.R. (2016). A systematic review of outcomes associated with participation in Housing First programs. *Housing Studies*, 31(3), pp.287-304. <https://doi.org/10.1080/02673037.2015.1080816>
- World Health Organisation. Self-care for health (2013). Available <http://apps.who.int/iris/bitstream/handle/10665/205887/B5084.pdf;jsessionid=9A88B65500EC3FF7EDF2F63A87D5844A?sequence=1>. Accessed 23 April 2018.

765 **Table 1: Pillars of self-care**

Pillars of Self-care	Example of topic guide prompts based on TDF
Self-awareness of physical and mental condition and use of health services	Awareness about their health and illness, use of healthcare services
Healthy eating	Seeking and consuming healthy food and balanced diet
Health literacy and seeking health related information	Whether participants actively seek health related information, ability to access and interpret information
Good hygiene	Maintenance of personal hygiene and associated barriers and facilitators
Physical activity and sleep	Physical activity levels, associated barriers and facilitators
Risk avoidance or mitigation	Substance misuse including drugs, alcohol and illicit substances
Rational and responsible use of medicines, services and products	Using medicines, services and products responsibly when necessary

766 Source: International Self Care Foundation (2018)

767
768

Table 2: Theoretical domains framework (TDF)

TDF domains and descriptions	
1. Knowledge	Knowledge of condition /scientific rationale, Procedural knowledge, Knowledge of task environment
2. Skills	Skills, skill development, Competence, Ability, Interpersonal skills, Practice Skill assessment
3. Social/ Professional Role and Identity	Professional identity, Professional role, Social identity, Identity, Professional boundaries, Professional confidence Group identity, Leadership, Organisational commitment
4. Beliefs about Capabilities	Self-confidence, Self-confidence Perceived competence Self-efficacy Perceived behavioural control Beliefs Self-esteem Empowerment Professional confidence
5. Optimism	Optimism Pessimism Unrealistic optimism, Identity
6. Beliefs about Consequences	Outcome expectancies, beliefs, anticipated regret, consequents
7. Reinforcement	Incentives, Rewards (proximal/distal, valued/not valued, probable/improbable), Incentives, Punishment, Consequents, Reinforcement, Contingencies, Sanctions
8. Intentions	Stability of intentions, Stages of change model, Trans. model/stages of change
9. Goals	Goals (distal/proximal), Goal priority, Goal / target setting, Goals (autonomous/controlled), Action planning Implementation intention
10. Memory, Attention and Decision Processes	Memory, attention, decision making, cognitive overload, tiredness
11. Environmental Context and Resources	Environmental stressors, Resources / material resources, Barriers and facilitators, Organisational culture /climate Person x environment interaction, Salient events / critical incidents
12. Social influences	Social pressure, Social norms, Group conformity, Social comparisons, Group norms, Social support, Intergroup, conflict, Power, Group identity, Alienation, Modelling
13. Emotion	Anxiety, Fear, Affect, Stress, Depression, Positive / negative affect, Burn-out,
14. Behavioural Regulation	Self-monitoring, Breaking habit, Action planning

Adapted from Cane *et al.* 2012, Atkins *et al.* 2017

Table 3: Participant demography

	Demographic Data	Number of Participants (%)
Sex (n=28)	Male	21 (75.0)
	Female	7 (25.0)
Age (n=28)	25-35 years old	9 (32.1)
	36-45 years old	10 (35.7)
	46-67 years old	9 (32.1)
Highest Level of Education (n=28)	Left school before 16	6 (21.4)
	Left school with GCSE/CSE/O-Level/Standard Grade or equivalent	14 (50.0)
	Left school with A-Level/Higher or equivalent	4 (14.3)
	University degree	1 (3.6)
	Other	3 (10.7)
Marital Status (n=28)	Single	16 (57.1)
	Divorced or separated	2 (7.1)
	Widowed	2 (7.1)
	Living with a partner (co-habiting)	6 (21.4)
	In a long term relationship	2 (7.1)
Where do you normally sleep? (n=28)	Hostel	5 (17.9)
	Council, housing association	11 (39.3)
	Sleeping rough	2 (7.1)
	Other such as with friends or relatives, B&B, Caravan	5 (17.9)
	Privately rented or owned accommodation	3 (10.7)
	Other	2 (7.1 (bedsit & shared house))
		20 (71.4)
Where do you normally obtain daily essentials? (n=27)	Mostly buys own food	4 (14.3)
	Churches	4 (14.3)
	Charity shelters or hostels	5 (17.9)
	Friends or relatives	1 (3.6)
	Begging	5 (17.9)
	Other	3 (10.7)
How did you become homeless? (n=28)	Alcohol misuse	17 (60.7)
	Drug misuse	2 (7.1)
	Gambling	310.7)
	Abusive situation	7 (25.0)
	Relationship breakdown	0 (0.0)
	Injury	2 (7.1)
	Loss of Job	8 (28.6)
	Mental Illness	9 (32.1)
	Other	5 (17.9)
		8 (28.6)
How long have you been homeless for? (n=28)	Less than 6 months	6 (21.4)
	6 months to a year	3 (10.7)
	1-2 years	6 (21.4)
	3-4 years	6 (21.4)
	5 or more years	8 (28.6)
How old were you when you first became homeless? (n=27)	Younger than 20 years old	6 (21.4)
	Between 20 – 30 years old	13 (46.4)
	Older than 30 years old	7 (25.0)
Responsible for any children? (n=28)	Yes	21 (75.0)
	No	20 (71.4)
Employment Status (n=28)	Unemployed and currently not looking for work	4 (14.3)
	Unemployed and currently looking for work	1 (3.6)
	Unemployed and student	2 (7.1)
	Employed full time	0 (0.0)
	Employed part time	1 (3.6)
	Retired	1 (3.6)
		1 (3.6)
How would you describe your general health? (n=28)	Very good	4 (14.3)
	Good	9 (32.1)
	Fair	12 (42.9)
	Bad	2 (7.1)
	Very bad	

*multiple choices were allowed

775
776
777

Table 4: Results of framework analysis

TDF^{18,19} domains	Self-awareness of physical and mental health	Healthy eating	Health literacy (Health information seeking behaviour)	Personal hygiene	Risk avoidance Rational and responsible use of products, services, diagnostics and medicines	Physical activity and rest (sleep)
1. Knowledge	Knowledge of personal health conditions, knowledge of the impact of homelessness on health	Knowledge (or lack of) about nutritional values of food; knowledge of the role of good food on health	Knowledge (or lack of) where to seek health information	Knowledge about facilities available in the locality for a wash	Knowledge of drug misuse about negative impact on health	Knowledge on the importance of physical activity or quality sleep to health Knowledge about medicines prescribed for better sleep quality
2. Skills		Lack of skills to prepare (cook) food	Ability (or lack of) to interpret health related information available online			
3. Social/ Professional Role and Identity	Being personally responsible for the homelessness and poor health		Identity as a patient, being an 'open book'		Social influence in taking up and giving up risky behaviours	Identity as a 'rough sleeper'
4. Beliefs about Capabilities	Losing self esteem	Poor health impacting on ability to eat healthily... not being able to 'open a tin.'	Self-confidence in asking health information from other individuals	'Nothing I could do' to maintain personal hygiene	Self-confidence in avoidance of risky behaviours	Disability, morbidity impacting Self-confidence or lack of in using physical exercise facilities
5. Optimism	Pessimistic about bringing positive change to health				Having already done 'irreversible' damage to health	Pessimistic about adopting better sleep pattern
6. Beliefs about Consequences	Consequences of prolonged homelessness on health	Consequences of not eating healthily			Consequences of illicit use of drugs, smoking and alcohol misuse to the health of	Consequences of good physical activity and sleep on health

7. Reinforcement	Positive health to enable job, work Lack of motivation to maintain health in temporary accommodation					
8. Intentions	Intention (or lack of) to maintain a good health such as keeping warm	Intention to eat healthily, e.g. through family, friends, charities	Intention (or lack of) to seek health information	Intention (or lack of) to remain clean	Intentions to come off drugs, smoking or alcohol	Lack of intentions to exercise Intentions to sleep well Seeking medications for better sleep quality
9. Goals	Good health a goal or in the 'back burner' Decision to make positive changes to health	Non-intention to spend on good food due to illicit drug habits Eating healthily a goal or not a goal		'More important things to worry about'	Goal setting in giving up risky behaviours	Physical activity not a goal or a priority Too tired to think about exercise, sleeping with 'one eye open'
10. Memory, Attention and Decision Processes						
11. Environmental Context and Resources	Vulnerable/ prone to poor health due to environmental hazards, lack of sleep Barriers of using health services including difficulties registering to the health care services and also lack of	Lack of facilities to store, cook or warm up food Lack of money to buy quality food Charity monetary resource to buy food Shoplifting to satisfy hunger	Resources including doctors, nurses and online sources or lack of seek health information	Lack of facilities for showers, Use of available facilities for shower, Use of limited facilities such as toilets for shower Charity facilitates to wash or dress clean	Importance of rehabilitation centres, methadone programmes and smoking cessation services on participants giving up of risky behaviours	Use of gym and exercises in temporary accommodation Weather having a big impact on sleep quality when rough sleeping

	Not being able to keep warm, Lack of place to store medicines					
Social influences	Alienation due to homelessness, lack of social support in maintaining health	Family and friends support to eating healthily Positive role of health care professional advice on healthy eating		Support from friends and family in maintaining personal hygiene	Social influence on taking up or giving up risky behaviours	Negative social attitude to rough sleeping, violence faced during rough sleeping
13. Emotion	Impact of stigma and discrimination Poor mental health, stress, depression, paranoia, suicidal ideation, feeling vulnerable, lack of self-esteem	Hunger often lasting several days	Mental health issues leading to fear and anxiety in learning new things Reluctance to speak about homelessness and its impact on health to HCPs they are not acquainted with	Fear of abuse due to poor hygiene	Illicit drug use as a coping mechanism	Personal worries disabling any sleep
14. Behavioural Regulation	Adoption (or non-adoption) of positive health behaviour; Adherence to the treatment to improve health	Discontinuation of drugs to eat healthily Attempting to eat as healthily as possible		Being able to maintain hygiene despite sleeping rough, actively seeking shower and clean up facilities	Determination (or lack of) giving up risky behaviours	Walking (instead of public transport) to improve health when no longer homeless

779
780
781
782