Ramsey properties of randomly perturbed graphs

Das, Shagnik; Treglown, Andrew

DOI:

[10.1017/S0963548320000231](https://doi.org/10.1017/S0963548320000231)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Das, S & Treglown, A 2020, 'Ramsey properties of randomly perturbed graphs: cliques and cycles', *Combinatorics, Probability and Computing*, vol. 29, no. 6, pp. 830-867.
<https://doi.org/10.1017/S0963548320000231>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This article has been published in a revised form in *Combinatorics, Probability and Computing* [<https://doi.org/10.1017/S0963548320000231>]. This version is free to view and download for private research and study only. Not for re-distribution or re-use. © The Author(s), 2020

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

RAMSEY PROPERTIES OF RANDOMLY PERTURBED GRAPHS: CLIQUEs AND CYCLES

SHAGNIK DAS AND ANDREW TREGLOWN

ABSTRACT. Given graphs H_1, H_2 , a graph G is (H_1, H_2) -Ramsey if for every colouring of the edges of G with red and blue, there is a red copy of H_1 or a blue copy of H_2 . In this paper we investigate Ramsey questions in the setting of *randomly perturbed graphs*: this is a random graph model introduced by Bohman, Frieze and Martin [8] in which one starts with a dense graph and then adds a given number of random edges to it. The study of Ramsey properties of randomly perturbed graphs was initiated by Krivelevich, Sudakov and Tetali [30] in 2006; they determined how many random edges must be added to a dense graph to ensure the resulting graph is with high probability (K_3, K_t) -Ramsey (for $t \geq 3$). They also raised the question of generalising this result to pairs of graphs other than (K_3, K_t) . We make significant progress on this question, giving a precise solution in the case when $H_1 = K_s$ and $H_2 = K_t$ where $s, t \geq 5$. Although we again show that one requires polynomially fewer edges than in the purely random graph, our result shows that the problem in this case is quite different to the (K_3, K_t) -Ramsey question. Moreover, we give bounds for the corresponding (K_4, K_t) -Ramsey question; together with a construction of Powerski [37] this resolves the (K_4, K_4) -Ramsey problem.

We also give a precise solution to the analogous question in the case when both $H_1 = C_s$ and $H_2 = C_t$ are cycles. Additionally we consider the corresponding multicolour problem. Our final result gives another generalisation of the Krivelevich, Sudakov and Tetali [30] result. Specifically, we determine how many random edges must be added to a dense graph to ensure the resulting graph is with high probability (C_s, K_t) -Ramsey (for odd $s \geq 5$ and $t \geq 4$).

To prove our results we combine a mixture of approaches, employing the container method, the regularity method as well as dependent random choice, and apply robust extensions of recent asymmetric random Ramsey results.

MSC2000: 5C55, 5C80, 5D10.

1. INTRODUCTION

Let G and H be graphs and $r \in \mathbb{N}$. We say that G is (H, r) -Ramsey if every r -colouring of G yields a monochromatic copy of H in G . More generally, for graphs $H_1, \dots, H_r$, a graph G is $(H_1, \dots, H_r)$ -Ramsey if for any r -colouring of G there is a copy of H_i in colour i for some $i \in [r]$. In the case $r = 2$, we shall take the first colour to be red and the second colour to be blue. Ramsey's classic theorem tells us that if $n \in \mathbb{N}$ is sufficiently large then K_n is $(H_1, \dots, H_r)$ -Ramsey. Whilst in general for given graphs $H_1, \dots, H_r$ it seems out of reach to determine the *smallest* $n =: R(H_1, \dots, H_r)$ with this property, much effort has gone into establishing good upper and lower bounds on $R(H_1, \dots, H_r)$ (particularly in the case when the H_i are cliques; see e.g. [12, 42]).

1.1. Ramsey properties of random graphs. There has also been significant interest in understanding Ramsey properties of the *random graph* $G(n, p)$. Recall that $G(n, p)$ has vertex set $[n] := \{1, \dots, n\}$ and each edge is present with probability p , independently of all other choices.

Date: February 19, 2020.

SD: Freie Universität Berlin, Germany, shagnik@mi.fu-berlin.de. Research supported by GIF grant G-1347-304.6/2016.

AT: University of Birmingham, United Kingdom, a.c.treglown@bham.ac.uk. Research supported by EPSRC grant EP/M016641/1.

Seminal work of Rödl and Ruciński [38, 39, 40] determined the *threshold* for the (H, r) -Ramsey property in $G(n, p)$ for all fixed H and $r \geq 2$. Before we state their result (in a slightly restricted form) we require two definitions. Given a graph H , set $d_2(H) := 0$ if $e(H) = 0$; $d_2(H) := 1/2$ when H is precisely an edge and define $d_2(H) := (e(H) - 1)/(v(H) - 2)$ otherwise. Then define $m_2(H) := \max_{H' \subseteq H} d_2(H')$ to be the *2-density* of H . We say that H is *strictly 2-balanced* if $m_2(H') < m_2(H)$ for all $H' \subsetneq H$. Rödl and Ruciński showed that the 2-density of H is the parameter that governs the threshold for the (H, r) -Ramsey property in $G(n, p)$.

Theorem 1.1 ([38, 39, 40]). *Let $r \geq 2$ be a positive integer and let H be a graph that is not a forest consisting of stars and paths of length 3. There are positive constants c, C such that*

$$\lim_{n \rightarrow \infty} \mathbb{P}[G(n, p) \text{ is } (H, r)\text{-Ramsey}] = \begin{cases} 0 & \text{if } p < cn^{-1/m_2(H)}; \\ 1 & \text{if } p > Cn^{-1/m_2(H)}. \end{cases}$$

We remark that a short proof of Theorem 1.1 was recently given in [35]. There has also been attention on the more general problem of determining the threshold of the $(H_1, \dots, H_r)$ -Ramsey property in $G(n, p)$. In particular, the focal question in the area is the following conjecture of Kohayakawa and Kreuter [25]. To state it, we need to introduce the *asymmetric density* of H_1, H_2 where $m_2(H_1) \geq m_2(H_2)$ via

$$(1.1) \quad m_2(H_1, H_2) := \max \left\{ \frac{e(H'_1)}{v(H'_1) - 2 + 1/m_2(H_2)} : H'_1 \subseteq H_1 \text{ and } e(H'_1) \geq 1 \right\}.$$

We say that H_1 is *strictly balanced with respect to $m_2(\cdot, H_2)$* if no $H'_1 \subsetneq H_1$ with at least one edge maximises (1.1).

Conjecture 1.2 ([25]). *For any graphs $H_1, \dots, H_r$ with $m_2(H_1) \geq \dots \geq m_2(H_r) > 1$, there are positive constants $c, C > 0$ such that*

$$\lim_{n \rightarrow \infty} \mathbb{P}[G(n, p) \text{ is } (H_1, \dots, H_r)\text{-Ramsey}] = \begin{cases} 0 & \text{if } p < cn^{-1/m_2(H_1, H_2)}; \\ 1 & \text{if } p > Cn^{-1/m_2(H_1, H_2)}. \end{cases}$$

Note that this conjectured threshold only depends on the ‘joint density’ of the densest two graphs H_1, H_2 . Further, notice $m_2(H_1) \geq m_2(H_1, H_2) \geq m_2(H_2)$ with equality if and only if $m_2(H_1) = m_2(H_2)$. Thus Conjecture 1.2 would generalise Theorem 1.1. The initial work on Conjecture 1.2 focused on the cases of cycles and cliques. (We take a similar approach in this paper when considering the analogous question in the perturbed setting.) In particular, Kohayakawa and Kreuter [25] confirmed Conjecture 1.2 when the H_i are cycles. When each H_i is a clique, the 0-statement was resolved by Marciniszyn, Skokan, Spöhel and Steger [32], who also observed that the approach used by Kohayakawa and Kreuter [25] implies the 1-statement of Conjecture 1.2 holds when H_1 is strictly 2-balanced *provided* the so-called KLR conjecture holds. This latter conjecture was proven by Balogh, Morris and Samotij [3], thereby proving the 1-statement of Conjecture 1.2 holds for strictly 2-balanced graphs H_1 . Hancock, Staden and Treglown [22] then proved a general result which implies (a resilient version of) the 1-statement in the case when $m_2(H_1) = m_2(H_2)$. Very recently, Mousset, Nenadov and Samotij [34] have shown the 1-statement is true without any assumptions regarding the balancedness of H_1 .

1.2. Ramsey properties of randomly perturbed graphs. Note that the results discussed above give us precise information about the Ramsey properties of *typical* graphs of a given density. Indeed, Theorem 1.1 implies that a typical graph of density $p = \omega(n^{-1/m_2(H)})$ is (H, r) -Ramsey whilst a typical graph of density $p = o(n^{-1/m_2(H)})$ is not (H, r) -Ramsey. In this paper we study the question of *how far away* a dense graph is from satisfying a given Ramsey property. The model of *randomly perturbed graphs*, introduced by Bohman, Frieze and Martin [8], provides a

framework for studying such questions. In their model one starts with a dense graph and then adds a fixed number of random edges to it. A natural problem in this setting is to determine how many random edges are required to ensure that the resulting graph w.h.p. satisfies a given property. For example Bohman, Frieze and Martin [8] proved that, given any n -vertex graph G of linear minimum degree, if one adds a linear number of random edges to G then, w.h.p., the resulting graph is Hamiltonian. In recent years, a whole host of results have been obtained concerning embedding spanning subgraphs into a randomly perturbed graph, as well as other properties of the model; see e.g. [4, 6, 7, 9, 10, 15, 24, 29, 30, 36]. The model has also been investigated in the setting of directed graphs and hypergraphs (see e.g. [5, 21, 28, 33]). Further, very recently an analogous model of randomly perturbed sets of integers has been studied [1].

The study of Ramsey properties of randomly perturbed graphs was initiated by Krivelevich, Sudakov and Tetali [30] in 2006. They considered the question of how many random edges one needs to add to any dense graph to ensure with high probability the resulting graph is (H_1, H_2) -Ramsey. Specifically, they resolved this question in the case when $H_1 = K_t$ and $H_2 = K_3$ for any $t \geq 3$. In this paper we give several generalisations of their result. Further, we look at the following more refined question: given any fixed $0 < d < 1$, how many random edges does one need to add to any graph G of density at least d to ensure with high probability the resulting graph is (H_1, H_2) -Ramsey? In order to present our results we define a threshold function below.

Definition 1.3 (Perturbed Ramsey threshold probability). *Given a density $0 < d < 1$, a number of colours $r \in \mathbb{N}$ and a sequence of graphs $(H_1, H_2, \dots, H_r)$, the perturbed Ramsey threshold probability $p(n; H_1, H_2, \dots, H_r, d)$ satisfies the following.*

- (i) *If $p = p(n) = \omega(p(n; H_1, H_2, \dots, H_r, d))$, then for any sequence $(G_n)_{n \in \mathbb{N}}$ of n -vertex graphs with density at least d , the graph $G_n \cup G(n, p)$ is $(H_1, H_2, \dots, H_r)$ -Ramsey with high probability.*
- (ii) *If $p = p(n) = o(p(n; H_1, H_2, \dots, H_r, d))$, for some sequence $(G_n)_{n \in \mathbb{N}}$ of n -vertex graphs with density at least d , the graph $G_n \cup G(n, p)$ is with high probability not $(H_1, H_2, \dots, H_r)$ -Ramsey.*

If it is the case that every sufficiently large graph of density at least d is $(H_1, H_2, \dots, H_r)$ -Ramsey then we define $p(n; H_1, H_2, \dots, H_r, d) := 0$. In the symmetric case, where $H_1 = H_2 = \dots = H_r = H$, we denote the threshold by $p(n; r, H, d)$.

We begin by observing some simple lower and upper bounds on the perturbed Ramsey threshold probability, which will serve as points of reference for our results.

Observation 1.4. *The following bounds on $p(n; H_1, H_2, d)$ hold:*

- (i) *If $d \leq 1 - 1/k$ and H_2 is not k -partite, then we may take G to be a complete balanced k -partite graph and colour all its edges blue. As long as $G(n, p)$ is H_1 -free, we may colour all its uncoloured edges red, and so $p(n; H_1, H_2, d)$ is at least the threshold probability for the appearance of H_1 in $G(n, p)$.*
- (ii) *If $G(n, p)$ alone is already (H_1, H_2) -Ramsey, then $G \cup G(n, p)$ will be as well. Since the 1-statement of Conjecture 1.2 is known to hold, it follows that for any d and graphs H_1 and H_2 with $m_2(H_1) \geq m_2(H_2) > 1$, we have $p(n; H_1, H_2, d) \leq n^{-1/m_2(H_1, H_2)}$.*
- (iii) *Suppose there is a $(k+1)$ -chromatic graph H that is (H_1, H_2) -Ramsey. If $d > 1 - 1/k$, the Erdős–Stone–Simonovits Theorem [17] implies any sufficiently large graph of density d will contain H , and thus will already be (H_1, H_2) -Ramsey before the addition of any random edges. Hence $p(n; H_1, H_2, d) = 0$ for $d > 1 - 1/k$.*

In particular, parts (i) and (ii) imply $n^{-2/(t-1)} \leq p(n; K_t, K_s, d) \leq n^{-2(ts+t-2s)/(t(t-1)(s+1))}$ for integers $t \geq s \geq 3$ and density $d \leq \frac{s-2}{s-1}$.

The aforementioned result of Krivelevich, Sudakov and Tetali [30] shows that the lower bound given by Observation 1.4(i) is in fact tight for the (K_t, K_3) -Ramsey problem.

Theorem 1.5 ([30]). *For $0 < d \leq 1/2$ and $t \geq 3$, we have $p(n; K_t, K_3, d) = n^{-2/(t-1)}$.*

1.3. Our results. Krivelevich, Sudakov and Tetali [30] raised the question of extending Theorem 1.5 to other pairs of graphs (H_1, H_2) . In this paper, we make the first progress on this question since it was raised in 2006, focusing on the cases where the H_i are cliques or cycles. In our first result we resolve their question in the case of cliques of size at least 5.

Theorem 1.6. *For $0 < d \leq 1/2$ and $t \geq s \geq 5$, we have $p(n; K_t, K_s, d) = n^{-1/m_2(K_t, K_{\lceil s/2 \rceil})}$.*

Recall that in the random graph setting one needs $\Theta(n^{2-1/m_2(K_t, K_s)})$ random edges to ensure $G(n, p)$ is with high probability (K_t, K_s) -Ramsey. On the other hand, Theorem 1.6 shows that adding just $\omega(n^{2-1/m_2(K_t, K_{\lceil s/2 \rceil})})$ random edges is already enough to make any dense n -vertex graph (K_t, K_s) -Ramsey. (In fact, in the proof we see that we only require $\Theta(n^{2-1/m_2(K_t, K_{\lceil s/2 \rceil})})$ random edges.)

To see that this is best possible, consider the complete balanced bipartite graph G on n vertices. Indeed, if $G(n, p)$ is such that there is a 2-colouring without a red K_t or blue $K_{\lceil s/2 \rceil}$, then by further colouring all the edges in G blue, one obtains a 2-colouring of $G \cup G(n, p)$ without a red K_t or blue K_s .

As one might expect, if one starts with an even denser graph G , one needs less randomness to be (K_t, K_s) -Ramsey, and we prove a stronger version of Theorem 1.6 that also gives exact (or log-asymptotically exact) results for larger densities d .

Theorem 1.7. *Given an integer $k \geq 2$, let d be such that $1 - 1/(k-1) < d \leq 1 - 1/k$, and let s and t be fixed integers with $2k+1 \leq s \leq t$. If*

- (i) $k = 2$ (that is, $0 < d \leq 1/2$), or
- (ii) $s \equiv 1 \pmod{k}$,

then $p(n; K_t, K_s, d) = n^{-1/m_2(K_t, K_{\lceil s/k \rceil})}$. Otherwise, if

- (iii) $k \geq 3$ and $s \not\equiv 1 \pmod{k}$,

we have $p(n; K_t, K_s, d) = n^{-(1-o(1))/m_2(K_t, K_{\lceil s/k \rceil})}$.

Comparing this result to the bounds in Observation 1.4, we note that in these cases, the thresholds lie strictly between the lower and upper bound, unlike when $s = 3$. Moreover, perhaps surprisingly, we see that if $s < s' \leq t$ and $\lceil s/k \rceil = \lceil s'/k \rceil$, the perturbed Ramsey threshold probabilities for the pairs (K_t, K_s) and $(K_t, K_{s'})$ are essentially the same.

In light of Theorems 1.5 and 1.6, in the setting of cliques, the Krivelevich–Sudakov–Tetali question now remains open only in the (K_t, K_4) case. As we will see in the next proposition though, this problem exhibits a different behaviour to the other cases.

Proposition 1.8. *For $k \geq 2$, let $1 - 1/(k-1) < d \leq 1 - 1/k$. If $k+2 \leq s \leq 2k$ and $t \geq s$, we have $n^{-2t/(t(t-1)+\lceil t/a \rceil)} \leq p(n; K_t, K_s, d) \leq n^{-2/t}$, where a is the smallest integer for which $R(K_{a+1}, K_{s-k}) > k$.*

It is worth noting that the formula in Theorem 1.7, had it also been valid for $k+2 \leq s \leq 2k$, would have implied the lower bound of $n^{-2/(t-1)}$ from Observation 1.4 is (essentially) correct. While Theorem 1.5 shows this to be the case for $s = 3$, the lower bound in Proposition 1.8 is higher, highlighting that the threshold probability truly does behave differently when $k+2 \leq s \leq 2k$. The upper bound in Proposition 1.8 also represents an improvement over the upper bound from Observation 1.4.

Remark. After making our manuscript available online, we learnt of the simultaneous and independent work of Powierski [37]. He proves Theorem 1.6 in the case $s = t \geq 5$ odd, and improves the lower bound of Proposition 1.8 when $k = 2$ and $s = t = 4$. In particular, combined with our upper bound from Proposition 1.8 this shows $p(n; K_4, K_4, d) = n^{-1/2}$ for $0 < d \leq 1/2$. Thus, the question from [30] is now resolved for the (K_t, K_t) -Ramsey problem for all $t \geq 3$. We suspect though that resolving the (K_t, K_4) -Ramsey problem for all $t \geq 5$ will be a significant challenge.

We next turn our attention to cycles, completely determining the perturbed Ramsey thresholds for all pairs of cycles and all densities.

Theorem 1.9. *Let $k, \ell \geq 3$ be integers such that either k is odd and ℓ is even, or they have the same parity and $k \leq \ell$, and let $d \in (0, 1)$. There exist $d_1 = d_1(k, \ell), d_2 = d_2(k, \ell) \in [0, 1]$ such that*

$$p(n; C_k, C_\ell, d) = \begin{cases} n^{-1} & \text{if } 0 < d \leq d_1, \\ n^{-2} & \text{if } d_1 < d \leq d_2, \\ 0 & \text{if } d_2 < d. \end{cases}$$

Moreover, the values of d_1 and d_2 are as given below.

	k even	k odd		
	ℓ even	ℓ even	$\ell = 3$	$\ell \geq 5$ odd
$d_1(k, \ell)$	0	1/2	1/2	1/2
$d_2(k, \ell)$	0	1/2	4/5	3/4

Theorem 1.9 shows there are at most three phases: an initial phase, where a linear number of random edges is needed to make a dense graph (C_k, C_ℓ) -Ramsey, an intermediate phase, where it suffices to add a large constant number of edges, and a supercritical phase, where the underlying graph is dense enough to already be (C_k, C_ℓ) -Ramsey. The parities and, in some cases, lengths of the two cycles in question determine at which densities (if at all) the transitions between these phases occur. In all cases though, our result demonstrates that one needs significantly fewer random edges for the perturbed (C_k, C_ℓ) -Ramsey question compared to the analogous result in the random graph setting [25].

We have thus far focused on the perturbed Ramsey thresholds for pairs of graphs, and our next observation explains why this is the case of greatest interest. Indeed, for the graph pairs we have studied, our results show that, when adding random edges to a graph of positive density, one needs significantly less randomness to make the graph (H_1, H_2) -Ramsey than in $G(n, p)$. However, this is not the case when there are three or more colours; since the probability threshold for $G(n, p)$ to be (H, r) -Ramsey does not depend on r , one needs a much denser base graph before requiring less randomness in the perturbed model. For simplicity we consider only the symmetric case, but similar remarks can be made in a more general setting.

Observation 1.10. *For a graph H and $r \geq 3$, set $k := \min\{\chi(F) : F \text{ is } (H, r-2)\text{-Ramsey}\}$. Then for $d \leq 1 - 1/(k-1)$, we have $p(n; r, H, d) = n^{-1/m_2(H)}$.*

Indeed, if $p \geq Cn^{-1/m_2(H)}$ for some constant C , Theorem 1.1 shows that $G(n, p)$ itself will be (H, r) -Ramsey, and hence this is an upper bound on the perturbed Ramsey threshold.

For the lower bound, take G to be a complete balanced $(k-1)$ -partite graph. By definition of k , we can $(r-2)$ -colour the edges of G without creating a monochromatic copy of H . We use the remaining two colours on the edges of $G(n, p)$. By Theorem 1.1, if $p \leq cn^{-1/m_2(H)}$ for some constant c , then with high probability $G(n, p)$ is not $(H, 2)$ -Ramsey. This thus gives an r -colouring of $G \cup G(n, p)$ without a monochromatic copy of H .

Despite this, in our next result we consider the symmetric multicoloured perturbed Ramsey thresholds for long cycles. As predicted by Observation 1.10, there is a large subcritical regime,

where the perturbed thresholds are the same as those from $G(n, p)$. However, once the underlying graph is very dense, we observe some similarities to Theorem 1.9. There remains one range of densities where we have been unable to determine the perturbed threshold.

Theorem 1.11. *For a number of colours $r \geq 3$ and odd cycle length $\ell \geq 2^r + 1$, we have*

$$p(n; r, C_\ell, d) \begin{cases} = n^{-1+1/(\ell-1)} & 0 \leq d \leq 1 - 2^{-r+2}, \\ \in [n^{-1}, n^{-1+1/(\ell-1)}] & 1 - 2^{-r+2} < d \leq 1 - 2^{-r+1}, \\ = n^{-2} & 1 - 2^{-r+1} < d \leq 1 - 2^{-r}, \\ = 0 & 1 - 2^{-r} < d \leq 1. \end{cases}$$

Finally, we combine these different settings, determining the perturbed Ramsey thresholds for odd cycles versus cliques. Our last result provides a different extension of Theorem 1.5, showing that the same perturbed Ramsey threshold remains valid if K_3 is replaced with any odd cycle.

Theorem 1.12. *For any clique size $t \geq 4$, odd cycle length $\ell \geq 5$, and density $0 < d \leq 1/2$, we have $p(n; K_t, C_\ell, d) = n^{-2/(t-1)}$.*

We note that the missing cases are covered by previous results; when $\ell = 3$, the threshold is given by Theorem 1.5, while the case $t = 3$ is covered by Theorem 1.9. Theorem 1.12 is perhaps of additional interest as it is the only example of the perturbed Ramsey question where we know the threshold precisely, but the analogous question in the setting of random graphs remains open.

1.4. Notation and organisation of the paper. Let G be a graph. We define $V(G)$ to be the vertex set of G and $E(G)$ to be the edge set of G . Define $v(G) := |V(G)|$ and $e(G) := |E(G)|$. For each $x \in V(G)$, we define the *neighbourhood of x in G* to be $N_G(x) := \{y \in V(G) : xy \in E(G)\}$ and define $d_G(x) := |N_G(x)|$. We define $\Delta(G)$ to be the *maximum degree of G* ; that is, the maximum value of $d_G(x)$ over all $x \in V(G)$. Given $X \subseteq V(G)$ we write $N_G(X) := \bigcap_{x \in X} N_G(x)$ for the *common neighbourhood of X in G* .

Let $X \subseteq V(G)$. Then $G[X]$ is the *graph induced by X on G* and has vertex set X and edge set $E_G(X) := \{xy \in E(G) : x, y \in X\}$. Similarly, if $A, B \subseteq V(G)$ are disjoint, we write $G[A, B]$ for the bipartite graph with vertex classes A and B and edge set $E_G(A, B) := \{xy \in E(G) : x \in A, y \in B\}$, and define $e_G(A, B) := |E_G(A, B)|$. We will often drop the subscript G from our notation if the graph under consideration is clear from context.

Given two graphs G, H on the same vertex set V we write $G \cup H$ for the graph with vertex set V and edge set $E(G) \cup E(H)$. We shall assume the vertex set of an n -vertex graph to be $[n] := \{1, 2, \dots, n\}$, unless otherwise specified.

Given a set A and $k \in \mathbb{N}$ we denote by A^k the set of all ordered k -tuples $(a_1, \dots, a_k)$ of elements from A , while $\binom{A}{k}$ denotes the set of all (unordered) k -element subsets $\{a_1, \dots, a_k\}$ of A .

Suppose $H_1, \dots, H_s$ and $H'_1, \dots, H'_t$ are graphs. We write $R(\{H_1, \dots, H_s\}, \{H'_1, \dots, H'_t\})$ for the smallest $n \in \mathbb{N}$ such that whenever K_n is 2-coloured, there is a red copy of *some* H_i (with $1 \leq i \leq s$) or there is a blue copy of *some* H'_j (with $1 \leq j \leq t$).

We write $0 < a \ll b \ll c < 1$ to mean that we can choose the constants a, b, c from right to left, with each constant sufficiently small with respect to those preceding it. More precisely, there exist non-decreasing functions $f : (0, 1] \rightarrow (0, 1]$ and $g : (0, 1] \rightarrow (0, 1]$ such that for all $b \leq f(c)$ and $a \leq g(b)$ our calculations and arguments in our proofs are correct. Thus $a \ll b$ implies that we may assume e.g. $a < b$ or $a < b^2$, as needed. Hierarchies of different lengths are defined similarly.

We will also use the standard asymptotic notation. Given two positive functions $\alpha, \beta : \mathbb{N} \rightarrow \mathbb{R}$, we write $\alpha = o(\beta)$ and $\beta = \omega(\alpha)$ if $\lim_{n \rightarrow \infty} \alpha/\beta = 0$. We emphasise that while in certain texts, $\alpha = o(\beta)$ and $\alpha \ll \beta$ are used interchangeably, that is not the case here.

The paper is organised as follows. In the next section we introduce useful tools concerning structures in dense, random, and randomly perturbed graphs. Then in Section 3 we give the proofs

of all of our main results. Some concluding remarks are given in Section 4. Finally, we include proofs of the less standard results from Section 2 in Appendix A.

2. USEFUL TOOLS

In this section we collect several of the tools we shall use in proving our results. We shall need some results for finding useful structures in the dense underlying graph G and others to analyse the random edges from $G(n, p)$.

2.1. Structures in dense graphs. When working with the dense graph, our main tool will be the famous Szemerédi Regularity Lemma [43]. We recall some of the key definitions and facts here, omitting many details that can be found in the survey of Komlós and Simonovits [26]. We start with that of an ε -regular pair, which is a pair of vertex sets that is very regular in terms of edge densities between linear-sized subsets, as is characteristic of bipartite random graphs.

Definition 2.1. *Given $\varepsilon > 0$, a graph G and two disjoint vertex sets $A, B \subset V(G)$, the pair (A, B) is ε -regular if for every $X \subseteq A$ and $Y \subseteq B$ with $|X| > \varepsilon|A|$ and $|Y| > \varepsilon|B|$, we have $|d(X, Y) - d(A, B)| < \varepsilon$, where $d(S, T) := e(S, T)/(|S||T|)$ for any vertex sets S and T .*

This strong regularity condition implies many desirable properties, some of which are collected in the following lemma. The first states that almost all ℓ -sets of vertices from the same side of an ε -regular pair will have a large common neighbourhood, while the second states that ε -regularity is essentially inherited by subsets. We omit the proofs of these standard facts.

Lemma 2.2. *Let (A, B) be an ε -regular pair in a graph G with $d(A, B) = d$.*

(i) *If $\ell \geq 1$ and $(d - \varepsilon)^{\ell-1} > \varepsilon$, then*

$$\left| \left\{ (x_1, x_2, \dots, x_\ell) \in A^\ell : |\cap_i N(x_i) \cap B| \leq (d - \varepsilon)^\ell |B| \right\} \right| \leq \ell \varepsilon |A|^\ell.$$

(ii) *If $\alpha > \varepsilon$, and $A' \subset A$ and $B' \subset B$ satisfy $|A'| \geq \alpha|A|$ and $|B'| \geq \alpha|B|$, then (A', B') is an ε' -regular pair of density d' , where $\varepsilon' := \max\{\varepsilon/\alpha, 2\varepsilon\}$ and $|d' - d| < \varepsilon$.*

We can now state the Regularity Lemma itself. We present the multicoloured version, as given in [26].

Theorem 2.3. *For any $\varepsilon > 0$ and $r, t \in \mathbb{N}$, there are $T = T(\varepsilon, r, t)$ and $n_0 = n_0(\varepsilon, r, t)$ such that if $n \geq n_0$ and the edges of an n -vertex graph G are r -coloured, with G_ℓ representing the ℓ th colour class, the vertex set $V(G)$ can be partitioned into sets $V_0, V_1, \dots, V_k$ for some $t \leq k \leq T$, such that $|V_0| < \varepsilon n$, $|V_1| = |V_2| = \dots = |V_k|$, and all but at most εk^2 pairs (V_i, V_j) , $1 \leq i < j \leq k$, are ε -regular pairs in each of the subgraphs G_ℓ simultaneously.*

Informally speaking, the lemma says that, apart from a small exceptional set, the vertices of G can be partitioned into a large but bounded number of parts, such that between almost all pairs of parts the edges of each colour seem randomly distributed. We shall often apply this in the form of the following corollary, which follows by combining the Regularity Lemma with Turán's Theorem [44] (see [26] for details).

Corollary 2.4. *For every $r \geq 1$ and $\varepsilon, \delta > 0$ with $\delta \geq 3\varepsilon$ there is some $\eta = \eta(\varepsilon, \delta, r) > 0$ and $n_0 = n_0(\varepsilon, \delta, r) \in \mathbb{N}$ such that the following holds for all $n \geq n_0$ and $k \geq 2$. If G is an r -coloured n -vertex graph of density at least $1 - 1/(k - 1) + \delta$, then there are pairwise disjoint vertex sets $V_1, V_2, \dots, V_k \subset V(G)$ such that $|V_1| = \dots = |V_k| \geq \eta n$, and, for each $1 \leq i < j \leq k$, there is some colour $c_{i,j} \in [r]$ for which the edges between V_i and V_j of colour $c_{i,j}$ form an ε -regular pair of density at least $\delta/(2r)$.*

We will use the structure from Corollary 2.4 to build monochromatic cliques. We will be able to do so because, as given by Lemma 2.2, almost all ℓ -tuples of vertices in V_i will have large common neighbourhoods in each of the other parts V_j . However, in some of our applications we will require *all* ℓ -tuples to have large common neighbourhoods. For that we make use of another powerful tool from extremal combinatorics, dependent random choice. The following lemma, proven in Appendix A.1, is a multipartite version of the basic lemma from the survey of Fox and Sudakov [18].

Lemma 2.5. *Given $s \in \mathbb{N}$ and $\delta > 0$, suppose $0 < \varepsilon \leq \min\{1/(2s), \delta/2\}$. Let $V_1, V_2, \dots, V_s$ be disjoint sets of vertices from a graph G , each of size m , such that for all $i \in [s-1]$, the pair (V_i, V_s) is ε -regular of density at least δ . For every $\beta > 0$ and $\ell \in \mathbb{N}$ there is some $\gamma = \gamma(s, \delta, \beta, \ell) > 0$ such that, if m is sufficiently large, there is a subset $U_s \subseteq V_s$ of size at least $m^{1-\beta}$ such that every set of ℓ vertices from U_s has at least γm common neighbours in each of the sets V_i , $i \in [s-1]$.*

When we are instead seeking slightly sparser structures within our ε -regular k -tuples, we shall use the following well-known result, a special case of the so-called Key Lemma (see [26]).

Lemma 2.6. *Let H be a fixed graph, $r \geq 2$, $0 < \varepsilon \ll d$, and let $m \in \mathbb{N}$ be sufficiently large. Suppose R is a graph on $[r]$ and suppose that G is a graph with vertex classes $V_1, \dots, V_r$ so that*

- (i) $|V_i| = m$ for all $i \in [r]$, and
- (ii) (V_i, V_j) forms an ε -regular pair of density at least d in G for all $1 \leq i < j \leq r$ with $\{i, j\} \in E(R)$.

If there is a homomorphism from H to R then G contains a copy of H .

2.2. Properties of random graphs. The previous lemma guaranteed the existence of fixed subgraphs within deterministic graphs. When we are instead dealing with the random graph $G(n, p)$, we shall apply the following result, a standard application of the Janson inequality [23, Theorem 2.14], whose proof is given in Appendix A.2.

Theorem 2.7. *Let H be a graph with $v \geq 2$ vertices and $e \geq 1$ edges. Let $[n]$ be the vertex set of $G(n, p)$, and, for some $\xi > 0$, let $\mathcal{H}$ be a collection of ξn^v possible copies of H supported on $[n]$. The probability that $G(n, p)$ does not contain any copy of H from $\mathcal{H}$ is at most $\exp(-\xi \mu_1 / (2^{v+1} v!))$, where $\mu_1 = \mu_1(H) := \min\{n^{v(F)} p^{e(F)} : F \subseteq H, e(F) \geq 1\}$.*

However, for our purposes, the most important properties of random graphs will be their Ramsey properties. As stated in the introduction, Conjecture 1.2 of Kohayakawa and Kreuter [25] suggests what the threshold for the random graph $G(n, p)$ being $(H_1, \dots, H_r)$ -Ramsey should be. While the 1-statement is now known to hold in general [34], the corresponding 0-statement is only known for cycles [25] and cliques [32].

Theorem 2.8 ([25, 32]). *Let $r \geq 2$, and let $H_1, H_2, \dots, H_r$ be graphs such that either every H_i is a complete graph or every H_i is a cycle, and $m_2(H_1) \geq m_2(H_2) \geq \dots \geq m_2(H_r)$. There is some constant $c > 0$ such that if $p \leq cn^{-1/m_2(H_1, H_2)}$, then with high probability $G(n, p)$ is not $(H_1, H_2, \dots, H_r)$ -Ramsey.*

In our applications, we require slightly stronger variants of the 1-statement, and we define the properties we shall need below.

Definition 2.9 (Robust and global Ramsey properties). *Let H_1 and H_2 be two fixed graphs, and let G be an n -vertex graph on the vertex set $[n]$.*

Given, for $i \in [2]$, families $\mathcal{F}_i \subseteq \binom{[n]}{v(H_i)}$ of forbidden subsets of $v(H_i)$ vertices, we say G is robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$ if every 2-colouring of G contains a red copy of H_1 or a blue copy of H_2 , such that the vertex set of the monochromatic subgraph is not forbidden.

Given $\mu > 0$, we say that G is μ -globally (H_1, H_2) -Ramsey if, for every 2-colouring of G and for every subset $U \subseteq [n]$ of at least μn vertices, $G[U]$ contains a red copy of H_1 or a blue copy of H_2 .

It turns out for a wide collection of pairs of graphs H_1, H_2 , above the threshold from Conjecture 1.2, $G(n, p)$ is not only (H_1, H_2) -Ramsey, but robustly and globally so. While [25] does not give explicit bounds on the error terms, making it a little harder to verify that this strengthening is possible, the more recent containers-based proofs given by Gugelmann, Nenadov, Person, Škorić, Steger and Thomas [20] and Hancock, Staden and Treglown [22] allow for the necessary extensions. In Appendix A.3 we prove Theorem 2.10, implementing the modifications that must be made to the existing proofs.

Theorem 2.10 ([20, 22]). *Let H_1 and H_2 be graphs such that $m_2(H_1) \geq m_2(H_2) \geq 1$, and*

- (a) $m_2(H_1) = m_2(H_2)$, or
- (b) H_1 is strictly balanced with respect to $m_2(\cdot, H_2)$.

The random graph $G(n, p)$ then has the following Ramsey properties:

- (i) *There are constants $\gamma = \gamma(H_1, H_2) > 0$ and $C_1 = C_1(H_1, H_2)$ such that if $p \geq C_1 n^{-1/m_2(H_1, H_2)}$ and, for $i \in [2]$, $\mathcal{F}_i \subseteq \binom{[n]}{v(H_i)}$ is a collection of at most $\gamma n^{v(H_i)}$ forbidden subsets, then $G(n, p)$ is with high probability robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$.*
- (ii) *For every $\mu > 0$ there is a constant $C_2 = C_2(H_1, H_2, \mu)$ such that if $p \geq C_2 n^{-1/m_2(H_1, H_2)}$, then $G(n, p)$ is with high probability μ -globally (H_1, H_2) -Ramsey.*
- (iii) *If we further have $m_2(H_2) > 1$, then there are constants $\beta_0 = \beta_0(H_1, H_2) > 0$ and $C_3 = C_3(H_1, H_2)$ such that if $0 \leq \beta \leq \beta_0$ and $p \geq C_3 n^{-(1-\beta)/m_2(H_1, H_2)}$, then with high probability $G(n, p)$ is $n^{-\beta}$ -globally (H_1, H_2) -Ramsey.*

Before we proceed, it is worth comparing the exponents in these different thresholds for pairs of cliques. For $t \geq 4$, we have $m_2(K_{t-1}, K_{t-1}) = \frac{t}{2} < \frac{t(t-1)}{2t-3} = m_2(K_t, K_3) < m_2(K_t, K_4) < \dots < m_2(K_t, K_t) = \frac{t+1}{2}$. In particular, when $G(n, p)$ is (K_t, K_3) -Ramsey, it will also be (K_k, K_ℓ) -Ramsey for $k, \ell < t$.

2.3. Properties of randomly perturbed graphs. While some of the previous results allowed us to find subgraphs within dense or random graphs, we will sometimes need the existence of certain graphs in the randomly perturbed model. Krivelevich, Sudakov and Tetali [30] showed that the threshold probabilities for this problem depend on the sparsest partitions of the desired graph, and the final result we shall make use of is their 1-statement.

Theorem 2.11 ([30]). *Given a graph F , define $\rho(F) := \max\{e(F')/v(F') : F' \subseteq F, v(F') \geq 1\}$, and, for $k \geq 2$, set*

$$\rho_k(F) := \min_{V(F) = \cup_i V_i} \max_i \rho(F[V_i]),$$

where the minimum is taken over all partitions into at most k parts. If $d > 1 - 1/(k-1)$, $p = \omega(n^{-1/\rho_k(F)})$, and G is an n -vertex graph of density d , then $F \subseteq G \cup G(n, p)$ with high probability.

3. PROOFS

With these tools at our disposal, we are now ready to prove our main results, establishing perturbed Ramsey thresholds for various pairs of graphs.

3.1. Proof of Theorem 1.7. Our first result establishes perturbed Ramsey thresholds for pairs of cliques that are not too small. Specifically, let $k \geq 2$ be such that $1 - 1/(k-1) < d \leq 1 - 1/k$, and suppose $2k+1 \leq s \leq t$. For convenience, we set $\ell := \lceil s/k \rceil$, and note that $\ell \geq 3$. We shall show $p(n; K_t, K_s, d) = n^{-1/m_2(K_t, K_\ell)}$ if $k = 2$ or $s \equiv 1 \pmod{k}$, and otherwise obtain this same threshold log-asymptotically. In all cases, the lower bound follows from the same argument, which we now present.

Proof of Theorem 1.7 (lower bound). Let G be the k -partite n -vertex Turán graph with vertex classes $V_1, V_2, \dots, V_k$, which has density at least $1 - 1/k$, and let $p \leq cn^{-1/m_2(K_t, K_\ell)}$, where the constant c is as in Theorem 2.8. We need to show that, with high probability, the edges of $G \cup G(n, p)$ can be 2-coloured without creating a red K_t or a blue K_s . We colour the edges of G blue, leaving only the edges of $G(n, p)[V_i]$, $i \in [k]$, uncoloured. By Theorem 2.8, with high probability $G(n, p)$ can be 2-coloured without creating a red K_t or a blue K_ℓ . This then gives the desired colouring — the connected components of the red subgraph lie within the red subgraphs of $G(n, p)[V_i]$, which are K_t -free, while, since the largest blue clique in each $G(n, p)[V_i]$ has at most $\ell - 1$ vertices, the largest blue clique in $G \cup G(n, p)$ has size at most $k(\ell - 1) < s$. Hence $p(n; K_t, K_s, d) \geq n^{-1/m_2(K_t, K_\ell)}$. $\square$

We divide the proof of the upper bound into three parts, treating each case in turn.

3.1.1. *Proof of Theorem 1.7 (upper bound), Case (i).* We start with the case $k = 2$. Let G be a graph of density $4d > 0$. By applying Corollary 2.4 with $r = 1$, we can find an ε -regular pair (U, W) of density at least $2d$, where $\varepsilon = \varepsilon(d) > 0$ is sufficiently small and $|U| = |W| = m := \eta n$ for some $\eta = \eta(d) > 0$.

Set $\xi := d^\ell/(2\ell)$, let $\mu := \xi d/2$, and recall that $m_2(K_t, K_\ell) > m_2(K_{t-1}, K_{t-1})$. Thus, if $C \geq \max\{C_2(K_t, K_\ell, \mu\eta), C_2(K_{t-1}, K_{t-1}, \mu\eta)\}$, it follows from Theorem 2.10 that if $p \geq Cn^{-1/m_2(K_t, K_\ell)}$, then with high probability,

(a) $G(n, p)$ is both $\mu\eta$ -globally (K_t, K_ℓ) -Ramsey and $\mu\eta$ -globally (K_{t-1}, K_{t-1}) -Ramsey.

Furthermore, set $\mathcal{F}_1 = \emptyset$, and let $\mathcal{F}_2 \subseteq \binom{U}{\ell}$ be those ℓ -sets with fewer than $d^\ell m$ common neighbours in W . By Lemma 2.2(i), it follows that $|\mathcal{F}_2| \leq \ell \varepsilon m^\ell$. Since ε was chosen to be sufficiently small, we can ensure $\ell \varepsilon < \gamma(K_t, K_\ell)$, where γ is as in Theorem 2.10. Since $G(n, p)[U] \sim G(m, p)$, provided $C \geq C_1(K_t, K_\ell) \eta^{-1/m_2(K_t, K_\ell)}$ as well, with high probability we also have, by Theorem 2.10,

(b) $G(n, p)[U]$ is robustly (K_t, K_ℓ) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$.

We may therefore assume $G(n, p)$ has Properties (a) and (b), and shall show that this implies $G \cup G(n, p)$ is (K_t, K_s) -Ramsey. Suppose for a contradiction that $G \cup G(n, p)$ has a 2-colouring with neither a red K_t nor a blue K_s . We first need the following claim.

Claim 3.1. *We may assume that no vertex in U has more than ξm red edges to W .*

Proof. First suppose $s \leq t - 1$. If we have a vertex $u \in U$ with at least ξm red neighbours in W , let $Y \subseteq W$ be the set of those red neighbours. By Property (a), we find a red K_{t-1} in $G(n, p)[Y]$, in which case we can add u to obtain the desired red K_t , or we find a blue K_{t-1} , which contains the desired blue s -clique.

Next suppose $s = t$. By symmetry, we may assume that at least half the edges of G between U and W are blue. In particular, since the density of (U, W) is at least $2d$, this means we can find a vertex $v \in W$ and a set $A \subset U$ of dm blue neighbours of v . Now suppose further that there is a vertex $u \in U$ with a set $B \subset W$ of ξm red neighbours of u .

By ε -regularity, there are at least $d|A||B|$ edges between A and B . If at least half of these are red, then we can find a vertex $b \in B$ with a set $A' \subseteq A$ of at least $d|A|/2$ red neighbours of b . Since $|A'| \geq \mu m$, by Property (a), $G(n, p)[A']$ has a red K_{t-1} , which together with b gives a red K_t , or a blue K_{t-1} , which together with v gives a blue K_t .

Hence at least half the edges between A and B are blue, which gives a vertex $a \in A$ with a set $B' \subseteq B$ of at least $d|B|/2$ blue neighbours of a . Again, $|B'| \geq \mu m$, and so by Property (a) we find in $G(n, p)[B']$ a red K_{t-1} , which extends via u to a red K_t , or a blue K_{t-1} , to which we can add a to obtain a blue K_t . $\square$

With the claim established, we complete our proof. By Property (b), $G(n, p)[U]$ contains a red K_t , in which case we are done, or a blue ℓ -clique whose vertex set S does not lie in $\mathcal{F}_2$. In particular, this implies that S has at least $d^\ell m$ common neighbours in W .

We discard those common neighbours with a least one red edge to S ; by Claim 3.1, we are left with a set Y of at least $(d^\ell m - \ell \xi m) = d^\ell m/2$ common blue neighbours of S . By Property (a), $G(n, p)[Y]$ has a red K_t , in which case we are done, or a blue K_ℓ , which together with S gives the desired blue s -clique.

Thus, when $k = 2$, we have $p(n; K_t, K_s, d) \leq n^{-1/m_2(K_t, K_\ell)}$, matching our lower bound. $\square$

3.1.2. Proof of Theorem 1.7 (upper bound), Case (ii). We next consider the case $k \geq 3$, where we provide the exact result if $s \equiv 1 \pmod{k}$. Specifically, we wish to show that if G is a graph of density $d = 1 - 1/(k-1) + \delta$, where $\delta > 0$, then if $p \geq Cn^{-1/m_2(K_t, K_\ell)}$ for some appropriately large constant C , with high probability $G \cup G(n, p)$ will be (K_t, K_s) -Ramsey.

Our strategy bears some resemblance to the previous proof. In Case (i), we built a blue K_s in two stages, finding half in each part of an ε -regular pair. When $k \geq 3$ and $s = k(\ell - 1) + 1$, we instead take k steps to build the blue clique within an ε -regular k -tuple, finding a blue $K_{\ell-1}$ in each of the first $k-1$ parts and a blue K_ℓ in the last part.

When we find a set of $\ell-1$ vertices from a part to add to the growing blue clique, we will need to ensure that they have a sufficiently large neighbourhood in each of the remaining parts, so that the process may continue. We will do this through repeated use of Lemma 2.5.¹ However, this leaves us with a sublinear set in which we will have to find a blue clique, and so we will need strong global Ramsey properties.

To that end, recall that if a pair (q, r) with $q \geq r$ precedes (t, ℓ) lexicographically (that is, either $q < t$, or $q = t$ and $r < \ell$), then $m_2(K_q, K_r) < m_2(K_t, K_\ell)$. We can therefore fix some $\beta > 0$ such that, for all such pairs (q, r) , we have $m_2(K_q, K_r) \leq (1 - \beta)m_2(K_t, K_\ell)$ and $\beta \leq \beta_0(K_q, K_r)$, where β_0 is as in Theorem 2.10.

We now define a sequence of constants that we shall use in the sequel. To start, let $\delta_0 := \delta$, and set $\delta_1 := \delta_0/4$. For $i \in [k-2]$, set $\delta_{i+1} := \delta_i/2$. For some soon-to-be-determined ε_1 , let $\eta := \eta(\delta_1, \varepsilon_1, 2)$, where η is as in Corollary 2.4, and set $m_1 := \eta n$. For $i \in [k-1]$, let $\gamma_i := \gamma(k-i+1, \delta_i, \beta/2, t-1)$, where γ is as in Lemma 2.5, and set $m_{i+1} := \lceil \gamma_i m_i \rceil$. Note that the m_i are linear in n , and set $\mu := m_k/n$.

Further, for $i \in [k-2]$, set $\varepsilon_{i+1} := \max\{2\varepsilon_i, \varepsilon_i/\gamma_i\}$. Noting that the δ_i and γ_i are independent of ε_1 , while the ε_i are linear in ε_1 , choose ε_1 sufficiently small so that $\varepsilon_i \leq \delta_i/2$ for all $i \in [k-1]$. We shall further assume that n is large enough for all following calculations and applications of the lemmas to be valid.

With these technicalities out of the way, we can proceed with our proof. We shall show that $G \cup G(n, p)$ is (K_t, K_s) -Ramsey, provided $G(n, p)$ satisfies the following properties.

- (a) $G(n, p)$ is $n^{-\beta}$ -globally (K_{t-1}, K_{s-1}) -Ramsey,
- (b) $G(n, p)$ is $n^{-\beta}$ -globally $(K_t, K_{\ell-1})$ -Ramsey,
- (c) $G(n, p)$ is μ -globally (K_t, K_ℓ) -Ramsey, and
- (d) if $s < t$, then $G(n, p)$ is $n^{-\beta}$ -globally $(K_{\lceil t/k \rceil}, K_s)$ -Ramsey.

Theorem 2.10 ensures Properties (a), (c), (d) and, if $\ell \geq 4$, (b) hold with high probability whenever $p \geq Cn^{-1/m_2(K_t, K_\ell)}$ for a large enough constant C . When $\ell = 3$, being $n^{-\beta}$ -globally (K_t, K_2) -Ramsey is equivalent to every induced subgraph on $n^{1-\beta}$ vertices containing a t -clique. Hence in this case we instead apply Theorem 2.7, taking a union bound over all such vertex subsets.

Now let G and $G(n, p)$ be as above, and suppose for contradiction there is a 2-colouring of $G \cup G(n, p)$ with neither a red K_t nor a blue K_s . Applying Corollary 2.4 with $r = 2$ to the coloured

¹In Case (i), we identified a forbidden set of vertex sets with small neighbourhoods, and then used the robust Ramsey properties of $G(n, p)$ to ensure the monochromatic K_ℓ had a large neighbourhood. However, in this setting the ε -regular pairs we have depend on the vertices chosen earlier, and our error bounds do not allow for so many applications of the robust Ramsey property.

graph G , we find pairwise disjoint vertex sets $V_1^{(1)}, V_2^{(1)}, \dots, V_k^{(1)}$, each of size at least m_1 , such that for each pair $\{i, j\} \in \binom{[k]}{2}$ there is some colour $c_{i,j}$ for which $(V_i^{(1)}, V_j^{(1)})$ is ε_1 -regular with density at least δ_1 in the colour $c_{i,j}$.

Claim 3.2. *All the colours $c_{i,j}$, $\{i, j\} \in \binom{[k]}{2}$, are the same.*

Proof. Suppose this were not the case. There are then i, j_1, j_2 such that c_{i,j_1} is red while c_{i,j_2} is blue. Apply Lemma 2.5 to the triple $V_{j_1}^{(1)}, V_{j_2}^{(1)}, V_i^{(1)}$ to find a set $U_i \subset V_i^{(1)}$ of size $m_1^{1-\beta/2} > n^{1-\beta}$ such that every $(t-1)$ -set in U_i has at least one common neighbour in both $V_{j_1}^{(1)}$ and $V_{j_2}^{(1)}$, using red and blue edges respectively (note that our conservative definition of γ_1 in fact guarantees us linearly many common neighbours in both $V_{j_1}^{(1)}$ and $V_{j_2}^{(1)}$, but here we only require one).

By Property (a), $G(n, p)[U_i]$ must contain a red K_{t-1} or a blue K_{s-1} . Extending this monochromatic clique with a common neighbour in $V_{j_1}^{(1)}$ or $V_{j_2}^{(1)}$ respectively gives a red K_t or a blue K_s in $G \cup G(n, p)$, a contradiction. $\square$

Let us first assume that $c_{i,j}$ is blue for every pair $\{i, j\}$ (if $s = t$, we may assume this without loss of generality). We shall either find a red K_t or build a blue K_s in k stages.

Suppose we have already run $a \geq 0$ stages of this process, thereby building a blue $K_{a(\ell-1)}$ on vertices $W = \cup_{i=1}^a W_{k+1-i}$, where W_j is a set of $\ell-1$ vertices from V_j . Further, for every $i \in [k-a]$, we have sets $V_i^{(a+1)}$ of size m_{a+1} contained in the common (blue) neighbourhood of W , such that each pair $(V_i^{(a+1)}, V_j^{(a+1)})$, $\{i, j\} \in \binom{[k-a]}{2}$, is ε_{a+1} -regular of density at least δ_{a+1} in blue.

If $a \leq k-2$, we apply Lemma 2.5 to the $(k-a)$ -tuple $V_1^{(a+1)}, V_2^{(a+1)}, \dots, V_{k-a}^{(a+1)}$. This gives us a set $U_{k-a} \subset V_{k-a}^{(a+1)}$ of size $m_{a+1}^{1-\beta/2} > n^{1-\beta}$ (since m_{a+1} is linear in n), such that any $(\ell-1)$ -set in U_{k-a} has at least m_{a+2} common (blue) neighbours in each $V_i^{(a+1)}$, $i \in [k-1-a]$.

By Property (b), $G(n, p)[U_{k-a}]$ contains a red K_t or a blue $K_{\ell-1}$. In the former case we are done, so we may assume there is a blue $(\ell-1)$ -clique on the vertices $W_{k-a} \subset U_{k-a}$. For each $i \in [k-1-a]$, let $V_i^{(a+2)} \subset V_i^{(a+1)}$ be a set of m_{a+2} common blue neighbours of W_{k-a} . By Lemma 2.2(ii), each pair $(V_i^{(a+2)}, V_j^{(a+2)})$, $\{i, j\} \in \binom{[k-1-a]}{2}$, is ε_{a+2} -regular with density at least $\delta_{a+1} - \varepsilon_{a+1} > \delta_{a+2}$.

In the final stage, when $a = k-1$, we simply set $U_1 := V_1^{(k)}$, which has size $m_k = \mu n$. By Property (c), $G(n, p)[U_1]$ contains a red K_t or a blue K_ℓ . We are again done in the former case, so we may assume the existence of a blue ℓ -clique on the vertices $W_1 \subset U_a$. This then gives a blue K_s on the vertices $\cup_{i=1}^k W_i$, contradicting our assumption that $G \cup G(n, p)$ has no blue K_s .

On the other hand, if $s < t$ and each colour $c_{i,j}$ is red instead, we follow a very similar process to that above, except in each U_i we use Property (d) to find either a blue K_s or a red $K_{\lceil t/k \rceil}$ instead. In this way, we either have a blue K_s or we build a red K_t in the k stages, obtaining the desired contradiction.

Hence we indeed have $p(n; K_t, K_s, d) = n^{-1/m_2(K_t, K_\ell)}$ in this case as well. $\square$

3.1.3. Proof of Theorem 1.7 (upper bound), Case (iii). The third and final case is when $k \geq 3$ and $k(\ell-1) + 2 \leq s \leq k\ell$. In this setting we can only match the lower bound on the perturbed Ramsey threshold log-asymptotically; that is, for any $\beta > 0$, we show that if G is a graph of density $d \geq 1 - 1/(k-1) + \delta$ and $p \geq Cn^{-(1-\beta)/m_2(K_t, K_\ell)}$, then with high probability $G \cup G(n, p)$ will be (K_t, K_s) -Ramsey.

The proof is essentially the same as in Case (ii), except when building the blue clique, we will find a blue K_ℓ in each part V_i , rather than just a $K_{\ell-1}$. Properties (a) and (d) hold as before. We can replace Properties (b) and (c) above with the following.

(b') $G(n, p)$ is $n^{-\beta}$ -globally (K_t, K_ℓ) -Ramsey.

Since we may freely assume that β is small enough to satisfy $\beta \leq \beta_0(K_t, K_\ell)$, Theorem 2.10 guarantees $G(n, p)$ satisfies Property (b') with high probability. We can then run the same process as before, picking up ℓ vertices from each U_i to add to the blue clique. Thus, after k stages, we would have built a blue $K_{k\ell} \supseteq K_s$.

This shows $p(n; K_t, K_s, d) \leq n^{-(1-\beta)/m_2(K_t, K_\ell)}$ for n sufficiently large. Since β can be taken to be arbitrarily small, we have $p(n; K_t, K_s, d) = n^{-(1-o(1))/m_2(K_t, K_\ell)}$, thereby completing the proof of Theorem 1.7. $\square$

3.2. Proof of Proposition 1.8. In Theorem 1.7, we required $s \geq 2k + 1$ or, equivalently, $\ell := \lceil s/k \rceil \geq 3$. This condition was necessary to apply Theorem 2.10, which asserts that $G(n, p)$ will be globally (K_t, K_ℓ) -Ramsey when $p = \omega(n^{-1/m_2(K_t, K_\ell)})$. Unfortunately, this is not true for $\ell = 2$. Indeed, being (K_t, K_2) -Ramsey is equivalent to containing a copy of K_t , and it is well known that the local property of the appearance of K_t in $G(n, p)$ occurs at a lower threshold probability than the global property of every large subset containing a t -clique.

This gives hope of improving the lower bound when we are dealing with a smaller clique: rather than our simplistic approach in Theorem 1.7, where all edges of the dense graph G received the same colour, we might hope to take advantage of the sparseness of the t -cliques in $G(n, p)$ to find a cleverer colouring of the edges of G , thus making it easier to avoid monochromatic copies of K_t and K_s when $s \leq 2k$. Proposition 1.8, despite falling short of determining $p(n; K_t, K_s, d)$, shows that this is indeed the case, and that one can improve upon both the obvious lower and upper bounds when $k + 2 \leq s \leq 2k$. We start with the former.

Proof of Proposition 1.8 (lower bound). We write $\ell := \lceil t/a \rceil$ for simplicity. Let G be the k -partite Turán graph with vertex classes $V_1, V_2, \dots, V_k$. We shall show that there is some constant $b > 0$ such that if $p \leq bn^{-2t/(t(t-1)+\ell)}$, then with high probability $G \cup G(n, p)$ is not (K_t, K_s) -Ramsey.

Using a result of Kreuter [27] concerning asymmetric vertex-Ramsey properties of random graphs, if $p \leq bn^{-2t/(t(t-1)+\ell)}$ for some constant $b = b(t, \ell)$, then we can with high probability partition the vertices $V(G(n, p)) = A \cup B$ such that $G(n, p)[A]$ is K_t -free and $G(n, p)[B]$ is K_ℓ -free. For $i \in [k]$, let $A_i := A \cap V_i$ and $B_i := B \cap V_i$.

For each $i \in [k]$, colour all edges within A_i and within B_i red. We further colour all edges from A_i to any other part blue. Now all that remains are the edges between B_i and B_j for $1 \leq i < j \leq k$.

Recall that we have $R(a + 1, s - k) > k$. Hence we can find a colouring $\varphi : \binom{[k]}{2} \rightarrow \{\text{red}, \text{blue}\}$ of K_k with no red clique of size $a + 1$ and no blue clique of size $s - k$. Then, for each $1 \leq i < j \leq k$, we colour all edges between B_i and B_j with the colour $\varphi(\{i, j\})$.

We claim that this colouring of $G \cup G(n, p)$ has neither a red K_t nor a blue K_s . First consider the red subgraph. Each part A_i is disconnected from the remainder of the graph, and since the only edges within A_i come from $G(n, p)$, we know that they are K_t -free. Any red component within B that corresponds to a clique in φ can involve at most a parts B_i . The largest clique within such a part has size at most $\ell - 1$, and so the largest red clique in B has size at most $a(\ell - 1) < t$. Hence there is indeed no red K_t .

Within the blue subgraph, the parts A_i and B_i are independent sets, and hence any blue clique K can contain at most one vertex from each part. Moreover, by the colouring φ , there can be at most $s - k - 1$ vertices from B in K . As there are only k parts in A , this shows that the largest blue clique has size at most $s - 1$, and hence there is no blue K_s either. This completes the proof of the lower bound. $\square$

We next establish the upper bound $p(n; K_t, K_s, d) \leq n^{-2/t}$.

Proof of Proposition 1.8 (upper bound). We start by defining sequences of constants that we shall require in the proof. Let $d := 1 - 1/(k-1) + \delta$ for some $\delta > 0$. We then set $\delta_1 := \delta/2$ and shall soon (implicitly) specify a sufficiently small ε_1 . Let $\eta = \eta(\varepsilon_1, \delta_1, 1)$ be as given by Corollary 2.4, and set $m_1 := \eta n$. Now, for $i \in [k-1]$, set $\delta_{i+1} := \delta_i/2$, $m_{i+1} := \delta_i^2 m_i/4$, and $\varepsilon_{i+1} := 4\varepsilon_i/\delta_i^2$. Observe that each m_i is linear in n , and that the ratios m_i/m_1 are independent of ε_1 .

Given these constants, choose $\alpha = \alpha(\delta) > 0$ sufficiently small to ensure $2k\alpha m_1 \binom{m_i}{t-1} \leq \frac{1}{4} \binom{m_i}{t}$ for all i , noting that this is independent of our choice of ε_1 . Finally, we choose ε_1 to be small enough that $\varepsilon_1 < \alpha$, $\delta_i \geq 3\varepsilon_i$ for each i , and $2k\varepsilon_i m_i^2 \binom{m_i}{t-2} \leq \frac{1}{4} \binom{m_i}{t}$ for all i . With these preliminaries sorted, we can now begin the proof.

We first seek a well-structured part of the deterministic graph G before exposing the random edges. Since G has density d , we can apply Corollary 2.4 with $r = 1$ to find sets $V_1, V_2, \dots, V_k$, each of size m_1 , such that each pair (V_i, V_j) is ε_1 -regular of density at least δ_1 .

Note that this is taking place in the uncoloured graph G , and hence the parts V_i are determined before we expose the random edges of $G(n, p)$. The edges in these regular pairs could later be coloured either red or blue, but Claim 3.3 will show that we may assume they are almost all blue. For this, we require the following properties of our random graph $G(n, p)$, where $\beta = \beta(t, s)$ is defined below:

- (a) $G(n, p)$ is $(\delta_1 \alpha \eta/4)$ -globally (K_{t-1}, K_{s-1}) -Ramsey, and
- (b) if $t > s$, $G(n, p)$ is $n^{-\beta}$ -globally $(K_{\lceil t/2 \rceil}, K_s)$ -Ramsey.

For the first property, we have $m_2(K_{t-1}, K_{s-1}) \leq m_2(K_{t-1}) = t/2$. Next, if $t > s$, let $s' := \min(s, \lceil t/2 \rceil)$ and $t' := \max(s, \lceil t/2 \rceil)$. Observe that $m_2(K_{t'}, K_{s'}) < t/2$, and so we can find some $\beta = \beta(t, s) > 0$ such that $m_2(K_{t'}, K_{s'})/(1-\beta) = t/2$. By Theorem 2.10, it follows that if $p = Cn^{-2/t}$ for some suitably large constant C , then $G(n, p)$ has both properties with high probability.

Claim 3.3. *If $G(n, p)$ has Properties (a) and (b) above, then we may assume that in any 2-colouring of $G \cup G(n, p)$ with neither a red K_t nor a blue K_s , for every pair $i \neq j$, the maximum red-degree in (V_i, V_j) is at most αm_1 .*

We shall prove Claim 3.3 in due course, but let us first see how it implies our desired upper bound on $p(n; K_t, K_s, d)$. Roughly speaking, we shall find t -cliques within each part V_i . Since there are no red t -cliques, each such t -clique must contain a blue edge. We shall choose the cliques to ensure that these blue edges combine to form a blue K_{2k} , contradicting our colouring being blue- K_s -free.

More precisely, let $p = Cn^{-2/t}$ where C is a sufficiently large constant, and assume (a) and (b) above hold. Consider any 2-colouring of $G \cup G(n, p)$, and recall that we assume the vertex set to be $[n]$, which we equip with its natural ordering.

Suppose for some $a \geq 0$, we have selected a set $S_a = \{s_1, s_2, \dots, s_{2a}\}$ of vertices from $\cup_{i=1}^a V_i$, such that they induce a blue K_{2a} in our colouring of $G \cup G(n, p)$ and have at least m_{a+1} common neighbours in each V_i for $a+1 \leq i \leq k$. When $a = 0$, the set $S_0 := \emptyset$ trivially satisfies these requirements.

For each $a+1 \leq i \leq k$, let $V_i^{(a+1)}$ be the first m_{a+1} common neighbours of S_a in V_i . Our goal is to find two vertices $s_{2a+1}, s_{2a+2} \in V_{a+1}^{(a+1)}$ to add to S_a in order to obtain a valid set S_{a+1} with which to proceed. In particular, the two new vertices should share a blue edge, all edges from them to S_a should also be blue, and s_{2a+1} and s_{2a+2} should have many common neighbours in each of the remaining parts. We therefore define $\mathcal{H}_{a+1}$ to be the collection of all copies of K_t whose vertex sets $T \subseteq V_{a+1}^{(a+1)}$ satisfy the following properties:

- (A) all edges between S_a and T are blue, and
- (B) every pair of vertices in T have at least m_{a+2} common neighbours in $V_j^{(a+1)}$ for each $a+2 \leq j \leq k$.

Let $R \subset V_{a+1}^{(a+1)}$ be those vertices that have a red edge to some vertex in S_a . By Claim 3.3, we have $|R| \leq \alpha m_1 |S_a| \leq 2k\alpha m_1$. Since every t -clique in $V_{a+1}^{(a+1)}$ violating Condition (A) must contain a vertex from R , there are at most $2k\alpha m_1 \binom{m_{a+1}}{t-1}$ such cliques. By our choice of α , this is at most $\frac{1}{4} \binom{m_{a+1}}{t}$ cliques.

By Lemma 2.2(ii), it follows that, for each $a+2 \leq j \leq k$, the pair $(V_{a+1}^{(a+1)}, V_j^{(a+1)})$ is ε_{a+1} -regular of density at least δ_{a+1} . We may therefore apply Lemma 2.2(i) to deduce that, for each given j , there are at most $2\varepsilon_{a+1} m_{a+1}^2$ pairs in $V_{a+1}^{(a+1)}$ with fewer than m_{a+2} common neighbours in $V_j^{(a+1)}$. There are thus a total of at most $2k\varepsilon_{a+1} m_{a+1}^2 \binom{m_{a+1}}{t-2}$ cliques of size t in $V_{a+1}^{(a+1)}$ violating Condition (B). Our choice of ε_1 ensures that this is again at most $\frac{1}{4} \binom{m_{a+1}}{t}$ cliques.

Thus $\mathcal{H}_{a+1}$ contains at least $\frac{1}{2} \binom{m_{a+1}}{t}$ sets. Suppose we find in $G(n, p)$ a t -clique $H \in \mathcal{H}_{a+1}$. If all edges of H are red, we have our desired red K_t , and hence we may assume there is some blue edge $e_{a+1} = \{s_{2a+1}, s_{2a+2}\}$ in H .

This pair of vertices has all the properties we required: they share the blue edge e_{a+1} , Property (A) of T ensures that all edges from e_{a+1} to S_a are blue, and Property (B) gives that, for each $j \geq a+2$, s_{2a+1} and s_{2a+2} have at least m_{a+2} common neighbours in $V_j^{(a+1)}$.

Hence, provided we can find a t -clique from the collection $\mathcal{H}_{a+1}$, we may set $S_{a+1} := S_a \cup \{s_{2a+1}, s_{2a+2}\}$ and proceed to the next iteration. We appeal to Theorem 2.7 to find the desired clique.

Indeed, we know $|\mathcal{H}_{a+1}| \geq \frac{1}{2} \binom{m_{a+1}}{t}$, which, since m_{a+1} is linear in n , is at least ξn^t for some constant $\xi = \xi(\delta) > 0$. Moreover, when $H := K_t$, we have $\mu_1 = \mu_1(H) = n^t p^{\binom{t}{2}}$. Recall $p = Cn^{-2/t}$, so $\mu_1 = C^{\binom{t}{2}} n$. Theorem 2.7 thus gives that the probability $G(n, p)$ does not contain a t -clique from a given collection $\mathcal{H}_{a+1}$ is at most $\exp(-C'n)$, where $C' := \xi C^{\binom{t}{2}} / (2^{t+1} t!)$.

However, the collection of t -cliques $\mathcal{H}_{a+1}$ depends on the colouring of the edges of G , which in turn could depend on the random graph $G(n, p)$ itself. To resolve this issue, we take a union bound over all collections of t -cliques $\mathcal{H}_{a+1}$ that could possibly arise. Observe that the sets $V_i^{(a+1)}$, for $a+1 \leq i \leq k$, are determined by the set S_a . This already specifies which cliques in $V_{a+1}^{(a+1)}$ fail to satisfy Property (B). To identify those violating Property (A), it suffices to identify the subset $R \subset V_{a+1}^{(a+1)}$ of vertices incident to a red edge from S_a .

The collection $\mathcal{H}_{a+1}$ is thus fully determined by the pair (S_a, R) , and there are fewer than $n^{2a} 2^{m_{a+1}}$ such pairs, which we can (wastefully) bound from above by 4^n . Hence, as C has been chosen sufficiently large (with respect to ξ and t), it follows from the union bound that with high probability, for each $0 \leq a \leq k-1$ and for every possible collection $\mathcal{H}_{a+1}$ that may arise, $G(n, p)$ contains a copy of K_t from $\mathcal{H}_{a+1}$.

We can thus repeat this process until we obtain a set S_k that induces a blue K_{2k} . Since $s \leq 2k$, this shows that with high probability $G \cup G(n, p)$ is indeed (K_t, K_s) -Ramsey, as desired. $\square$

It remains to prove Claim 3.3.

Proof of Claim 3.3. We begin with a straightforward observation: we cannot have a vertex u , a vertex v , and a set U of $\delta_1 \alpha m_1 / 4$ common neighbours of u and v , such that all edges from u to U are red and all edges from v to U are blue. Indeed, by Property (a), there is a red K_{t-1} or a blue K_{s-1} in U . Extending this monochromatic clique by u or v respectively gives either a red K_t or a blue K_s , contradicting our assumption on the colouring of $G \cup G(n, p)$.

Let us first consider the case $t > s$. Suppose for contradiction we have some $u \in V_i$ with a set R of αm_1 red neighbours in V_j , for some $j \neq i$. Let $S := \{v \in V_i : |N_G(v) \cap R| \geq (\delta_1 - \varepsilon_1)|R|\}$. By the ε_1 -regularity of (V_i, V_j) , it follows that $|S| \geq (1 - \varepsilon_1)m_1$.

By our earlier observation, each $v \in S$ can have at most $\delta_1 \alpha m_1 / 4$ blue edges to R , and hence there are at least $(1 - \varepsilon_1)(\delta_1 - \varepsilon_1 - \delta_1/4)|R|m_1 \geq \delta_1|R|m_1/8$ red edges between S and R . It follows from dependent random choice (see [18]) that we can find a subset $U \subset S$ of size $n^{1-\beta}$, such that every subset of $\lceil t/2 \rceil$ vertices from U has at least $n^{1-\beta}$ common red neighbours in R .

By Property (b), we find a red $K_{\lceil t/2 \rceil}$ or a blue K_s in U . In the latter case, we are done, so we may assume the former. Let A be the set of vertices of this $\lceil t/2 \rceil$ -clique, and let $W \subseteq R$ be a set of $n^{1-\beta}$ common red neighbours of A . Applying Property (b) to W , we again find a blue K_s , and are done, or find a set B of $\lceil t/2 \rceil$ vertices inducing a red clique. In this latter case, $A \cup B$ gives rise to a red clique on at least t vertices, and hence we have the desired contradiction.

This leaves us with the case $t = s$, where by symmetry we may assume that the majority of edges in $G \cup G(n, p)$ are coloured blue. For a contradiction, we suppose without loss of generality that there is some $u \in V_1$ with a set R of αm_1 red neighbours in V_2 . Since we assumed blue was the more popular colour, we can find some (ordered) pair (V_i, V_j) , $j \neq 2$, where at least one-third of the edges are blue. By averaging, this gives some vertex $v \in V_i$ and a set $S \subseteq V_j$ of αm_1 blue neighbours of v .

Now note that (V_2, V_j) is an ε_1 -regular pair, and so there are at least $(\delta_1 - \varepsilon_1)\alpha^2 m_1^2$ edges between R and S . If at least half of these edges were coloured red, then by averaging, we would find a vertex $u' \in R$ with a set U of at least $\delta_1 \alpha m_1 / 4$ red neighbours in S . Then the vertices u' and v , together with the set U , violate our initial observation. On the other hand, if half of the edges between R and S are blue, then we find a vertex $v' \in S$ with a set U' of at least $\delta_1 \alpha m_1 / 4$ blue neighbours in R . Then u, v' and U' violate our initial observation instead.

Hence, if the colouring of $G \cup G(n, p)$ has neither a red K_t nor a blue K_s , we may indeed assume that the maximum red-degree in each pair (V_i, V_j) is at most αm_1 . $\square$

3.3. Proof of Theorem 1.9. First we restate Theorem 1.9 in the following equivalent form.

Theorem 3.4. *Let $\ell, k \geq 3$ be integers.*

- (i) *If $k, \ell \in 2\mathbb{N}$ then $p(n; C_k, C_\ell, d) = 0$ for all $d > 0$.*
- (ii) *If $k \in 2\mathbb{N} + 1$ then $p(n; C_k, C_\ell, d) = 1/n$ for all $d \in (0, 1/2]$.*
- (iii) *If $k \in 2\mathbb{N} + 1$ and $\ell \in 2\mathbb{N}$ then $p(n; C_k, C_\ell, d) = 0$ for all $d > 1/2$.*
- (iv)

$$p(n; C_3, C_3, d) = \begin{cases} 1/n^2 & \text{if } d \in (1/2, 4/5] \\ 0 & \text{if } d > 4/5. \end{cases}$$

- (v) *If $k \in 2\mathbb{N} + 1$ and $\ell \in 2\mathbb{N} + 3$ then*

$$p(n; C_k, C_\ell, d) = \begin{cases} 1/n^2 & \text{if } d \in (1/2, 3/4] \\ 0 & \text{if } d > 3/4. \end{cases}$$

3.3.1. Proof of Theorem 3.4(i). Let $r := R_{\text{bip}}(k, \ell)$, the bipartite Ramsey number for $K_{k,k}$ and $K_{\ell,\ell}$. Then $K_{r,r}$ is a bipartite graph that is $(K_{k,k}, K_{\ell,\ell})$ -Ramsey, and therefore (C_k, C_ℓ) -Ramsey as well. By Observation 1.4(iii), it follows that $p(n; C_k, C_\ell, d) = 0$ for $d > 0$; that is, any sufficiently large dense graph will already be (C_k, C_ℓ) -Ramsey before any random edges are added. $\square$

3.3.2. Proof of Theorem 3.4(ii). The lower bound here is the trivial lower bound from Observation 1.4, since C_k is not bipartite and for $p = o(1/n)$, $G(n, p)$ is with high probability C_ℓ -free. Thus $p(n; C_k, C_\ell, d) \geq 1/n$ for all $d \in (0, 1/2]$.

Let $d > 0$ and set $0 < \eta \ll \varepsilon \ll \gamma \ll \delta \ll d, 1/k, 1/\ell$. Suppose that G is a sufficiently large n -vertex graph. Consider $G \cup G(n, p)$ where $p = \omega(1/n)$. By Corollary 2.4 (with $r = 1$) there exist disjoint $A, B \subseteq V(G)$ so that $|A| = |B| \geq \eta n$ and $(A, B)_G$ is ε -regular with density at least δ . We call an ordered $(k + \ell)$ -tuple $(x_1, \dots, x_k, y_1, \dots, y_\ell) \in A^{k+\ell}$ *good* if $|(\cap_i N_G(x_i)) \cap (\cap_i N_G(y_i)) \cap B| \geq \gamma|B|$. Lemma 2.2(i) implies that all but at most $\gamma|A|^{k+\ell}$ ordered $(k + \ell)$ -tuples in $A^{k+\ell}$ are good.

Using Theorem 2.7, we deduce that with high probability there is a good $(k + \ell)$ -tuple such that in $G(n, p)$, $x_1 x_2 \dots x_k x_1$ forms a k -cycle and $y_1 y_2 \dots y_\ell y_1$ gives an ℓ -cycle. Indeed, letting H be the vertex-disjoint union of C_k and C_ℓ , our above argument shows that the collection $\mathcal{H}$ of potential copies of H with good supports has size $\xi n^{k+\ell}$ for some $\xi > 0$. Moreover, we have $\mu_1 = (np)^{\min(k, \ell)} = \omega(1)$, and so the probability of having no copy of H supported on a good $(k + \ell)$ -tuple tends to zero as n grows. Therefore w.h.p. in $G(n, p)[A]$ we have a disjoint k -cycle X and ℓ -cycle Y together with a set $N \subseteq B$ where $|N| \geq \gamma|B|$ and $N \subseteq N_G(X \cup Y)$.

Consider any 2-colouring of $G \cup G(n, p)$. Given any $b \in B$ there are $2^{k+\ell}$ possible ways to colour those edges incident to b with an endpoint in $X \cup Y$. Thus, there is a set $N_1 \subseteq N$ of size at least $\gamma|B|/2^{k+\ell}$ where each vertex in N_1 has the same ‘colour profile’ (i.e. every vertex in N_1 has the same red neighbourhood in $X \cup Y$ and therefore the same blue neighbourhood in $X \cup Y$).

Colour each vertex $v \in X \cup Y$ with the colour of the edges it receives from N_1 . So we now have a red/blue colouring of the vertices and edges of $X \cup Y$. Suppose there is a red vertex $u \in X \cup Y$ and a blue vertex $v \in X \cup Y$. Theorem 2.10(ii) implies that w.h.p. $G(n, p)[N_1]$ is $(P_{k-1}, P_{\ell-1})$ -Ramsey. If there is a red copy of P_{k-1} in N_1 , then together with u we obtain our desired red copy of C_k in $G \cup G(n, p)$; otherwise we obtain a blue copy of $P_{\ell-1}$ in N_1 and thus together with v we obtain a blue copy of C_ℓ .

Thus, we may assume that every vertex in $X \cup Y$ is coloured the same. Suppose that they are all red. If the edges of Y are all blue, we obtain the desired blue copy of C_ℓ . So we may assume that there is at least one red edge in Y . As all the edges between $X \cup Y$ and N_1 are red, we can extend this red edge to a red copy of C_k in $G \cup G(n, p)$. The case when every vertex in $X \cup Y$ is blue is similar (and in fact easier if ℓ is even), and so in all cases we obtain a desired monochromatic cycle. $\square$

3.3.3. Proof of Theorem 3.4(iii). Let $d > 1/2$ and define $\delta > 0$ so that $d > 1/2 + \delta$. Set $0 < \varepsilon \ll \delta$. Let $\eta > 0$ be obtained by applying Corollary 2.4 with input $r = 2$. Consider any sufficiently large n -vertex graph G of density at least d , and consider any 2-colouring of G . By Corollary 2.4 we have that there are disjoint sets V_1, V_2, V_3 in G so that $|V_1| = |V_2| = |V_3| \geq \eta n$ and for each $1 \leq i < j \leq 3$, there is some colour $c_{i,j}$ for which the edges between V_i and V_j of colour $c_{i,j}$ form an ε -regular pair of density at least $\delta/4$. Suppose one of these colours $c_{i,j}$ is blue. Then Lemma 2.6 implies G contains a blue copy of C_ℓ . Otherwise all the $c_{i,j}$ are red and then since C_k is 3-partite, Lemma 2.6 implies G contains a red copy of C_k . $\square$

3.3.4. Proof of Theorem 3.4(iv). Let G_n denote the 5-partite Turán graph on n vertices. Since K_5 has a 2-colouring without a monochromatic copy of C_3 , so does G_n . Let $p = o(1/n^2)$. Then w.h.p. $G(n, p)$ is empty. Thus, w.h.p. $G_n \cup G(n, p)$ is not (C_3, C_3) -Ramsey. This shows that $p(n, C_3, C_3; d) \geq 1/n^2$ for all $d \leq 4/5$.

Next suppose that $d > 1/2$, and let $p = \omega(1/n^2)$. Since K_6 is (C_3, C_3) -Ramsey, to prove that $p(n; C_3, C_3, d) = 1/n^2$, it suffices to show that given any n -vertex graph G of density d , w.h.p. $K_6 \subseteq G \cup G(n, p)$. This follows immediately from Theorem 2.11 with $k = 3$, since $\rho_3(K_6) = 1/2$. So indeed $p(n; C_3, C_3, d) = 1/n^2$.

Let $d > 4/5$ and suppose that G is any sufficiently large graph with density at least d . Then by Turán’s theorem G contains a copy of K_6 . Since K_6 is (C_3, C_3) -Ramsey, any 2-colouring of G yields a monochromatic copy of C_3 . Thus, $p(n; C_3, C_3, d) = 0$. $\square$

3.3.5. *Proof of Theorem 3.4(v).* To prove this result we will need two additional lemmas.

Lemma 3.5. $R(\{C_3\}, \{C_3, C_5\}) = 5$.

Proof. To see that $R(\{C_3\}, \{C_3, C_5\}) > 4$, consider a 2-colouring of K_4 whose red edges induce a copy of P_4 and whose blue edges induce a copy of P_4 .

Next consider any 2-colouring of K_5 . If there is a vertex incident to at least three red edges then we must have a red or blue copy of C_3 . The same conclusion holds if there is a vertex incident to at least three blue edges. Thus, we may assume that every vertex has red and blue degree two. In particular, the blue subgraph is 2-regular and so is a copy of C_5 , as desired. $\square$

Let H_m be the graph formed by taking disjoint vertex sets $V_1, \dots, V_5$ each of size m , and with edge set as follows: H_m contains a perfect matching between V_1 and V_2 ; a perfect matching between V_3 and V_4 ; between all other pairs of distinct V_i there are all possible edges. The next result proves that finding H_m in a graph G (for m sufficiently large) ensures G is (C_k, C_ℓ) -Ramsey.

Lemma 3.6. *Given $k \in 2\mathbb{N} + 1$ and $\ell \in 2\mathbb{N} + 3$, there exists an $m_0 = m_0(k, \ell)$ such that if $m \geq m_0$, then H_m is (C_k, C_ℓ) -Ramsey.*

Proof. Consider any 2-colouring c of H_m . Given an edge $xy \in E(H_m)$, write $c(xy)$ for the colour of xy in c . We first build an auxiliary complete bipartite graph B with classes B_1 and B_2 . The vertices of B_1 are the edges $\{v_{1,i}v_{2,i} : i \in [m]\}$ of the perfect matching between V_1 and V_2 in H_m ; the vertices of B_2 are the edges $\{v_{3,i}v_{4,i} : i \in [m]\}$ of the perfect matching between V_3 and V_4 in H_m .

Next we 16-colour the edges of B as follows: for all $i, j \in [m]$, we colour the edge from $v_{1,i}v_{2,i}$ to $v_{3,j}v_{4,j}$ in B with the 4-tuple $(c(v_{1,i}v_{3,j}), c(v_{1,i}v_{4,j}), c(v_{2,i}v_{3,j}), c(v_{2,i}v_{4,j}))$. Since m is sufficiently large, the bipartite Ramsey theorem implies the existence of a monochromatic copy K of $K_{4(k+\ell), 4(k+\ell)}$ in B . Let $(c_{1,3}, c_{1,4}, c_{2,3}, c_{2,4})$ be the colour of K . Let $V'_i \subseteq V_i$ denote the set of vertices in V_i that are ‘present’ in K ; for example, $v_{1,j} \in V'_1$ precisely if $v_{1,j}v_{2,j}$ is a vertex in K . It follows that, for every $i \in \{1, 2\}$, $j \in \{3, 4\}$, all edges in the complete bipartite graphs $H_m[V'_i, V'_j]$ have the colour $c_{i,j}$.

Now consider the vertices in V_5 . There are $2^{16(k+\ell)}$ possible ways the edges between a vertex $v \in V_5$ and the vertices in $\cup_{i=1}^4 V'_i$ can be coloured. Hence, we can find a set $V_5'' \subseteq V_5$ of at least $m2^{-16(k+\ell)} \geq k + \ell$ vertices that all have the same colour profile.

Next consider the matching between V'_1 and V'_2 in H_m . For each edge v_1v_2 in this matching, there are four possible ways the edges from $\{v_1, v_2\}$ to V_5'' can be coloured. Hence, there are subsets $V_1'' \subseteq V'_1$ and $V_2'' \subseteq V'_2$ of size $k + \ell$ such that there is a perfect matching in $H_m[V_1'', V_2'']$, and for each $i \in [2]$, all edges between V_i'' and V_5'' have colour $c_{i,5}$. Similarly, there are subsets $V_3'' \subseteq V'_3$ and $V_4'' \subseteq V'_4$ of size $k + \ell$ such that there is a perfect matching in $H_m[V_3'', V_4'']$, and for each $i \in \{3, 4\}$, all edges between V_i'' and V_5'' have colour $c_{i,5}$.

In summary, we have an induced subgraph H'_m of H_m on $\cup_{i=1}^5 V_i''$ where $|V_i''| \geq k + \ell$; a perfect matching in $H'_m[V_1'', V_2'']$ and in $H'_m[V_3'', V_4'']$; for all other pairs $i < j$, a monochromatic complete bipartite graph of colour $c_{i,j}$ between V_i'' and V_j'' in H'_m .

Fix an edge x_1x_2 in $H'_m[V_1'', V_2'']$ and denote its colour in c by $c_{1,2}$. Similarly let x_3x_4 be an edge in $H'_m[V_3'', V_4'']$ with colour $c_{3,4}$. Now consider an auxiliary copy of K_5 with vertex set $[5]$, colouring each edge ij with colour $c_{i,j}$. By Lemma 3.5 we find a red C_3 , blue C_3 or blue C_5 .

Case 1: There is a monochromatic copy of C_3 on $\{i, j, r\}$ in K_5 . Suppose this C_3 is red; the blue case is analogous. Choose vertices $y_1 \in V_i''$, $y_2, y_4, \dots, y_{k-1} \in V_j''$ and $y_3, y_5, \dots, y_k \in V_r''$. These choices can be made arbitrarily unless $\{1, 2\} \subseteq \{i, j, r\}$ in which case we set $i := 1$, $j := 2$, $y_1 := x_1$ and $y_2 := x_2$; or if $\{3, 4\} \subseteq \{i, j, r\}$, in which case we set $i := 3$, $j := 4$, $y_1 := x_3$ and $y_2 := x_4$.

Note that $y_1 y_2 y_3 \dots y_k y_1$ is a red copy of C_k in $H'_m \subseteq H_m$, as desired.

Case 2: There is a blue copy $\pi_1 \pi_2 \pi_3 \pi_4 \pi_5 \pi_1$ of C_5 in K_5 . We may assume that $\pi_5 = 5$. Choose vertices $y_1 \in V''_{\pi_1}$, $y_2 \in V''_{\pi_2}$, $y_3 \in V''_{\pi_3}$, $y_4, y_6, \dots, y_{\ell-1} \in V''_{\pi_4}$ and $y_5, y_7, \dots, y_{\ell} \in V''_5$ such that $y_i = x_{\pi_i} \in V''_{\pi_i}$ for each $i \in [4]$, with the other vertices chosen arbitrarily. Then $y_1 y_2 y_3 \dots y_{\ell} y_1$ is a blue copy of C_{ℓ} in $H'_m \subseteq H_m$, as desired. $\square$

With Lemma 3.6 at hand it is now straightforward to prove Theorem 3.4(v).

Proof of Theorem 3.4(v) Let G_n denote the 4-partite Turán graph on n vertices. Colour the edges of K_4 so that the red edges induce a P_4 , and the blue edges also a P_4 . Lift this to a 2-colouring of G_n . Since the red (blue) subgraph of G_n is bipartite we do not have a red (blue) copy of any odd cycle. Let $p = o(1/n^2)$. Then w.h.p. $G(n, p)$ is empty. Thus, w.h.p. $G_n \cup G(n, p)$ is not (C_k, C_{ℓ}) -Ramsey. This shows that $p(n; C_k, C_{\ell}, d) \geq 1/n^2$ for all $d \leq 3/4$.

Next suppose that $d > 1/2$. Let $p = \omega(1/n^2)$, and let $m \geq m_0$ as in Lemma 3.6. Let G be a sufficiently large n -vertex graph of density at least d . The 3-partition $V(H_m) = (V_1 \cup V_2) \cup (V_3 \cup V_4) \cup V_5$ shows $\rho_3(H_m) \leq 1/2$, and so by Theorem 2.11, w.h.p. $H_m \subseteq G \cup G(n, p)$, and therefore $G \cup G(n, p)$ is (C_k, C_{ℓ}) -Ramsey. So indeed $p(n; C_k, C_{\ell}, d) = 1/n^2$ for all $d \in (1/2, 3/4]$.

Finally, suppose that $d > 3/4$ and let G be a sufficiently large graph of density d . Since H_m is 5-partite, the Erdős–Stone–Simonovits theorem [17] implies that $H_m \subseteq G$. Lemma 3.6 implies that G is (C_k, C_{ℓ}) -Ramsey. So $p(n; C_k, C_{\ell}, d) = 0$ for all $d > 3/4$. $\square$

3.4. Proof of Theorem 1.11. We next turn to our multicolour result, where we seek the threshold at which any r -colouring of $G \cup G(n, p)$ will contain a monochromatic copy of C_{ℓ} , where $\ell \geq 2^r + 1$ is odd. In the proof of the theorem we will repeatedly make use of the following simple property of 2^i -partite graphs.

Fact 3.7. *The edge set of any 2^i -partite graph H can be partitioned into i bipartite graphs.*

The 2^i is best possible, as shown by the following Ramsey result, which can be proven by induction on r .

Fact 3.8. *Given any $r \geq 1$, and any r -colouring of K_{2^r+1} , there is a monochromatic odd cycle in K_{2^r+1} .*

3.4.1. The case when $0 \leq d \leq 1 - 2^{-r+2}$. When the underlying graph G is not very dense, we can appeal to Observation 1.10. By Fact 3.7, any graph that is $(C_{\ell}, r-2)$ -Ramsey must have chromatic number at least $2^{r-2} + 1$, as otherwise its edges can be partitioned into $r-2$ bipartite (and hence C_{ℓ} -free) subgraphs. Thus, by Observation 1.10, if $d \leq 1 - 2^{-r+2}$, we have $p(n; r, C_{\ell}, d) = n^{-1/m_2(C_{\ell})} = n^{-1+1/(\ell-1)}$.

3.4.2. The case when $1 - 2^{-r+2} < d \leq 1 - 2^{-r+1}$. For each $n \in \mathbb{N}$, let G_n denote the 2^{r-1} -partite Turán graph on n -vertices. Let $p = o(n^{-1})$. By Fact 3.7 we can $(r-1)$ -colour G_n so that each colour class is bipartite, thus avoiding monochromatic copies of C_{ℓ} . A simple application of Markov's inequality yields that w.h.p. $G(n, p)$ does not contain a copy of C_{ℓ} , and thus its edges can be coloured with the remaining colour. Together this implies that w.h.p. $G_n \cup G(n, p)$ is not (C_{ℓ}, r) -Ramsey. Hence, $p(n; r, C_{\ell}, d) \geq n^{-1}$ for all $d \leq 1 - 2^{-r+1}$. As in the previous case, the upper bound follows from Theorem 1.1, since $n^{-1+1/(\ell-1)}$ is the threshold for $G(n, p)$ itself to be (C_{ℓ}, r) -Ramsey. This shows $p(n; r, C_{\ell}, d) \in [n^{-1}, n^{-1+1/(\ell-1)}]$ for all $1 - 2^{-r+2} < d \leq 1 - 2^{-r+1}$.

3.4.3. The case when $1 - 2^{-r+1} < d \leq 1 - 2^{-r}$. For each $n \in \mathbb{N}$, let G_n denote the 2^r -partite Turán graph on n -vertices. Let $p = o(n^{-2})$. By Fact 3.7 we can r -colour G_n so that there are no monochromatic copies of C_{ℓ} . Further w.h.p. $G(n, p)$ is empty. So w.h.p. $G_n \cup G(n, p)$ is not (C_{ℓ}, r) -Ramsey. Hence, $p(n; r, C_{\ell}, d) \geq n^{-2}$ for all $1 - 2^{-r+1} < d \leq 1 - 2^{-r}$.

To finish this case off, we require the following generalisation of the graph H_m introduced in the proof of Theorem 1.9. Let $H_{m,r}$ be the graph formed by taking disjoint vertex sets $V_1, \dots, V_{2^r+1}$ each of size m , and with edge set as follows: $H_{m,r}$ contains a perfect matching between V_{2i-1} and V_{2i} for each $1 \leq i \leq 2^{r-1}$; between all other distinct pairs V_i, V_j there are all possible edges. Note that $H_m = H_{m,2}$.

Lemma 3.9. *Given $\ell \geq 2^r + 1$, there exists an $m_0 = m_0(r, \ell)$ such that if $m \geq m_0$, then $H_{m,r}$ is (C_ℓ, r) -Ramsey.*

Proof. Let m be sufficiently large and consider any r -colouring c of $H_{m,r}$. By repeatedly arguing as in the proof of Lemma 3.6 we can obtain subsets $V_i^* \subseteq V_i$ (for each $i \leq 2^r$) of size $r^3 \ell$ so that:

- There is a perfect matching from V_{2i-1}^* to V_{2i}^* for every $1 \leq i \leq 2^{r-1}$;
- For all other pairs (V_i^*, V_j^*) (with $i < j \leq 2^r$), all edges from V_i^* to V_j^* have the same colour $c_{i,j}$.

To construct such sets V_i^* we first construct the auxiliary bipartite graph B precisely as in Lemma 3.6; this yields sets $V_1', \dots, V_4'$ (whose sizes will now still be huge). Then we construct an analogous auxiliary graph for the pair $((V_1', V_2'), (V_5, V_6))$, and so on, until we have considered every pair of tuples of such vertex classes.

By further restricting to the most popular colour within each of the perfect matchings, we can now assume that each such set has size $r^2 \ell$ and

- There is a *monochromatic* perfect matching from V_{2i-1}^* to V_{2i}^* for every $1 \leq i \leq 2^{r-1}$. Denote this colour by $c_{2i-1,2i}$.

Now consider the vertices in V_{2^r+1} . There are $r^{r^2 \ell 2^r}$ possible ways the edges between a vertex $v \in V_{2^r+1}$ and the vertices in $\cup_{i=1}^{2^r} V_i^*$ can be coloured. Hence, we can find a set $V_{2^r+1}^{**} \subseteq V_{2^r+1}$ of size at least $m r^{-r^2 \ell 2^r} \geq \ell$ vertices that all have the same colour profile.

Next consider the matching between V_{2i-1}^* and V_{2i}^* in $H_{m,r}$ for all $1 \leq i \leq 2^{r-1}$. For each edge $v_{2i-1} v_{2i}$ in this matching, there are r^2 possible ways the edges from $\{v_{2i-1}, v_{2i}\}$ to $V_{2^r+1}^{**}$ can be coloured. Hence, for each $1 \leq j \leq 2^r$, there are subsets $V_j^{**} \subseteq V_j^*$ of size ℓ such that all edges between V_j^{**} and $V_{2^r+1}^{**}$ have colour $c_{j,2^r+1}$, and such that there is a perfect matching in $H_{m,r}[V_{2i-1}^{**}, V_{2i}^{**}]$ for each $1 \leq i \leq 2^{r-1}$.

In summary, we have an induced subgraph $H_{m,r}^*$ of $H_{m,r}$ on $\cup_{j=1}^{2^r+1} V_j^{**}$ with $|V_j^{**}| \geq \ell$, a monochromatic perfect matching of colour $c_{2i-1,2i}$ in $H_{m,r}^*[V_{2i-1}^{**}, V_{2i}^{**}]$ for all $1 \leq i \leq 2^{r-1}$, and, for all other pairs $1 \leq i < j \leq 2^r + 1$, a monochromatic complete bipartite graph of colour $c_{i,j}$ between V_i^{**} and V_j^{**} in $H_{m,r}^*$.

Now consider an auxiliary copy of K_{2^r+1} with vertex set $[2^r + 1]$, colouring each edge ij with colour $c_{i,j}$. By Fact 3.8 we find a monochromatic odd cycle $C_{\ell'} = x_1 \dots x_{\ell'} x_1$ in K_{2^r+1} where $2 \leq \ell' \leq 2^r + 1 \leq \ell$. It is easy to see that there is a homomorphism ϕ from C_ℓ to $C_{\ell'}$ so that for all but one of the edges $x_i x_{i+1}$, precisely one edge is mapped onto $x_i x_{i+1}$ by ϕ . We can thus ensure that all edges corresponding to parts with a perfect matching between them are mapped to exactly once, and it is then easy to lift the $C_{\ell'}$ to a monochromatic copy of C_ℓ in $H_{m,r}^*$, as desired. $\square$

Returning to Theorem 1.11, suppose that $d > 1 - 2^{-r+1}$. Let $p = \omega(1/n^2)$, and let $m \geq m_0$ as in Lemma 3.9. Let G be a sufficiently large n -vertex graph of density at least d . Pairing up the parts of $H_{m,r}$ joined by a perfect matching gives a $(2^{r-1} + 1)$ -partition of $V(H_{m,r})$ that shows $\rho_{2^{r-1}+1}(H_{m,r}) \leq 1/2$. Theorem 2.11 then implies that w.h.p. $G_n \cup G(n, p)$ contains a copy of $H_{m,r}$. Thus, by Lemma 3.9, w.h.p. $G_n \cup G(n, p)$ is (C_ℓ, r) -Ramsey. So indeed $p(n; r, C_\ell, d) = n^{-2}$ for all $1 - 2^{-r+1} < d \leq 1 - 2^{-r}$.

3.4.4. *The case when $1 - 2^{-r} < d \leq 1$.* Suppose $d > 1 - 2^{-r}$, and G is a sufficiently large graph of density d . Since the graph $H_{m,r}$ from Lemma 3.9 is $(2^r + 1)$ -partite, it follows from the Erdős–Stone–Simonovits theorem [17] that $H_{m,r} \subseteq G$. Hence, by Lemma 3.9, G is (C_ℓ, r) -Ramsey, and thus $p(n; r, C_\ell, d) = 0$ for $d > 1 - 2^{-r}$, completing the proof of the theorem.

3.5. Proof of Theorem 1.12. In our final result, we establish the perturbed Ramsey thresholds for cliques versus odd cycles by showing $p(n; K_t, C_\ell, d) = n^{-2/(t-1)}$ for $t \geq 4$, odd $\ell \geq 5$ and $d \in (0, 1/2]$.

Proof of Theorem 1.12. First note that the lower bound follows from Observation 1.4(i). Indeed, we can take G to be the balanced complete bipartite n -vertex graph, which is C_ℓ -free and of density at least $1/2$, and colour all its edges blue. If $p = o(n^{-2/(t-1)})$, then with high probability $G(n, p)$ is K_t -free, and so we can colour the remaining edges of $G(n, p)$ red, obtaining an edge-colouring of $G \cup G(n, p)$ without any red K_t or blue C_ℓ .

For the upper bound, let $p = \omega(n^{-2/(t-1)})$. Apply Corollary 2.4 with $\delta := d, k := 2, r = 1$ and $\varepsilon := (d/4)^{t+\ell}$ to the uncoloured graph G of density d , obtaining an ε -regular pair (V_1, V_2) of density at least $d/2$, with $|V_1| = |V_2| = \eta n$ for some $\eta > 0$. We shall show that $G \cup G(n, p)$ is (K_t, C_ℓ) -Ramsey provided $G(n, p)$ has the three properties given below, where H is the vertex-disjoint union of K_t and C_ℓ , and $\alpha := (d/8)^{t+\ell}$:

- (a) $G(n, p)[V_1]$ contains a copy of H with at least $\alpha 2^{t+\ell} |V_2|$ common neighbours in V_2 ,
- (b) $G(n, p)$ is $\alpha\eta$ -globally $(K_{t-1}, P_{\ell-1})$ -Ramsey, and
- (c) $G(n, p)$ is $\alpha\eta$ -globally (K_{t-2}, C_ℓ) -Ramsey.

We first show that (a) holds with high probability. By Lemma 2.2(i), we know that at least half of all $(t + \ell)$ -sets of vertices in V_1 have at least $(d/2 - \varepsilon)^{t+\ell} |V_2|$ common neighbours in V_2 , and $(d/2 - \varepsilon)^{t+\ell} \geq \alpha 2^{t+\ell}$. Let $\mathcal{H}$ be the collection of all possible copies of H supported on such $(t + \ell)$ -sets of V_1 ; it follows that $|\mathcal{H}| \geq \xi n^{t+\ell}$ for some constant $\xi = \xi(t, \ell, d) > 0$. Furthermore, as $p = \omega(n^{-2/(t-1)})$, we have $\mu_1(H) = n^t p^{\binom{t}{2}} = \omega(1)$ (attained by taking $F = K_t$ in the definition of μ_1). Thus, by Theorem 2.7, the probability that $G(n, p)[V_1]$ does not contain a suitable copy of H is at most $\exp(-\xi \omega(1)/(2^{t+\ell+1}(t + \ell)!)) = o(1)$.

For (b), we appeal to Theorem 2.10. Observe that $m_2(K_{t-1}) \geq m_2(P_{\ell-1}) = 1$, K_{t-1} is strictly balanced with respect to $m_2(\cdot, P_{\ell-1})$, and that $m_2(K_{t-1}, P_{\ell-1}) = (t - 1)/2$. Thus there is some constant $C' = C'(\alpha\eta, K_{t-1}, P_{\ell-1})$ such that for $p' \geq C' n^{-2/(t-1)}$, $G(n, p')$ satisfies (b) with high probability (so certainly our choice of p satisfies (b) with high probability).

For (c), the case $t \geq 5$ can be handled with Theorem 2.10 just as above. If $t = 4$, though, being (K_{t-2}, C_ℓ) -Ramsey is equivalent to containing C_ℓ . For this, we can apply Theorem 2.7 instead. For every set $U \subset V(G(n, p))$ of size $\alpha \nu n$, let $\mathcal{H}_U$ be the possible copies of C_ℓ within U , observing that $|\mathcal{H}_U| = (\ell - 1)! \binom{|U|}{\ell} / 2 = \xi' n^\ell$ for some appropriate constant $\xi' > 0$. Moreover, as $p = \omega(n^{-2/3})$, we have $\mu_1(C_\ell) = n^\ell p^\ell = \omega(n^{\ell/3}) \geq n^{5/3}$. By Theorem 2.7, the probability that $G(n, p)$ does not contain a copy of C_ℓ from $\mathcal{H}_U$ is at most $\exp(-\xi' n^{5/3}/(2^{\ell+1}\ell!))$. Taking a union bound over all possible choices of U , we see that with high probability, $G(n, p)$ will contain an ℓ -cycle in each subset U , thus satisfying (c).

To finish, let us see how these properties imply that $G \cup G(n, p)$ is (K_t, C_ℓ) -Ramsey. Consider any 2-colouring of $G \cup G(n, p)$. By (a), we can find a copy H_0 of H within $G(n, p)[V_1]$ with a set $W \subset V_2$ of at least $\alpha 2^{t+\ell} |V_2|$ common neighbours of $V(H_0)$. For each $w \in W$, there are $2^{t+\ell}$ ways the edges between w and $V(H_0)$ can be coloured, and so we can find a subset $U \subset W$ of $\alpha |V_2|$ common neighbours such that each vertex in U has the same colour profile to $V(H_0)$ (i.e. the red neighbourhood in $V(H_0)$ of each $u \in U$ is the same; as is the blue neighbourhood).

First suppose there are $x, y \in V(H_0)$ such that the edges from x to U are all red while those from y to U are all blue. By (b), we find a red K_{t-1} or a blue $P_{\ell-1}$ in $G(n, p)[U]$. Extending by either x or y respectively, we find either a red K_t or a blue C_ℓ in $G \cup G(n, p)$, and are done.

Next, suppose all edges between $V(H_0)$ and U are red. Recall that H_0 contains C_ℓ as a subgraph. If all edges of this ℓ -cycle are blue, we are done, and hence there must be a red edge $\{x, y\}$. By (c), U must contain either a red K_{t-2} or a blue C_ℓ . In the latter case, we are done, while in the former, extending the clique by x and y gives the desired red K_t .

Thus all edges between $V(H_0)$ and U must be blue. In this case, recall that H_0 contains K_t as a subgraph. If all its edges are red, then we are done, and hence there is a blue edge $\{u_0, u_1\}$. Writing $\ell = 2k + 1$, let $\{u_2, u_3, \dots, u_k\}$ be a set of $k - 1$ other vertices from $V(H_0)$, and let $\{v_1, v_2, \dots, v_k\}$ be an arbitrary set of k vertices from U . We then have a blue ℓ -cycle $(u_0, u_1, v_1, u_2, v_2, \dots, u_k, v_k, u_0)$ in $G \cup G(n, p)$, as required. $\square$

4. CONCLUDING REMARKS AND OPEN PROBLEMS

In this paper we have determined how many random edges one must add to a dense graph to ensure w.h.p. the resulting graph is (K_t, K_s) -Ramsey for all $t \geq s \geq 5$; more precisely $p(n; K_t, K_s, d) = n^{-1/m_2(K_t, K_{\lceil s/2 \rceil})}$ for all $0 < d \leq 1/2$ (see Theorem 1.7). This complements work of Krivelevich, Sudakov and Tetali [30] who determined the corresponding threshold when $s = 3$. This leaves only one open case for this range of densities; when $s = 4$. This is perhaps the most interesting open question resulting from our work. As mentioned previously, our work combined with Powierski's [37] settles this problem when $t = 4$. The construction from [37] can be extended to the (K_t, K_4) -Ramsey question as well, and improves the lower bound from Proposition 1.8 for $t \in \{5, 6\}$. It would be of great interest to determine the correct threshold for $t \geq 5$.

Note Theorem 1.7 does not consider all possible densities d ; indeed, it is still an open problem to determine how many random edges one must add to a graph G of density d to ensure it is w.h.p. (K_t, K_s) -Ramsey if $d > 1 - 2/(s - 1)$ and $s \leq t$. Notice though that if $d > 1 - \frac{1}{r-1}$ where $r := R(K_t, K_s)$, then by Turán's theorem G contains a copy of K_r and thus is (K_t, K_s) -Ramsey. Further, setting $r' := \lceil R(K_t, K_s)/2 \rceil$, Theorem 2.11 implies that if $d > 1 - \frac{1}{r'-1}$ then one only requires $p = \omega(n^{-2})$ to ensure $G \cup G(n, p)$ w.h.p. contains K_r and thus is (K_t, K_s) -Ramsey.

We have completely resolved the question of how many random edges one must add to a graph of fixed density to ensure w.h.p. the resulting graph is (C_k, C_ℓ) -Ramsey for all $k, \ell \geq 3$ (see Theorem 1.9). Additionally we made some progress towards the analogous question for more colours via Theorem 1.11; it would be interesting to close the one remaining gap in this statement (i.e. fully resolve the case when $1 - 2^{-r+1} < d \leq 1 - 2^{-r+1}$). Our work here is also related to a conjecture of Erdős and Graham [16] from 1973. Indeed, recall that Theorem 1.11 only considers cycles C_ℓ that are sufficiently large compared to the number of colours r (i.e. $\ell \geq 2^r + 1$). This is because we apply Fact 3.8: any r -colouring of K_{2^r+1} yields a monochromatic odd cycle. In particular, this monochromatic cycle could have length $2^r + 1$. Erdős and Graham [16] asked how large can the smallest monochromatic odd cycle in an r -colouring of K_{2^r+1} be. Thus, getting non-trivial upper bounds on this question would yield a strengthening of Theorem 1.11. We are not aware of such progress on this problem, though Day and Johnson [14] did prove the smallest such monochromatic cycle is unbounded as r grows, thereby answering a question of Chung [11].

Note that when the density d of the graph G is large, our upper bounds on $p(n; H_1, H_2, d)$ often come from the existence of a fixed-size subgraph that is (H_1, H_2) -Ramsey. However, for small densities, the upper bounds use more global arguments, either using results on the Ramsey properties of the random graph, or finding large common neighbourhoods and then finding subgraphs there. It would be interesting to see to what extent this dichotomy extends to other cases of the Ramsey problem in randomly perturbed graphs.

Finally, there has been significant interest in Ramsey properties of random hypergraphs (see, for example, [13, 19, 20]). It would be interesting to obtain analogues of our results in the setting of randomly perturbed hypergraphs.

ACKNOWLEDGEMENTS

We are grateful to Patrick Morris, whose advice was of great help in proving Claim A.9 from the appendix. We are also grateful to the referee for a careful and helpful review.

REFERENCES

- [1] E. Aigner-Horev and Y. Person, Monochromatic Schur triples in randomly perturbed dense sets of integers, *SIAM J. Discrete Math.* **33** (2019), 2175–2180.
- [2] N. Alon and J. Spencer, *The Probabilistic Method* (John Wiley & Sons, 2015).
- [3] J. Balogh, R. Morris and W. Samotij, Independent sets in hypergraphs, *J. Am. Math. Soc.* **28** (2015), 669–709.
- [4] J. Balogh, A. Treglown and A.Z. Wagner, Tilings in randomly perturbed dense graphs, *Comb. Probab. Comput.* **28** (2019), 159–176.
- [5] W. Bedenknecht, J. Han, Y. Kohayakawa and G. O. Mota, Powers of tight Hamilton cycles in randomly perturbed hypergraphs, *Random Struct. Algor.* **55** (2019), 795–807.
- [6] P. Bennett, A. Dudek and A. Frieze, Adding random edges to create the square of a Hamilton cycle, *arXiv:1710.02716* (2017).
- [7] T. Bohman, A. Frieze, M. Krivelevich and R. Martin, Adding random edges to dense graphs, *Random Struct. Algor.* **24** (2004), 105–117.
- [8] T. Bohman, A. Frieze and R. Martin, How many edges make a dense graph Hamiltonian?, *Random Struct. Algor.* **22** (2003), 33–42.
- [9] J. Böttcher, J. Han, Y. Kohayakawa, R. Montgomery, O. Parczyk and Y. Person, Universality for bounded degree spanning trees in randomly perturbed graphs, *Random Struct. Algor.* **55** (2019), 854–864.
- [10] J. Böttcher, R. Montgomery, O. Parczyk and Y. Person, Embedding spanning bounded degree graphs in randomly perturbed graphs, *Mathematika*, to appear.
- [11] F. Chung, Open problems of Paul Erdős in graph theory, *J. Graph Theor.* **25** (1997), 3–36.
- [12] D. Conlon, A new upper bound for diagonal Ramsey numbers, *Ann. Math.* **170** (2009), 941–960.
- [13] D. Conlon and W.T. Gowers, Combinatorial theorems in sparse random sets, *Ann. Math.* **84** (2016), 367–454.
- [14] A.N. Day and J.R. Johnson, Multicolour Ramsey numbers of odd cycles, *J. Comb. Theory Ser. B* **124** (2017), 56–63.
- [15] A. Dudek, C. Reiher, A. Ruciński, M. Schacht, Powers of Hamiltonian cycles in randomly augmented graphs, *Random Struct. Algor.* **56** (2020), 122–141.
- [16] P. Erdős and R.L. Graham, On Partition Theorems for Finite Graphs, *Colloq. Math. Soc. J. B.* **10** (1973), 515–527.
- [17] P. Erdős and A.H. Stone, On the structure of linear graphs, *B. Am. Math. Soc.* **52** (12) (1946), 1087–1091.
- [18] J. Fox and B. Sudakov, Dependent random choice, *Random Struct. Algor.* **38** (2011), 68–99.
- [19] E. Friedgut, V. Rödl and M. Schacht, Ramsey properties of random discrete structures, *Random Struct. Algor.* **37** (2010), 407–436.
- [20] L. Gugelmann, R. Nenadov, Y. Person, N. Škorić, A. Steger and H. Thomas, Symmetric and asymmetric Ramsey properties in random hypergraphs, *Forum Math., Sigma* **5**, Cambridge University Press (2017).
- [21] J. Han and Y. Zhao, Hamiltonicity in randomly perturbed hypergraphs, *J. Comb. Theory Ser. B*, to appear.
- [22] R. Hancock, K. Staden and A. Treglown, Independent sets in hypergraphs and Ramsey properties of graphs and the integers, *SIAM J. Discrete Math.* **33** (2019), 153–188.
- [23] S. Janson, T. Łuczak and A. Ruciński, *Random Graphs* (Wiley, New York, 2000).
- [24] F. Joos and J. Kim, Spanning trees in randomly perturbed graphs, *Random Struct. Algor.* **56** (2020), 169–219.
- [25] Y. Kohayakawa and B. Kreuter, Threshold functions for asymmetric Ramsey properties involving cycles, *Random Struct. Algor.* **11** (1997), 245–276.
- [26] J. Komlós and M. Simonovits, Szemerédi’s Regularity Lemma and its applications in graph theory, *Combinatorics, Paul Erdős is Eighty* (Volume 2), Keszthely (Hungary), 1993, (D. Miklós, V. T. Sós, T. Szönyi eds.), Bolyai Math. Stud., Budapest (1996), 295–352.
- [27] B. Kreuter, Threshold functions for asymmetric Ramsey properties with respect to vertex colorings, *Random Struct. Algor.* **9** (1996), 335–348.
- [28] M. Krivelevich, M. Kwan and B. Sudakov, Cycles and matchings in randomly perturbed digraphs and hypergraphs, *Comb. Probab. Comput.* **25** (2016), 909–927.

- [29] M. Krivelevich, M. Kwan and B. Sudakov, Bounded-degree spanning trees in randomly perturbed graphs, *SIAM J. Discrete Math.* **31** (2017), 155–171.
- [30] M. Krivelevich, B. Sudakov and P. Tetali, On smoothed analysis in dense graphs and formulas, *Random Struct. Algor.* **29** (2006), 180–193.
- [31] T. Łuczak, A. Ruciński and B. Voigt, Ramsey properties of random graphs, *J. Comb. Theory Ser. B* **56** (1992), 55–68.
- [32] M. Marciniszyn, J. Skokan, R. Spöhel and A. Steger, Asymmetric Ramsey properties of random graphs involving cliques, *Random Struct. Algor.* **34** (2009), 419–453.
- [33] A. McDowell and R. Mycroft, Hamilton ℓ -cycles in randomly perturbed hypergraphs, *Electron. J. Comb.* **25** (2018), P4.36.
- [34] F. Mousset, R. Nenadov and W. Samotij, Towards the Kohayakawa–Kreuter conjecture on asymmetric Ramsey properties, *arXiv:1808.05070* (2018).
- [35] R. Nenadov and A. Steger, A short proof of the random Ramsey theorem, *Comb. Probab. Comput.* **25** (2016), 130–144.
- [36] R. Nenadov and M. Trujić, Sprinkling a few random edges doubles the power, *arXiv:1811.09209* (2018).
- [37] E. Powierski, Ramsey properties of randomly perturbed dense graphs, *arXiv:1902.02197* (2019).
- [38] V. Rödl and A. Ruciński, Lower bounds on probability thresholds for Ramsey properties, *Combinatorics, Paul Erdős is Eighty* (Volume 1), Keszthely (Hungary), 1993, (D. Miklós, V. T. Sós, T. Szőnyi eds.), Bolyai Math. Stud., Budapest (1993), 317–346.
- [39] V. Rödl and A. Ruciński, Random graphs with monochromatic triangles in every edge coloring, *Random Struct. Algor.* **5** (1994), 253–270.
- [40] V. Rödl and A. Ruciński, Threshold functions for Ramsey properties, *J. Am. Math. Soc.* **8** (1995), 917–942.
- [41] D. Saxton and A. Thomason, Hypergraph containers, *Invent. Math.* **201** (2015), 925–992.
- [42] J.H. Spencer, Ramsey’s theorem – a new lower bound, *J. Comb. Theory Ser. A* **18** (1975), 108–115.
- [43] E. Szemerédi, Regular partitions of graphs, *Colloques Internationaux C.N.R.S. No 260 – Problèmes Combinatoires et Théorie des Graphes*, Orsay (1976), 399–401.
- [44] P. Turán, On an extremal problem in graph theory (in Hungarian), *Math. Fiz. Lapok* **48** (1941), 436–452.

APPENDIX A. PROOFS OF USEFUL TOOLS

In this appendix we prove some of the results from Section 2 namely; Lemma 2.5, a multipartite version of the dependent random choice lemma; Theorem 2.7, which guarantees the existence of subgraphs within $G(n, p)$; Theorem 2.10, which shows that when random graphs have certain Ramsey properties, they have them robustly and globally. These follow from either fairly standard arguments or minor modifications of existing proofs, but we include these results here for the sake of completeness.

A.1. Proof of Lemma 2.5. In Lemma 2.5, copied below, we extend the basic lemma of dependent random choice from [18], showing that if a set of vertices V_s is in ε -regular pairs with other sets $V_1, V_2, \dots, V_{s-1}$, then we can find a relatively large subset $U \subseteq V_s$ such that all small subsets of U have many common neighbours in each of the other parts V_i .

Lemma 2.5. *Given $s \in \mathbb{N}$ and $\delta > 0$, suppose $0 < \varepsilon \leq \min\{1/(2s), \delta/2\}$. Let $V_1, V_2, \dots, V_s$ be disjoint sets of vertices from a graph G , each of size m , such that for all $i \in [s-1]$, the pair (V_i, V_s) is ε -regular of density at least δ . For every $\beta > 0$ and $\ell \in \mathbb{N}$ there is some $\gamma = \gamma(s, \delta, \beta, \ell) > 0$ such that, if m is sufficiently large, there is a subset $U_s \subseteq V_s$ of size at least $m^{1-\beta}$ such that every set of ℓ vertices from U_s has at least γm common neighbours in each of the sets V_i , $i \in [s-1]$.*

Proof. For $i \in [s-1]$, let $T_i \subset V_i$ be the subset of vertices obtained when selecting t vertices uniformly and independently (with repetition), where t is to be determined. Let $W_s := \{v \in V_s : \cup_i T_i \subseteq N(v)\}$ be those vertices in V_s adjacent to all chosen vertices. For fixed $v \in V_s$, we have

$$\mathbb{P}(v \in W_s) = \prod_{i=1}^{s-1} \left(\frac{|N(v) \cap V_i|}{m} \right)^t.$$

24

By Lemma 2.2(i), with $\ell = 1$, the number of vertices in V_s with at most $(\delta - \varepsilon)m$ neighbours in V_i for some $i \in [s - 1]$ is at most $(s - 1)\varepsilon m$. Hence for at least $(1 - (s - 1)\varepsilon)m$ vertices in V_s we have $|N(v) \cap V_i| > (\delta - \varepsilon)m$ for all $i \in [s - 1]$. Thus $\mathbb{E}[|W_s|] \geq (1 - (s - 1)\varepsilon)m(\delta - \varepsilon)^{(s-1)t} \geq (m/2)(\delta/2)^{(s-1)t}$.

We now set $\gamma := (\delta/2)^{2(s-1)\ell/\beta}$. For every ℓ -set $S \subseteq W_s$ with fewer than γm common neighbours in some V_i , remove one vertex of S from W_s , and let U_s be the set of remaining vertices. Clearly every ℓ -subset of U_s has the requisite number of common neighbours, and so we need only show that U_s is sufficiently large.

Observe that if $S \subseteq W_s$, then we must have $T_i \subseteq N(S) \cap V_i$ for each $i \in [s - 1]$. Thus, if S has fewer than γm common neighbours in some part V_i , we can bound $\mathbb{P}(S \subseteq W_s) \leq \gamma^t$. In expectation, the number of vertices that needs to be removed from W_s is thus at most $\binom{m}{\ell} \gamma^t$. This leaves us with

$$\mathbb{E}[|U_s|] \geq \frac{m}{2} \left(\frac{\delta}{2}\right)^{(s-1)t} - \binom{m}{\ell} \gamma^t \geq \frac{m}{2} \left(\frac{\delta}{2}\right)^{(s-1)t} - m^\ell \gamma^t.$$

Thus, if we set $t := \lfloor \log(4m^{-\beta}) / ((s - 1) \log(\delta/2)) \rfloor$, the first term will be at least $2m^{1-\beta}$, while our choice of γ ensures $m^\ell \gamma^t = o(1)$. Hence we can find some set $U_s \subset V_s$ of size at least $m^{1-\beta}$ with the desired property. $\square$

A.2. Proof of Theorem 2.7. The next result is a standard, straightforward, and very useful application of the Janson inequality [23].

Theorem 2.7. *Let H be a graph with $v \geq 2$ vertices and $e \geq 1$ edges. Let $[n]$ be the vertex set of $G(n, p)$, and, for some $\xi > 0$, let $\mathcal{H}$ be a collection of ξn^v possible copies of H supported on $[n]$. The probability that $G(n, p)$ does not contain any copy of H from $\mathcal{H}$ is at most $\exp(-\xi \mu_1 / (2^{v+1} v!))$, where $\mu_1 = \mu_1(H) := \min\{n^{v(F)} p^{e(F)} : F \subseteq H, e(F) \geq 1\}$.*

We separate a calculation that will be needed in later proofs as well. The quantity Δ bounds the expected number of pairs of distinct edge-intersecting copies of H , with the first copy from $\mathcal{H}$, that appear in $G(n, p)$, a quantity closely linked to the variance of the number of copies of H from $\mathcal{H}$ in $G(n, p)$.

Lemma A.1. *Let H be a graph with v vertices and e edges, let $\mathcal{K}_n(H)$ be the collection of all copies of H in the complete graph K_n , and let $\mathcal{H} \subseteq \mathcal{K}_n(H)$ be a subcollection thereof. We then have*

$$\Delta := \sum_{\substack{H' \in \mathcal{H}, H'' \in \mathcal{K}_n(H): \\ H' \neq H'', e(H' \cap H'') \neq 0}} \mathbb{P}(H' \cup H'' \subseteq G(n, p)) \leq 2^v v! |\mathcal{H}| n^v p^{2e} / \mu_0,$$

where $\mu_0 = \mu_0(H) := \min\{n^{v(F)} p^{e(F)} : F \subsetneq H, e(F) \geq 1\}$.

Proof. We write $H' \sim H''$ if H', H'' are distinct copies of H that share at least an edge, with $H' \in \mathcal{H}$ and $H'' \in \mathcal{K}_n(H)$. Thus Δ denotes the expected number of (ordered) pairs $H' \sim H''$ with both H' and H'' appearing in $G(n, p)$, which we want to bound from above.

Given $H' \in \mathcal{H}$, let us estimate its contribution to Δ . There are fewer than 2^v ways to choose a subset $S \subseteq V(H')$ of the vertices of H' that are shared with H'' , and at most $v!$ assignments φ of these vertices to the vertices of H'' . This then determines the subgraph $F = F(S, \varphi) \subset H$ in which H' and H'' intersect. Let $\mathcal{F} := \{(S, \varphi) : e(F(S, \varphi)) \geq 1\}$ denote the set of viable pairs (S, φ) . For each such pair, letting $F = F(S, \varphi)$, there are at most $n^{v-v(F)}$ choices for the remaining vertices of H'' , and each such copy H'' introduces a further $e - e(F)$ edges. Hence the probability that both H' and H'' appear in $G(n, p)$ is $p^{2e-e(F)}$. This gives

$$\Delta \leq \sum_{H' \in \mathcal{H}} \sum_{(S, \varphi) \in \mathcal{F}} n^{v-v(F)} p^{2e-e(F)} = n^v p^{2e} \sum_{H' \in \mathcal{H}} \sum_{(S, \varphi) \in \mathcal{F}} n^{-v(F)} p^{-e(F)}.$$

The summand is, by definition of μ_0 , at most μ_0^{-1} . Since there are at most $2^v v!$ choices of (S, φ) in the inner sum, and $|\mathcal{H}|$ choices of H' in the outer sum, we get the desired bound of

$$\Delta \leq 2^v v! |\mathcal{H}| n^v p^{2e} / \mu_0. \quad \square$$

Theorem 2.7 follows almost immediately from the lemma.

Proof of Theorem 2.7. Let $X_{\mathcal{H}}$ be the random variable counting the number of copies of H from $\mathcal{H}$ that appear in $G(n, p)$, so that we seek to bound $\mathbb{P}(X_{\mathcal{H}} = 0)$ from above. We clearly have $\mu := \mathbb{E}[X_{\mathcal{H}}] = |\mathcal{H}| p^e = \xi n^v p^e$. Thus, letting Δ and μ_0 be as in Lemma A.1, we have $\Delta \leq 2^v v! \mu^2 / (\xi \mu_0)$. Furthermore, we have $\mu_1 = \min\{\mu/\xi, \mu_0\}$.

By the Janson inequality (see [2, Theorem 8.1.1]), if $\Delta \leq \mu$, then

$$\mathbb{P}(X_{\mathcal{H}} = 0) \leq \exp(-\mu/2) \leq \exp(-\xi \mu_1 / (2^{v+1} v!)).$$

Otherwise, when $\Delta > \mu$, we can instead apply the extended Janson inequality [2, Theorem 8.1.2]:

$$\mathbb{P}(X_{\mathcal{H}} = 0) \leq \exp(-\mu^2 / (2\Delta)) \leq \exp(-\xi \mu_0 / (2^{v+1} v!)) \leq \exp(-\xi \mu_1 / (2^{v+1} v!)). \quad \square$$

A.3. Proof of Theorem 2.10. We finally justify our strengthening of the 1-statement of the asymmetric random Ramsey problem, showing that above the threshold, $G(n, p)$ is both robustly and globally (H_1, H_2) -Ramsey. We recall the precise statement below.

Theorem 2.10 ([20, 22]). *Let H_1 and H_2 be graphs such that $m_2(H_1) \geq m_2(H_2) \geq 1$, and*

- (a) $m_2(H_1) = m_2(H_2)$, or
- (b) H_1 is strictly balanced with respect to $m_2(\cdot, H_2)$.

The random graph $G(n, p)$ then has the following Ramsey properties:

- (i) *There are constants $\gamma = \gamma(H_1, H_2) > 0$ and $C_1 = C_1(H_1, H_2)$ such that if $p \geq C_1 n^{-1/m_2(H_1, H_2)}$ and, for $i \in [2]$, $\mathcal{F}_i \subseteq \binom{[n]}{v(H_i)}$ is a collection of at most $\gamma n^{v(H_i)}$ forbidden subsets, then $G(n, p)$ is with high probability robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$.*
- (ii) *For every $\mu > 0$ there is a constant $C_2 = C_2(H_1, H_2, \mu)$ such that if $p \geq C_2 n^{-1/m_2(H_1, H_2)}$, then $G(n, p)$ is with high probability μ -globally (H_1, H_2) -Ramsey.*
- (iii) *If we further have $m_2(H_2) > 1$, then there are constants $\beta_0 = \beta_0(H_1, H_2) > 0$ and $C_3 = C_3(H_1, H_2)$ such that if $0 \leq \beta \leq \beta_0$ and $p \geq C_3 n^{-(1-\beta)/m_2(H_1, H_2)}$, then with high probability $G(n, p)$ is $n^{-\beta}$ -globally (H_1, H_2) -Ramsey.*

A.3.1. Containers for nearly- H -free graphs and Ramsey supersaturation. As stated earlier, we shall prove this result by retracing the proofs of Hancock, Staden and Treglown [22] and of Gugelmann, Nenadov, Person, Škorić, Steger and Thomas [20], making minor modifications along the way. One commonality between the two proofs is the use of the hypergraph containers theorems of Balogh, Morris and Samotij [3] and of Saxton and Thomason [41].

Indeed, if $G(n, p)$ is not (H_1, H_2) -Ramsey, then its edges can be partitioned into a red H_1 -free subgraph and a blue H_2 -free subgraph. The aforementioned containers theorems show there are only a small number of “containers” that these H_i -free subgraphs must belong to, and one may use the properties of these containers to show it is very unlikely that $G(n, p)$ admits such a partition.

However, in the robust Ramsey setting, it is no longer true that the red subgraph must be H_1 -free or that the blue subgraph must be H_2 -free, as they may contain some forbidden copies of these subgraphs. Fortunately, the containers do not only capture H_i -free graphs, but also graphs with relatively few copies of H_i . To make this precise, we use the formulation of Saxton and Thomason [41], for which we require the following definition concerning r -uniform hypergraphs (or r -graphs).

Definition A.2 (Co-degree function). *Let Γ be a vertex-transitive d -regular r -graph, and let $\tau > 0$. Given a set S of at most r vertices, define $d(S)$ to be the number of edges of Γ containing S . For each $2 \leq j \leq r$, we define δ_j by the equation*

$$\delta_j \tau^{j-1} d = \max_{S \in \binom{V(\Gamma)}{j}} d(S).$$

The co-degree function $\delta(\Gamma, \tau)$ is then defined by

$$\delta(\Gamma, \tau) := 2^{\binom{r}{2}-1} \sum_{j=2}^r 2^{-\binom{j-1}{2}} \delta_j.$$

Note that the notion of co-degree can be defined more generally (i.e. not just for vertex-transitive r -graphs); we only state the definition in this form to simplify things a little. In the case of vertex-transitive graphs, Corollary 3.6 in [41] reads as below.

Corollary A.3 (Saxton–Thomason [41]). *Let Γ be a vertex-transitive r -graph on vertex set $[N]$. Let $0 < \varepsilon, \tau < 1/2$. Suppose that τ satisfies $\delta(\Gamma, \tau) \leq \varepsilon/12r!$. Then there exists a constant $c = c(r)$ and a function $\psi : \mathcal{P}([N])^q \rightarrow \mathcal{P}([N])$, where $q \leq c \log(1/\varepsilon)$, with the following properties. Let $\mathcal{T} := \{(T^{(1)}, \dots, T^{(q)}) \in \mathcal{P}([N])^q : |T^{(j)}| \leq c\tau N, j \in [q]\}$, and let $\mathcal{C} := \{\psi(T) : T \in \mathcal{T}\}$. Then*

- (a) *for every set $I \subset [N]$ for which $e(\Gamma[I]) \leq 24\varepsilon r! r\tau^r e(\Gamma)$, there exists $T = (T^{(1)}, \dots, T^{(q)}) \in \mathcal{T} \cap \mathcal{P}(I)^q$ with $I \subset \psi(T) \in \mathcal{C}$,*
- (b) *$e(\Gamma[\chi]) \leq \varepsilon e(\Gamma)$ for all $\chi \in \mathcal{C}$, and*
- (c) *$\log |\mathcal{C}| \leq c \log(1/\varepsilon) N \tau \log(1/\tau)$.*

Given a graph H , we wish to apply Corollary A.3 to build containers for n -vertex graphs that are nearly H -free. To that end, we set $N := \binom{n}{2}$ and $r := e(H)$, and take Γ to be the r -graph whose vertices correspond to the edges of K_n and whose hyperedges are the edge-sets of all copies of H in K_n . It is clear that Γ is vertex-transitive, and, writing $v := v(H)$ and $e := e(H)$, each vertex in Γ has degree

$$d := \binom{n}{v} \frac{v!}{|\text{Aut}(H)|} \frac{e}{\binom{n}{2}} = \Theta(n^{v-2}).$$

We next must bound $d(S)$ for sets $S \in \binom{V(\Gamma)}{j}$, where $2 \leq j \leq r$. Clearly, if the set S of edges spans a vertices of H , then $d(S) = \Theta(n^{v-a})$. Consequently,

$$\max_{S \in \binom{V(\Gamma)}{j}} d(S) = \Theta(n^{v-a_j}),$$

where a_j is the minimum number of vertices that a set of j edges of H can span. Hence

$$\delta_j := \max_{S \in \binom{V(\Gamma)}{j}} d(S) / (\tau^{j-1} d) = \Theta(\tau^{1-j} n^{2-a_j}),$$

and so $\delta(\Gamma, \tau) = \Theta(\max_j \tau^{1-j} n^{2-a_j})$. In order to apply Corollary A.3, we need the co-degree function to satisfy $\delta(\Gamma, \tau) \leq \varepsilon/12r!$, where ε will be a small constant, and therefore we cannot let τ be too small. Indeed, we need $\tau = \Omega(n^{-(a_j-2)/(j-1)})$ for all j , which is equivalent to saying that for every subgraph $F \subseteq H$, we need $\tau = \Omega(n^{-(v(F)-2)/(e(F)-1)})$. Recalling the definition of the 2-density $m_2(H)$, this implies that for every $0 < \varepsilon < 1/2$, we may apply Corollary A.3 with $\tau = O_\varepsilon(n^{-1/m_2(H)})$. Substituting this choice of τ into Corollary A.3 gives the following.

Corollary A.4. *Given a graph H and a constant $0 < \varepsilon < 1/2$, there are positive constants $q = q(\varepsilon, H)$, $\beta = \beta(\varepsilon, H)$ and $\kappa = \kappa(\varepsilon, H)$ such that for every $n \in \mathbb{N}$, there is a function $\psi : \mathcal{P}(E(K_n))^q \rightarrow \mathcal{P}(E(K_n))$ with the following properties. Let $\mathcal{T} := \{(T^{(1)}, \dots, T^{(q)}) \in \mathcal{P}(E(K_n))^q :$*

$|T^{(j)}| \leq \beta n^{2-1/m_2(H)}$, $j \in [q]$, and let $\mathcal{C} := \{\psi(T) : T \in \mathcal{T}\}$ be the family of corresponding subgraphs of K_n . Then

- (i) for every subgraph $F \subseteq K_n$ with at most $\kappa n^{v(H)-e(H)/m_2(H)}$ copies of H , there is some $T \in \mathcal{T} \cap \mathcal{P}(E(F))^q$ with $F \subseteq \psi(T) \in \mathcal{C}$, and
- (ii) each subgraph $\psi \in \mathcal{C}$ contains at most $\varepsilon \binom{n}{v(H)}$ copies of H .

Another result we shall need is a supersaturated version of Ramsey's Theorem — when colouring the edges of a large enough clique, we do not just see a monochromatic copy of a given graph, but we see many such copies. This follows from the classic Ramsey's Theorem by a standard averaging argument, the details of which can be found in [20].

Proposition A.5 (Folklore). *Given $r \geq 2$ and graphs $F_1, F_2, \dots, F_r$, there are positive constants $n_0 = n_0(F_1, \dots, F_r)$ and $\pi = \pi(F_1, \dots, F_r)$ such that if $n \geq n_0$, then every r -colouring of K_n has some $i \in [r]$ for which there are at least $\pi \binom{n}{v(F_i)}$ monochromatic copies of F_i in colour i .*

A.3.2. Random graphs are robustly Ramsey: Case (a). We now prove the robust Ramsey result under the assumption (a); that is, $m_2(H_1) = m_2(H_2) \geq 1$. Recall that we then have $m_2(H_1, H_2) = m_2(H_1)$ as well. We wish to show that there are constants $C, \gamma > 0$ such that if $p \geq Cn^{-1/m_2(H_1)}$ and, for $i \in [2]$, $\mathcal{F}_i$ is a collection of at most $\gamma n^{v(H_i)}$ forbidden copies of H_i , then $G(n, p)$ is with high probability robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$. By monotonicity, we may assume $p = Cn^{-1/m_2(H_1)}$ and, for each i , $|\mathcal{F}_i| = \gamma n^{v(H_i)}$.

Recall that the only copies of H_i that may be monochromatic in their respective colours are the forbidden ones from $\mathcal{F}_i$. We first claim that there are few forbidden copies of each H_i in $G(n, p)$.

Claim A.6. *If $p = Cn^{-1/m_2(H_i)}$ for some constant $C \geq 1$, then for each $i \in [2]$, with high probability $G(n, p)$ contains at most $2\gamma C^{e(H_i)} n^{v(H_i)-e(H_i)/m_2(H_i)}$ copies of H_i from $\mathcal{F}_i$.*

Proof. We begin with a preliminary calculation. Since $p = Cn^{-1/m_2(H_i)} \geq n^{-1}$, for any subgraph $F \subseteq H_i$ with at least one edge we have

$$(A.1) \quad n^{v(F)} p^{e(F)} = n^2 p \cdot n^{v(F)-2} p^{e(F)-1} \geq C^{e(F)-1} n^2 p = \Omega(n).$$

Now let X_i denote the number of forbidden copies of H_i from $\mathcal{F}_i$ that appear in $G(n, p)$. We clearly have $\mu(H_i) := \mathbb{E}[X_i] = |\mathcal{F}_i| p^{e(H_i)} = \gamma n^{v(H_i)} p^{e(H_i)} = \gamma C^{e(H_i)} n^{v(H_i)-e(H_i)/m_2(H_i)}$. Moreover, taking $F = H_i$ in (A.1), we have $\mu(H_i) = \Omega(n)$.

We next wish to estimate the variance of X_i . Since edge-disjoint copies of H_i appear independently of each other, we can bound

$$\text{Var}(X_i) \leq \sum_{\substack{F_1, F_2 \in \mathcal{F}_i: \\ e(F_1 \cap F_2) \geq 1}} \mathbb{P}(F_1 \cup F_2 \subseteq G(n, p)).$$

The diagonal terms, when $F_1 = F_2$, contribute a total of $\mu(H_i)$, which, since $\mu(H_i) = \Omega(n)$, is $O(\mu(H_i)^2/n)$. On the other hand, the sum over ordered pairs (F_1, F_2) with $F_1 \neq F_2$ can be bounded by Δ from Lemma A.1, applied with $H := H_i$ and $\mathcal{H} := \mathcal{F}_i$. The lemma gives $\Delta \leq 2^{v(H_i)} v(H_i)! |\mathcal{F}_i| n^{v(H_i)} p^{2e(H_i)} / \mu_0(H_i) = \Theta(\mu(H_i)^2 / \mu_0(H_i))$. By (A.1) we have $\mu_0(H_i) = \Omega(n)$, from which we can deduce $\Delta = O(\mu(H_i)^2/n)$.

Thus $\text{Var}(X_i) = O(\mu(H_i)^2/n)$. Chebyshev's inequality now gives the desired result, since

$$\mathbb{P}\left(X_i \geq 2\gamma C^{e(H_i)} n^{v(H_i)-e(H_i)/m_2(H_i)}\right) = \mathbb{P}(X_i \geq 2\mu(H_i)) \leq \frac{\text{Var}(X_i)}{\mu(H_i)^2} = O\left(\frac{1}{n}\right). \quad \square$$

Define $\mathcal{E}_0$ to be the event that $G(n, p)$ contains more than $2\gamma C^{e(H_i)} n^{v(H_i)-e(H_i)/m_2(H_i)}$ copies of H_i from $\mathcal{F}_i$, for some $i \in [2]$. Claim A.6 shows that $\mathcal{E}_0$ almost surely does not occur, and we shall henceforth assume it does not.

Now suppose $G(n, p)$ is not robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$. This means the edges of $G(n, p)$ may be 2-coloured so that the red copies of H_1 and the blue copies of H_2 all belong to $\mathcal{F}_1$ and $\mathcal{F}_2$ respectively. In light of Claim A.6, the red subgraph has few copies of H_1 , and hence by Corollary A.4 must lie within one of the containers for nearly- H_1 -free graphs. Similarly, the blue subgraph must lie within one of the containers for nearly- H_2 -free graphs. Since there are very few containers, and the containers are relatively small, this is very unlikely.

To be more precise, let $\pi = \pi(H_1, H_2, K_2)$ be the constant from Proposition A.5. For $i \in [2]$, we then apply Corollary A.4 with $\varepsilon := \pi/2$ and H_i to get constants q_i, β_i and κ_i , and a map ψ_i from a collection of sequences $\mathcal{T}_i := \{(T_i^{(1)}, \dots, T_i^{(q_i)}) \in \mathcal{P}(E(K_n))^{q_i} : |T_i^{(j)}| \leq \beta_i n^{2-1/m_2(H_i)}, j \in [q_i]\}$ to the set of containers $\mathcal{C}_i := \{\psi_i(T_i) : T_i \in \mathcal{T}_i\}$. Given a choice of the constant C , which we shall implicitly specify at the end, we set $\gamma := \min\{\kappa_i/(2C^{e(H_i)}) : i \in [2]\}$.

Now fix a 2-colouring of $G(n, p)$ where all the red copies of H_1 and blue copies of H_2 belong to $\mathcal{F}_1$ and $\mathcal{F}_2$ respectively. By Claim A.6, the red subgraph G_1 has at most $2\gamma C^{e(H_1)} n^{v(H_1)-e(H_1)/m_2(H_1)}$ copies of H_1 , which, by our choice of γ , is at most $\kappa_1 n^{v(H_1)-e(H_1)/m_2(H_1)}$. By Corollary A.4, there is some $T_1 \in \mathcal{T}_1 \cap \mathcal{P}(E(G_1))^{q_1}$ such that $G_1 \subseteq C_1 := \psi_1(T_1) \in \mathcal{C}_1$. A similar argument for the blue subgraph G_2 gives $T_2 \in \mathcal{T}_2 \cap \mathcal{P}(E(G_2))^{q_2}$ with $G_2 \subseteq C_2 := \psi_2(T_2) \in \mathcal{C}_2$. Since $G(n, p) = G_1 \cup G_2$, we have $G(n, p) \subseteq C_1 \cup C_2$.

Corollary A.4 also asserts that, for $i \in [2]$, C_i contains at most $\pi \binom{n}{v(H_i)}/2$ copies of H_i . By Proposition A.5, it follows that there are at least $\pi \binom{n}{2}$ edges in $R(T_1, T_2) := E(K_n) \setminus (C_1 \cup C_2)$, all of which must be missing from $G(n, p)$.

Given $T_1 \in \mathcal{T}_1$ and $T_2 \in \mathcal{T}_2$, we define the shorthand $T_1 \cup T_2 := \bigcup_{i \in [2]} \left\{ \bigcup_{j \in [q_i]} T_i^{(j)} \right\}$. Now let $\mathcal{E}(T_1, T_2)$ be the event that all edges in $T_1 \cup T_2$ are in $G(n, p)$ while all edges in $R(T_1, T_2)$ are missing. Since $|R(T_1, T_2)| \geq \pi \binom{n}{2}$, we have

$$\mathbb{P}(\mathcal{E}(T_1, T_2)) = p^{|T_1 \cup T_2|} (1-p)^{|R(T_1, T_2)|} \leq p^{|T_1 \cup T_2|} \exp\left(-\pi \binom{n}{2} p\right).$$

Our above discussion shows that if $G(n, p)$ does not have the desired robust Ramsey property, then either $\mathcal{E}_0$ or one of the events $\mathcal{E}(T_1, T_2)$ must occur. By the union bound, the probability of $G(n, p)$ not being robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$ is at most

$$\begin{aligned} \sum_{T_1 \in \mathcal{T}_1, T_2 \in \mathcal{T}_2} \mathcal{E}(T_1, T_2) + \mathbb{P}(\mathcal{E}_0) &\leq \exp\left(-\pi \binom{n}{2} p\right) \sum_{T_1 \in \mathcal{T}_1, T_2 \in \mathcal{T}_2} p^{|T_1 \cup T_2|} + o(1) \\ &= \exp\left(-\pi \binom{n}{2} p\right) \sum_{S \subseteq E(K_n)} \sum_{\substack{T_1 \in \mathcal{T}_1, T_2 \in \mathcal{T}_2: \\ T_1 \cup T_2 = S}} p^{|S|} + o(1). \end{aligned}$$

Now recall that $T_i = (T_i^{(1)}, \dots, T_i^{(q_i)})$, where $|T_i^{(j)}| \leq \beta_i n^{2-1/m_2(H_i)}$, and so $|T_1 \cup T_2| \leq M_n := (\beta_1 q_1 + \beta_2 q_2) n^{2-1/m_2(H_1)}$. Moreover, given a set S of at most this size, for each edge $e \in S$, we must choose which of the sets $T_i^{(j)}$ it belongs to, and hence there are at most $2^{(q_1+q_2)|S|}$ pairs (T_1, T_2) corresponding to S . Therefore

$$\sum_{S \subseteq E(K_n)} \sum_{\substack{T_1 \in \mathcal{T}_1, T_2 \in \mathcal{T}_2: \\ T_1 \cup T_2 = S}} p^{|S|} \leq \sum_{\substack{S \subseteq E(K_n): \\ |S| \leq M_n}} (2^{q_1+q_2} p)^{|S|} \leq \sum_{s=0}^{M_n} \binom{n^2}{s} (2^{q_1+q_2} p)^s \leq \sum_{s=0}^{M_n} \left(\frac{2^{q_1+q_2} e n^2 p}{s} \right)^s.$$

Substituting $p = C n^{-1/m_2(H_1)}$, we find that the summand is $(2^{q_1+q_2} C e n^{2-1/m_2(H_1)}/s)^s$. Since the function $f(x) = (A/x)^x$ increases to a maximum at $x = A/e$, it follows that, if C is sufficiently

large, the largest summand corresponds to $s = M_n$. We thus have

$$\sum_{T_1 \in \mathcal{T}_1, T_2 \in \mathcal{T}_2} p^{|T_1 \cup T_2|} \leq (M_n + 1) \left(\frac{2^{q_1+q_2} C e n^{2-1/m_2(H_1)}}{M_n} \right)^{M_n} = (M_n + 1) \exp(L_C n^{2-1/m_2(H_1)}),$$

where $L_C := (\beta_1 q_1 + \beta_2 q_2) \ln(2^{q_1+q_2} C e / (\beta_1 q_1 + \beta_2 q_2))$. This gives

$$\begin{aligned} \mathbb{P}(G(n, p) \text{ is not robustly Ramsey}) &\leq (M_n + 1) \exp \left(L_C n^{2-1/m_2(H_1)} - \pi \binom{n}{2} p \right) + o(1) \\ &\leq (M_n + 1) \exp \left((L_C - C\pi/4) n^{2-1/m_2(H_1)} \right) + o(1) \\ (A.2) \quad &\leq \exp \left(-C\pi n^{2-1/m_2(H_1)}/8 \right) + o(1), \end{aligned}$$

where the final inequality holds when we choose C to be sufficiently large, since L_C is logarithmic in C , and if n is large enough, since M_n is only polynomial in n . This shows $G(n, p)$ is with high probability robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$. $\square$

A.3.3. Random graphs are robustly Ramsey: Case (b). We next prove the robust Ramsey result under the assumption (b), when H_1 is strictly balanced with respect to $m_2(\cdot, H_2)$. When $m_2(H_1) > m_2(H_2)$ and $p = C n^{-1/m_2(H_1, H_2)}$, there are too many nearly- H_1 -free containers for the union bound calculation of Case (a) to work.

We will instead exploit the fact that nearly all copies of H_1 in $G(n, p)$ are edge-disjoint. To that end, call a copy of H_1 in $G(n, p)$ *isolated* if it is edge-disjoint from all other copies of H_1 , and call it *non-isolated* otherwise. Our first claim shows that there are few non-isolated copies of H_1 .

Claim A.7. *Let H_1 be strictly balanced with respect to $m_2(\cdot, H_2)$, and let $p = C n^{-1/m_2(H_1, H_2)}$ for some constant $C \geq 1$. There is a constant $\delta = \delta(H_1, H_2)$ such that the number of non-isolated copies of H_1 in $G(n, p)$ is with high probability at most $n^{2-1/m_2(H_2)-\delta}$.*

Proof. If a copy H' of H_1 is non-isolated, then there is some other copy $H'' \neq H'$ of H_1 such that $e(H' \cap H'') \geq 1$. The number of non-isolated copies of H_1 in $G(n, p)$ is thus bounded from above by the expected number of such pairs $\{H', H''\}$, which is precisely the quantity Δ from Lemma A.1, when $H := H_1$ and $\mathcal{H} := \mathcal{K}_n(H_1)$. The lemma gives

$$\Delta \leq \frac{2^{v(H_1)} v(H_1)! |\mathcal{K}_n(H_1)| n^{v(H_1)} p^{2e(H_1)}}{\mu_0(H_1)} \leq \frac{2^{v(H_1)} v(H_1)! (n^{v(H_1)} p^{e(H_1)})^2}{\mu_0(H_1)},$$

where $\mu_0(H_1) = \min\{n^{v(F)} p^{e(F)} : F \subsetneq H_1, e(F) \geq 1\}$.

For each $F \subseteq H_1$ with $e(F) \geq 1$, define δ_F such that

$$m_2(H_1, H_2) = \frac{e(H_1)}{v(H_1) - 2 + 1/m_2(H_2)} =: \frac{e(F)}{v(F) - 2 + 1/m_2(H_2) - 2\delta_F}.$$

It then follows that for every $F \subseteq H_1$ we have

$$(A.3) \quad n^{v(F)} p^{e(F)} = C^{e(F)} n^{v(F)-e(F)(v(H_1)-2+1/m_2(H_2))/e(H_1)} = C^{e(F)} n^{2-1/m_2(H_2)+2\delta_F}.$$

We clearly have $\delta_{H_1} = 0$. On the other hand, since H_1 is strictly balanced with respect to $m_2(\cdot, H_2)$, we have $\delta_F > 0$ in all other cases. Letting $\delta = \delta(H_1, H_2) := \min\{\delta_F : F \subsetneq H_1, e(F) \geq 1\}$, it follows that $\mu_0(H_1) \geq n^{2-1/m_2(H_2)+2\delta}$.

By using these values for δ_{H_1} and $\mu_0(H_1)$, we can bound the expected number of pairs of intersecting copies of H_1 by

$$\Delta = O \left(\frac{(n^{v(H_1)} p^{e(H_1)})^2}{\mu_0(H_1)} \right) = O \left(\frac{(n^{2-1/m_2(H_2)})^2}{n^{2-1/m_2(H_2)+2\delta}} \right) = O \left(n^{2-1/m_2(H_2)-2\delta} \right).$$

Using Markov's inequality, the probability that the number of pairs of intersecting copies of H_1 , and thus the number of non-isolated copies of H_1 , in $G(n, p)$ is more than $n^{2-1/m_2(H_2)-\delta}$ is $O(n^{-\delta})$, establishing the claim. $\square$

Let $\mathcal{E}_1$ be the event that $G(n, p)$ has more than $n^{2-1/m_2(H_2)-\delta}$ non-isolated copies of H_1 , which Claim A.7 shows we may assume does not occur. We denote by L the set of edges in non-isolated copies of H_1 . Note that any edge $e \in E(G(n, p)) \setminus L$ is contained in at most one copy of H_1 .

Now suppose that $G(n, p)$ is not robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$, and fix a 2-colouring of its edges where the only red copies of H_1 and blue copies of H_2 in $G(n, p)$ are forbidden (i.e. in $\mathcal{F}_1$ or $\mathcal{F}_2$ respectively). We may safely assume that any edge of $G(n, p)$ that is not contained in a copy of H_1 is red, since if not, recolouring it red does not create a red copy of H_1 . Furthermore, we may assume that every isolated non-forbidden copy of H_1 has exactly one blue edge — since it is not in $\mathcal{F}_1$, it cannot be monochromatic red, and if it has more than one blue edge, then since it is an isolated copy of H_1 , recolouring all but one of them red cannot create a red copy of H_1 .

Let B be the subgraph of $G(n, p)$ consisting of all edges not in L that are coloured blue. Note that each edge of B must be contained in a copy of H_1 , and moreover these copies are pairwise edge-disjoint. We thus call a set E of edges in $G(n, p)$ *viable* if each $e \in E$ is contained in some copy W_e of H_1 such that the copies $\{W_e : e \in E\}$ are pairwise edge-disjoint. We further call such a set of copies a *witness* for the viability of E .

Note that a set of edges E can only be contained in B if E is viable. The next claim bounds the likelihood of viability.

Claim A.8. *If $p = Cn^{-1/m_2(H_1, H_2)}$ for some positive constant C , then a set E of s edges is viable with probability at most ρ^s , where $\rho := C^{e(H_1)}v(H_1)^2n^{-1/m_2(H_2)}$.*

Proof. We take a union bound over all possible witnesses for E . Since the copies W_e of H_1 in a witness are edge-disjoint, each witness requires exactly $e(H_1)s$ edges to appear in $G(n, p)$, and hence appears with probability $p^{e(H_1)s}$.

To estimate the number of possible witnesses, note that for each edge $e \in E$, we have fewer than $v(H_1)^2$ ways to assign vertices of H_1 to the vertices of e in the copy W_e . There are then at most a further $n^{v(H_1)-2}$ choices for the remaining vertices in W_e . This shows there are at most $(v(H_1)^2n^{v(H_1)-2})^s$ possible witnesses for E .

The expected number of witnesses for E , which bounds from above the probability of E being viable, is therefore at most

$$\left(v(H_1)^2n^{v(H_1)-2}\right)^s p^{e(H_1)s} = \left(v(H_1)^2n^{v(H_1)-2}p^{e(H_1)}\right)^s = \rho^s,$$

where in the last equality we use the fact that $p^{e(H_1)} = C^{e(H_1)}n^{2-v(H_1)-1/m_2(H_2)}$. $\square$

Only forbidden copies of H_2 can be monochromatically blue. Furthermore, a copy of H_2 in B must also be viable. Claim A.9 asserts there cannot be many viable forbidden copies of H_2 . We will defer its proof for the moment, first seeing how it implies the robust Ramsey result.

Claim A.9. *The number of viable forbidden copies of H_2 from $\mathcal{F}_2$ is with high probability at most $\gamma K n^{v(H_2)-e(H_2)/m_2(H_2)}$ for some constant $K = K(C, H_1, H_2)$.*

We now lay some groundwork for the rest of the proof, including specifying our choice of the constant γ . Let $\pi = \pi(H_1, H_2, K_2)$ be the constant from Proposition A.5. We now apply Corollary A.4 with $H := H_2$ and $\varepsilon := \pi/2$. This gives us constants q, β and κ , together with a collection $\mathcal{T} := \{(T^{(1)}, \dots, T^{(q)}) \in \mathcal{P}(E(K_n))^q : |T^{(j)}| \leq \beta n^{2-1/m_2(H_2)}, j \in [q]\}$ and a set of containers $\mathcal{C} := \{\psi(T) : T \in \mathcal{T}\}$ for nearly- H_2 -free graphs. Finally, for whatever sufficiently large constant we

will later choose C to be, we set $\gamma := \min\{\kappa/K, \pi/(2v(H_1)^{v(H_1)})\}$, where $K = K(C, H_1, H_2)$ is the constant from Claim A.9.

Now let $\mathcal{E}_0$ be the event that there are more than $\kappa n^{v(H_2)-e(H_2)/m_2(H_2)}$ viable forbidden copies of H_2 , which, by Claim A.9 and the fact that $\kappa \geq \gamma K$, we may assume does not occur. Since all copies of H_2 in B must be viable and forbidden, it follows that B has at most $\kappa n^{v(H_2)-e(H_2)/m_2(H_2)}$ copies of H_2 . By Corollary A.4, there is then some $T \in \mathcal{T} \cap \mathcal{P}(E(B))^q$ with $B \subseteq \psi(T)$. Let $\mathcal{E}_2(T)$ be the event that the set of edges $E(T) = \cup_{j=1}^q T^{(j)}$ is viable, which must hold for them to be in B . By Claim A.8, we have

$$(A.4) \quad \mathbb{P}(\mathcal{E}_2(T)) \leq \rho^{|E(T)|}.$$

Next define $R = R(T, L) := E(K_n) \setminus (\psi(T) \cup L)$, and consider the 3-colouring of $E(K_n)$ where all edges in R receive colour 1, edges in $\psi(T)$ have colour 2 and those in $L \setminus \psi(T)$ are coloured 3. Corollary A.4 guarantees that there are at most $\pi \binom{n}{v(H_2)}/2$ copies of H_2 coloured 2, while Claim A.7 guarantees that there are fewer than $\pi \binom{n}{2}$ copies of K_2 in colour 3. By Proposition A.5, we must therefore have at least $\pi \binom{n}{v(H_1)}$ copies of H_1 in R . Let $\mathcal{H}$ be those copies of H_1 in R that are not forbidden, and note that since $|\mathcal{F}_1| \leq \gamma n^{v(H_1)}$, we have $|\mathcal{H}| \geq \pi \binom{n}{v(H_1)}/2 \geq \xi n^{v(H_1)}$ for $\xi := \pi/(2v(H_1)^{v(H_1)})$.

Let $\mathcal{E}_3(T, L)$ be the event that none of the copies of H_1 in $\mathcal{H}$ appear in $G(n, p)$, and note that this must hold. Otherwise, the copy of $H_1 \in \mathcal{H}$, which is not forbidden, must be monochromatic red, since all blue edges of $G(n, p)$ lie either in L or in $B \subseteq \psi(T)$, contradicting our assumption that the only red copies of H_1 are forbidden.

By Theorem 2.7, we have $\mathbb{P}(\mathcal{E}_3(T, L)) \leq \exp(-\xi \mu_1(H_1)/(2^{v(H_1)+1} v(H_1)!))$. As we saw in (A.3), since H_1 is strictly balanced with respect to $m_2(\cdot, H_2)$, it follows that $\mu_1(H_1) = n^{v(H_1)} p^{e(H_1)} = C^{e(H_1)} n^{2-1/m_2(H_2)}$. Thus, setting $\xi' := \xi/(2^{v(H_1)+1} v(H_1)!)$, we have

$$(A.5) \quad \mathbb{P}(\mathcal{E}_3(T, L)) \leq \exp(-\xi C^{e(H_1)} n^{2-1/m_2(H_2)} / (2^{v(H_1)+1} v(H_1)!)) = \exp(-\xi' C^{e(H_1)} n^{2-1/m_2(H_2)}).$$

To summarise, in order for $G(n, p)$ to have a 2-colouring where all the red copies of H_1 and blue copies of H_2 are forbidden, we either need $\mathcal{E}_0$ or $\mathcal{E}_1$ to hold, or, if they do not, for $\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)$ to hold for some choice of $T \in \mathcal{T}$ and some specified set L of edges in non-isolated copies of H_1 . We can thus estimate the probability that $G(n, p)$ is not robustly Ramsey by taking a union bound over these results:

$$(A.6) \quad \mathbb{P}(G(n, p) \text{ is not robustly } (H_1, H_2)\text{-Ramsey}) \leq \mathbb{P}(\mathcal{E}_0) + \mathbb{P}(\mathcal{E}_1) + \sum_{T, L} \mathbb{P}(\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)).$$

Since $\mathcal{E}_2(T)$ — that all of the edges from $E(T)$ appear in B — is a monotone increasing graph property, while $\mathcal{E}_3(T, L)$ — that none of the copies of H_1 from $R(T, L)$ appear in $G(n, p)$ — is a monotone decreasing graph property, it follows from the FKG Inequality (see [2, Theorem 6.3.3]) that $\mathbb{P}(\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)) \leq \mathbb{P}(\mathcal{E}_2(T)) \mathbb{P}(\mathcal{E}_3(T, L))$. Using the upper bounds from (A.4) and (A.5), we can bound the sum by

$$\sum_{T, L} \mathbb{P}(\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)) \leq \exp\left(-\xi' C^{e(H_1)} n^{2-1/m_2(H_2)}\right) \sum_L \sum_T \rho^{|E(T)|}.$$

There are at most $n^{2-1/m_2(H_2)-\delta}$ non-isolated copies of H_1 (as otherwise we are covered by $\mathcal{E}_1$), and hence the sum L only runs over sets of at most $e(H_1) n^{2-1/m_2(H_2)-\delta}$ edges. Thus

$$\sum_{T, L} \mathbb{P}(\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)) \leq \exp\left(-\xi' C^{e(H_1)} n^{2-1/m_2(H_2)}\right) \left(\sum_{\ell=0}^{e(H_1) n^{2-1/m_2(H_2)-\delta}} \binom{\binom{n}{2}}{\ell} \right) \sum_T \rho^{|E(T)|}$$

$$(A.7) \quad \leq \exp \left(-\xi' C^{e(H_1)} n^{2-1/m_2(H_2)} + o \left(n^{2-1/m_2(H_2)} \right) \right) \sum_T \rho^{|E(T)|}.$$

We now bound the sum over T as we did in Case (a). From Corollary A.4, we know $s = |E(T)| \leq M_n := q\beta n^{2-1/m_2(H_2)}$. We group the q -tuples T by the size of $E(T)$, observing that each set of edges $E(T)$ corresponds to at most $2^{q|E(T)|}$ q -tuples in $\mathcal{T}$. Recalling the definition of ρ from Claim A.8, we have

$$\sum_T \rho^{|E(T)|} \leq \sum_{s=0}^{M_n} \sum_{\substack{T \in \mathcal{T}: \\ |E(T)|=s}} \rho^s \leq \sum_{s=0}^{M_n} \binom{n^2}{s} (2^q \rho)^s \leq \sum_{s=0}^{M_n} \left(\frac{v(H_1)^2 e 2^q C^{e(H_1)} n^{2-1/m_2(H_2)}}{s} \right)^s.$$

As before, this summand is maximised when $s = M_n$, yielding, for $\lambda := v(H_1)^2 e 2^q / (q\beta)$,

$$\sum_T \rho^{|E(T)|} \leq \left(q\beta n^{2-1/m_2(H_2)} + 1 \right) \left(\lambda C^{e(H_1)} \right)^{q\beta n^{2-1/m_2(H_2)}}.$$

Making the substitution in (A.7), we find

$$(A.8) \quad \sum_{T,L} \mathbb{P}(\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)) \leq \exp \left(\left(-\xi' C^{e(H_1)} + q\beta \ln \left(\lambda C^{e(H_1)} \right) + o(1) \right) n^{2-1/m_2(H_2)} \right).$$

Thus, if we choose C to be a sufficiently large constant (with respect to ξ', q, β and λ), this sum is $o(1)$. By Claims A.9 and A.7 respectively, so are $\mathbb{P}(\mathcal{E}_0)$ and $\mathbb{P}(\mathcal{E}_1)$. Hence, by (A.6), it follows that $G(n, p)$ is with high probability robustly (H_1, H_2) -Ramsey with respect to $(\mathcal{F}_1, \mathcal{F}_2)$. $\square$

To complete the proof, we prove Claim A.9.

Proof of Claim A.9. In order for a forbidden copy $H \in \mathcal{F}_2$ of H_2 to be viable, it must have a witness — that is, a collection of pairwise edge-disjoint copies W_e of H_1 containing each edge $e \in E(H)$. We shall bound the number of witnesses for forbidden copies of H_2 , thereby obtaining an upper bound on the number of viable forbidden copies of H_2 .

Note that there are various non-isomorphic types of witnesses, depending on how the copies W_e of H_1 are attached to the edges of the forbidden copy of H_2 , and whether or not they share vertices. Let $\Gamma_1, \dots, \Gamma_m$ represent the different isomorphism classes, where $m := m(H_1, H_2)$ is some constant. Although these witnesses can have different numbers of vertices, up to a maximum of $v(H_2) + e(H_2)(v(H_1) - 2)$, the edge-disjointness of the W_e ensures they each have exactly $e(H_1)e(H_2)$ edges.

We will show that for $p = Cn^{-1/m_2(H_1, H_2)}$, if we set $K' := 2\gamma C^{e(H_1)e(H_2)}$, then for each $i \in [m]$ the probability that there are at most $\gamma K' n^{v(H_2) - e(H_2)/m_2(H_2)}$ copies of Γ_i with the central copy of H_2 coming from $\mathcal{F}_2$ is $O(1/n)$. Then, taking $K := mK'$, a union bound over the different types of witnesses Γ_i shows that the probability of there being more than $\gamma K n^{v(H_2) - e(H_2)/m_2(H_2)}$ viable forbidden copies of H_2 is also $O(1/n)$.

Let us fix some witness Γ_i , where $i \in [m]$, and let X_i denote the number of copies of Γ_i in $G(n, p)$ with the central copy of H_2 coming from $\mathcal{F}_2$. If we first select this central copy, and then the remaining vertices of Γ_i , we see there are at most $|\mathcal{F}_2| n^{v(\Gamma_i) - v(H_2)} = \gamma n^{v(\Gamma_i)}$ possible witnesses of this form, each appearing with probability $p^{e(H_1)e(H_2)}$. Thus $\mathbb{E}[X_i] \leq \gamma n^{v(\Gamma_i)} p^{e(H_1)e(H_2)}$.

We now differentiate between two cases.

Case I: $v(\Gamma_i) \leq v(H_2) + e(H_2)(v(H_1) - 2) - 1$. In this case we have

$$\mathbb{E}[X_i] \leq \gamma n^{v(H_2) + e(H_2)(v(H_1) - 2) - 1} p^{e(H_1)e(H_2)} = \gamma n^{v(H_2) - 1} \left(n^{v(H_1) - 2} p^{e(H_1)} \right)^{e(H_2)}.$$

Since $n^{v(H_1) - 2} p^{e(H_1)} = C^{e(H_1)} n^{-1/m_2(H_2)}$, we have $\mathbb{E}[X_i] \leq \gamma C^{e(H_1)e(H_2)} n^{v(H_2) - e(H_2)/m_2(H_2) - 1}$. By Markov's inequality, $\mathbb{P}(X_i > \gamma K' n^{v(H_2) - e(H_2)/m_2(H_2)}) = O(1/n)$, as desired.

Case II: $v(\Gamma_i) = v(H_2) + e(H_2)(v(H_1) - 2)$. In this case the copies $\{W_e : e \in E(H_2)\}$ of H_1 in Γ_i are all vertex-disjoint except for where a pair intersect in precisely one vertex in the central copy of H_2 .

The calculation above shows $\mathbb{E}[X_i] \leq \gamma C^{e(H_1)e(H_2)} n^{v(H_2)-e(H_2)/m_2(H_2)}$. We can again use Lemma A.1 to bound $\text{Var}(X_i)$, applying the lemma with $H := \Gamma_i$ and $\mathcal{H}$ the collection of all possible witnesses of this type for forbidden copies of H_2 . As in the proof of Claim A.6, this gives

$$\begin{aligned} \text{Var}(X_i) &\leq \Delta + \mathbb{E}[X_i] \leq \frac{\gamma 2^{v(\Gamma_i)} v(\Gamma_i)! (n^{v(\Gamma_i)} p^{e(\Gamma_i)})^2}{\mu_0(\Gamma_i)} + \gamma n^{v(\Gamma_i)} p^{e(\Gamma_i)} \\ &= O\left(\frac{(n^{v(\Gamma_i)} p^{e(\Gamma_i)})^2}{\mu_1(\Gamma_i)}\right) = O\left(\frac{(n^{v(H_2)-e(H_2)/m_2(H_2)})^2}{\mu_1(\Gamma_i)}\right), \end{aligned}$$

where $\mu_1(\Gamma_i)$ is as defined in Theorem 2.7. The penultimate equality above then follows from the fact that $\mu_1(\Gamma_i) = \min\{\mu_0(\Gamma_i), n^{v(\Gamma_i)} p^{e(\Gamma_i)}\}$. We shall shortly show that $\mu_1(\Gamma_i) = \Omega(n)$. Chebyshev's inequality then gives the desired bound:

$$\begin{aligned} \mathbb{P}\left(X_i > K' n^{v(H_2)-e(H_2)/m_2(H_2)}\right) &\leq \mathbb{P}\left(X_i - \mathbb{E}[X_i] > \gamma C^{e(H_1)e(H_2)} n^{v(H_2)-e(H_2)/m_2(H_2)}\right) \\ &\leq \frac{\text{Var}(X_i)}{(\gamma C^{e(H_1)e(H_2)} n^{v(H_2)-e(H_2)/m_2(H_2)})^2} = O\left(\frac{1}{n}\right). \end{aligned}$$

To finish, we establish a lower bound for $\mu_1(\Gamma_i) = \min\{n^{v(F)} p^{e(F)} : F \subseteq \Gamma_i, e(F) \geq 1\}$, and let F_0 be the subgraph of Γ_i that minimises this expression. Recall that for each edge h in the central forbidden copy $H \in \mathcal{F}_2$ of H_2 , we have a copy W_h of H_1 containing h . Let $F_0(h)$ be the subgraph of H_1 induced by F_0 on W_h , and let $w_h := e(F_0(h))$ be the number of edges it contains. We then have $e(F_0) = \sum_{h \in E(H_2)} w_h$.

When we count the vertices of F_0 , we observe that the vertices from the central H_2 will belong to several of the $F_0(h)$. Thus, for each $h \in E(H_2)$, we let $v_h := v(F_0(h))$, and let $r_h \in \{0, 1, 2\}$ denote the number of the vertices from h that are in F_0 . Let F_0^* denote the subgraph of the central H_2 contained within F_0 ; in particular, we have $r_h = 2$ if and only if $h \in E(F_0^*)$. Moreover, $v(F_0) = v(F_0^*) + \sum_{h \in E(H_2)} (v_h - r_h)$.

Thus we have

$$\mu_1(\Gamma_i) = n^{v(F_0)} p^{e(F_0)} = n^{v(F_0^*)} \prod_{h \in E(H_2)} n^{v_h - r_h} p^{w_h} \geq C^{e(F_0)} n^{v(F_0^*)} \prod_{\substack{h \in E(H_2): \\ w_h \geq 1}} n^{v_h - r_h - w_h/m_2(H_1, H_2)}.$$

For each h in the final product, since $F_0(h) \subseteq H_1$, the definition of $m_2(H_1, H_2)$ implies $w_h \leq (v_h - 2 + 1/m_2(H_2))m_2(H_1, H_2)$. Therefore

$$\mu_1(\Gamma_i) \geq C^{e(F_0)} n^{v(F_0^*)} \prod_{\substack{h \in E(H_2): \\ w_h \geq 1}} n^{2 - r_h - 1/m_2(H_2)} \geq C^{e(F_0)} n^{v(F_0^*) + |\{h \in E(H_2) : w_h \geq 1, r_h = 0\}| - e(F_0^*)/m_2(H_2)},$$

where the last inequality is due to the fact that, since $m_2(H_2) \geq 1$, the exponent $2 - r_h - 1/m_2(H_2)$ is at least 1 if $r_h = 0$, is non-negative when $r_h = 1$, and is $-1/m_2(H_2)$ when $r_h = 2$ or, equivalently, $h \in E(F_0^*)$.

If $e(F_0^*) \geq 1$ (and therefore $v(F_0^*) \geq 2$), then by definition of $m_2(H_2)$ we have $e(F_0^*) \leq (v(F_0^*) - 2)m_2(H_2) + 1$. Substituting this into the exponent, we find $n^{v(F_0^*) - e(F_0^*)/m_2(H_2)} \geq n^{2-1/m_2(H_2)} \geq n$, and hence $\mu_1(\Gamma_i) = \Omega(n)$.

Therefore we may assume $e(F_0^*) = 0$, in which case $\mu_1(\Gamma_i) = \Omega(n^{v(F_0^*) + |\{h \in E(H_2) : w_h \geq 1, r_h = 0\}|})$. This exponent is always at least 1, since either $v(F_0^*) \geq 1$, or $r_h = 0$ for all $h \in E(H_2)$ and,

since $e(F_0) \geq 1$, we must have $w_h \geq 1$ for some h as well. Thus we indeed have $\mu_1(\Gamma_i) = \Omega(n)$, completing the proof of this claim. $\square$

A.3.4. Random graphs are globally Ramsey. Having proven part (i) of Theorem 2.10, we finally deduce the global Ramsey properties of $G(n, p)$ from the above proofs, thereby establishing the remaining parts (ii) and (iii). Here we wish to show that, for appropriate probabilities, $G(n, p)$ is μ -globally (H_1, H_2) -Ramsey, for $\mu > 0$ constant and $\mu = n^{-\beta}$ respectively.

We begin with the assumption (a), that $m_2(H_1) = m_2(H_2)$. For (ii), let $\mu > 0$ be a fixed constant, and set $C_2 := C_1 \mu^{-1/m_2(H_1, H_2)}$, where C_1 is the constant from part (i). To show that $G(n, p)$ is μ -globally (H_1, H_2) -Ramsey when $p \geq C_2 n^{-1/m_2(H_1, H_2)}$, it will suffice to apply the union bound over all subsets $U \subseteq [n]$ of $n_0 := \mu n$ vertices.

Indeed, for any such set U , we have $G(n, p)[U] \sim G(n_0, p)$, where $p = C_2 n^{-1/m_2(H_1, H_2)} = C_1 n_0^{-1/m_2(H_1, H_2)}$. By (A.2), the probability that $G(n, p)[U]$ is not (H_1, H_2) -Ramsey is at most $\exp\left(-C_1 \pi n_0^{2-1/m_2(H_1)} / 8\right) + o(1)$, where $\pi = \pi(H_1, H_2, K_2)$ from Proposition A.5.

However, the $o(1)$ error term came from the event $\mathcal{E}_0$, where many forbidden copies of H_i appeared in $G(n_0, p)$. Since we do not have forbidden copies in our current setting, we do not incur this error. Thus, recalling that $m_2(H_1) \geq 1$, and assuming, as we may freely do, $C_1 \geq 8\pi\mu^{-1}$, we see that the probability that $G(n, p)[U]$ is not (H_1, H_2) -Ramsey is at most $\exp(-n)$. We may therefore take a union bound over all such sets U , of which there are fewer than 2^n , which shows that $G(n, p)$ is μ -globally (H_1, H_2) -Ramsey with high probability.

For part (iii), we instead set $n_0 := n^{1-\beta}$, and set $C_3 := C_1$. Letting $U \subseteq [n]$ be a set of n_0 vertices, we have $G(n, p)[U] \sim G(n_0, p)$, where $p \geq C_1 n_0^{-1/m_2(H_1, H_2)}$. As above, the probability that $G(n, p)[U]$ is not (H_1, H_2) -Ramsey is at most $\exp(-C_1 \pi n_0^{2-1/m_2(H_1)} / 8)$.

We can now take a union bound over all such sets U , of which there are $\binom{n}{n_0} \leq \exp(n_0 \ln(ne/n_0)) = \exp(n_0^{1+o(1)})$. Hence the probability that there is a non-Ramsey subgraph induced on some U is at most $\exp(n_0^{1+o(1)} - C_1 \pi n_0^{2-1/m_2(H_1)} / 8)$. Since $m_2(H_1) > 1$, it follows that $G(n, p)$ is $n^{-\beta}$ -globally (H_1, H_2) -Ramsey with high probability.

The argument under the assumption (b), that H_1 is strictly balanced with respect to $m_2(\cdot, H_2)$, is very similar. We will again run a union bound over all sets U of size $n_0 := \mu n$ (for constant $\mu > 0$ and $\mu = n^{-\beta}$), using the fact that $G(n, p)[U] \sim G(n_0, p)$ and choosing the constants C_2 and C_3 such that $p \geq C_1 n_0^{-1/m_2(H_1, H_2)}$.

In (A.6) we bounded from above the probability of $G(n, p)$ not being robustly (H_1, H_2) -Ramsey. Since we do not have collections of forbidden copies of H_1 and H_2 , we can again omit the error term from the event $\mathcal{E}_0$ of there being too many viable forbidden copies of H_2 . We then have

$$\mathbb{P}(G(n, p)[U] \text{ is not } (H_1, H_2)\text{-Ramsey}) \leq \mathbb{P}(\mathcal{E}_1) + \sum_{T, L} \mathbb{P}(\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)).$$

Recall that $\mathcal{E}_1$ is the event that we have more than $n^{2-1/m_2(H_2)-\delta}$ non-isolated copies of H_1 . In Claim A.7 we only obtained polynomially small bounds on its probability, and so we cannot afford to take a union bound over this event holding for each induced subgraph $G(n, p)[U]$. Instead, we replace the event by $\mathcal{E}'_1$, the event that $G(n, p)$ has more than $n_0^{2-1/m_2(H_2)-\delta}$ non-isolated copies of H_1 . Since a non-isolated copy of H_1 in $G(n, p)[U]$ must be non-isolated in $G(n, p)$ as well, this modified event would suffice for our purposes, and we can avoid the union bound for $\mathcal{E}_1$.

To see that $\mathcal{E}'_1$ holds with vanishingly small probability, observe that in Claim A.7, the expected number of pairs of intersecting copies of H_1 , which we bounded by $O(n^{2-1/m_2(H_1, H_2)-2\delta})$, was polynomial in n and p . Hence when we increase p to either $C_2 n^{-1/m_2(H_1, H_2)}$ or $C_3 n^{-(1-\beta)/m_2(H_1, H_2)}$,

provided $\beta < \beta_0(H_1, H_2)$, this expected number will grow to at most $O(n^{2-1/m_2(H_1, H_2)-5\delta/3})$. Similarly, since $n_0 \geq n^{1-\beta_0}$, we have $n_0^{2-1/m_2(H_2)-\delta} = \Omega(n^{2-1/m_2(H_2)-4\delta/3})$. Thus we can still apply Markov's inequality to deduce $\mathbb{P}(\mathcal{E}'_1) = o(1)$.

This leaves us with the sum $\sum_{T,L} \mathbb{P}(\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L))$, which we saw in (A.8) can be bounded by $\exp(-\Omega(n^{2-1/m_2(H_2)}))$. As under the assumption (a), we can afford to take a union bound over all $\binom{n}{n_0}$ choices for the set U , and still have the probability that there is some induced subgraph $G(n, p)[U]$ where $\mathcal{E}_2(T) \wedge \mathcal{E}_3(T, L)$ holds be $o(1)$. Hence it follows that $G(n, p)$ is with high probability μ -globally (H_1, H_2) -Ramsey in this case as well, completing the proof of Theorem 2.10. $\square$