Vertical profiles of biogenic volatile organic compounds as observed online at a tower in Beijing

Zhang, Huina; Zhang, Yanli; Huang, Zhonghui; Acton, W. Joe F.; Wang, Zhaoyi; Nemitz, Eiko; Langford, Ben; Mullinger, Neil; Davison, Brian; Shi, Zongbo; Liu, Di; Song, Wei; Yang, Weiqiang; Zeng, Jianqiang; Wu, Zhenfeng; Fu, Pingqing; Zhang, Qiang; Wang, Xinming

DOI:

[10.1016/j.jes.2020.03.032](https://doi.org/10.1016/j.jes.2020.03.032)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Zhang, H, Zhang, Y, Huang, Z, Acton, WF, Wang, Z, Nemitz, E, Langford, B, Mullinger, N, Davison, B, Shi, Z, Liu, D, Song, W, Yang, W, Zeng, J, Wu, Z, Fu, P, Zhang, Q & Wang, X 2020, 'Vertical profiles of biogenic volatile organic compounds as observed online at a tower in Beijing', *Journal of Environmental Sciences*, vol. 95, pp. 33-42. <https://doi.org/10.1016/j.jes.2020.03.032>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

**Vertical profiles of biogenic volatile organic compounds as observed
online at a tower in Beijing**

Huina Zhang^{1,3}, Yanli Zhang^{1,2,*}, Zhonghui Huang^{1,4}, W. Joe F. Acton⁵, Zhaoyi Wang¹,
Eiko Nemitz⁶, Ben Langford⁶, Neil Mullinger⁶, Brian Davison⁵, Zongbo Shi^{7,8}, Di Liu⁷,
Wei Song¹, Weiqiang Yang¹, Jianqiang Zeng^{1,2}, Zhenfeng Wu^{1,3}, Pingqing Fu^{8,9}, Qiang
Zhang¹⁰, Xinming Wang^{1,2,3}

1. State Key Laboratory of Organic Geochemistry and Guangdong Key Laboratory of
Environmental Protection and Resources Utilization, Guangzhou Institute of
Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

2. Center for Excellence in Regional Atmospheric Environment Institute of Urban
Environment, Chinese Academy of Sciences, Xiamen 361021, China

3. University of Chinese Academy of Sciences, Beijing 100049, China

4. State Environmental Protection Key Laboratory of Environmental Pollution Health
Risk Assessment and Guangdong Key Laboratory of Water and Air Pollution Control,
South China Institute of Environmental Science, Ministry of Ecology and Environment,
Guangzhou 510655, China

5. Lancaster Environment Centre, Lancaster University, Lancaster LA14YQ, UK

6. Centre for Ecology and Hydrology, Edinburgh, EH26 0QB, UK

7. School of Geography Earth and Environmental Sciences, University of Birmingham,
Birmingham B15 2TT, UK

8. Institute of Surface-Earth System Science, Tianjin University, Tianjin 300072, China

9. Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029,
China

10. Beijing Key Laboratory of Green Chemical Reaction Engineering and Technology,
Department of Chemical Engineering, Tsinghua University, Beijing 100084, China

Abstract

Vertical profiles of isoprene and monoterpenes were measured by a proton transfer reaction-time of flight-mass spectrometry (PTR-ToF-MS) at heights of 3, 15, 32, 64, and 102 m above the ground on the Institute of Atmospheric Physics (IAP) tower in central Beijing during the winter of 2016 and the summer of 2017. Isoprene mixing ratios were larger in summer due to much stronger local emissions whereas monoterpenes were lower in summer due largely to their consumption by much higher levels of ozone. Isoprene mixing ratios were the highest at the 32 m in summer (1.64 ± 0.66 ppbV) and at 15 m in winter (1.41 ± 0.64 ppbV) with decreasing concentrations to the ground and to the 102 m, indicating emission from the tree canopy of the surrounding parks. Monoterpene mixing ratios were the highest at the 3 m height in both the winter (0.71 ± 0.42 ppbV) and summer (0.16 ± 0.10 ppbV) with a gradual decreasing trend to 102 m, indicating an emission from near the ground level. The lowest isoprene and monoterpene mixing ratios all occurred at 102 m, which were 0.71 ± 0.42 ppbV (winter) and 1.35 ± 0.51 ppbV (summer) for isoprene, and 0.42 ± 0.22 ppbV (winter) and 0.07 ± 0.06 ppbV (summer) for monoterpenes. Isoprene in the summer and monoterpenes in the winter, as observed at the five heights, showed significant mutual correlations. In the winter monoterpenes were positively correlated with combustion tracers CO and acetonitrile at 3 m, suggesting possible anthropogenic sources.

Keywords:

Isoprene

Monoterpenes

Vertical profiles

Proton transfer reaction-time of flight-mass spectrometry (PTR-ToF-MS)

Biogenic volatile organic compounds (BVOCs)

55 *Corresponding author. E-mail: zhang_yl86@gig.ac.cn (Yanli Zhang)

56

Introduction

Volatile organic compounds (VOCs) from both anthropogenic and biogenic sources play crucial roles in the formation of ozone (O_3) and secondary organic aerosols (SOA) in the atmosphere (Henze et al., 2006). Biogenic VOCs (BVOCs) are mainly emitted from plant leaves and they account for ~90% of global annual VOC emissions (Guenther et al., 2012). Due to their relatively higher reactivity with atmospheric oxidants (Ryerson et al., 2001; Hallquist et al., 2009), BVOCs like isoprene have exceedingly strong ozone formation potential (OFP) compared with many anthropogenic VOCs (Atkinson, 2000; Calfapietra et al., 2013). In addition, on a global scale, it is estimated that SOA derived from biogenic sources greatly exceed that from anthropogenic sources (Hallquist et al., 2009).

As BVOCs, such as isoprene (C_5H_8) and monoterpenes ($C_{10}H_{16}$), are largely emitted from vegetation, the majority of field observations have been carried out in natural ecosystems (Baker et al., 2005; Eerdekens et al., 2009; Bai et al., 2017). However, model results suggest that BVOCs could contribute as much as 15% of ground level O_3 pollution formed in some of the metropolitan areas of Europe (Curci et al., 2009). Recent studies have also showed that anthropogenic pollutants including SO_2 and oxides of nitrogen (NO_x , $NO + NO_2$) can largely promote SOA formation from BVOCs (Shilling et al., 2013; Xu et al., 2015). Monoterpenes are estimated to be the largest source of summer organic aerosol in the southeastern United States (Zhang et al., 2018), and BVOC-derived SOA tracers have been widely observed to occur in atmospheric aerosols over cities (Fu et al., 2009; Ding et al., 2012; Lin et al., 2013; Martinsson et al., 2017). Thus, the effect of BVOCs in the urban environment is an important but currently understudied area of research.

Measurements of the vertical distribution of pollutants offer insight into their

sources, and enable the evaluation of chemical transport models and assessing indoor pollution at different heights (Jo and Kim, 2002; Caputi et al., 2019). Aircrafts, tethered balloons and towers are the most common platforms used to assess vertical gradients of pollutants. Yet, aircrafts are typically limited to higher altitude measurements (> 200 m) (Reeves et al., 2010; He et al., 2012), and tethered balloons are typically confined to rural areas due to space requirements (Greenberg et al., 1999; Sun et al., 2018). Tower-based measurements are, therefore, the most suitable option for use in the urban environment (Hollaway et al., 2019). While, many previous tower-based studies have been conducted in forests (Hemig et al., 1998; Kesselmeier et al., 2002; Yanez-Serrano et al., 2015), we believe the results presented here represent some of the first vertical profiles of BVOCs measured above a major urban conurbation.

Like many cities in developing countries, Beijing has air quality problems including very high levels of O₃ and fine particulate matter. Numerous field campaigns have been conducted to characterize anthropogenic VOCs in Beijing (Liu et al., 2017; Yang et al., 2018), but relatively few studies have focused on the measurement of biogenic compounds such as isoprene and monoterpenes and those that have typically been confined to offline analytical techniques (Duan et al., 2008; Cheng et al., 2018; Mo et al., 2018). Since monoterpenes are highly reactive with O₃, inefficient removal of O₃ during sampling could lead to the loss of monoterpenes and an underestimation of their abundance (Fick et al., 2001; Arnts et al., 2008). Furthermore, wall adsorption and storage time may also influence the determination of BVOCs in bags and canisters samples (Ahn et al., 2016). In contrast, online instruments such as proton transfer reaction-time of flight-mass spectrometry (PTR-ToF-MS) overcome these shortcomings by offering real-time monitoring with a sample resolution on the order of seconds (de Gouw and Warneke, 2007; Liu et al., 2016; Huang et al., 2017; Yuan et al.,

2017).

In this study, mixing ratios of isoprene and monoterpenes were measured by a PTR-ToF-MS in the winter of 2016 and the summer of 2017 at 5 different heights from the 325-meter the Institute of Atmospheric Physics (IAP) tower in central Beijing. The aims of this study were (i) to obtain vertical profiles of isoprene and monoterpenes in urban Beijing, (ii) to explore the factors influencing their abundance, and (iii) to evaluate if these predominantly biogenic species also have a significant contribution from anthropogenic sources within Beijing.

1. Material and methods

1.1 Vertical Profile Measurements

Field measurements were conducted as part of the joint UK-China Atmospheric Pollution and Human Health in a Developing Megacity (APHH-Beijing) research programme. The monitoring site (39°58'33" N, 116°22'41" E) is located in the Institute of Atmospheric Physics, Chinese Academy of Sciences (IAP, CAS), which is an urban site, between the North 3rd and 4th Ring Road in Beijing (Shi et al., 2019). Although in a central location, the sampling site is surrounded by four parks containing a variety of grassed and forested areas (Fig. 1). The dominant plants are evergreen trees, deciduous trees, evergreen shrubs and deciduous shrubs in the urban area of Beijing (Ghirardo et al., 2016). These trees are also the main species in parks near the sampling site.

Online monitoring was conducted at the 325 m tall tower located within the grounds of the IAP tower section. For each hour, ~15 minutes were spent measuring vertical profiles of isoprene and monoterpenes mixing ratios by sequentially sampling

air from the five measurement heights at 3, 15, 32, 64, and 102 m. Online measurements of ambient BVOCs were jointly conducted with colleagues from the United Kingdom who measured eddy covariance BVOC fluxes also using data from the same PTR-ToF-MS. Air samples were drawn from inlets at different heights to the ground station with high-volume pumps to minimize the residence time in the sampling lines to be less than 1 second. Then the automatic switching valves were used to achieve gradient switching. Sample air was drawn through individual 1/4 inch outer diameter (O.D.) Polytetrafluoroethylene (PTFE) sample lines at a flow rate of 3.3 L/min, with 0.3 L/min of this airflow diverted into heated (30°C) 10 L stainless steel canisters. The turnover time of air within each of the five canisters was ~ 30 min, which meant the PTR-ToF-MS could sample sequentially from each canister for just 170 sec and provide mixing ratios that were representative of the average during the previous half an hour. A further 5 minutes was spent measuring the instrument background and the remaining 40 min of each hour were for eddy covariance flux measurements. All air samples entered the PTR-ToF-MS via a 1 m long 1/4 inch O.D. PTFE line followed by a 20 cm long 1/16 inch O.D. polyetheretherketone (PEEK) inlet tube. A PTFE filter (Mitex, Merck KGaA, Ireland) was installed in front of the inlet to remove particulate matters. Trichlorobenzene was introduced into the inlet flow via diffusion through a needle valve to provide a high mass compound for mass calibration. The VOCs were measured by a PTR-ToF-MS 2000 (Ionicon Analytik GmbH, Innsbruck, Austria) housed in an air-conditioned container from November 23rd to December 12th, 2016 during the winter campaign and from June 10th to June 25th, 2017 during the summer campaign.

1.2 Instrument setup

The basic principles of PTR-ToF-MS have been described elsewhere in detail (Jordan et al., 2009; Huang et al., 2016). Briefly, the instrument consists of a hollow cathode ion source that generates a pure H_3O^+ reagent ion stream, a drift tube is used to ionize VOCs, and a high resolution time-of-flight mass spectrometer separates the ions according to their mass-to-charge ratio (m/z).

During the campaigns, the PTR-ToF-MS was operated under conditions of 1.9 mbar drift tube pressure, 60°C of inlet and drift temperature, and a drift tube voltage of 484 V, and E/N of 130 Td (where, E is electric field strength and N is the number density of a neutral gas; 1 Td is 10^{-17} V cm²).

Multipoint calibrations were performed twice every week using a VOC standard mixture (UK National Physical Laboratory; $\sim 1 \pm 0.10$ ppmV) by dynamic dilution with two mass flow controllers (Model 8500, KOFLOC, Japan; F-201CV, Bronkhorst, Germany), which had been calibrated before use by a flowmeter (Gilian Cilibrator 2, Sensidyne, USA). Calibration curves and method detection limit (MDLs) of isoprene and α -pinene in this campaign were presented in Appendix A Fig. S1. The MDLs of isoprene and α -pinene were 58 and 64 pptV, respectively (Huang et al., 2017). It is worth noting that isoprene (m/z 69) may be overestimated due to interference from Furan and the fragmentation of 2-methyl-3-butene-2-ol (MBO) (de Gouw et al., 2003; Yuan et al., 2017). Monoterpenes (m/z 137) were quantified by the sensitivity of α -pinene and may be underestimated because of a small fragment in m/z 87 (Warneke et al., 2003).

1.3 Trace gases and meteorological data

Data of other pollutants such as carbon monoxide (CO) and O_3 were obtained from

the Beijing Air Quality Monitoring Network with online monitoring at the site Chaoyang Olympic Sports Center (39°59'01" N, 116°23'56" E). The meteorological data including wind direction (WD), wind speed (WS), relative humidity (RH), and temperature (T) were obtained from the IAP tower where these parameters were monitored at three heights (8, 120 and 240 m); solar radiation (SR) was obtained from monitored at ground level on the tower; planetary boundary layer (PBL) height was calculated online from NOAA's READY Archived Meteorology program (<http://ready.arl.noaa.gov/READYamet.php>).

During the winter campaign (November 23rd to December 12th, 2016), air temperature ranged from -3.8 to 11.5°C with an average of 3.4°C; relative humidity ranged 11%-92% with an average of 43%; PBL height ranged 50-2047 m with an average of 225 m; solar radiation was 70.2 W/m² on average with a maximum of 625.8 W/m². During the summer campaign (June 10th to June 25th, 2017), air temperature ranged from 17.8 to 39.9°C with an average of 27.4°C; relative humidity ranged 14%-100% with an average of 54%; PBL height ranged 50-3078 m with an average of 676 m; solar radiation was 211.4 W/m² on average with a maximum of 1156.0 W/m².

2. Results and discussion

2.1 Overview about mixing ratios of isoprene and monoterpenes

Figs. 2 and **3** showed the time series of mixing ratios of isoprene and monoterpenes as well as that of meteorological parameters (wind speed and direction, temperature, relative humidity, PBL and solar radiation) during the winter and the summer campaign, respectively.

During the winter campaign, measured mixing ratios of isoprene at the five heights ranged from 0.01 to 3.11 ppbV with averages from 0.71 ± 0.42 ppbV (102 m) to $1.41 \pm$

0.64 ppbV (15 m). Our measured isoprene mixing ratios (1.18 ± 0.53 ppbV) at the 3 m height was comparable with that of 1.00 ± 0.48 ppbV (Table 1) measured 2.5 m above ground also during November 2016 by PTR-ToF-MS at an urban site about 10 km away from the IAP tower (Sheng et al., 2018). However, measured mixing ratios of isoprene at 15 m (1.41 ± 0.64 ppbV) or at other heights in this study were much higher than those previously reported based on canister-sampling offline measurements in Beijing during the cold non-growing seasons, for example 0.22 ± 0.13 ppbV observed ~15 m above ground at an urban site in Beijing during January-February 2015 (Li et al., 2019), 0.14 ± 0.14 ppbV at another urban site ~12 m above the ground in Beijing during January 2015 (Cheng et al., 2018), or even 0.04 ± 0.04 ppbV at a suburban site ~12 m above the ground in Beijing during November-December, 2014 (Li et al., 2019). The mixing ratios on average were also much higher than that of 0.05 ppbV reported at a forest site 532 m above sea level in Cyprus during March 2015 (Debevec et al., 2018).

During the winter campaign, observed mixing ratios of monoterpenes ranged from below the MDL to 1.31 ppbV with similar averages at the five heights from 0.57 ± 0.20 ppbV at 3 m (the highest) to 0.42 ± 0.22 ppbV at 102 m (the lowest). The average mixing ratios of monoterpenes at 15 m was 0.54 ± 0.18 ppbV. These averages were also about an order of magnitude higher when compared to those based on canister-sampling offline measurements in Beijing, such as 0.06 ± 0.06 ppbV at an urban site ~15 m above the ground during January-February 2015 or 0.04 ± 0.04 ppbV at a suburban site ~12 m above the ground during November-December 2014 (Table 1; Li et al., 2019). They were also substantially much higher than that measured in situ (0.28 ± 0.31 ppbV) with

an online gas chromatography-flame ionization detector (GC-FID) system at the forest site in Cyprus during March 2015 (Debevec et al., 2018).

During the summer campaign, which captured the growing seasons of the local vegetation, observed mixing ratios of isoprene were between 0.44 and 2.51 ppbV with an average from 1.35 ± 0.51 ppbV at 102 m to 1.64 ± 0.66 ppbV at 32 m (Table 1). These are higher than those measured during the winter campaign. This is not surprising because isoprene is typically emitted from vegetation as a function of light and temperature (Guenther et al., 2006, 2012). They are also higher than those previously observed using offline canister-sampling measurements in Beijing during the growing seasons. For example, Xie et al. (2018) reported concentrations of 0.89 ± 0.55 ppbV at an urban site ~20 m above ground during August-September 2006, Li et al. (2019) reported concentrations of 0.93 ± 0.53 ppbV at another urban site ~8 m above ground during July-August 2014, and Gong et al. (2018) observed concentrations of 0.29 ± 0.03 ppbV at a forest site 1690 above sea level in TianJian Mountain in south China during July-August 2016 (Table 1).

During the summer campaign, mixing ratios of monoterpenes ranged from below detection limits to 0.62 ppbV as observed at the five heights with an average from 0.07 ± 0.06 ppbV at 102 m to 0.16 ± 0.10 ppbV at 3 m (Table 1). They are even lower than that of 0.28 ± 0.31 ppbV at the forest site in Cyprus during March 2015 (Debevec et al., 2018). It is worth noting that the monoterpenes concentrations at the five heights during the summer campaign were much lower than those during the winter campaign (Table 1). Given that biogenic monoterpene emissions are temperature and/or light dependent,

this finding is somewhat counterintuitive. This is likely the result of the combination of atmospheric chemistry and boundary layer dynamics. Evergreen trees and shrubs, which account for ~50% of Beijing's vegetation distribution (Ghirardo et al., 2016) can emit a small amount of monoterpenes, even in wintertime (Guenther et al., 2012). However, PBL height (225 m) during the winter campaign was much lower than that during the summer campaign (676 m). This would facilitate the accumulation of monoterpenes in the winter. In addition, monoterpenes have a much shorter atmospheric lifetime with respect to ozone than compared with isoprene (Seinfeld and Pandis, 1998). During the summer, the O₃ concentration (62.8 ppbV) during the summer campaign was about 7-fold that of 8.6 ppbV during the winter campaign, resulting in a much shorter chemical lifetime in the summer (Shi et al., 2019). Thus, both the strong chemical sink and the higher boundary layer contributed to the higher measured monoterpene concentrations in the summer despite stronger emission rates (Acton et al., 2018).

2.2 Vertical profiles of isoprene and monoterpenes

2.2.1 Gradient distributions

Tower-based and airborne measurements in Amazonia indicate that concentrations of BVOCs declined strongly with altitude (Kuhn et al., 2007). As showed in Fig. 4 and Table 1, the highest average mixing ratios of isoprene (1.41 ± 0.61 ppbV) in the winter campaign was observed at 15 m, which is at a similar height as the tree canopy. Average mixing ratios of isoprene at 3 m was 1.18 ± 0.53 ppbV, just below that at 15 m. This was probably influenced by deposition towards the ground, as revealed in an

Amazonian rainforest when strong gradients towards the ground level from canopy height were also observed (Yanez-Serrano et al., 2015). From 15 to 102 m, isoprene mixing ratios showed a gradual decreasing trend, with the minimum of 0.71 ± 0.42 ppbV at 102 m, about 50% of that at 15 m. In the summer campaign, isoprene mixing ratio peaked at 32 m (1.64 ± 0.66 ppbV), and decreased to 15 m (1.61 ± 0.68 ppbV) and 3 m (1.49 ± 0.64 ppbV). This is probably also due to deposition towards the ground. From 32 to 102 m, it also showed a downward trend with the lowest level at 102 m (1.35 ± 0.51 ppbV), about 82% of that at 32 m. The contrast between the maximum and minimum of the averages at the five heights for isoprene was within 50% in the winter campaign but only 20% in the summer campaign, largely due to better near-surface mixing in the summer.

For monoterpenes, the highest mixing ratios were observed at 3 m during both the winter and the summer campaigns, and the average mixing ratios of monoterpenes all showed a downward trend from 3 to 102 m (Table 1; Fig. 4). During the winter campaign, the average mixing ratios of monoterpenes decrease by about 27% from 0.57 ± 0.20 ppbV at 3 m to 0.42 ± 0.22 ppbV at 102 m; whereas during the summer campaign it decrease by about 56% from 0.16 ± 0.10 to 0.07 ± 0.06 ppbV. This larger contrast in mixing ratios of monoterpenes between 3 and 102 m in the summertime reflects the influence of ozone on the scavenging of monoterpenes as discussed above.

The highest isoprene mixing ratios were observed on November 26th, 2016 in the winter campaign and June 16th, 2017 in the summer campaign. The highest mixing ratios of monoterpenes appeared were observed on November 26th, 2016 during the

winter campaign and on June 11th and 13th, 2017 in the summer campaign. As showed in **Fig. 4**, the vertical profiles of the isoprene and monoterpenes during these days were similar to that of the campaign averages, but the average mixing ratios during these particular days were about 50% higher.

2.2.2 Correlation of BVOCs between different heights

To indicate if the mixing ratios of BVOCs at the five heights changed in a similar pattern, the correlation analysis between BVOCs at the height with the highest average concentration and those at other heights and were performed (**Fig. 5**). During the winter campaign, the mixing ratios of isoprene at 102, 64, 32 and 3 m were only modestly correlated with those at 15 m, with correlation coefficient R^2 of 0.31-0.60. In the summer, however, the mixing ratios of isoprene at other heights showed significant correlations with those at 32 m with R^2 of 0.91-0.98. This is probably due to the dominant contribution of local emission from plant leaves and better near-ground mixing conditions in the summer. In contrary to isoprene, monoterpenes at 3 m showed a much better correlation with those at other heights in the winter ($R^2 = 0.54- 0.75$) than in the summer($R^2 = 0.78-0.95$). The relatively poorer correlations of the mixing ratios of monoterpenes between the different heights and the decrease in R^2 with heights, might be resulted from their reaction with ozone at a much higher rate than in the winter during upward transport.

2.2.3 Diurnal variations at different heights

As shown in **Fig. 6**, similar diurnal variations were observed at the five heights. During the winter campaign, isoprene showed no obvious diurnal changes while

monoterpenes had the higher concentrations at night and the lowest ones occurred at midday, consistent with the results by Hellen et al. (2012) and Cheng et al. (2018). Unlike isoprene, monoterpenes are emitted not only directly by plants' synthesis under light but also by pool storage (Oderbolz et al., 2013). For some tree species, terpenes may be formed and preserved in mesophyll or glandular cells during the night and evaporates as the temperature increases (Loreto and Schnitzler, 2010). Overall, the diurnal variation of BVOCs is a balance between the higher emission during daytime under elevated temperature or solar radiation and enhanced accumulation during nighttime with lower PBL heights.

During the summer campaign, isoprene mixing ratios peaked at 15:00-16:00, coincident with the maxima in solar radiation, and decreased from 18:00 until the following morning. In the summer, the stronger emissions of BVOCs during the daytime overtook the influence of changing PBL heights, thus they all had higher mixing ratios during daytime than nighttime.

2.3 Influencing factors and source implications

As shown in Appendix A Fig. S2, monoterpenes were positively correlated with relatively humidity (RH, $R^2 = 0.41-0.47$) during the winter campaign. Previous studies demonstrated that humidity could increase monoterpenes emission rates under dry conditions (Lamb et al., 1985; Schade et al., 1999). A recent study in the Mediterranean also suggested that higher relative humidity and rainfall will promote the emission of plant BVOCs in the dry season (Debevec et al., 2018). During the winter campaign, mixing ratios of isoprene and monoterpenes were both elevated on November 25th and

December 3rd-4th, largely due to their accumulation in a shallow boundary layer heights and lower wind speeds (Fig. 2); the higher relative humidity might also be a contributing factor to elevated mixing ratios during these two episodes. Except humidity, no further significant correlations between isoprene and monoterpenes and other meteorological parameters during the winter campaign.

The mixing ratios of isoprene were significantly correlated with temperature ($R^2 = 0.73-0.86$) but negatively correlated with relative humidity ($R^2 = 0.22-0.39$). In the summer campaign, isoprene mixing ratios were the highest on June 14th-17th and 20th-21st when the temperature and solar radiation were higher (Fig. 3). Although biogenic emissions of isoprene are known to be both light and temperature dependent (Guenther et al., 2006), no correlation between isoprene and light were observed during either measurement campaign. No significant correlations between the monoterpenes and meteorological parameters were found in the summer.

To further examine if there were sources of BVOCs other than emission from plant leaves, the relationship between BVOCs and other trace gases were also investigated. Monoterpenes showed significant correlations with CO and acetonitrile in the winter as observed at the 3 m height with R^2 of 0.54 and 0.71, respectively (Fig. 7). Since CO is a typical tracer of incomplete combustion of biomass or fossil fuels (Parrish et al., 2009; Zhang et al., 2015) and acetonitrile was a marker of biomass burning (Fang et al., 2017), this good correlation between monoterpenes and combustion tracers at 3 m near the ground suggests that combustion sources, particularly biomass burning, might have contributed to monoterpenes at the IAP site. Indeed, biomass burning has been reported

as an anthropogenic source of monoterpenes (Andreae and Merlet, 2001; Stockwell et al., 2015). In addition, wintertime heating in Beijing started on November 15th, and coal consumption and residential biomass were still the main fuels for central heating in Beijing (Beijing Municipal Bureau of Statistics, 2017; Yang et al., 2018). Thus, it was possible that combustion processes contributed to the emission of monoterpenes in the winter. Isoprene did not show a significant correlation with combustion tracers.

3. Conclusions

The mixing ratios of isoprene and total monoterpenes mixing ratios were measured online with a PTR-ToF-MS at five heights in central Beijing during two contrasting seasons (winter 2016 and summer 2017). Observed mixing ratios of isoprene and monoterpenes based on online measurements at the five heights were much higher than those previous measured by offline canister measurements. At the lowest height of 3 m, the average mixing ratios of isoprene reached 1.49 ± 0.64 ppbV in the summer and 1.18 ± 0.53 ppbV in the winter, suggesting potential important contribution to ozone formation by BVOCs even in the urban areas of Beijing.

Among the five heights, average mixing ratios of isoprene were highest at 32 m (1.64 ± 0.66 ppbV) in the summer and 15 m (1.41 ± 0.64 ppbV) in the winter. From the height with the highest average isoprene mixing ratios, isoprene showed a downward trend towards the ground and towards the height of 102 m. Mixing ratios of monoterpenes were highest at 3 m both in the winter (0.71 ± 0.42 ppbV) and the summer (0.16 ± 0.10 ppbV), and they decreased with altitude during both the winter and summer campaigns. The main reason for the lower ambient mixing ratios of monoterpenes

observed in the summer, when higher emissions occur, is much higher levels of O₃ and reduced the atmospheric lifetime of the monoterpenes.

At the five heights, isoprene observed in summer and monoterpenes observed in winter, showed highly significant mutual correlations, while isoprene observed in winter or monoterpenes observed in the summer showed poorer mutual correlations, largely due to different roles play by emissions, atmospheric oxidation and dispersion conditions in the two seasons. Isoprene showed significant correlations with temperature in the summer while monoterpenes with relative humidity in the winter. In addition, in winter monoterpenes showed significantly positive correlations with CO and acetonitrile, suggesting possible emissions from anthropogenic sources, particularly combustion sources such as biomass burning and coal combustion.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (Nos. 41571130031, 41673116, 41703112, 41603070), the Natural Environment Research Council (Nos. NE/N006992/1 and NE/N006976/1), Theme-based Research Scheme (No. T24-504/17-N), Youth Innovation Promotion Association of the Chinese Academy of Sciences (No. 2017406) and Natural Science Foundation of Guangdong Province (No. 2016A030310117). ZS acknowledges financial support from Natural Environment Research Council (NE/N007190/1). We acknowledge the support from Pingqing Fu, Zifa Wang, Jie Li and Yele Sun from IAP for hosting the Atmospheric Pollution and Human Health in a Developing Megacity (APHH-Beijing) campaign at IAP. We thank Zongbo Shi, Di Liu, Roy Harrison, Tuan Vu and Bill Bloss from the

University of Birmingham, Siyao Yue, Liangfang Wei, Hong Ren, Qiaorong Xie, Wanyu Zhao, Linjie Li, Ping Li, Shengjie Hou, Qingqing Wang from IAP, Rachel Dunmore, Ally Lewis and James Lee from the University of York, Kebin He and Xiaoting Cheng from Tsinghua University, and James Allan and Hugh Coe from the University of Manchester for providing logistic and scientific support for the field campaigns.

Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version at xxxxxx.

References

- Acton, W.J.F., Jud, W., Ghirardo, A., Wohlfahrt, G., Hewitt, C.N., Taylor, J.E., et al. 2018. The effect of ozone fumigation on the biogenic volatile organic compounds (BVOCs) emitted from *Brassica napus* above- and below-ground. *PLoS ONE* 13(12), e0208825.
- Ahn, J.H., Deep, A., Kim, K.H., 2016. The storage stability of biogenic volatile organic compounds (BVOCs) in polyester aluminum bags. *Atmos. Environ.* 141, 430-434.
- Andreae, M.O., Merlet, P., 2001. Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966.
- Arnts, R.R., 2008. Reduction of biogenic VOC sampling losses from ozone via trans-2-Butene addition. *Environ. Sci. Technol.* 42 (20), 7663-7669.
- Atkinson, R., 2000. Atmospheric chemistry of VOCs and NO_x. *Atmos. Environ.* 34 (12-14), 2063-2101.
- Bai, J.H., Guenther, A., Turnipseed, A., Duhl, T., Greenberg, J., 2017. Seasonal and interannual variations in whole-ecosystem BVOC emissions from a subtropical plantation in China. *Atmos. Environ.* 161, 176-190.
- Baker, B., Bai, J.H., Johnson, C., Cai, Z.T., Li, Q.J., Wang, Y.F., et al., 2005. Wet and

dry season ecosystem level fluxes of isoprene and monoterpenes from a southeast Asian secondary forest and rubber tree plantation. *Atmos. Environ.* 39 (2), 381-390.

Beijing Municipal Bureau of Statistics (BMBS): Beijing Statistical Yearbook 2017, China Statistics Press, Beijing, 2017.

Cheng, X., Li, H., Zhang, Y.J., Li, Y.P., Zhang, W.Q., Wang, X.Z., et al., 2018. Atmospheric isoprene and monoterpenes in a typical urban area of Beijing: Pollution characterization, chemical reactivity and source identification. *J. Environ. Sci.* 71, 150-167.

Curci, G., Beekmann, M., Vautard, R., Smiatek, G., Steinbrecher, R., Theloke, J., et al., 2009. Modelling study of the impact of isoprene and terpene biogenic emissions on European ozone levels. *Atmos. Environ.* 43 (7), 1444-1455.

de Gouw, J., Warneke, C., 2007. Measurements of volatile organic compounds in the earth's atmosphere using proton-transfer-reaction mass spectrometry. *Mass Spectrom. Rev.* 26 (2), 223-257.

de Gouw, J.A., Goldan, P.D., Warneke, C., Kuster, W.C., Roberts, J.M., Marchewka, M., et al., 2003. Validation of proton transfer reaction-mass spectrometry (PTR-MS) measurements of gas-phase organic compounds in the atmosphere during the New England Air Quality Study (NEAQS) in 2002. *J. Geophys. Res.* 108 (D21).

Debevec, C., Sauvage, S., Gros, V., Sellegri, K., Sciare, J., Pikridas, M., et al., 2018. Driving parameters of biogenic volatile organic compounds and consequences on new particle formation observed at an eastern Mediterranean background site. *Atmos. Chem. Phys.* 18 (19), 14297-14325.

Ding, X., Wang, X.M., Gao, B., Fu, X.X., He, Q.F., Zhao, X.Y., et al., 2012. Tracer-based estimation of secondary organic carbon in the Pearl River Delta, south China. *J. Geophys. Res.* 117, D05313.

Duan, J.C., Tan, J.H., Yang, L., Wu, S., Hao, J.M., 2008. Concentration, sources and ozone formation potential of volatile organic compounds (VOCs) during ozone episode in Beijing. *Atmos. Res.* 88 (1), 25-35.

Eerdekens, G., Ganzeveld, L., de Arellano, J.V.-G., Kluepfel, T., Sinha, V., Yassaa, N.,

- et al., 2009. Flux estimates of isoprene, methanol and acetone from airborne PTR-MS measurements over the tropical rainforest during the GABRIEL 2005 campaign. *Atmos. Chem. Phys.* 9 (13), 4207-4227.
- Fang, Z., Deng, W., Zhang, Y.L., Ding, X., Tang, M.J., Liu, T.Y., et al., 2017. Open burning of rice, corn and wheat straws: primary emissions, photochemical aging, and secondary organic aerosol formation. *Atmos. Chem. Phys.* 17 (24), 14821-14839.
- Fick, J., Pommer, L., Andersson, B., Nilsson, C., 2001. Ozone removal in the sampling of parts per billion levels of terpenoid compounds: An evaluation of different scrubber materials. *Environ. Sci. Technol.* 35 (7), 1458-1462.
- Fu, P., Kawamura, K., Chen, J., Barrie, L.A., 2009. Isoprene, Monoterpene, and Sesquiterpene oxidation products in the high Arctic aerosols during late Winter to early Summer. *Environ. Sci. Technol.* 43 (11), 4022-4028.
- Ghirardo, A., Xie, J., Zheng, X., Wang, Y., Grote, R., Block, K., et al., 2016. Urban stress-induced biogenic VOC emissions and SOA-forming potentials in Beijing. *Atmos. Chem. Phys.* 16 (5), 2901-2920.
- Gong, D.C., Wang, H., Zhang, S.Y., Wang, Y., Liu, S.C., Guo, H., et al., 2018. Low-level summertime isoprene observed at a forested mountaintop site in southern China: implications for strong regional atmospheric oxidative capacity. *Atmos. Chem. Phys.* 18 (19), 14417-14432.
- Greenberg, J.P., Guenther, A., Zimmerman, P., Baugh, W., Geron, C., Davis, K., et al., 1999. Tethered balloon measurements of biogenic VOCs in the atmospheric boundary layer. *Atmos. Environ.* 33 (6), 855-867.
- Guenther, A.B., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., Geron, C., 2006. Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210.
- Guenther, A.B., Jiang, X., Heald, C.L., Sakulyanontvittaya, T., Duhl, T., Emmons, L.K., et al., 2012. The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): An extended and updated framework for modeling biogenic

emissions. *Geosci. Model Dev.* 5 (6), 1471-1492.

Hallquist, M., Wenger, J.C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., et al., 2009. The formation, properties and impact of secondary organic aerosol: current and emerging issues. *Atmos. Chem. Phys.* 9 (14), 5155-5236.

He, H., Li, C., Loughner, C.P., Li, Z.Q., Krotkov, N.A., Yang, K., et al., 2012. SO₂ over central China: Measurements, numerical simulations and the tropospheric sulfur budget. *J. Geophys. Res.* 117, D00K37.

Hellen, H., Tykka, T., Hakola, H., 2012. Importance of monoterpenes and isoprene in urban air in northern Europe. *Atmos. Environ.* 59, 59-66.

Helmig, D., Balsley, B., Davis, K., Kuck, L.R., Jensen, M., Bognar, J., et al., 1998. Vertical profiling and determination of landscape fluxes of biogenic nonmethane hydrocarbons within the planetary boundary layer in the Peruvian Amazon. *J. Geophys. Res.* 103 (D19), 25519-25532.

Henze, D.K., Seinfeld, J.H., 2006. Global secondary organic aerosol from isoprene oxidation. *Geophys. Res. Lett.* 33 (9).

Hollaway, M., Wild, O., Yang, T., Sun, Y., Xu, W., Xie, C., et al., 2019. Photochemical impacts of haze pollution in an urban environment. *Atmos. Chem. Phys.* 2019 (19) 9699-9714.

Huang, Z.H., Zhang, Y.L., Yan, Q., Zhang, Z., Wang, X.M., 2016. Real-time monitoring of respiratory absorption factors of volatile organic compounds in ambient air by proton transfer reaction time-of-flight mass spectrometry. *J. Hazard. Mater.* 320, 547-555.

Huang, Z.H., Zhang, Y.L., Yan, Q., Wang, Z.Y., Zhang, Z., Wang, X.M., 2017. Decreased human respiratory absorption factors of aromatic hydrocarbons at lower exposure levels: The dual effect in reducing ambient air toxics. *Environ. Sci. Technol. Lett.* 4 (11), 463-469.

Jo, W.K., Kim, K.Y., 2002. Vertical variability of volatile organic compound (VOC) levels in ambient air of high-rise apartment buildings with and without occurrence of surface inversion. *Atmos. Environ.* 36, 5645-5652.

Jordan, A., Haidacher, S., Hanel, G., Hartungen, E., Maerk, L., Seehauser, H., et al.,

2009. A high resolution and high sensitivity proton-transfer-reaction time-of-flight mass spectrometer (PTR-TOF-MS). *Int. J. Mass spectrom.* 286, 122-128.
- Kesselmeier, J., Kuhn, U., Rottenberger, S., Biesenthal, T., Wolf, A., Schebeske, G., et al., 2002. Concentrations and species composition of atmospheric volatile organic compounds (VOCs) as observed during the wet and dry season in Rondonia (Amazonia). *J. Geophys. Res.* 107 (D20), 8053.
- Kuhn, U., Andreae, M.O., Ammann, C., Araujo, A.C., Brancaleoni, E., Ciccioli, P., et al., 2007. Isoprene and monoterpene fluxes from Central Amazonian rainforest inferred from tower-based and airborne measurements, and implications on the atmospheric chemistry and the local carbon budget. *Atmos. Chem. Phys.* 7 (11), 2855-2879.
- Lamb, B., Westberg, H., Allwine, G., Quarles, T., 1985. Biogenic hydrocarbon emissions from deciduous and coniferous trees in United-States. *J. Geophys. Res.* 90 (ND1), 2380-2390.
- Li, K., Li, J.L., Tong, S.R., Wang, W.G., Huang, R.J., Ge, M.F., 2019. Characteristics of wintertime VOCs in suburban and urban Beijing: concentrations, emission ratios, and festival effects. *Atmos. Chem. Phys.* 19 (12), 8021-8036.
- Lin, Y.H., Zhang, H., Pye, H.O.T., Zhang, Z., Marth, W.J., Park, S., et al., 2013. Epoxide as a precursor to secondary organic aerosol formation from isoprene photooxidation in the presence of nitrogen oxides. *Proc. Natl. Acad. Sci. USA* 110 (17), 6718-6723.
- Liu, C.T., Ma, Z.B., Mu, Y.J., Liu, J.F., Zhang, C.L., Zhang, Y.Y., et al., 2017. The levels, variation characteristics, and sources of atmospheric non-methane hydrocarbon compounds during wintertime in Beijing, China. *Atmos. Chem. Phys.* 17 (17), 10633-10649.
- Liu, C.T., Mu, Y.J., Zhang, C.L., Zhang, Z.B., Zhang, Y.Y., Liu, J.F., et al., 2016. Development of gas chromatography-flame ionization detection system with a single column and liquid nitrogen-free for measuring atmospheric C₂-C₁₂ hydrocarbons. *J. Chromatogr. A* 1427, 134-141.
- Loreto, F., Schnitzler, J.-P., 2010. Abiotic stresses and induced BVOCs. *Trends Plant*

Sci. 15 (3), 154-166.

Martinsson, J., Monteil, G., Sporre, M.K., Hansen, A.M.K., Kristensson, A., Stenstrom, K.E., et al., 2017. Exploring sources of biogenic secondary organic aerosol compounds using chemical analysis and the FLEXPART model. *Atmos. Chem. Phys.* 17 (18), 11025-11040.

Mo, Z.W., Shao, M., Wang, W.J., Liu, Y., Wang, M., Lu, S.H., 2018. Evaluation of biogenic isoprene emissions and their contribution to ozone formation by ground-based measurements in Beijing, China. *Sci. Total Environ.* 627, 1485-1494.

Oderbolz, D.C., Aksoyoglu, S., Keller, J., Barmpadimos, I., Steinbrecher, R., Skjoth, C.A., et al., 2013. A comprehensive emission inventory of biogenic volatile organic compounds in Europe: Improved seasonality and land-cover. *Atmos. Chem. Phys.* 13 (4), 1689-1712.

Parrish, D.D., Kuster, W.C., Shao, M., Yokouchi, Y., Kondo, Y., Goldan, P.D., et al., 2009. Comparison of air pollutant emissions among mega-cities. *Atmos. Environ.* 43 (40), 6435-6441.

Reeves, C.E., Formenti, P., Afif, C., Ancellet, G., Attie, J.L., Bechara, J., et al., 2010. Chemical and aerosol characterisation of the troposphere over West Africa during the monsoon period as part of AMMA. *Atmos. Chem. Phys.* 10 (16), 7575-7601.

Ryerson, T.B., Trainer, M., Holloway, J.S., Parrish, D.D., Huey, L.G., Sueper, D.T., et al., 2001. Observations of ozone formation in power plant plumes and implications for ozone control strategies. *Science*, 292 (5517), 719-723.

Schade, G.W., Goldstein, A.H., Lamanna, M.S., 1999. Are monoterpene emissions influenced by humidity? *Geophys. Res. Lett.* 26 (14), 2187-2190.

Seinfeld, J.H., Pandis, S.N., 1998. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Changes*. Wiley, New York, USA.

Sheng, J.J., Zhao, D.L., Ding, D.P., Li, X., Huang, M.Y., Gao, Y., et al., 2018. Characterizing the level, photochemical reactivity, emission, and source contribution of the volatile organic compounds based on PTR-TOF-MS during winter haze period in Beijing, China. *Atmos. Res.* 212, 54-63.

Shi, Z.B., Vu, T., Kotthaus, S., Harrison, R.M., Grimmond, S., Yue, S.Y., et al., 2019.

- Introduction to the special issue "In-depth study of air pollution sources and processes within Beijing and its surrounding region (APHH-Beijing)". *Atmos. Chem. Phys.* 19 (11), 7519-7546.
- Shilling, J.E., Zaveri, R.A., Fast, J.D., Kleinman, L., Alexander, M.L., Canagaratna, M.R., et al., 2013. Enhanced SOA formation from mixed anthropogenic and biogenic emissions during the CARES campaign. *Atmos. Chem. Phys.* 13 (4), 2091-2113.
- Stockwell, C.E., Veres, P.R., Williams, J., Yokelson, R.J., 2015. Characterization of biomass burning emissions from cooking fires, peat, crop residue, and other fuels with high-resolution proton-transfer-reaction time-of-flight mass spectrometry. *Atmos. Chem. Phys.* 15 (2), 845-865.
- Sun, J., Wang, Y.S., Wu, F.K., Tang, G.Q., Wang, L.L., Wang, Y.H., et al., 2018. Vertical characteristics of VOCs in the lower troposphere over the North China Plain during pollution periods. *Environ. Pollut.* 236, 907-915.
- Warneke, C., De Gouw, J.A., Kuster, W.C., Goldan, P.D., Fall, R., 2003. Validation of atmospheric VOC measurements by proton-transfer-reaction mass spectrometry using a gas-chromatographic pre-separation method. *Environ. Sci. Technol.* 37 (11), 2494-2501.
- Xie, X., Shao, M., Liu, Y., Lu, S.H., Chang, C.-C., Chen, Z.-M., 2008. Estimate of initial isoprene contribution to ozone formation potential in Beijing, China. *Atmos. Environ.* 42 (24), 6000-6010.
- Xu, L., Guo, H.Y., Boyd, C.M., Klein, M., Bougiatioti, A., Cerully, K.M., et al., 2015. Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the southeastern United States. *Proc. Natl. Acad. Sci. USA.* 112 (1), 37-42.
- Yanez-Serrano, A.M., Noelscher, A.C., Williams, J., Wolff, S., Alves, E., Martins, G.A., et al., 2015. Diel and seasonal changes of biogenic volatile organic compounds within and above an Amazonian rainforest. *Atmos. Chem. Phys.* 15 (6), 3359-3378.
- Yang, W.Q., Zhang, Y.L., Wang, X.M., Li, S., Zhu, M., Yu, Q.Q., et al., 2018. Volatile organic compounds at a rural site in Beijing: influence of temporary emission

control and wintertime heating. *Atmos. Chem. Phys.* 18 (17), 12663-12682.

Yuan, B., Koss, A.R., Warneke, C., Coggon, M., Sekimoto, K., de Gouw, J.A., 2017. Proton-Transfer-Reaction Mass Spectrometry: Applications in atmospheric sciences. *Chem. Rev.* 117 (21), 13187-13229.

Zhang, H.F., Yee, L.D., Lee, B.H., Curtis, M.P., Worton, D.R., Isaacman-VanWertz, G., et al., 2018. Monoterpenes are the largest source of summertime organic aerosol in the southeastern United States. *Proc. Natl. Acad. Sci. USA* 115 (9), 2038-2043.

Zhang, Z., Zhang, Y.L., Wang, X.M., Lu, S.J., Huang, Z.H., Huang, X.Y., et al., 2016. Spatiotemporal patterns and source implications of aromatic hydrocarbons at six rural sites across China's developed coastal regions. *J. Geophys. Res.* 121 (11), 6669-6687.

Table 1 Comparison of average mixing ratios of isoprene and monoterpenes (average $\pm$ standard deviation) observed at IAP tower in our study with those reported in Beijing and other forest sites.

Sampling site	Height (m)	Type	Isoprene (ppbV)	Monoterpenes (ppbV)	Study periods
PKU, Beijing ^a	~20	Urban	0.89 $\pm$ 0.55	NA	Aug.-Sep., 2006
NCNST, Beijing ^b	~15	Urban	0.22 $\pm$ 0.13	0.06 $\pm$ 0.06	Jan.-Feb., 2015
YUFA, Beijing ^b	~12	Rural	0.64 $\pm$ 0.44	NA	Jan.-Feb., 2015
UCAS, Beijing ^b	~12	Suburban	0.04 $\pm$ 0.04	0.04 $\pm$ 0.04	Nov.-Dec., 2014
CRAES, Beijing ^c	8	Urban	0.93 $\pm$ 0.53	NA	Jul.-Aug., 2014
			0.14 $\pm$ 0.14	NA	Jan., 2015
BAAF, Beijing ^d	2.5	Urban	1.00 $\pm$ 0.48	NA	Dec., 2016
Cyprus ^e	532 ^h	Forest	0.05 $\pm$ 0.01	0.28 $\pm$ 0.31	Mar., 2015
Mt. TianJing, China ^f	1690 ^h	Forest	0.29 $\pm$ 0.03	NA	Jul.-Aug., 2016
	3		1.18 $\pm$ 0.53	0.57 $\pm$ 0.20	
	15		1.41 $\pm$ 0.64	0.54 $\pm$ 0.18	
IAP, Beijing ^g	32	Urban	0.99 $\pm$ 0.44	0.52 $\pm$ 0.17	Nov.-Dec., 2016
	64		0.83 $\pm$ 0.43	0.49 $\pm$ 0.17	
	102		0.71 $\pm$ 0.42	0.42 $\pm$ 0.22	
	3		1.49 $\pm$ 0.64	0.16 $\pm$ 0.10	
	15		1.61 $\pm$ 0.68	0.14 $\pm$ 0.09	
IAP, Beijing ^g	32	Urban	1.64 $\pm$ 0.66	0.11 $\pm$ 0.08	Jun., 2017
	64		1.37 $\pm$ 0.62	0.09 $\pm$ 0.07	
	102		1.35 $\pm$ 0.51	0.07 $\pm$ 0.06	

^aXie et al., 2008, ^bLi et al., 2019, ^cCheng et al., 2018, ^dSheng et al., 2018, ^eDebevec et al., 2018, ^fGong et al., 2018, ^gThis study; ^hMeters above sea level. NA: not available; PKU: Peking University; NCNST: the National Center for Nanoscience and Technology of China; YUFA: Yu Fa Town; UCAS: the University of Chinese Academy of Sciences; CRAES: the Chinese Research Academy of Environmental Sciences; BAAF: the Beijing Academy of Agriculture and Forestry; Mt. TianJing: TianJing Mountain.

Fig. 1 A map showing the surroundings of the Institute of Atmospheric Physics (IAP) tower (left), a diagram showing the sampling heights in the IAP tower (middle) and the location of IAP tower in Beijing (upper-right corner).

Fig. 2 Time series of (a) wind speed and wind direction, (b) temperature (T), (c) relative humidity (RH), (d) solar radiation (SR) and planetary boundary layer height (PBL), (e) mixing ratios of isoprene and (f) monoterpenes during the winter campaign in 2016.

Fig. 3 Time series of (a) wind speed and wind direction, (b) temperature (T), (c) relative humidity (RH), (d) solar radiation (SR) and planetary boundary layer height (PBL), (e) mixing ratios of isoprene and (f) monoterpenes during the summer campaign in 2017.

Fig. 4 Vertical profiles for mixing ratios of isoprene and monoterpenes during the winter and the summer campaigns. Range bars represent the 25% and 75% percentiles. Solid lines represents the averages during the whole summer or winter campaign; dashed lines are the days (2016/11/26, 2017/06/11, 2017/06/13 and 2017/06/16) when higher levels of isoprene or monoterpenes occurred.

Fig. 5 (a) Correlations between the isoprene concentrations at 15 m and those at other heights in the winter; (b) correlations between the isoprene mixing ratios at 32 m and those at other heights in the summer; correlations between the mixing ratios of monoterpenes at 3 m and those at other heights (c) in the winter and (d) in the summer.

Fig. 6 Diurnal variations of isoprene and monoterpenes observed at the five heights in the winter 2016 campaign and the summer 2017 campaign. Range bars are expressed the 25% and 75% percentiles.

655

656 **Fig. 7** Correlations of monoterpenes with CO and acetonitrile at 3 m during the 2016
 657 winter campaign.