Rationale for CD40 pathway blockade in autoimmune rheumatic disorders

Pucino, Valentina; Gardner, David; Fisher, Benjamin

DOI:

[10.1016/S2665-9913\(20\)30038-2](https://doi.org/10.1016/S2665-9913(20)30038-2)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Pucino, V, Gardner, D & Fisher, B 2020, 'Rationale for CD40 pathway blockade in autoimmune rheumatic disorders', *The Lancet Rheumatology*, vol. 2, no. 5, pp. e292-e301. [https://doi.org/10.1016/S2665-9913\(20\)30038-2](https://doi.org/10.1016/S2665-9913(20)30038-2)

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Rationale for CD40 pathway blockade in autoimmune rheumatic disorders

Valentina Pucino^{1,2} PhD; David H. Gardner¹ PhD; Benjamin A. Fisher^{1,2} MD(Res)

1. Institute of Inflammation and Ageing, College of Medical and Dental Sciences, University of Birmingham (UK)
2. National Institute for Health Research (NIHR) Birmingham Biomedical Research Centre and Department of Rheumatology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK.

* Correspondence to: Dr Benjamin Fisher

Institute of Inflammation and Ageing
College of Medical and Dental Sciences
University of Birmingham
Queen Elizabeth Hospital, Birmingham, B15 2WB
Direct Line: +44 (0) 7956617385
Email: B.Fisher@bham.ac.uk

Abstract

CD40 and CD40L (CD154) belong to the tumour necrosis factor receptor superfamily and are expressed by immune and non-immune cells. CD40L plays a central role in co-stimulation and regulation of the immune response via activation of CD40-expressing cells. Imbalance of the CD40/CD40L costimulatory pathway is reported in many autoimmune diseases including systemic lupus erythematosus (SLE), rheumatoid arthritis (RA) and Sjögren's syndrome (SS) thus supporting its role in the breach of immune tolerance typical of these diseases. Targeting CD40/CD40L signalling may represent a novel therapeutic option for several autoimmune disorders.

Key points:

1. CD40/CD40L signalling pathway regulates immune and non-immune cell responses
2. CD40/CD40L signalling is altered in autoimmune diseases such as SLE, RA, and SS
3. Early clinical trials programmes targeting CD40L were halted due to thrombotic adverse events
4. New therapeutic approaches targeting CD40 or using modified molecules against CD40L are currently being tested
5. Targeting CD40/CD40L signalling is a promising novel therapeutic strategy for reducing inflammation in autoimmune rheumatic disorders

1. Introduction

CD40L/CD40 interactions exert profound effects on immune and non-immune cells (1-3).

In a pathogenic setting, the deregulation of CD40 signalling has been observed in multiple autoimmune diseases (1-3).

Conversely, therapeutic up-regulation of the CD40 pathway in cancer may have potent anti-tumor effects. Thus the CD40 pathway has long been an attractive therapeutic target for treating autoimmune diseases; however, early clinical trials of monoclonal antibodies blocking CD40 ligand (CD40L) were halted due to platelet-related thromboembolic complications (4).

CD40 is a transmembrane type I glycoprotein which belongs to the tumour necrosis factor (TNF) gene superfamily and behaves as a co-stimulatory molecule. It is constitutively expressed by B cells and antigen presenting cells including monocytes, neutrophils and dendritic cells, and may be expressed on other cell types such as epithelial cells, endothelial cells, smooth muscle cells, fibroblasts, and platelets (2). Its ligand, CD40L (CD154), is a type II transmembrane protein which exists in soluble (sCD40L) or membrane-bound form. CD40L is mainly expressed on activated T cells, B cells, natural killer, platelets, endothelial, epithelial, and smooth muscle cells (1, 2). Soluble CD40L is mainly produced by platelets (5) and activated T cells (6) and is functional, being capable of enhancing platelet activation and B cell proliferation (7). The wide expression of this co-stimulatory machinery indicates the pivotal roles they play in different cellular immune processes.

In this review, we will first outline the role of CD40–CD40L in the biology of immune cells. We will then review published data on the role of the CD40–CD40L signalling pathway in autoimmune rheumatic diseases and, finally, novel approaches to targeting the pathway for clinical efficacy.

2. Co-stimulation

The requirement for B and T cell co-stimulation during the initiation of adaptive immune responses acts as a checkpoint that is involved in maintaining immune tolerance. A failure to regulate these

signals can underlie the development of autoimmunity which makes them an attractive therapeutic target. Similarly, in cancer therapy, the inhibition of regulatory pathways such as CTLA-4 and PD-1 has been utilised successfully with the aim of propagating anti-tumor immunity.

The requirement for CD28 is of fundamental importance to the initiation of adaptive immune responses in that naïve T cells are dependent upon CD28-signalling for activation and proliferation. The CD28 pathway has been effectively targeted in the treatment of rheumatoid arthritis (RA) with abatacept [Orencia; CTLA-4Ig; (8)]. Although patient responses to therapy can be variable, a subpopulation of patients appear to respond particularly well to treatment. Nevertheless, a key problem associated with CD28-blockade is the inability to effectively time the treatment. Specifically, patients are unlikely to present in the clinic until many years after tolerance mechanisms were initially bypassed and autoimmune responses are initiated. At this stage, activated/memory T cells become less dependent upon CD28-costimulation and are located at sites of inflammation where features of the local microenvironment diminish the efficacy of therapy. Indeed, attention has turned towards treating patients at earlier stages of the disease (9), even so far as to treat individuals who are at risk of RA development prior to the onset of active synovitis (10).

Although, CD28 can continue to play a role in an ongoing immune response, the upregulation of other costimulatory pathways implies some redundancy. These alternate co-stimulatory pathways include other members of the CD28 family and wider Immunoglobulin superfamily and TNF-receptor superfamily members which cooperate with/take over from CD28 in driving the maintenance, differentiation and effector function of activated T cell populations (11).

This hierarchical arrangement in the relative contributions of various costimulatory and inhibitory pathways brings additional targets for biological therapies in autoimmunity with the capacity to modulate features of lymphocyte effector function which are the basis of immune mediated pathologies (**Figure 1**). For example, ICOS (Inducible T cell co-stimulator) expression is driven by CD28 signaling in activated T cells (12). This mediates signalling that is important to the germinal centre reaction via the maintenance of T follicular helper cells (Tfh) (13, 14) which places the ICOS-

ICOS-L pathway at the center of B/T Cell crosstalk and therefore an appealing target in B cell mediated immune pathologies. Indeed, ICOS-L has been targeted via a fully human monoclonal antibody (AMG-557; prezalumab) although a phase 2a clinical trial to evaluate the safety and efficacy of AMG 557 in subjects with primary Sjögren's syndrome showed no efficacy. It is possible that this is associated with redundancy created by the presence of additional costimulatory pathways.

The CD40-CD40L pathway fulfils multiple roles within this co-stimulation hierarchy. By driving Antigen Presenting Cell (APC) activation, including the expression of the CD28 ligands CD80 and CD86 (15, 16), CD40 signaling could be seen to sit above CD28 in the hierarchy by licensing effective CD28 co-stimulation. However, the CD40-CD40-L also forms an integral part of T cell effector function through a key role that is played in B/T cell crosstalk. This multifaceted role that is played in the generation of an adaptive immune responses has generated significant interest in the pathway as a target for therapy in autoimmune/inflammatory diseases.

3. CD40L/CD40 signalling in immune cells

The engagement of CD40 by CD40L promotes an intracellular signalling cascade, including the recruitment of proteins called TNFR-associated factors (TRAFs) and the activation of the nuclear factor κ B (NF κ B)-signaling pathway which culminates in the activation of transcription factors and cytokine production (17).

CD40 co-stimulation, in concert with cytokines or other stimuli, promotes B cell activation, proliferation, differentiation and survival via the upregulation of co-stimulatory molecules CD80 and CD86 which interact with the B cell receptor (BCR) on human peripheral and tissue resident naïve and memory B cells (**Figure 2**). CD40 signals also support the differentiation of B cells into antibody-secreting plasma cells and, in combination with additional signals, drive switching to various antibody isotypes (18-21). CD40 signals have been implicated in the development of germinal centres (**Figure 2**, GCs) (22). Indeed, lymph nodes from patients with X-linked hyper IgM syndrome

(HIGM), arising due to genetic defects in CD40L, have normal primary follicles but GCs are largely absent (23). Interestingly patients with HIGM, despite a high incidence of infections due to impairment of the immune system, are prone to autoimmune manifestations, especially hematologic abnormalities, arthritis, and inflammatory bowel disease. The mechanisms by which HIGM is associated with autoimmunity are not completely elucidated. A defective development of regulatory T cells as well as an impaired peripheral B-cell tolerance checkpoint may be important (24). Conversely, the CD40 pathway contributes to the ‘licencing’ of dendritic cells through the upregulation of other co-stimulation molecules and cytokine production. The presentation of peptide by an ‘unlicensed’ APC to a cognate T cell receptor leads to T cell non-responsiveness or deletion, helping to maintain peripheral tolerance (25).

CD40-CD40L interactions have been shown to influence the behaviour of other immune cells including T cells (26), natural killer cells (27), dendritic cells (28) and macrophages (29) favouring their maturation, survival and effector functions (**Figure 2**). For instance, recent evidence has shown that macrophages and neutrophils from CD40L-deficient patients show a decreased oxidative burst and microbicide activity (30, 31).

CD4 help for CD8 T cell responses also involves CD40–CD40L interactions, since the ‘licensing’ of dendritic cells by CD40L-expressing CD4 T cells is required to drive CD8 responses (**Figure 2**). Additionally, stimulations by an agonist anti-CD40 Ab have been proven to be sufficient to induce efficient CD8 responses in the absence of CD4 T cell help *in vitro* and *in vivo* (32, 33).

4. CD40/CD40L signalling in autoimmunity

4.1 Systemic Lupus Erythematosus (SLE)

SLE is a systemic autoimmune disease characterized by autoantibodies to nuclear antigens and immune complex deposits in small blood vessels, affecting the skin, joints, lungs, heart, brain, and kidney (34). Several studies have shown a dysregulation of CD40/CD40L signalling in SLE.

CD40L is up-regulated on several immune cells isolated from SLE patients with active disease (35-38). B cells expressing CD40L from SLE patients spontaneously produce antibodies *in vitro* (35) and transgenic mice overexpressing CD40L develop a lupus-like disease with age, suggesting a key role of CD40L in promoting autoimmunity (39). Treatment with rituximab, an anti-CD20 mAb which depletes B cells, decreases the frequency of B cells expressing CD40 and T cells expressing CD40L, suggesting that some of the benefit of this drug in treating lupus patients may be secondary to decreased activation of the CD40 signalling pathway (40, 41). SLE patients also have elevated levels of sCD40L in serum which correlates with disease activity (42).

Altered CD40 signals have been associated with kidney involvement as confirmed by CD40 upregulation in the kidney of SLE patients which in turn leads to the production of the pro-fibrotic cytokine TGF- β , which may contribute to kidney disease (43).

SLE patients display increased apoptosis of CD34⁺ hematopoietic progenitor cells, contributing to the cytopenias often seen in lupus, and which may, in part, be driven by activation of the CD40 pathway via the upregulation of the death receptor Fas (44).

A role for the CD40 pathway in pathogenesis has been further validated in spontaneous mouse models of SLE. Treatment with anti-CD40L antibody prior to onset of symptoms reduces proteinuria, prolongs survival, ameliorates kidney disease and decreases anti-dsDNA Ab titres (45-47). Treatment commenced after the onset of moderate to severe proteinuria also ameliorates kidney disease and immune complex deposition as well as prolongs survival. Interestingly, greater benefits have been observed when combining CD40 pathway blockade with CTLA4-Ig especially on survival, anti-dsDNA Ab production, and kidney disease (47).

Multiple genome-wide linkage analyses have identified regions in humans and in mice which are associated with SLE (48). The CD40 gene lies on human chromosome 20q11.2-13.1, a region with

linkage to SLE incidence (48, 49). A missense SNP, rs11086998 G, which results in an amino acid substitution within the cytoplasmic tail of CD40 and three residues upstream of the TRAF6 binding site (CD40-P227A), has been associated with risk of SLE (50). Similarly the rs481085 SNP major allele G is increased in SLE compared to controls, while the minor allele T is associated with reduced CD40 expression and is under-represented in patients with SLE (51).

4.2 Rheumatoid arthritis (RA)

RA is a chronic inflammatory disease of synovial joints. It is characterized by accumulation of adaptive and innate immune cells within the synovium, autoantibody production [anti-citrullinated protein antibodies (ACPA)] and proliferation of resident stromal cells leading to degradation of the underlying cartilage and bone (52). GWAS studies have shown an association between the CD40 locus in both juvenile and adult arthritis (53-57). Indeed, RA patients homozygous for the risk allele rs4810485 have a third more CD40 expression on CD19⁺ B cells in peripheral blood than those homozygous for the non-risk allele (56). Studies have also identified an association between polymorphisms in the CD40 locus and increased rate of joint destruction in patients with ACPA-positive RA (58) and response to TNF inhibitor treatments (59).

From the immunological point of view, CD40 is functionally expressed on several stromal and synovial immune cell populations including fibroblasts, B cells, T cells and monocytes (60-62). Once activated, CD40 signalling stimulates immune cells to proliferate, to upregulate adhesion molecules and to secrete pro-inflammatory cytokines and chemokines (60, 63). In addition, CD40 signalling in fibroblast-like synovial cells induces the expression of RANKL, involved in the osteoclast-mediated bone resorption (64) suggesting a possible role of CD40 signalling in the onset of bone erosions. CD40 expression on immune cells is upregulated by pro-inflammatory cytokines, including IFN- γ and TNF- α (60-62).

Therapeutically, treatment with anti-CD40 blocking antibody reduced TNF- α production from RA synovial fibroblasts *in vitro*, whether in the presence or absence of other immune cells, (62).

Regarding CD40L, RA synovial and peripheral T cells and B cells express CD40L at higher levels than peripheral cells isolated from healthy controls (62) and treatment with an anti-CD40L mAb prevented or ameliorated arthritis in pre-clinical models (65). However, the drug did not reverse the disease when arthritis was already established (65) indicating a more pivotal role for blocking this pathway in the earliest phase of the disease.

By contrast, a transcriptomic study of human RA synovium taken at different stages of disease found an increased expression of CD40L related gene signature in both pre-RA (arthralgia and undifferentiated arthritis) and RA samples (early RA, and established RA) suggesting that the pathway is active in both early and established human disease (61).

CD40L expression levels were found to be positively correlated with disease activity index (DAS28), and C-reactive protein (CRP) and treatment with TNF- α inhibitors decreased the expression of CD40L, ameliorated disease activity, as well as reduced ACPA antibody production by RA peripheral blood mononuclear cells (PBMCs) (66).

Although there is some conflicting data (67) the effect of activating the CD40 pathway with agonist antibodies in animal models resulted in earlier onset and more severe disease (68), supporting the hypothesis of an adjuvant property of CD40 on the initiating pathogenic mechanisms.

Altogether, these data suggest that blocking CD40/CD40L may be crucial at the initial stages of arthritis but with more variable and unpredictable efficacy in later stages.

4.3 Sjögren's syndrome (SS)

SS is a systemic autoimmune disease with a female-to-male predominance of 9:1 that primarily involves the salivary and lacrimal glands with dry eyes and mouth as common symptoms. It can also involve other major organs, and is associated with an increased risk of developing B cell lymphomas

(69). Around 25% of SS patients develop GCs within salivary gland (SG) lymphocytic foci and which are associated with more severe disease (70). The very definition of SS as a disease characterised by ectopic lymphocytic structures associated with B cell hyperactivity, together with organisation of the lymphocytic structures and germinal centre formation, makes the CD40 pathway an attractive target for this population. Indeed, the presence of both CD40 and CD40L has been confirmed within SS lymphocytic foci (42).

Constitutive expression of CD40, CD40L, and other co-stimulatory markers (i.e. CD80 and CD86) has also been described on SG ductal epithelial and endothelial cells in SS thus suggesting the potential to activate effector immune cells (71-73). The expression of CD40 on SG is regulated by pro-inflammatory stimuli such as interferon-gamma (IFN- γ) and IL-1 β . In turn CD40 enhances the surface expression of the adhesion molecule intercellular adhesion molecule-1 (ICAM-1)/CD54 on SG epithelial cells (73) promoting leukocyte recruitment and the establishment of inflammation. Evidence suggests a possible role of CD40 signalling pathway in promoting SS epithelial cell apoptosis but this mechanism needs to be further elucidated (74). Increased levels of sCD40L have also been reported in SS (42).

Therapeutic treatment with systemic anti-CD40L antibody reduced sialadenitis, inhibited ectopic lymphoid structure formation and autoantibody production, as well as decreased the frequency of SG antibody-secreting cells in mouse models (75)

4.4 Systemic sclerosis (SSc)

Systemic sclerosis is an immune-mediated rheumatic disease that is characterised by fibrosis of the skin and internal organs as well as vasculopathy and systemic complications such as scleroderma renal crisis, pulmonary arterial hypertension, digital ulceration, and gastro-oesophageal reflux. Treatments for specific complications have emerged and a growing evidence base supports the use of immune suppression for the treatment of skin and lung fibrosis (76). Increased expression of

CD40L has been reported on activated CD4⁺ T lymphocytes in systemic sclerosis and blocking anti-CD40 antibody reduced the expression of the co-stimulatory molecule, CD80, in SSc activated monocytes (77).

Soluble CD40L concentration is higher in systemic sclerosis than in controls and correlates with clinical and laboratory features (78, 79).

The blockade of CD40/CD40L interactions by anti-CD40L monoclonal antibody significantly reduced cutaneous fibrosis, anti-topoisomerase I autoantibody and normalised B lymphocyte activation, as evidenced by reduced levels of immunoglobulin, in a mouse model (TSK/+ mice) (79).

CD40 mRNA was found to be constitutively expressed in both SSc and normal human fibroblasts but CD40 protein expression was higher on SSc fibroblasts. In addition, ligation of CD40 by recombinant human CD40L resulted in increased production of IL-6, IL-8, and monocyte chemoattractant protein-1 in SSc but not normal fibroblasts in a dose-dependent manner. The co-stimulatory molecule CD80, was also induced on SSc fibroblasts by CD40 ligation (80). Although CD40/CD40L may to be implicated in the pathogenesis of SSs, there have been no clinical trials to date.

5. Clinical trials

Given the key role that the CD40 pathway may have in the pathogenesis of autoimmune disease, it has long been considered an attractive therapeutic target. Several clinical trials targeting the CD40/CD40L signalling pathway have been completed or are ongoing. Early trials using anti-CD40L mAb were discontinued due to thrombotic side effects. New molecules targeting CD40 or novel engineering approaches to CD40L have been identified to minimize the collateral effects (**Table 1**).

5.1 Targeting CD40/CD40L in SLE

Ruplizumab (Hu5c8, BG9588), a CD40L-specific humanized IgG1 mAb, was one of the first molecules generated to target the CD40 pathway. Treatment with ruplizumab resulted in a significant decrease in anti-dsDNA antibody level and haematuria as well as a significant increase in complement C3 concentration in an open label Phase 2 study in patients with proliferative lupus nephritis (81). CD38^{high} B cells and IgM and IgG anti-dsDNA secreting plasma cells were decreased (82). In addition, the reduction of anti-dsDNA antibodies was also associated with improvement in the SLE disease activity index (SLEDAI) (82).

However, despite this promising evidence of clinical effect, further development of hu5c8 was discontinued because of treatment-emergent cardiovascular thrombotic events (TEs) (83). Numerous TEs including pulmonary vascular thrombi and vasculopathy were subsequently found after administration of hu5c8 in Rhesus monkeys (4, 83).

Toralizumab (IDEC-131), a humanized anti-CD40L mAb, was evaluated in a number of early clinical trials, including Phase 1 and Phase 2 studies in SLE (84, 85). Even at higher doses of 0.05-15.0 mg/kg toralizumab was demonstrated to be safe and well tolerated with no adverse event in both studies (85). Unfortunately, the efficacy of IDEC-131 compared to placebo was not demonstrated in SLE patients with mild to moderate disease activity (85). Similar to ruplizumab, further development of this agent was stopped due to increased thrombosis in other trials (86).

The TEs of ruplizumab and toralizumab appear to require a functional Fc portion (87, 88). Activated platelets express CD40L and *in vitro* analyses have shown that immune complexes consisting of sCD40L and an anti-CD40L monoclonal antibody can trigger platelet aggregation (87, 88). Inhibition of platelet Fc receptors can block this antibody mediated platelet aggregation.

One approach to reducing the TE risk is to remove the Fc portion, and an anti-CD40L Fab' antibody fragment conjugated to polyethylene glycol (PEG), **dapirolizumab pegol (CDP7657)**, was subsequently developed and showed no evidence of pro-thrombotic effects in pre-clinical studies (4). Dapirolizumab was found to inhibit humoral immune responses in monkeys without thromboembolic complications (4). In a Phase I study, dapirolizumab pegol showed no serious treatment-emergent

adverse events. Exploratory analyses of patients with high disease activity at baseline indicated the potential for clinical improvement and reduction in anti-dsDNA antibodies, with down-modulation of peripheral blood plasma cell and B cell genes, particularly those related to immunoglobulins. (89). A Phase 2 dose-ranging study (NCT02804763) in adults with moderately-to-severely active SLE failed to achieve the primary end point of establishing a dose response at week 24 ($P = 0.06$), although numerical improvements in clinical endpoints were observed with dapirolizumab pegol in comparison to placebo, as were improvements in complement levels and reductions in anti-dsDNA antibodies. Four TEs were observed but three of these were in the placebo-treated arm (90).

5.2 Targeting CD40/CD40L in RA

VIB4920 (MEDI4920) is a novel CD40L binding protein comprised of two Tn3 proteins derived from fibronectin type III domain, fused to human serum albumin. VIB4920 targets CD40L but does not possess an Fc domain and therefore is unlikely to induce thrombotic effects. A phase 1, randomized, blinded, placebo-controlled, single-ascending dose study confirmed safety and tolerability of MEDI4920 in healthy adults (91). In a Phase 1b, randomized, double-blind, placebo-controlled, multiple-ascending dose study, VIB4920 significantly reduced RA disease activity, as measured by DAS28-CRP, at day 85 as well as additional clinical parameters including tender/swollen joint counts, CRP, and patient and physician global assessments (3, 86). Improvements in clinical activity in this trial were accompanied by reductions in rheumatoid factor of approximately 50% at the higher doses and in other circulating biomarkers associated with pathways that drive RA disease activity (86). Lower doses of VIB4920 were associated with a high prevalence of anti-drug antibodies. Although these were not detected at the highest dose, consistent with effective suppression of the CD40 pathway, the implications of this finding for future clinical utility remain to be determined (86). A Phase 2 study to evaluate the efficacy, safety, and pharmacokinetics is currently ongoing (NCT04163991).

An alternative approach to reducing TE risk is to utilise blocking antibodies to the more widely expressed CD40. **BI-655064** is a humanized antagonistic IgG1 mAb that binds CD40 and is modified to reduce Fc mediated effector functions including cytotoxicity and platelet activation (92). BI-655064 demonstrated safety and was well tolerated in a cohort of healthy volunteers (92). In a double-blind, randomized Phase 2a trial (NCT01751776), patients with RA received either weekly BI-655064 (120 mg) or placebo as add-on therapy to methotrexate (93). The primary endpoint of an ACR20 response at week 12 was seen in 68.2% in the active arm (n=44) compared to 45.5% in placebo (p=0.06). Interestingly, a significant difference in efficacy between active drug and placebo was seen in those patients with a disease duration less than 2.5 years (p=0.009). Treatment with BI-655064 also reduced the frequency of activated B cells, specifically class switched (CD19⁺IgD⁻CD27⁺CD95⁺), pre-switched (CD19⁺IgD⁺CD27⁺CD95⁺) and double-negative (CD19⁺IgD⁻CD27⁻CD95⁺) cells. Reductions in IgG and IgA rheumatoid factor, total IgG and IgM, IL6, metalloproteinase (MMP)-3 and RANKL levels were also observed. Amongst T allele carriers of the CD40 SNP rs4810485, ACR50 responses were greater in the active treatment arm compared with placebo (54.5% vs 16.7%; p=0.04) (93). Pharmacokinetic variability was observed with steady state only being reached towards the end of the 12 week intervention period, raising the possibility of inadequate dosing. A Phase 2 trial to evaluate the long term efficacy and safety of different doses of BI 655064 versus placebo as add-on therapy to Standard of Care (SOC) during maintenance therapy for lupus nephritis is currently ongoing (NCT03385564).

5.3 Targeting CD40/CD40L in SS

Iscalimab (CFZ533), is a human non-agonistic anti-CD40 monoclonal IgG1 blocking antibody, containing a modified Fc domain that renders it unable to mediate Fcγ-dependent effector functions and is therefore non-depleting (94). Iscalimab reduces humoral responses as well as GC formation in monkeys following kidney transplantation (95). In a Phase 1/2 study in de novo renal transplant,

iscalimab in combination with mycophenolate mofetil (MMF) and corticosteroids (CS) demonstrated comparable efficacy to tacrolimus, MMF and CS in terms of graft rejection, graft loss or death. However, compared to tacrolimus, iscalimab demonstrated improved renal function with fewer serious adverse events and infectious complications (96) as well as normal histology in the allografts (97)

Iscalimab also demonstrated efficacy in a Phase 2a randomized controlled trial (NCT02291029) in patients with primary SS (98). Forty-four SS patients were enrolled: 8 patients received 3 mg/kg s.c. iscalimab and 4 placebo in cohort 1 and 21 received 10 mg/kg i.v. iscalimab and 11 placebo in cohort 2. Iscalimab was safe and well tolerated. In cohort 1, high target-mediated drug disposition (TMDD) was observed with correspondingly low plasma PK values and no evidence of a treatment difference between iscalimab and placebo on EULAR Sjögren's Syndrome Disease Activity Index (ESSDAI) scores [baseline-adjusted reduction from placebo at week 12 was only 0.41 points (95% CI -2.89 to 3.70)]. However in cohort 2, there was a statistically significant and clinically meaningful improvement in clinical disease activity as measured by improvement in ESSDAI in patients receiving iscalimab with a baseline-adjusted mean reduction in ESSDAI at week 12 of 5.21 points (95% CI 0.96 to 9.46; one-sided $p=0.009$). The high-dose iscalimab arm showed statistically significant improvements in physician's global assessment compared with placebo, and other measures such as the EULAR syndrome patient reported index (ESSPRI), multidimensional fatigue inventory (MFI), and patient's global assessment showed trends to improvement. Data suggest that the chemokine CXCL13 may be a biomarker of germinal centre formation and, in keeping with the histological definition of SS as a focal lymphocytic sialadenitis, we have previously found that serum levels of CXCL13 correlate with the extent of histological SG inflammation (99). It is therefore of interest that iscalimab 10 mg/kg was associated with a marked reduction in CXCL13 levels. The TMDD observed with subcutaneous dosing, which may also be relevant to other anti-CD40 mAbs, can be overcome with either i.v. or s.c loading with corresponding evidence of clinical efficacy and reduction in symptoms and CXCL13, albeit from an open-label cohort (100).

Any beneficial effect of anti-CD40–CD40L blockade on lymphoma risk in SS remains to be seen.

6. Conclusions

Progressive disability, systemic complications and early death are still a reality leading to socioeconomic costs and unmet needs for several rheumatic autoimmune disorders. Indeed, some patients still fail to respond to current conventional and biologic disease modifying therapies.

The CD40-CD40L axis modulates many immune responses, and alterations in this signalling pathway have been reported in several autoimmune rheumatic disorders.

Early attempts to modulate this pathway through CD40L blockade were terminated due to thromboembolic events. Recent developments have focussed on targeting CD40 or on modifying anti-CD40L molecules to prevent Fc receptor binding. Several molecules blocking CD40 or CD40L have been tested in humans, with encouraging data emerging and new clinical trials ongoing in multiple autoimmune diseases. More data is required to be certain that newer approaches to CD40L blockade do not increase risk for thromboembolic events. Aside from this safety consideration, in making the choice between targeting CD40 or CD40L, one key distinction is the more widespread nature of CD40 expression and susceptibility to TMDD, necessitating higher dosing, with implications for cost, and more extensive receptor occupancy.

The use of monoclonal antibodies in autoimmune disease is often effective through tuning down excessive target signalling, rather than by completely eliminating it. However, given the importance and multi-faceted roles of CD40-CD40L interactions, it is plausible that persistent pathway blockade may be associated with infectious adverse events. Effective CD40 pathway blockade would also be anticipated to impair vaccination responses. Given the potential roles of the CD40 pathway in facilitating immune responses to cancer, it is unknown if persistent pathway blockade might increase cancer risk. Although the safety data from recent trials has been encouraging, ongoing study is required as programmes progress.

It is still not clear at which stage of disease this therapy will be most efficacious. Many pre-clinical studies reviewed here suggest greater efficacy in the very early stages of autoimmunity. Indeed, short-term deep pathway blockade in this setting might hypothetically allow efficacy with a lower overall risk infection compared with long-term therapy. However, the results of iscalimab in established SS are encouraging, and may reflect disruption of pathogenically important B-T cell interactions within glandular lymphocytic foci/ectopic germinal centres. Results require confirmation in larger studies. It remains to be determined if stratification by pathotype in RA or by CD40 genotype may identify patients more likely to benefit. Given the important role of CD40/CD40L at multiple levels within the immune system, further studies should also determine the mechanisms of action most relevant to human autoimmune disease modification with this promising therapeutic approach.

Contributors

All authors contributed to the conception and content of the Review. All authors critically revised and edited the first draft and approved the final version.

Acknowledgements

VP and BF have received support from the NIHR Birmingham Biomedical Research Centre. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

Conflict of Interest

BF is paid instructor/consultant for: Novartis, Roche, BMS and Servier.

Search strategy and selection criteria

References for this Review were identified through searches of PubMed and ClinicalTrials.gov with the search terms “CD40 and CD40L in SLE, RA, SS, SSc, psoriasis” from 1991 until 2019. We

largely selected publications from the past 5 years, but did not exclude commonly referenced and highly regarded older publications. We also searched the reference lists of articles identified by this search strategy and selected those we judged relevant.

Figure 1 – Co-stimulatory pathways relevant to current drug development in rheumatic autoimmune disorders

Only those immune co-stimulation targets referred to in the review are illustrated. Antigen-specific interaction between TCR and MHC expressed on APC provides the first signal for adequate T cell activation. Different stimulatory (green) or inhibitory (red) co-stimulatory signals may influence T cell-dependent immune responses. Dysregulation of these pathways is reported in several autoimmune diseases and novel pharmacological agents (e.g. Iscalimab, AMG 557) targeting molecules involved in the co-stimulatory machinery are being developed.

Abbreviations: APC, antigen presenting cell; CTLA4, cytotoxic T-lymphocyte-associated protein 4; ICOS, inducible co-stimulator; MHC, major histocompatibility complex; PD-1, programmed cell death-1; TCR, T cell receptor

Figure 2 - CD40/CD40L in autoimmunity

CD40L expressed on T cells interacts with CD40 on APCs and regulates different effector functions. CD40–CD40L interaction regulates T-cell co-stimulation and differentiation of conventional and regulatory T cells as well as contributes to the activation of macrophages, dendritic cells (DCs) and B cells. On B cells, the interaction between CD40 and CD40L induces their differentiation to memory B cells and antibody-producing plasma cells. All these events may contribute to the formation of germinal centres (GCs) and in the development of autoimmunity.

References

1. Toubi E, Shoenfeld Y. The role of CD40-CD154 interactions in autoimmunity and the benefit of disrupting this pathway. *Autoimmunity*. 2004;37(6-7):457-64.
2. Elgueta R, Benson MJ, de Vries VC, Wasiuk A, Guo Y, Noelle RJ. Molecular mechanism and function of CD40/CD40L engagement in the immune system. *Immunol Rev*. 2009;229(1):152-72.
3. Karnell JL, Rieder SA, Ettinger R, Kolbeck R. Targeting the CD40-CD40L pathway in autoimmune diseases: Humoral immunity and beyond. *Adv Drug Deliv Rev*. 2019;141:92-103.
4. Shock A, Burkly L, Wakefield I, Peters C, Garber E, Ferrant J, et al. CDP7657, an anti-CD40L antibody lacking an Fc domain, inhibits CD40L-dependent immune responses without thrombotic complications: an in vivo study. *Arthritis Res Ther*. 2015;17:234.
5. Henn V, Slupsky JR, Grafe M, Anagnostopoulos I, Forster R, Muller-Berghaus G, et al. CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature*. 1998;391(6667):591-4.
6. Matthies KM, Newman JL, Hodzic A, Wingett DG. Differential regulation of soluble and membrane CD40L proteins in T cells. *Cell Immunol*. 2006;241(1):47-58.
7. Pietravalle F, Lecoanet-Henchoz S, Blasey H, Aubry JP, Elson G, Edgerton MD, et al. Human native soluble CD40L is a biologically active trimer, processed inside microsomes. *J Biol Chem*. 1996;271(11):5965-7.
8. Schiff M. Abatacept treatment for rheumatoid arthritis. *Rheumatology (Oxford)*. 2011;50(3):437-49.
9. Emery P, Burmester GR, Bykerk VP, Combe BG, Furst DE, Barre E, et al. Evaluating drug-free remission with abatacept in early rheumatoid arthritis: results from the phase 3b, multicentre, randomised, active-controlled AVERT study of 24 months, with a 12-month, double-blind treatment period. *Ann Rheum Dis*. 2015;74(1):19-26.
10. Al-Laith M, Jasenecova M, Abraham S, Bosworth A, Bruce IN, Buckley CD, et al. Arthritis prevention in the pre-clinical phase of RA with abatacept (the APIPPRA study): a multi-centre, randomised, double-blind, parallel-group, placebo-controlled clinical trial protocol. *Trials*. 2019;20(1):429.
11. Chen L, Flies DB. Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat Rev Immunol*. 2013;13(4):227-42.
12. McAdam AJ, Chang TT, Lumelsky AE, Greenfield EA, Boussiotis VA, Duke-Cohan JS, et al. Mouse inducible costimulatory molecule (ICOS) expression is enhanced by CD28 costimulation and regulates differentiation of CD4+ T cells. *J Immunol*. 2000;165(9):5035-40.
13. Warnatz K, Bossaller L, Salzer U, Skrabl-Baumgartner A, Schwinger W, van der Burg M, et al. Human ICOS deficiency abrogates the germinal center reaction and provides a monogenic model for common variable immunodeficiency. *Blood*. 2006;107(8):3045-52.
14. Weber JP, Fuhrmann F, Feist RK, Lahmann A, Al Baz MS, Gentz LJ, et al. ICOS maintains the T follicular helper cell phenotype by down-regulating Kruppel-like factor 2. *J Exp Med*. 2015;212(2):217-33.
15. Caux C, Massacrier C, Vanbervliet B, Dubois B, Van Kooten C, Durand I, et al. Activation of human dendritic cells through CD40 cross-linking. *J Exp Med*. 1994;180(4):1263-72.
16. Pinchuk LM, Polacino PS, Agy MB, Klaus SJ, Clark EA. The role of CD40 and CD80 accessory cell molecules in dendritic cell-dependent HIV-1 infection. *Immunity*. 1994;1(4):317-25.
17. Bishop GA, Moore CR, Xie P, Stunz LL, Kraus ZJ. TRAF proteins in CD40 signaling. *Adv Exp Med Biol*. 2007;597:131-51.

18. Ettinger R, Sims GP, Fairhurst AM, Robbins R, da Silva YS, Spolski R, et al. IL-21 induces differentiation of human naive and memory B cells into antibody-secreting plasma cells. *J Immunol.* 2005;175(12):7867-79.
19. Rousset F, Garcia E, Banchereau J. Cytokine-induced proliferation and immunoglobulin production of human B lymphocytes triggered through their CD40 antigen. *J Exp Med.* 1991;173(3):705-10.
20. Spriggs MK, Armitage RJ, Strockbine L, Clifford KN, Macduff BM, Sato TA, et al. Recombinant human CD40 ligand stimulates B cell proliferation and immunoglobulin E secretion. *J Exp Med.* 1992;176(6):1543-50.
21. Defrance T, Vanbervliet B, Durand I, Briolay J, Banchereau J. Proliferation and differentiation of human CD5+ and CD5- B cell subsets activated through their antigen receptors or CD40 antigens. *Eur J Immunol.* 1992;22(11):2831-9.
22. Kawabe T, Naka T, Yoshida K, Tanaka T, Fujiwara H, Suematsu S, et al. The immune responses in CD40-deficient mice: impaired immunoglobulin class switching and germinal center formation. *Immunity.* 1994;1(3):167-78.
23. Facchetti F, Appiani C, Salvi L, Levy J, Notarangelo LD. Immunohistologic analysis of ineffective CD40-CD40 ligand interaction in lymphoid tissues from patients with X-linked immunodeficiency with hyper-IgM. Abortive germinal center cell reaction and severe depletion of follicular dendritic cells. *J Immunol.* 1995;154(12):6624-33.
24. Jesus AA, Duarte AJ, Oliveira JB. Autoimmunity in hyper-IgM syndrome. *J Clin Immunol.* 2008;28 Suppl 1:S62-6.
25. Quezada SA, Jarvinen LZ, Lind EF, Noelle RJ. CD40/CD154 interactions at the interface of tolerance and immunity. *Annu Rev Immunol.* 2004;22:307-28.
26. Macatonia SE, Hosken NA, Litton M, Vieira P, Hsieh CS, Culpepper JA, et al. Dendritic cells produce IL-12 and direct the development of Th1 cells from naive CD4+ T cells. *J Immunol.* 1995;154(10):5071-9.
27. Carbone E, Ruggiero G, Terrazzano G, Palomba C, Manzo C, Fontana S, et al. A new mechanism of NK cell cytotoxicity activation: the CD40-CD40 ligand interaction. *J Exp Med.* 1997;185(12):2053-60.
28. Ma DY, Clark EA. The role of CD40 and CD154/CD40L in dendritic cells. *Semin Immunol.* 2009;21(5):265-72.
29. Suttles J, Stout RD. Macrophage CD40 signaling: a pivotal regulator of disease protection and pathogenesis. *Semin Immunol.* 2009;21(5):257-64.
30. Cabral-Marques O, Ramos RN, Schimke LF, Khan TA, Amaral EP, Barbosa Bomfim CC, et al. Human CD40 ligand deficiency dysregulates the macrophage transcriptome causing functional defects that are improved by exogenous IFN-gamma. *J Allergy Clin Immunol.* 2017;139(3):900-12.e7.
31. Cabral-Marques O, Franca TT, Al-Sbiei A, Schimke LF, Khan TA, Feriotti C, et al. CD40 ligand deficiency causes functional defects of peripheral neutrophils that are improved by exogenous IFN-gamma. *J Allergy Clin Immunol.* 2018;142(5):1571-88.e9.
32. Meunier S, Rapetti L, Beziaud L, Pontoux C, Legrand A, Tanchot C. Synergistic CD40 signaling on APCs and CD8 T cells drives efficient CD8 response and memory differentiation. *J Leukoc Biol.* 2012;91(6):859-69.
33. Bennett SR, Carbone FR, Karamalis F, Flavell RA, Miller JF, Heath WR. Help for cytotoxic-T-cell responses is mediated by CD40 signalling. *Nature.* 1998;393(6684):478-80.
34. Lisnevskaja L, Murphy G, Isenberg D. Systemic lupus erythematosus. *Lancet.* 2014;384(9957):1878-88.
35. Grammer AC, Slota R, Fischer R, Gur H, Girschick H, Yarboro C, et al. Abnormal germinal center reactions in systemic lupus erythematosus demonstrated by blockade of CD154-CD40 interactions. *J Clin Invest.* 2003;112(10):1506-20.

36. Katsiari CG, Liossis SN, Souliotis VL, Dimopoulos AM, Manoussakis MN, Sfikakis PP. Aberrant expression of the costimulatory molecule CD40 ligand on monocytes from patients with systemic lupus erythematosus. *Clin Immunol.* 2002;103(1):54-62.
37. Desai-Mehta A, Lu L, Ramsey-Goldman R, Datta SK. Hyperexpression of CD40 ligand by B and T cells in human lupus and its role in pathogenic autoantibody production. *J Clin Invest.* 1996;97(9):2063-73.
38. Koshy M, Berger D, Crow MK. Increased expression of CD40 ligand on systemic lupus erythematosus lymphocytes. *J Clin Invest.* 1996;98(3):826-37.
39. Higuchi T, Aiba Y, Nomura T, Matsuda J, Mochida K, Suzuki M, et al. Cutting Edge: Ectopic expression of CD40 ligand on B cells induces lupus-like autoimmune disease. *J Immunol.* 2002;168(1):9-12.
40. Tokunaga M, Saito K, Kawabata D, Imura Y, Fujii T, Nakayamada S, et al. Efficacy of rituximab (anti-CD20) for refractory systemic lupus erythematosus involving the central nervous system. *Ann Rheum Dis.* 2007;66(4):470-5.
41. Tamimoto Y, Horiuchi T, Tsukamoto H, Otsuka J, Mitoma H, Kimoto Y, et al. A dose-escalation study of rituximab for treatment of systemic lupus erythematosus and Evans' syndrome: immunological analysis of B cells, T cells and cytokines. *Rheumatology (Oxford).* 2008;47(6):821-7.
42. Goules A, Tzioufas AG, Manousakis MN, Kirou KA, Crow MK, Routsias JG. Elevated levels of soluble CD40 ligand (sCD40L) in serum of patients with systemic autoimmune diseases. *J Autoimmun.* 2006;26(3):165-71.
43. Delmas Y, Viallard JF, Solanilla A, Villeneuve J, Pasquet JM, Belloc F, et al. Activation of mesangial cells by platelets in systemic lupus erythematosus via a CD154-dependent induction of CD40. *Kidney Int.* 2005;68(5):2068-78.
44. Pyrovolaki K, Mavroudi I, Sidiropoulos P, Eliopoulos AG, Boumpas DT, Papadaki HA. Increased expression of CD40 on bone marrow CD34+ hematopoietic progenitor cells in patients with systemic lupus erythematosus: contribution to Fas-mediated apoptosis. *Arthritis Rheum.* 2009;60(2):543-52.
45. Wang X, Huang W, Schiffer LE, Mihara M, Akkerman A, Hiromatsu K, et al. Effects of anti-CD154 treatment on B cells in murine systemic lupus erythematosus. *Arthritis Rheum.* 2003;48(2):495-506.
46. Quezada SA, Eckert M, Adeyi OA, Schned AR, Noelle RJ, Burns CM. Distinct mechanisms of action of anti-CD154 in early versus late treatment of murine lupus nephritis. *Arthritis Rheum.* 2003;48(9):2541-54.
47. Wang X, Huang W, Mihara M, Sinha J, Davidson A. Mechanism of action of combined short-term CTLA4Ig and anti-CD40 ligand in murine systemic lupus erythematosus. *J Immunol.* 2002;168(4):2046-53.
48. Wakeland EK, Liu K, Graham RR, Behrens TW. Delineating the genetic basis of systemic lupus erythematosus. *Immunity.* 2001;15(3):397-408.
49. Gaffney PM, Langefeld CD, Graham RR, Ortmann WA, Williams AH, Rodine PR, et al. Fine-mapping chromosome 20 in 230 systemic lupus erythematosus sib pair and multiplex families: evidence for genetic epistasis with chromosome 16q12. *Am J Hum Genet.* 2006;78(5):747-58.
50. Peters AL, Plenge RM, Graham RR, Altshuler DM, Moser KL, Gaffney PM, et al. A novel polymorphism of the human CD40 receptor with enhanced function. *Blood.* 2008;112(5):1863-71.
51. Vazgiourakis VM, Zervou MI, Choulaki C, Bertsiyas G, Melissourgaki M, Yilmaz N, et al. A common SNP in the CD40 region is associated with systemic lupus erythematosus and correlates with altered CD40 expression: implications for the pathogenesis. *Ann Rheum Dis.* 2011;70(12):2184-90.

52. McInnes IB, Schett G. The pathogenesis of rheumatoid arthritis. *N Engl J Med*. 2011;365(23):2205-19.
53. Thompson SD, Moroldo MB, Guyer L, Ryan M, Tombragel EM, Shear ES, et al. A genome-wide scan for juvenile rheumatoid arthritis in affected sibpair families provides evidence of linkage. *Arthritis Rheum*. 2004;50(9):2920-30.
54. Orozco G, Eyre S, Hinks A, Ke X, Wilson AG, Bax DE, et al. Association of CD40 with rheumatoid arthritis confirmed in a large UK case-control study. *Ann Rheum Dis*. 2010;69(5):813-6.
55. Raychaudhuri S, Remmers EF, Lee AT, Hackett R, Guiducci C, Burt NP, et al. Common variants at CD40 and other loci confer risk of rheumatoid arthritis. *Nat Genet*. 2008;40(10):1216-23.
56. Li G, Diogo D, Wu D, Spoonamore J, Dancik V, Franke L, et al. Human genetics in rheumatoid arthritis guides a high-throughput drug screen of the CD40 signaling pathway. *PLoS Genet*. 2013;9(5):e1003487.
57. Fang H, De Wolf H, Knezevic B, Burnham KL, Osgood J, Sanniti A, et al. A genetics-led approach defines the drug target landscape of 30 immune-related traits. *Nat Genet*. 2019;51(7):1082-91.
58. van der Linden MP, Feitsma AL, le Cessie S, Kern M, Olsson LM, Raychaudhuri S, et al. Association of a single-nucleotide polymorphism in CD40 with the rate of joint destruction in rheumatoid arthritis. *Arthritis Rheum*. 2009;60(8):2242-7.
59. Spiliopoulou A, Colombo M, Plant D, Nair N, Cui J, Coenen MJ, et al. Association of response to TNF inhibitors in rheumatoid arthritis with quantitative trait loci for CD40 and CD39. *Ann Rheum Dis*. 2019;78(8):1055-61.
60. Rissoan MC, Van Kooten C, Chomarat P, Galibert L, Durand I, Thivolet-Bejui F, et al. The functional CD40 antigen of fibroblasts may contribute to the proliferation of rheumatoid synovium. *Clin Exp Immunol*. 1996;106(3):481-90.
61. Guo Y, Walsh AM, Fearon U, Smith MD, Wechalekar MD, Yin X, et al. CD40L-Dependent Pathway Is Active at Various Stages of Rheumatoid Arthritis Disease Progression. *J Immunol*. 2017;198(11):4490-501.
62. Liu MF, Chao SC, Wang CR, Lei HY. Expression of CD40 and CD40 ligand among cell populations within rheumatoid synovial compartment. *Autoimmunity*. 2001;34(2):107-13.
63. Harigai M, Hara M, Kawamoto M, Kawaguchi Y, Sugiura T, Tanaka M, et al. Amplification of the synovial inflammatory response through activation of mitogen-activated protein kinases and nuclear factor kappaB using ligation of CD40 on CD14+ synovial cells from patients with rheumatoid arthritis. *Arthritis Rheum*. 2004;50(7):2167-77.
64. Lee BO, Haynes L, Eaton SM, Swain SL, Randall TD. The biological outcome of CD40 signaling is dependent on the duration of CD40 ligand expression: reciprocal regulation by interleukin (IL)-4 and IL-12. *J Exp Med*. 2002;196(5):693-704.
65. Durie FH, Fava RA, Foy TM, Aruffo A, Ledbetter JA, Noelle RJ. Prevention of collagen-induced arthritis with an antibody to gp39, the ligand for CD40. *Science*. 1993;261(5126):1328-30.
66. Tung CH, Lu MC, Lai NS, Wu SF. Tumor necrosis factor-alpha blockade treatment decreased CD154 (CD40-ligand) expression in rheumatoid arthritis. *PLoS One*. 2017;12(8):e0183726.
67. Mauri C, Mars LT, Londei M. Therapeutic activity of agonistic monoclonal antibodies against CD40 in a chronic autoimmune inflammatory process. *Nat Med*. 2000;6(6):673-9.
68. Tellander AC, Michaelsson E, Brunmark C, Andersson M. Potent adjuvant effect by anti-CD40 in collagen-induced arthritis. Enhanced disease is accompanied by increased production of collagen type-II reactive IgG2a and IFN-gamma. *J Autoimmun*. 2000;14(4):295-302.

69. Brito-Zeron P, Baldini C, Bootsma H, Bowman SJ, Jonsson R, Mariette X, et al. Sjogren syndrome. *Nat Rev Dis Primers*. 2016;2:16047.
70. Risselada AP, Looije MF, Kruize AA, Bijlsma JW, van Roon JA. The role of ectopic germinal centers in the immunopathology of primary Sjogren's syndrome: a systematic review. *Semin Arthritis Rheum*. 2013;42(4):368-76.
71. Tsunawaki S, Nakamura S, Ohyama Y, Sasaki M, Ikebe-Hiroki A, Hiraki A, et al. Possible function of salivary gland epithelial cells as nonprofessional antigen-presenting cells in the development of Sjogren's syndrome. *J Rheumatol*. 2002;29(9):1884-96.
72. Nakamura H, Kawakami A, Tominaga M, Migita K, Kawabe Y, Nakamura T, et al. Expression of CD40/CD40 ligand and Bcl-2 family proteins in labial salivary glands of patients with Sjogren's syndrome. *Lab Invest*. 1999;79(3):261-9.
73. Dimitriou ID, Kapsogeorgou EK, Moutsopoulos HM, Manoussakis MN. CD40 on salivary gland epithelial cells: high constitutive expression by cultured cells from Sjogren's syndrome patients indicating their intrinsic activation. *Clin Exp Immunol*. 2002;127(2):386-92.
74. Ping L, Ogawa N, Sugai S. Novel role of CD40 in Fas-dependent apoptosis of cultured salivary epithelial cells from patients with Sjogren's syndrome. *Arthritis Rheum*. 2005;52(2):573-81.
75. Wieczorek G, Bigaud M, Pfister S, Ceci M, McMichael K, Afatsawo C, et al. Blockade of CD40-CD154 pathway interactions suppresses ectopic lymphoid structures and inhibits pathology in the NOD/ShiLtJ mouse model of Sjogren's syndrome. *Ann Rheum Dis*. 2019;78(7):974-8.
76. Denton CP, Khanna D. Systemic sclerosis. *Lancet*. 2017;390(10103):1685-99.
77. Valentini G, Romano MF, Naclerio C, Bisogni R, Lamberti A, Turco MC, et al. Increased expression of CD40 ligand in activated CD4+ T lymphocytes of systemic sclerosis patients. *J Autoimmun*. 2000;15(1):61-6.
78. Allanore Y, Borderie D, Meune C, Lemarechal H, Weber S, Ekindjian OG, et al. Increased plasma soluble CD40 ligand concentrations in systemic sclerosis and association with pulmonary arterial hypertension and digital ulcers. *Ann Rheum Dis*. 2005;64(3):481-3.
79. Komura K, Sato S, Hasegawa M, Fujimoto M, Takehara K. Elevated circulating CD40L concentrations in patients with systemic sclerosis. *J Rheumatol*. 2004;31(3):514-9.
80. Fukasawa C, Kawaguchi Y, Harigai M, Sugiura T, Takagi K, Kawamoto M, et al. Increased CD40 expression in skin fibroblasts from patients with systemic sclerosis (SSc): role of CD40-CD154 in the phenotype of SSc fibroblasts. *Eur J Immunol*. 2003;33(10):2792-800.
81. Boumpas DT, Furie R, Manzi S, Illei GG, Wallace DJ, Balow JE, et al. A short course of BG9588 (anti-CD40 ligand antibody) improves serologic activity and decreases hematuria in patients with proliferative lupus glomerulonephritis. *Arthritis Rheum*. 2003;48(3):719-27.
82. Huang W, Sinha J, Newman J, Reddy B, Budhai L, Furie R, et al. The effect of anti-CD40 ligand antibody on B cells in human systemic lupus erythematosus. *Arthritis Rheum*. 2002;46(6):1554-62.
83. Kawai T, Andrews D, Colvin RB, Sachs DH, Cosimi AB. Thromboembolic complications after treatment with monoclonal antibody against CD40 ligand. *Nat Med*. 2000;6(2):114.
84. Davis JC, Jr., Totoritis MC, Rosenberg J, Sklenar TA, Wofsy D. Phase I clinical trial of a monoclonal antibody against CD40-ligand (IDEC-131) in patients with systemic lupus erythematosus. *J Rheumatol*. 2001;28(1):95-101.
85. Kalunian KC, Davis JC, Jr., Merrill JT, Totoritis MC, Wofsy D. Treatment of systemic lupus erythematosus by inhibition of T cell costimulation with anti-CD154: a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum*. 2002;46(12):3251-8.
86. Karnell JL, Albulescu M, Drabic S, Wang L, Moate R, Baca M, et al. A CD40L-targeting protein reduces autoantibodies and improves disease activity in patients with autoimmunity. *Sci Transl Med*. 2019;11(489).

87. Robles-Carrillo L, Meyer T, Hatfield M, Desai H, Davila M, Langer F, et al. Anti-CD40L immune complexes potently activate platelets in vitro and cause thrombosis in FCGR2A transgenic mice. *J Immunol*. 2010;185(3):1577-83.
88. Langer F, Ingersoll SB, Amirkhosravi A, Meyer T, Siddiqui FA, Ahmad S, et al. The role of CD40 in CD40L- and antibody-mediated platelet activation. *Thromb Haemost*. 2005;93(6):1137-46.
89. Chamberlain C, Colman PJ, Ranger AM, Burkly LC, Johnston GI, Otoul C, et al. Repeated administration of dapirolizumab pegol in a randomised phase I study is well tolerated and accompanied by improvements in several composite measures of systemic lupus erythematosus disease activity and changes in whole blood transcriptomic profiles. *Ann Rheum Dis*. 2017;76(11):1837-44.
90. Furie R, Bruce, I.N., Dörner, T., Leon, M.G., Leszczynski, P., Urowitz, M.B., Haier, B., Jimenez, T., Barbey, C., Liu, J., Stach, C. EFFICACY AND SAFETY OF DAPIROLIZUMAB PEGOL (DZP) IN PATIENTS WITH MODERATELY TO SEVERELY ACTIVE SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): A RANDOMISED, PLACEBO (PBO)-CONTROLLED STUDY. *Ann Rheum Dis* 2019;volume 78, supplement 2 page A775.
91. Li J GM, Ly N, Godwood A, Howe D, Albulescu M, Butler LH, Arbaugh K, Faggioni R. Pharmacokinetics, Pharmacodynamics, and Immunogenicity of MEDI4920, a Novel, Engineered CD40 Ligand Antagonist, in Healthy Volunteers [abstract]. *Arthritis Rheumatol* 2016;68 (suppl 10). .
92. Albach FN, Wagner F, Huser A, Igel J, Joseph D, Hilbert J, et al. Safety, pharmacokinetics and pharmacodynamics of single rising doses of BI 655064, an antagonistic anti-CD40 antibody in healthy subjects: a potential novel treatment for autoimmune diseases. *Eur J Clin Pharmacol*. 2018;74(2):161-9.
93. Visvanathan S, Daniluk S, Ptaszynski R, Muller-Ladner U, Ramanujam M, Rosenstock B, et al. Effects of BI 655064, an antagonistic anti-CD40 antibody, on clinical and biomarker variables in patients with active rheumatoid arthritis: a randomised, double-blind, placebo-controlled, phase IIa study. *Ann Rheum Dis*. 2019;78(6):754-60.
94. Ristov J, Espie P, Ulrich P, Sickert D, Flandre T, Dimitrova M, et al. Characterization of the in vitro and in vivo properties of CFZ533, a blocking and non-depleting anti-CD40 monoclonal antibody. *Am J Transplant*. 2018;18(12):2895-904.
95. Cordoba F, Wieczorek G, Audet M, Roth L, Schneider MA, Kunkler A, et al. A novel, blocking, Fc-silent anti-CD40 monoclonal antibody prolongs nonhuman primate renal allograft survival in the absence of B cell depletion. *Am J Transplant*. 2015;15(11):2825-36.
96. Nashan B, Tedesco H, van den Hoogen M, Berger S, Cibrik D, Mulgaonkar S, et al. CD40 Inhibition with CFZ533 - A New, Fully Human, Non-Depleting, Fc Silent mAb - Improves Renal Allograft Function While Demonstrating Comparable Efficacy vs. Tacrolimus in De-Novo CNI-Free Kidney Transplant Recipients. 2018;102:S366.
97. Farkash E NA, Tedesco-Silva H, Nashan B, Witzke O, Hoogen Mvanden, Berger S, Cibrik DM, Mulgaonkar S, Leeser DB, Alloway R, Patel A, Pratchke J, Sommerer C, Wiseman A, Zuilen AVan, Laessing U, Rush J, Haraldsson B. . CNI-Free Therapy with Iscalimab (anti-CD40 mAb) Preserves Allograft Histology Compared to Standard of Care after Kidney Transplantation [abstract]. *Am J Transplant*. 2019;19 (suppl 3).
98. Fisher B, Szántó A, Ng WF, Bombardieri M, Posch M, Papas A, et al. Safety, tolerability, efficacy, and pharmacokinetics of the anti-CD40 antibody iscalimab in patients with primary Sjögren's syndrome: a multi-center, randomised, double-blind, placebo-controlled, parallel group proof-of-concept study *Lancet Rheumatology*. 2020;In press.
99. Colafrancesco S, Priori R, Smith CG, Minniti A, Iannizzotto V, Pipi E, et al. CXCL13 as biomarker for histological involvement in Sjogren's syndrome. *Rheumatology (Oxford)*. 2019.

100. Fisher B SA, Ng WF, et al. FRI0174 SUBCUTANEOUS DOSING OF THE NOVEL ANTI-CD40 ANTIBODY ISCALIMAB ACHIEVES TARGET DRUG EXPOSURE AND CLINICAL EFFICACY IN PRIMARY SJÖGREN'S SYNDROME; RESULTS OF A PHASE IIA RANDOMISED OPEN LABEL TWO ARM PARALLEL GROUP TRIAL. *Annals of the Rheumatic Diseases* 2019;78:760-1.

1 Table 1 CD40/CD40L targeting molecules

Target	Commercial name	Molecule	Diseases	Clinical trial stage	Outcome in rheumatic disorders	References
CD40L	Ruplizumab (BG9588, hu5c8)	Humanized IgG1 mAb	SLE, transplantation	Stopped after phase 2 (thromboembolism)	Reduced disease activity (SLE)	(83)
	Toralizumab	Humanized IgG1 mAb	SLE, AT, CD, MS	Stopped after phase 2 (thromboembolism)	No superiority compared to placebo (SLE)	(84, 85, 87)
	Dapirolizumab	Fab' fragment	SLE	Phase 2b completed (NCT02804763)	Clinical response rate higher in dapirolizumab group vs placebo (P=0.06)	(90)
	Letolizumab (BMS-986004)	Human IgG1 fusion protein	Transplantation, AT	Phase 1 and 2 completed (AT, NCT02273960) Phase 1 and 2 ongoing (GVHD, NCT02273960)		
	VIB4920 (MEDI4920)	Tn3 fusion protein	RA, SS, transplantation	Phase 1b completed (RA, NCT02780388) Phase 2 ongoing (RA, NCT04163991) Phase 2 ongoing (SS, NCT04129164) Phase 2 ongoing (transplantation, NCT04046549, NCT04174677)	Clinical and laboratory response (NCT02780388)	(86)
CD40	Isclimab (CFZ533)	Fc-modified human IgG1 mAb	SS, RA, SLE, MG, transplantation	Phase 1 completed (RA) (NCT02089087) Phase 2a completed (SS) (NCT02291029) Phase 2 completed (GD) (NCT02713256) Phase 2 completed (MG) (NCT02565576)	Safe and tolerated (NCT02089087) Safe and tolerated. Clinical and laboratory response (NCT02291029)	(98)

				Phase 2b ongoing (SS) (NCT03905525) Phase 2 ongoing (SLE, LN) (NCT03656562, NCT03610516) Phase 2 ongoing (LT, KT) (NCT03781414, NCT03663335) Phase 2 ongoing (T1DM) NCT04129528		
	Bleselumab	Human IgG4 mAb	Psoriasis, transplantation, FSGS	Phase 1 completed (KT) (NCT01279538) Phase 2a completed (psoriasis) Phase 2 ongoing (KT and FSGS) (NCT02921789)		
	BI-655064	Humanized IgG1 mAb	RA, AT, SLE	Phase 2 completed (RA, AT) (NCT01751776, NCT02009761) Phase 2 ongoing (SLE) (NCT03385564)	Safe and tolerated. Clinical and laboratory response (NCT01751776)	(92)
	Ch5D12	Human IgG4 mAb	CD	Phase 1 and 2 completed		
	FFP104	Human IgG4 mAb	CD, PBC	Phase 1 and 2 ongoing (PBC) (NCT02193360) Phase 2 ongoing (CD) (NCT02465944)		

2

3 **Abbreviations:** AT, autoimmune thrombocytopenia; CD, Crohn's disease; DM, Type 1 Diabetes; FSGS, focal segmental glomerulosclerosis; GD,
4 Graves' disease; KT, kidney transplant; LN, lupus nephritis; LT, liver transplant; MG, myasthenia gravis; MS, multiple sclerosis; PBC, primary biliary
5 cirrhosis; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus; SS, Sjögren's syndrome