

Temporal trends in radiometrically dated sediment cores from English lakes show polybrominated diphenyl ethers correlate with brominated but not mixed bromo/chloro dioxins and furans

Peters, Leon I.; Rose, Neil L.; Yang, Handong; Klánová, Jana; Moehring, Thomas; Harrad, Stuart

DOI:
[10.1016/j.scitotenv.2020.143118](https://doi.org/10.1016/j.scitotenv.2020.143118)

License:
Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version
Peer reviewed version

Citation for published version (Harvard):
Peters, LI, Rose, NL, Yang, H, Klánová, J, Moehring, T & Harrad, S 2021, 'Temporal trends in radiometrically dated sediment cores from English lakes show polybrominated diphenyl ethers correlate with brominated but not mixed bromo/chloro dioxins and furans', *Science of the Total Environment*, vol. 762, 143118.
<https://doi.org/10.1016/j.scitotenv.2020.143118>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Download date: 29. Nov. 2021

1 **Temporal Trends in Radiometrically Dated Sediment Cores**
2 **from English Lakes Show Polybrominated Diphenyl Ethers**
3 **Correlate with Brominated but not Mixed Bromo/Chloro**
4 **Dioxins and Furans**

5
6 Leon I. Peters^{a#}, Neil L. Rose^b, Handong Yang^b, Jana Klánová^c, Thomas Mochring^d, Stuart
7 Harrad^{a*}
8

9 ^aSchool of Geography, Earth, and Environmental Sciences, University of Birmingham,
10 Birmingham, B15 2TT, U.K.

11 ^bEnvironmental Change Research Centre, Department of Geography, University College London,
12 London, WC1E 6BT, U.K.

13 ^cRECETOX, Masaryk University, Brno, Czech Republic

14 ^dThermo Fisher Scientific (GmbH) Bremen, Hanna-Kunath-Str. 11, 28199 Bremen, Germany

15
16 [#]Deceased

17 ^{*}Email address: S.J.Harrad@bham.ac.uk
18

19 **ABSTRACT**

20 This paper reports concentrations between ~1950 and present, of polybrominated diphenyl ethers
21 (PBDEs) and polybrominated dibenzo-*p*-dioxins and furans (PBDD/Fs), in radiometrically-dated
22 sediment cores from three English lakes. Mixed bromo/chloro dibenzo-*p*-dioxins and furans
23 (PXDD/Fs) were measured in two of the same lakes. Concentrations of PXDD/Fs decreased over
24 time to the present. To our knowledge, this is the first report of temporal trends of PXDD/Fs in the
25 environment. In contrast, concentrations of PBDEs increased towards the present and were
26 significantly correlated ($R= 0.88-0.98$; $p<0.05$) with concentrations of PBDFs in all three lakes.
27 These observations suggest that the sources of PXDD/Fs are not related to PBDEs and differ from
28 those of PBDFs. We also report for the first time the presence of octabromodibenzofuran (OBDF)
29 in the two most recent core slices at one lake. The source of OBDF in these samples is unclear.
30 While OBDF has been reported previously as a significant contaminant of some commercial
31 formulations of Deca-BDE, it is also present in Octa-BDE products and in emissions from a variety
32 of combustion activities. Overall, while the positive correlation between PBDEs and PBDFs
33 suggests increased use of PBDEs has contributed substantially to environmental contamination
34 with PBDFs; examination of PBDF homologue patterns implies emissions from combustion
35 activities are likely also important.

36 **Keywords**

37 PBDD/Fs; PXDD/Fs; PBDEs; Time Trends; Sources; Lakes

38 **Introduction**

39 Polybrominated diphenyl ethers (PBDEs) are industrial chemicals that have found extensive global
40 use as flame retardants added to a wide range of consumer items such as electrical and electronic
41 equipment and furniture fabrics and foams. As a consequence of their environmental persistence,
42 potential for long-range atmospheric transport and bioaccumulation, and adverse effects on
43 humans and/or wildlife, PBDEs are listed under the United Nations Environment Program's
44 Stockholm Convention on Persistent Organic Pollutants (POPs). Moreover, manufacture and new
45 use of PBDEs is either banned or restricted by many jurisdictions. Against this backdrop, there is a
46 clear need to evaluate the efficacy of such actions; for example, by monitoring temporal trends in
47 environmental contamination. Several studies worldwide have previously delineated the increase
48 in concentrations of PBDEs in the environment from the onset of their widespread use in the 1980s
49 to mid-2000s when the first restrictions on them were introduced (Kohler et al, 2008; Zegers et al,
50 2003; Vane et al, 2010; Webster et al, 2008). Subsequent trends up to the present, point to
51 concentrations levelling off and even declining as actions to eliminate PBDEs take effect (Yang et
52 al, 2016). One concern with respect to PBDEs is that commercial PBDE products have been shown
53 to contain polybrominated dibenzo-*p*-dioxins and furans (PBDD/Fs) as contaminants at levels that
54 amount to a substantial mass (0.43 - 2.2 t globally) of PBDD/Fs (Hanari et al, 2006; Ren et al,
55 2011). This is consistent with a recent report that temporal trends of PBDFs correlate with those of
56 PBDEs in sediment cores from Tokyo Bay (Goto et al, 2017). This is concerning owing to evidence
57 of the toxicity of PBDFs that is considered by the WHO to be on a par with that of their chlorinated
58 analogs (Van den Berg et al, 2013). Moreover, there exists emerging concern about environmental
59 contamination with mixed bromo-/chloro-dibenzo-*p*-dioxins and furans (PXDD/Fs) (Wall et al,
60 2015). Although little is known about the toxicity of PXDD/Fs, their structural similarity to

61 PBDD/Fs means that establishing the level at which they are present in the environment and
62 elucidating their sources is desirable. Studies exist that demonstrate that combustion activities such
63 as iron ore sintering and waste incineration are sources of PXDD/Fs (Chatkittikunwong and
64 Creaser, 1994; Weber et al, 2003), with their presence in air in Japan (Hayakawa et al, 2004) as
65 well as soil impacted by a recycling plant fire (Myers et al, 2012) and by informal e-waste handling
66 reported (Leung et al, 2007; Ma et al, 2008; Tue et al, 2013; Yu et al, 2008; Zennegg et al, 2009).
67 Currently however, very few data exist on concentrations of PXDD/Fs in sediments.
68 Concentrations of Σ PXDD/Fs were between 0.03-0.1 ng/g dry weight in freshwater sediment taken
69 from a former chlor-alkali plant in the USA, with monobromoheptachloro dibenzo-*p*-dioxin the
70 only congener detected (Kannan et al, 1998). In Osaka Bay, Japan, a wider range of PXDD/Fs
71 were detected in marine sediment, most prominently 2-Br-3,7,8-ClCDD detected between 0.84-6.5
72 pg/g dry weight (Ohta et al, 2002), while concentrations of monobromo-polychlorinated
73 dibenzo-*p*-dioxins/dibenzofurans which ranged between below detection limit and 1.8 ng/g dry
74 weight were reported for marine surficial sediments from Hong Kong and Korea (Terauchi et al,
75 2009).

76 In this study we exploit the fact that sediment deposited in lakes over time can provide a reliable
77 record of contaminant inputs into lacustrine systems. We thus collected sediment cores from three
78 English lakes for which data on concentrations of PBDEs, hexabromocyclododecane (HBCDD),
79 and polychlorinated biphenyls (PCBs) have been measured previously (Yang et al, 2016) and used
80 radiometric techniques to assign dates to core slices representing different depths. Concentrations
81 of PBDEs, PBDD/Fs, and PXDD/Fs were measured in individual core slices and the data used to
82 test the hypothesis that these contaminant classes will display similar temporal trends.

83 **2. Experimental**

84 *2.1 Sampling locations*

85 A map of our sampling locations is given as supplementary data (Figure SD-1), with additional
86 information about individual sites supplied in Table SD-1. Locations studied were 3 of the same
87 seepage lakes for which we have previously reported temporal trends in concentrations of PBDEs,
88 HBCDD, and PCBs in sediment cores (Yang et al, 2016). These were: Edgbaston Pool, Holt Hall
89 Lake, and Wake Valley Pond. These lakes were selected from locations with a range of population
90 densities as a surrogate indicator of anthropogenic inputs. One sediment core from each lake was
91 collected between 23rd – 26th June 2015, covering sedimentation from at least ~1950 to the date of
92 sampling. Consistent with our previous study of these lakes (Yang et al, 2016), each core was
93 collected from a flat area of the lake basin near the deepest point.

94 *2.2 Sampling and sectioning of lake sediment cores*

95 Cores were sampled from a flat area close to the maximum depth at each site and were collected
96 from a purpose built pontoon to a depth of between 0.75-0.95 m below the benthic surface using a
97 large diameter sediment core apparatus ('Big-Ben') (Patmore et al, 2014). The sediment corer,
98 piston and core covers were all thoroughly decontaminated with hexane before and after use. The
99 'Big-Ben' corer having an internal diameter of 140 mm, is considerably larger than conventional
100 piston corers (typically 50 - 80 mm i.d.) and provides sediment cores with a cross-sectional area of
101 154 cm² resulting in far greater sample for analysis, and facilitating the analysis of PBDD/Fs and
102 PXDD/Fs at the anticipated ultra-trace levels. Sediment cores were extruded vertically in the field
103 at 10 mm intervals, with each sample stored at -20° C in individually sealed Whirl-PackTM
104 sampling bags until extraction analysis. Sample contamination derived from use of Whirl-PackTM
105 sampling bags manufactured from low density polyethylene (LDPE) was controlled for with the
106 use of sampling blanks, which consisted of 30 g pre- cleaned Na₂SO₄ spiked with 10 µL

107 $^{13}\text{C}_{12}$ -BDE-138. Sampling control blanks were opened to the atmosphere for approximately 30 min
108 to allow the sampling spike solvent to evaporate, before being homogenised and sealed until
109 analysis. Three sampling blanks were employed per site and treated analogously to sediment
110 samples, including extraction and analysis. Recoveries of $^{13}\text{C}_{12}$ -BDE-138 ranged between 50 - 110
111 % with a mean of 85.4 ± 35.2 % (Mean \pm 1SD) and all cases (n= 9) yielded BFR concentrations
112 below limits of quantification confirming that the sample collection procedure did not contribute to
113 sediment contamination with PBDEs.

114 Each sediment sample was divided into two during the core-sectioning procedure: two thirds kept
115 for analysis of our target contaminants; with the remaining one third used for radiometric dating
116 and determination of sediment water and total organic carbon (TOC) content. The latter two
117 metrics were determined gravimetrically by mass loss from a 2 g (wet weight) sample, oven dried
118 at 105 °C for 3 h to determine water content and a further 2 h at 550 °C to determine TOC by
119 loss-on-ignition. After freeze-drying, each 10 mm core slice was homogenized. Thereafter, based
120 on information from our radiometric dating results, we combined 10 mm core slices from the same
121 core to generate a series of pooled core slice samples that each represented around five years of
122 sedimented material. In total, 31 such pooled samples were prepared for analysis. A further core
123 slice was analyzed from the bottom of the core taken from each of our lakes, to provide an
124 indication of concentrations of our target contaminants prior to ~1950.

125 *Analytical methods*

126 Details of methods used to radiometrically date (using ^{210}Pb) sediment core slices are given as SD,
127 with sediment chronologies and sedimentation rates provided as Tables SD-2-SD-4 and Figures
128 SD-2-SD-7. For determination of concentrations of target contaminants in sediment samples, 5 g
129 each of both hydromatrix and sodium sulfate (both pre-cleaned) were loaded into pre-cleaned

130 stainless steel extraction cells, along with an aliquot of freeze-dried and homogenized sediment
131 core slice (typically 5 g, accurately weighed). This was treated with ¹³C-labeled internal standards
132 (details below) and subjected to pressurized liquid extraction using an ASE-350 (Dionex, CA).
133 Extraction was first with hexane:dichloromethane (3:2, v/v; 2 cycles), followed by toluene (2
134 cycles) at 90°C, 1500 psi and hold time= 5 min. Following concentration to ~ 1 mL using a
135 Turbovap solvent evaporator, crude sediment extracts were purified prior to instrumental analysis
136 via elution through a sequence of acid silica and activated carbon columns (Cape Technologies,
137 Maine, USA). This yielded two concentrated extracts (F1b and F2b) for instrumental analysis of
138 PBDEs (F1b) and PBDD/Fs + PXDD/Fs (F2b) for each sediment core slice. Detailed descriptions
139 of the complex procedures used to purify and fractionate sediment core extracts prior to
140 instrumental analysis are provided as SD (Figures SD-8-10 and accompanying text).

141 Commercially available standards were used for the analysis of BFRs. These comprised native
142 PBDEs and ¹³C₁₂-BDEs 28, 47, 99, 100, 153, 154, 183, 207, and 209 (Wellington Laboratories)
143 and native and ¹³C₁₂- analogues of the following PBDD/Fs (Cambridge Isotope Laboratories):
144 2,3,7,8-TBDD, 1,2,3,7,8-PeBDD, 1,2,3,4,7,8-HxBDD, 1,2,3,6,7,8-HxBDD, 1,2,3,7,8,9-HxBDD,
145 1,2,3,4,6,7,8-HpBDD, OBDD, 2,3,7,8-TBDF, 2,4,6,8-TBDF, 1,2,3,7,8-PeBDF, 2,3,4,7,8-PBDF,
146 1,2,3,4,7,8-HxBDF, 1,2,3,4,6,7,8-HpBDF, and OBDF. The following native standards of
147 PXDD/Fs were kindly provided by Dr. Alwyn Fernandes, FERA, UK: 2-Br-7,8-CDD,
148 2-Br-3,7,8-CDD, 2,3-Br-7,8-CDD, 1-Br-2,3,7,8-CDD+2-Br-1,3,7,8-CDD, 2-Br-3,6,7,8,9-CDD,
149 2-Br-7,8-CDF, 2-Br-6,7,8-CDF+3-Br,2,7,8-CDF, and 1-Br-2,3,7,8-CDF. In the absence of
150 ¹³C₁₂-labeled PXDD/Fs, ¹³C₁₂-2,4,6,8-TBDF was used as internal (surrogate) standard for
151 quantification of PBDD/Fs.

152 *Instrumental Analysis*

153 Quantification of target compounds and congener groups was conducted on a Thermo Scientific
154 GC Q Exactive GC/orbitrap MS. Concentrations of PBDEs were determined via injection of 2 μ L
155 of F1 onto a Thermo Fisher Scientific Trace Gold Dioxin capillary column (12 m x 0.25 mm x 0.1
156 μ m), with the GC-MS operated in selected ion monitoring (SIM) mode. Concentrations of
157 PBDD/Fs and PXDD/Fs were determined by injecting 2 μ L of F2 onto the same GC column, with
158 the GC-MS operated in full scan mode. For both F1 and F2, extracts were injected via a PTV
159 injector operating in splitless mode. A programmed temperature of injection of 120 $^{\circ}$ C followed by
160 a ramp profile from 150 $^{\circ}$ C to 320 $^{\circ}$ C at 14.5 $^{\circ}$ C/sec was deployed. A constant He flow of 1.3
161 mL/min was deployed with a temperature program of: 120 $^{\circ}$ C held for 3 min before ramping to 250
162 $^{\circ}$ C at 6.5 $^{\circ}$ C/min, followed by a second ramp from 250 $^{\circ}$ C to 305 $^{\circ}$ C at 8 $^{\circ}$ C/min held for 7 min. The
163 orbitrap MS was operated at 60,000 mass resolution, which facilitated confirmation of analyte
164 identity via accurate mass. In all cases, where a standard for a given compound was included in our
165 calibration standard, identification of that compound was via comparison of retention time to the
166 standard, while quantification of that compound was made using its response factor relative to the
167 appropriate internal (surrogate) standard. For PBDD/F and PXDD/F compounds for which we did
168 not have standards, quantification was achieved using an averaged relative response factor for
169 compounds of the same homologue group. For example, hexabromo-PBDFs were quantified using
170 the relative response factor for 1,2,3,4,7,8-HxBDF. Full details of mass spectrometric parameters
171 are provided as SD.

172 *QA/QC*

173 Recoveries for all target analyte internal standards ranged between 50 and 110 %, with the
174 exception of $^{13}\text{C}_{12}$ -OBDD and $^{13}\text{C}_{12}$ -OBDF for which recoveries in some samples were as low as
175 40%. Method accuracy was assessed for PBDEs by replicate (n=7) analysis of NIST SRM1944

176 (New York/ New Jersey Waterway Sediment). This revealed good agreement between
177 concentrations measured in this study with the certified values reported by NIST. Full details are
178 provided in SD (Table SD-5). While certified or indicative values have not previously been
179 reported for PBDD/Fs and PXDD/Fs, they are also provided as SD (Tables SD-6 and SD-7) for
180 comparison with future studies.

181 *Statistical methods*

182 All statistics in this study were conducted using IBM SPSS Statistics 21, and Microsoft Excel
183 2007.

184 **Results and discussion**

185 *Concentrations of PBDEs*

186 Concentrations of all target PBDEs and PBDD/Fs in each individual core slice from each sampling
187 location are provided as supporting data (Tables SD-8 – SD-13). Table 1 provides a summary of
188 concentrations of Σ PBDEs detected in this study together with those reported in selected relevant
189 previous studies for comparison. Most pertinently, concentrations of Σ PBDEs reported for surficial
190 sediments for our three lakes match closely with those reported previously by our group for
191 surficial sediments (albeit dating from 6-7 years earlier) for the same lakes (Yang et al, 2016).

192 While the lake situated in the local authority with the highest population density (Edgbaston Pool)
193 displayed the greatest concentrations of Σ PBDEs, concentrations at Wake Valley Pond and Holt
194 Hall Lake are similar despite Wake Valley Pond being located in a more densely populated local
195 authority area (Table SD-1 – i.e. Edgbaston Pool 2,500 people/km² >Wake Valley Pond 250-499
196 people/km² >Holt Hall Lake 100-249 people/km²). This is partly consistent with previous findings
197 showing higher concentrations of PBDEs in air and soil from more densely populated areas due to

198 emissions from the built environment (Harrad and Hunter, 2006). Concentrations of Σ PBDEs in
199 this study at Edgbaston Pool ca 3 km from the center of the city of Birmingham are close to the
200 average reported for surficial sediments collected in 2011 from 45 locations along the River
201 Thames in the UK (Ganci et al, 2019). By comparison, those at Holt Hall Lake and Wake Valley
202 Pond are around an order of magnitude lower than at Edgbaston Pool. When compared to Σ PBDE
203 concentrations in surficial sediments collected in 2002-2003 as part of two studies conducted in
204 industrial areas of the River Clyde in Scotland (Vane et al, 2010; Webster et al, 2008),
205 concentrations at all three lakes in our study are substantially lower.

206 *Relative Abundance of PBDE Congeners*

207 The PBDE congener pattern in all our samples is dominated by BDE-209. This is consistent with
208 industry figures that in Europe, Deca-BDE production far exceeded that of both Penta- and
209 Octa-BDE (BSEF, 2003). It is also in line with previous reports for the same lakes (Yang et al,
210 2016), with sediment cores from the Clyde Estuary in Scotland, UK (Vane et al, 2010; Webster et
211 al, 2008), and with surficial sediments from the River Thames running through London, UK (Ganci
212 et al, 2019). Consistent with the Clyde Estuary, the relative contribution of the nona-BDEs-206,
213 -207, and -208 to Σ PBDEs in this study exceeds that observed in commercial formulations of
214 Deca-BDE. Specifically, in the top three core slices from all three locations, Σ nona-BDEs
215 comprises 8-27% of Σ BDEs. This compares to 2.5 and 9.3% respectively in the commercial
216 Deca-BDE formulations Saytex 102E and Bromkal 82-0DE (La Guardia et al, 2006). While the
217 contribution of Σ nona-BDEs to Σ BDEs is higher in the commercial Octa-BDE products DE-79 and
218 Bromkal 79-8DE at 12.1 and 18.9% respectively (La Guardia et al, 2006); the contribution of the
219 main indicator congener for Octa-BDE (BDE-183) to Σ PBDEs in our sediments is much lower
220 (0.45-2.1% in the top three slices at all lakes) than in DE-79 and Bromkal 79-8DE (42% and 12.6%

221 respectively). On this evidence, the presence of the three nona-BDEs in this study seem most likely
222 attributable to debromination of BDE-209. This is supported by the fact that BDE-209 and
223 Σ nona-BDEs are positively correlated ($p < 0.05$ at Edgbaston Pool; $p < 0.1$ at Wake Valley Pond and
224 Holt Hall Lake) at each lake studied here. Similar conclusions were drawn based on the
225 observation that nona-BDEs were the second most abundant homologue group after BDE-209 in
226 river sediments from China (Mai et al, 2005; Zhang et al, 2009). However, we note a report of
227 elevated abundances of BDE-208 relative to BDE-209 in TV/PC display casings, and PC
228 components sampled in South China. In this study the authors attributed to the decomposition of
229 higher brominated PBDEs during the process of manufacturing use and/or recycling of
230 PBDE-containing products (Chen et al., 2010). Specifically, the ratios of average concentrations
231 of BDE-208: average concentrations of BDE-209 in TV and PC housing were 5.4% and 2.8%
232 respectively. By comparison, BDE-208:BDE-209 ratios in the uppermost 4 core slices in our
233 study ranged from 0.2% to 0.6% at Edgbaston Pool; 1.3% to 2.0% at Wake Valley Pond; and 1.3%
234 to 9.5% at Holt Hall Lake. It is thus possible that the nona-BDEs detected in our study have
235 arisen at least in part to emissions from treated products as well as any post-emission
236 debromination (Law et al., 2008).

237 *Temporal Trends in PBDEs in English Lake Sediments*

238 Figure 1 plots the temporal trends in concentrations of Σ PBDEs at all three locations. Given the
239 predominant contribution of BDE-209 to Σ PBDEs, the trends in Σ PBDEs reflect those of
240 BDE-209. Importantly, BDE-209 concentrations in the cores from Edgbaston Pool and Wake
241 Valley Pond do not peak in the uppermost core slice (2015); instead the highest concentrations are
242 found in the slice immediately below (2012 and 2009 for Edgbaston Pool and Wake Valley Pond
243 respectively). The situation differs for Holt Hall Lake, where the highest BDE-209 concentration is

244 in the surficial sediment (2015). Overall, this suggests that while the full impact of recent
245 restrictions on the manufacture and use of Deca-BDE has yet to manifest itself at the lakes studied
246 here; there are tentatively encouraging signs that contamination is levelling off.

247 *Concentrations, Relative Abundance, and Possible Sources of PBDD/Fs in English Lake*
248 *Sediments*

249 PBDD/Fs were detected in all samples in this study at concentrations two orders of magnitude
250 below those of PBDEs. Table 1 compares Σ PBDD/F concentrations in this study compared to those
251 reported in a variety of previous studies from other locations. Concentrations in sediment core
252 slices from the less urbanised Wake Valley Pond and Holt Hall Lake are in line with those reported
253 for Swedish lakes and rivers (Hagberg et al, 2005; Lundstedt, 2016). However, those at Edgbaston
254 Pool exceed those reported elsewhere. Similar to the situation for PBDEs, while the lake situated in
255 the local authority with the highest population density (Edgbaston Pool) has noticeably the highest
256 Σ PBDD/F concentrations; levels of these contaminants are similar at Wake Valley Pond and Holt
257 Hall Lake despite the former being located in a more densely populated local authority area.
258 PBDFs were overwhelmingly dominant, with OBDD the only PBDD detected and that only in the
259 uppermost two layers in the cores taken at Wake Valley Pond and Holt Hall Lake. This is
260 consistent with previous observations for surficial sediments and atmospheric deposition (Goto et
261 al, 2017; Hayakawa et al, 2004), as well as in emissions from waste incinerators (Wang et al,
262 2010), and iron ore sintering (Drage et al, 2014). It is also pertinent to note that while OBDD and to
263 a far lesser extent penta- and tetra-BDDs have been detected in commercial PBDE formulations
264 (Ren et al, 2011), PBDFs – especially OBDF and 1,2,3,4,6,7,8-HpBDF - were in comparison
265 detected in such products at far higher concentrations (Hanari et al, 2006; Ren et al, 2011).

266 Figure 2 shows the homologue profiles of PBDFs in sediment core slices from each lake. It is
267 interesting to note the spatial variation in these profiles, for example TBDFs are far less abundant at
268 Edgbaston Pool than at the other two lakes. A striking observation is that OBDF was detected at
269 Wake Valley Pond - albeit only in the two uppermost core slices (2015 and 2009). To our
270 knowledge, this is the first report of OBDF in sediments. The low detection frequency of OBDF in
271 sediments is perhaps surprising given previous data that reports OBDF to be an impurity in both
272 Octa-BDE and Deca-BDE commercial formulations (Hanari et al, 2006; Ren et al, 2011).
273 However, this may be because OBDF has been reported to be easily debrominated (Goto et al,
274 2017). Overall, the PBDF homologue pattern in our sediment cores (expressed as a percentage of
275 Σ PBDFs) more closely resembles that reported for the Octa-BDE formulation which comprises 0.3
276 % TBDFs, 1.8 % PeBDFs, 44.9 % HxBDFs, 27.1 % HpBDFs, and 25.8 % OBDF, than the
277 Deca-BDE formulation which comprises 0.1% for each of TBDFs, PeBDFs, and HxBDFs, 3.6%
278 for HpBDFs, and 96.1% OBDF. While this might suggest the Octa-BDE commercial formulation
279 is a more significant source of the PBDFs observed in our sediments than the Deca-BDE product;
280 as noted above, the major PBDE formulation used in the UK was Deca-BDE. Moreover, we note
281 that concentrations of Σ PBDFs are more strongly correlated with those of BDE-209 ($R=0.96$,
282 $p<0.05$ and $R=0.85$, $p<0.1$) than those of BDE-183 at both Edgbaston Pool and Wake Valley Pond
283 ($R=0.48$, $p>0.1$ and $R=0.64$, $p>0.1$) - no meaningful inference could be drawn in this regard at Holt
284 Hall Lake as there were too few samples in which PBDFs and both BDEs-183 and -209 were
285 detected.

286 In addition to the potential contribution of PBDE use, both OBDF and 1,2,3,4,6,7,8-HpBDF (the
287 predominant HpBDF detected in this study) have been reported to be emitted from combustion
288 processes (Wang et al, 2015). We also note that we did not target

289 2,7-/2,8-dibromodibenzo-p-dioxins or 1,3,7-/1,3,8-tribromodibenzo-p-dioxins, which have been
290 identified as being of biogenic origin (Goto et al, 2017) and can therefore not rule out possible
291 contributions from biogenic sources to the burden of PBDD/Fs in our lakes. In summary, drawing
292 firm conclusions about PBDD/F source attribution based on homologue and congener patterns in
293 environmental matrices like lake sediments, is complicated by post-emission modification of such
294 profiles by weathering processes that favour some PBDD/Fs more than others. Notwithstanding
295 this, the evidence of congener/homologue profiles presented here, suggests that the PBDD/Fs
296 detected in this study are predominantly a complex integral of combustion source emissions and
297 their presence as contaminants of the Deca-BDE formulation.

298 *Temporal Trends in concentrations of Σ PBDFs and PBDF homologue patterns*

299 Temporal trends in concentrations of Σ PBDFs are shown in Figure 1. Overall, they reveal a steady
300 rise in concentrations from the 1980s onwards, peaking at the surface (2015) at Edgbaston Pool and
301 Wake Valley Pond, and in the second most recent core slice (dated to 2009) at Holt Hall Lake. Prior
302 to the onset of rising concentrations in the 1980s, concentrations were much lower. Inspection of
303 Figure 1 suggests temporal trends in Σ PBDEs and Σ PBDFs are positively correlated and indeed the
304 correlation coefficient for the three sites is significant, ranging between 0.88-0.98 ($p < 0.05$ at each
305 site). With respect to changes in the homologue pattern over time, Figure 2 reveals no clear trend at
306 Holt Hall Lake. In contrast, there is a marked shift from HpBDFs to HxBDFs in more recent
307 sediment layers at Edgbaston Pool. Meanwhile at Wake Valley Pond, there is a noticeable decline
308 in the abundance of TBDFs, along with increased abundance of HpBDFs in more recent core
309 slices, as well as the aforementioned presence of OBDF in the top two layers (dated to 2009 and
310 2015) only. These variable homologue patterns and temporal trends suggest that changes in the
311 relative contributions of different sources of PBDFs over time differ between our three lakes. To

312 our knowledge there is only one study with which our data on PBDF homologue patterns in
313 sediment cores can be compared. In this study of a sediment core from Osaka Bay in Japan that
314 covered the period 1904-2000, OBDF was not detected and in declining order of abundance the
315 other homologues followed the order HpBDF>HxBDF>PeBDF>TBDF (Takigami et al, 2005).
316 This homologue pattern most closely resembled that observed in this study at Holt Hall Lake.

317

318 *Concentrations, Relative Abundance, and Temporal Trends of PXDD/Fs in English Lake*

319 *Sediments*

320 Table 2 reports concentrations of individual PXDD/Fs and homologue groups in each sediment
321 core slice analyzed in this study. Also included in Table 2 are concentrations of 2-Br-3,7,8-CDD,
322 2-Br-3,6,7,8,9-CDD, 2-Br-2,7,8-CDF, and 1-Br-2,3,7,8-CDF reported previously in surficial
323 sediments collected from Osaka Bay, Japan (Ohta et al, 2002). Concentrations of these PXDD/Fs
324 in our study are of a similar range to those reported for Osaka Bay. As observed for both PBDEs
325 and PBDD/Fs, concentrations of Σ PXDD/Fs at the two lakes studied for these contaminants do not
326 correspond with the population density of the local authority within which the lake was submitted
327 (Table SD-1). Specifically, t-test comparison shows PXDD/F concentrations at Wake Valley Pond
328 to be statistically indistinguishable ($p>0.05$) from those at Holt Hall Lake. For PXDD/Fs, the
329 concentrations detected were typically two orders of magnitude below those of PBDD/Fs. PXDFs
330 were more abundant than PXDDs in every core slice analysed from both locations where PXDD/Fs
331 were determined, with PXDFs typically 2-3 times more abundant. There was no clear temporal
332 trend in the ratio of PXDFs:PXDDs at either lake. Moreover, the homologue pattern is broadly
333 similar at both lakes and does not appear subject to temporal variation.

334 Temporal trends in concentrations of PXDD/Fs differ between the two lakes in which they were
335 measured. Concentrations at Wake Valley Pond peak in the oldest slice (1954) analyzed at this
336 location and then steadily decrease through to the present (2015). In slight contrast, concentrations
337 at Holt Hall Lake rise from 1935 to 1976, before declining through to the uppermost layer (2015).
338 In contrast to the significant positive correlation between Σ PBDEs and Σ PBDFs at all three lakes,
339 concentrations of Σ PBDEs are not significantly correlated ($p>0.05$) with those of Σ PXDD/Fs.
340 Likewise, Σ PBDD/Fs are not correlated with Σ PXDD/Fs ($p>0.05$). Overall, this suggests that the
341 sources of PXDD/Fs are different to those of PBDD/Fs, and that the manufacture and use of
342 PBDEs does not appear to be a source of PXDD/Fs.

343

344 **Conclusions**

345 Concentrations of PXDD/Fs in radiometrically-dated sediment cores from 3 English lakes
346 decreased over time to the present. In contrast, concentrations of PBDEs increased towards the
347 present and were significantly correlated with concentrations of PBDFs in all three lakes. These
348 findings suggest that PBDEs do not appear to be a source of PXDD/Fs and that the sources of
349 PXDD/Fs are different to those of PBDFs. Moreover, while the correlation between PBDEs and
350 PBDFs implies that increased use of PBDEs has contributed substantially to environmental
351 contamination with PBDFs; examination of PBDF homologue patterns implies emissions from
352 combustion activities such as waste incineration and iron ore sintering are likely also important.
353 Concentrations of PBDEs in all samples are dominated by BDE-209 which is the main constituent
354 of the Deca-BDE formulation most widely used in the UK. At two of the three lakes studied, the
355 concentrations of BDE-209 were highest in the sediment layers dated to 2012 and 2009, rather than
356 in the surface layer dated to 2015. This may indicate that recent bans and restrictions on the

357 manufacture and use of Deca-BDE may be starting to take effect and reduce concentrations in the
358 environment.

359

360 **Acknowledgements**

361 The research leading to these results has received funding from the Marie Curie Actions of the
362 European Union's FP7 Programme under REA grant agreement # 606857 (ELUTE project). The
363 analytical work was supported by the RECETOX research infrastructure (LM2018121). We thank
364 Ian Patmore for his help in the field, Aristide Ganci for field and lab-work assistance, and Alwyn
365 Fernandes for the kind gift of PXDD/F standards. This paper is dedicated to the memory of Dr
366 Leon Peters.

367

368 **Supplementary Data**

369 Supplementary data to this article can be found online at

370

371 **References**

- 372 1. Bromine Science and Environmental Forum (BSEF) 2003. Major Brominated Flame
373 Retardants Volume Estimates: Total Market Demand by Region in 2001, www.bsef.com
374 (accessed 21st January 2003).
- 375 2. Chattkittikunwong, W., Creaser, C. S. 1994. Bromo-, Bromochloro- and
376 Chloro-Dibenzo-p-Dioxins and Dibenzofurans in Incinerator Flyash. *Chemosphere* 29,
377 559-566.

- 378 3. Chen, S. J., Ma, Y. J., Wang, J., Tian, M., Luo, X-J., Chen, D., Mai, B-X. 2010 Measurement
379 and human exposure assessment of brominated flame retardants in household products from
380 South China. *J. Hazard Mater.* 176, 979–984.
- 381 4. Choi, J. W., Fujimaki, S., Kitamura, K., Hashimoto, S., Ito, H., Sakurai, T., Suzuki, N.,
382 Nagasaka, H., Tanabe, K., Sakai, S., Morita, M. 2003. Historical trends of PBDD/Fs,
383 PBDEs, PCDD/Fs, and dioxin - like PCBs in sediment cores from Tokyo Bay.
384 *Organohalogen Compd.* 61, 119-122.
- 385 5. Drage, D. S., Aries, E., Harrad., S. 2014. Studies into the formation of PBDEs and PBDD/Fs
386 in the iron ore sintering process. *Sci. Tot. Environ.* 485-486, 497-507.
- 387 6. Ganci, A. P., Vane, C. H., Abdallah, M. A.-E., Moehring, T., Harrad, S. 2019. Legacy PBDEs
388 and NBRFRs in sediments of the tidal River Thames using liquid chromatography coupled to a
389 high resolution accurate mass Orbitrap mass spectrometer. *Sci. Tot. Environ.* 658, 1355–
390 1366.
- 391 7. Goto, A., Tue, N. M., Someya, M., Isobe, T., Takahashi, S., Tanabe, S., Kinisue, T. 2017.
392 Spatio-temporal trends of polybrominated dibenzo-*p*-dioxins and dibenzofurans in archived
393 sediments from Tokyo Bay, Japan. *Sci.Tot. Environ.* 599-600, 340-347.
- 394 8. Hagberg, J., Grahn, E., van Bavel, B., Lindström, G. 2005. Occurrence and levels of
395 PCDD/Fs and PBDD/Fs in two Swedish lake sediments. *Organohalogen Compd.* 67, 2030-
396 2032.
- 397 9. Hanari, N., Kannan, K., Okazawa, T., Kodavanti, P. R. S., Aldous, K. M., Yamashita, N.
398 2006. Occurrence of Polybrominated Biphenyls, Polybrominated Dibenzo-*p*-dioxins, and

- 399 Polybrominated Dibenzofurans as Impurities in Commercial Polybrominated Diphenyl Ether
400 Mixtures. *Environ. Sci. Technol.* 40, 4400–4405.
- 401 10. Harrad, S., Hunter, S. 2006. Concentrations of Polybrominated Diphenyl Ethers in Air and
402 Soil on a Rural-Urban Transect Across a Major UK Conurbation. *Environ. Sci. Technol.* 40,
403 4548-4553.
- 404 11. Hayakawa, K., Takatsuki, H., Watanabe, I., Sakai, S. 2004. Polybrominated diphenyl ethers
405 (PBDEs), polybrominated dibenzo-p-dioxins/dibenzofurans (PBDD/Fs) and
406 monobromo-polychlorinated dibenzo-p-dioxins/dibenzofurans (MoBPXDD/Fs) in the
407 atmosphere and bulk deposition in Kyoto, Japan. *Chemosphere* 57, 343–56.
- 408 12. Kannan, K., Watanabe, I., Giesy, J. P. 1998. Congener profile of polychlorinated/brominated
409 dibenzo-p-dioxins and dibenzofurans in soil and sediments collected at a former chlor-alkali
410 plant. *Toxicol. Environ. Chem.* 67, 135-146.
- 411 13. Law, R. J.; Herzke, D.; Harrad, S.; Morris, S.; Bersuder, P.; Allchin, C. R. 2008. Levels and
412 trends of brominated flame retardants in the European environment. *Chemosphere* 73, 223–
413 241.
- 414 14. Kohler, M., Zennegg, M., Bogdal, C., Gerecke, A. C., Schmid, P. V., Heeb, N., Sturm, M.,
415 Vonmont, H., Kohler, H-P. E., Giger, W. 2008. Temporal Trends, Congener Patterns, and
416 Sources of Octa-, Nona-, and Decabromodiphenyl Ethers (PBDE) and
417 Hexabromocyclododecanes (HBCD) in Swiss Lake Sediments. *Environ. Sci. Technol.* 42,
418 6378–6384.

- 419 15. La Guardia, M. J., Hale, R. C., Harvey, E. 2006. Detailed Polybrominated Diphenyl Ether
420 (PBDE) Congener Composition of the Widely Used Penta-, Octa-, and Deca-PBDE Technical
421 Flame-retardant Mixtures. *Environ. Sci. Technol.* 40, 6247–6254.
- 422 16. Leung, A. O. W., Luksemburg, W. J., Wong, A. S., Wong, M. H. 2007. Spatial Distribution
423 of Polybrominated Diphenyl Ethers and Polychlorinated Dibenzo-*p*-dioxins and
424 Dibenzofurans in Soil and Combusted Residue at Guiyu, an Electronic Waste Recycling Site
425 in Southeast China. *Environ. Sci. Technol.* 41, 2730–2737.
- 426 17. Lundstedt, S. 2016. Sources and levels of PBDD/Fs in the Swedish environment
427 <https://www.diva-portal.org/smash/get/diva2:945535/FULLTEXT01.pdf>. (accessed March 3rd
428 2020).
- 429 18. Ma, J., Kannan, K., Cheng, J., Hori, Y., Wu, Q., Wang, W. 2008. Concentrations, profiles,
430 and estimated human exposures for polychlorinated dibenzo - *p* - dioxins and dibenzofurans
431 from electronic waste recycling facilities and a chemical industrial complex in Eastern China.
432 *Environ. Sci. Technol.* 42, 8252 - 8259.
- 433 19. Ma, J., Addink, R., Yun, S. H., Cheng, J. P., Wang, W. H., Kannan K. 2009. Polybrominated
434 dibenzo-*p* - dioxins/dibenzofurans and polybrominated diphenyl ethers in soil, vegetation,
435 workshop - floor dust, and electronic shredder residue from an electronic waste recycling
436 facility and in soils from a chemical industrial complex in Eastern China. *Environ. Sci.*
437 *Technol.* 43, 7350 - 7356.

- 438 20. Mai, B., Chen, S., Luo, X., Chen, L., Yang, Q., Sheng, G., Peng, P., Fu, J., Zeng, E. Y. 2005.
439 Distribution of Polybrominated Diphenyl Ethers in Sediments of the Pearl River Delta and
440 Adjacent South China Sea. *Environ. Sci. Technol.* 39, 3521-3527.
- 441 21. Myers, A. L., Mabury, S. A., Reiner, E. J. 2012. Analysis of mixed halogenated
442 dibenzo-*p*-dioxins and dibenzofurans (PXDD/PXDFs) in soil by gas chromatography tandem
443 mass spectrometry (GC-MS/MS). *Chemosphere* 87, 1063-1069.
- 444 22. Naturvårdsverket. 2011. Recycling and disposal of electronic waste - health hazards and
445 environmental impacts. 6417.
446 <http://naturvardsverket.se/Documents/publikationer6400/978-91-620-6417-4.pdf> (accessed
447 March 3rd 2020)
- 448 23. Ohta, S., Nakao, T., Nishimura, H., Okumura, T., Aozasa, O., Miyata, H. 2002.
449 Contamination levels of PBDEs, TBBPA, PCDD/Fs, PBDD/Fs and PXDD/Fs in the
450 environment of Japan. *Organohalogen Compd.* 57, 57-60.
- 451 24. Patmore, I. R., Sayer, C. D., Goldsmith, B., Davidson, T. A., Rawcliffe, R., Salgado, J. 2014.
452 Big Ben: A new wide-bore piston corer for multi-proxy palaeolimnology. *J. Paleolimnol.* 51,
453 79-86.
- 454 25. Ramu, K., Isobe, T., Takahashi, S., Subramanian, A., Parthasarathy, P., Tanabe S. 2008.
455 Brominated flame retardants and dioxins in soil from electronic waste recycling sites in India.
456 *Organohalogen Compd.* 70, 2058 - 2061.
- 457 26. Ren, M., Peng, P., Cai, Y., Chen, D., Zhou, L., Chen, P., Hu, J. 2011. PBDD/F impurities in
458 some commercial deca-BDE. *Environ. Pollut.* 159, 1375-1380.

- 459 27. Takigami, H., Sakai, S., Brouwer, A. 2005. Bio/chemical Analysis of Dioxin-like Compounds
460 in Sediment Samples from Osaka Bay, Japan. *Environ. Technol.* 26, 459–70.
- 461 28. Terauchi, H., Takahashi, S., Lam, P. K. S., Min, B-Y., Tanabe, S. 2009. Polybrominated,
462 polychlorinated and monobromo-polychlorinated dibenzo-*p*-dioxins/dibenzofurans and
463 dioxin-like polychlorinated biphenyls in marine surface sediments from Hong Kong and
464 Korea. *Environ. Pollut.* 157, 724-730.
- 465 29. Tue, N. M., Takahashi, S., Subramanian, A., Sakai, S., Tanabe, S. 2013. Environmental
466 contamination and human exposure to dioxin-related compounds in e-waste recycling sites of
467 developing countries. *Environ. Sci. Proc. Imp.* 15, 1326–1331.
- 468 30. Van den Berg, M., Denison, M. S., Birnbaum, L. S., DeVito, M. J., Fiedler, H., Falandysz, J.,
469 Rose, M., Schrenk, D., Safe, S., Tohyama, C., Tritscher, A., Tysklind, M., Peterson, R. E.
470 2013. Polybrominated Dibenzo-*p*-Dioxins, Dibenzofurans, and Biphenyls: Inclusion in the
471 Toxicity Equivalency Factor Concept for Dioxin-Like Compounds. *Toxicol. Sci.* 133, 197-
472 208.
- 473 31. Vane, C. H., Ma, Y-J., Chen, S-J., Mai, B-X. Increasing polybrominated diphenyl ether
474 (PBDE) contamination in sediment cores from the inner Clyde Estuary, UK. 2010. *Environ.*
475 *Geochem. Hlth.* 32, 13–21.
- 476 32. Wall, R. J., Fernandes, A., Rose, M., Bell, D. R., Mellor, I. R. 2015. Characterisation of
477 chlorinated, brominated and mixed halogenated dioxins, furans and biphenyls as potent and
478 as partial agonists of the Aryl hydrocarbon receptor. *Environ. Int.* 76, 49–56.
- 479 33. Wang, L-C., Hsi, H-C., Wang, Y-F., Lin, S-L, Chang-Chien, G-P. 2010. Distribution of
480 polybrominated diphenyl ethers (PBDEs) and polybrominated dibenzo-*p*-dioxins and

- 481 dibenzofurans (PBDD/Fs) in municipal solid waste incinerators. *Environ. Pollut.* 158, 1595–
482 602.
- 483 34. Wang, M., Liu, G., Jiang, X., Liu, W., Li, L., Li, S., Zheng, M., Zhan, J. 2015. Brominated
484 dioxin and furan stack gas emissions during different stages of the secondary copper smelting
485 process. *Atmos. Pollut. Res.* 6, 464-468.
- 486 35. Weber, R., Kuch, B. 2003. Relevance of BFRs and thermal conditions on the formation
487 pathways of brominated and brominated–chlorinated dibenzodioxins and dibenzofurans.
488 *Environ. Int.* 29, 699– 710.
- 489 36. Webster, L., Russell, M., Adefehinti, F., Dalgarno, E. J., Moffat, C. F. 2008. Preliminary
490 assessment of polybrominated diphenyl ethers (PBDEs) in the Scottish aquatic environment,
491 including the Firth of Clyde. *J. Environ. Monit.* 10, 463–473.
- 492 37. Yang, C., Rose, N.L., Turner, S.D., Yang, H., Goldsmith, B., Losada, S., Barber, J. L., Harrad,
493 S. 2016. Hexabromocyclododecanes, polybrominated diphenyl ethers, and polychlorinated
494 biphenyls in radiometrically dated sediment cores from English lakes, ~ 1950–present. *Sci.*
495 *Tot. Environ.* 541, 721–728.
- 496 38. Yu, X., Zennegg, M., Engwall, M., Rotander, A., Larsson, M., Wong, M. H., Weber, R. 2008.
497 E-Waste Recycling Heavily Contaminates a Chinese City with Chlorinated, Brominated and
498 Mixed Halogenated Dioxins. *Organohalogen Compd.* 70, 813-816.
- 499 39. Zegers, B. N., Lewis, W. E., Booij, K., Smittenberg, R. H., Boer, W., de Boer, J., Boon, J. P.
500 2003. Levels of polybrominated diphenyl ether flame retardants in sediment cores from
501 Western Europe. *Environ. Sci. Technol.* 37, 3803–3807.

- 502 40. Zennegg, M., Yu, X., Wong, M. H., Weber, R. 2009. Fingerprints of Chlorinated, Brominated
503 and Mixed Halogenated Dioxins at Two E-Waste Recycling Sites in Guiyu/China.
504 *Organohalogen Compd.* 71, 2248-2252.
- 505 41. Zhang, X-L., Luo, X-J., Chen, S-J., Wu, J-P., Mai, B-X. 2009. Spatial distribution and
506 vertical profile of polybrominated diphenyl ethers, tetrabromobisphenol A, and
507 decabromodiphenylethane in river sediment from an industrialized region of South China.
508 *Environ. Pollut.* 157, 1917–1923
- 509 42. Zhou, L., Li, H., Yu, Z., Ren, M., Zeng, X., Peng, P., Sheng, G., Fu, J. 2012. Chlorinated and
510 brominated dibenzo-p-dioxins and dibenzofurans in surface sediment from Taihu Lake, China.
511 *J. Environ. Monit.* 14, 1935-1942.

512 **Table 1: Concentrations of Σ PBDEs and Σ PBDD/Fs (ng/g dry weight) in surficial sediments**
 513 **from this study with those in soil and sediments from selected other studies**

Matrix, Location	Σ PBDD/F	Σ PBDE	Reference
Fresh water sediments			
Edgbaston Pool, UK	7.1	210	This Study
Holt Hall Lake, UK	0.49	18	This Study
Wake Valley Pond, UK	0.49	21	This Study
Edgbaston Pool, UK ^a	-	120	Yang et al, 2016
Holt Hall Lake, UK ^a	-	10	Yang et al, 2016
Wake Valley Pond, UK ^a	-	9.8	Yang et al, 2016
Surficial Sediments, River Thames, UK	-	180 (mean)	Ganci et al, 2019
Clyde Estuary, UK (core slices)	-	1-2,600	Vane et al, 2010
Scotland, UK (core slices)	-	2.3-98,000	Webster et, 2008
Rural/urban lakes, Sweden	0.44-0.54	-	Hagberg et al, 2005
Urban river, Sweden	0.41-1.7	29-62	Lundstedt, 2016
Rural lake Sweden	0.082-0.085	4.4-16	Lundstedt, 2016
Urban lake, China	0.00048-0.0057	-	Zhou et al, 2012
Stream at dump site, Peru	0.012-0.074	3.7-6.1	Naturvårdsverket, 2011
Lake, industrial area, Thailand	0.037-1.5	3.4-58	Naturvårdsverket, 2011
Marine sediments			
Hong Kong/Korea	nd.-0.46	-	Terauchi et al, 2009
Tokyo Bay, Japan	0.0052-0.070	10-78,050	Choi et al, 2003
Osaka, Japan	0.0041-0.077	8.0-352	Ohta et al, 2002
Osaka, Japan	0.0024-0.59	53-910	Takigami et al, 2005
Coastal and offshore, Sweden	0.050-10	-	Lundstedt, 2016
Rural Soil			
Lanna, Sweden	0.028-0.054	0.065-1.3	Lundstedt, 2016
Urban soil			
Umeå and Norrköping, Sweden	0.0011-0.22	0.18-66	Lundstedt, 2016
Bangalore and Chennai, India	0.0060-0.31	-	Ramu et al, 2008
Kyoto, Japan	0.28	-	Hayakawa et al, 2004
Industrial area, China	nd.-0.43	2.03-269	Ma et al, 2008; Ma et al, 2009
Industrial area, Thailand	0.019-0.16	1.8-13	Naturvårdsverket, 2011
Dump site, Peru	0.0086-0.32	3.6-92	Naturvårdsverket, 2011

514 ^aData for core slices dated to 2008 for Edgbaston Pool and for 2009 for both Holt Hall Lake and
 515 Wake Valley Pond.

516
517

Table 2: Concentrations (pg/g OC) of PXDD/Fs in Sediment Core Slices from Wake Valley Pond and Holt Hall Lake with Comparative Data from Osaka Bay, Japan


Location	Congener/ Homologue/ Year	2015	2009	2005	2004	1999	1993	1985	1977	1969	1965	1954	1935
Wake Valley Pond	2-Br-7,8-CDD	0.023	0.16	0.24		0.32	0.4	0.035			0.079	<0.032	
Holt Hall Lake	2-Br-7,8-CDD	1.1	2.8		2.6	3.4		1.1	<0.025	<0.024		<0.021	<0.021
Wake Valley Pond	Total Br-2CDD	0.73	1.9	1.9		2.7	2.3	3.6			0.8	9.3	
Holt Hall Lake	Total Br-2CDD	1.8	4.1		4.0	5.3		2.2	11	7.1		2.3	<0.021
Wake Valley Pond	2-Br-3,7,8-CDD	0.081	<0.028	<0.027		0.12	0.22	0.22			0.29	<0.047	
Holt Hall Lake	2-Br-3,7,8-CDD	0.02	0.049		0.054	0.16		0.08	1.4	<0.035		<0.031	<0.03
Osaka Bay, Japan ^a	2-Br-3,7,8-CDD	0.84-6.5											
Wake Valley Pond	Total Br-3CDD	1.5	2.2	2.8		3.8	3	3.5			6.2	9.3	
Holt Hall Lake	Total Br-3CDD	0.77	2.1		2.4	2.7		2.4	18	<0.027		<0.019	<0.018
Wake Valley Pond	2,3-Br-7,8-CDD	<0.015	<0.017	<0.016		<0.016	<0.017	<0.016			<0.016	<0.029	
Holt Hall Lake	2,3-Br-7,8-CDD	<0.013	0.095		0.062	<0.018		0.034	<0.022	<0.021		<0.019	<0.018
Osaka Bay, Japan ^a	2,3-Br-7,8-CDD	Not detected											
Wake Valley Pond	Total 2Br-2CDD	0.48	0.64	0.87		1.2	1.5	1.8			1.2	4.4	
Holt Hall Lake	Total 2Br-2CDD	0.33	0.79		0.82	1.1		0.97	5.0	2.2		0.62	<0.017
Wake Valley Pond	1-Br-2,3,7,8-CDD + 2-Br-1,3,7,8-CDD ^b	<0.018	<0.021	0.37		<0.02	<0.021	<0.02			<0.021	<0.032	
Holt Hall Lake	1-Br-2,3,7,8-CDD + 2-Br-1,3,7,8-CDD ^b	<0.016	<0.019		<0.02	<0.023		<0.027	<0.028	<0.027		<0.024	<0.023
Osaka Bay, Japan ^a	1-Br-2,3,7,8-CDD	Not detected											
Wake Valley Pond	Total Br-4CDD	1.7	1.8	2.8		3.8	5.9	4.6			7	5.7	
Holt Hall Lake	Total Br-4CDD	0.47	1.9		1.0	1.9		<0.021	<0.022	<0.021		0.71	<0.018
Wake Valley Pond	2-Br-3,6,7,8,9-CDD	<0.015	<0.017	<0.016		<0.016	<0.017	<0.016			<0.016	<0.029	
Holt Hall Lake	2-Br-3,6,7,8,9-CDD	<0.013	<0.015		<0.016	<0.018		0.6	<0.022	<0.021		<0.019	<0.018
Osaka Bay, Japan ^a	2-Br-3,6,7,8,9-CDD	Not detected - 0.32											
Wake Valley Pond	Total Br-5CDD	1	0.82	3.5		2.2	5.1	7			7.5	15	
Holt Hall Lake	Total Br-5CDD	<0.018	0.11		<0.01	<0.012		0.6	<0.022	<0.021		<0.019	<0.018

Location	Congener/ Homologue/ Year	2015	2009	2005	2004	1999	1993	1985	1977	1969	1965	1954	1935
Wake Valley Pond	2-Br-7,8-CDF	0.77	1	0.88		<0.01	2	2.7			<0.01	<0.018	
Holt Hall Lake	2-Br-7,8-CDF	<0.008	<0.01		<0.01	<0.012		0.53	<0.014	<0.013		<0.012	<0.012
Wake Valley Pond	Total Br-2CDF	6.9	16	16		26	17	25			22	62	
Holt Hall Lake	Total Br-2CDF	4.7	7.9		7.7	12		9.6	88	89		15	9.5
Wake Valley Pond	2-Br-6,7,8-CDF + 3-Br-2,7,8-CDF*	<0.012	0.13	0.17		0.3	0.65	0.46			0.076	<0.023	
Holt Hall Lake	2-Br-6,7,8-CDF + 3-Br-2,7,8-CDF*	0.07	0.28		0.18	0.40		1.3	5.1	<0.017		0.63	<0.015
Osaka Bay, Japan ^a	3-Br-2,7,8-CDF	Not detected-2.0											
Wake Valley Pond	Total Br-3CDF	4.3	6	6.4		8.4	8.7	10			11	57	
Holt Hall Lake	Total Br-3CDF	2.5	4.8		5.3	8.8		5.0	44	49		6.6	3.1
Wake Valley Pond	1-Br-2,3,7,8-CDF	0.022	<0.014	0.047		0.042	0.66	<0.012			<0.012	<0.021	
Holt Hall Lake	1-Br-2,3,7,8-CDF	<0.009	<0.011		<0.012	<0.014		<0.016	<0.017	<0.016		<0.014	<0.014
Osaka Bay, Japan ^a	1-Br-2,3,7,8-CDF	Not detected											
Wake Valley Pond	Total Br-4CDF	1.5	1.6	2.6		3.6	6.4	5.8			2	20	
Holt Hall Lake	Total Br-4CDF	1.8	5.0		2.4	5.2		0.81	7.3	<0.016		3.5	4.4
Wake Valley Pond	∑PXDD	5.4	7.5	12		14	18	20			23	43	
Holt Hall Lake	∑PXDD	3.3	9.1		8.2	11		6.0	34	9.3		7.1	<0.03
Wake Valley Pond	∑PXDF	13	24	25		38	32	41			34	140	
Holt Hall Lake	∑PXDF	9.0	18		15	26		15	140	140		25	17
Wake Valley Pond	∑PXDD/F	18	31	37		51	50	61			57	180	
Holt Hall Lake	∑PXDD/F	12	27		24	37		21	170	150		32	17

518 ^aRange (pg/g dry weight) surficial sediments from Osaka, Japan (n=6) (Ohta et al, 2002) – detection limits not reported


519

520 **Figure 1: Temporal Trends in Concentrations (ng/g OC) of Σ PBDE, Σ PBDF, and Σ PXDD/F in 3 English Lake Sediment Cores**
 521
 522




526
527

Figure 2: Temporal Trends in Relative Contributions of PBDF Homologues to Σ PBDF Concentrations in 3 English Lake Sediment Cores



528
529



530

