

Global prevalence and types of complementary and alternative medicines use amongst adults with diabetes

Alzahrani, Abdulaziz Saud S; Price, Malcolm; Greenfield, Sheila; Paudyal, Vibhu

DOI:

[10.1007/s00228-021-03097-x](https://doi.org/10.1007/s00228-021-03097-x)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Alzahrani, ASS, Price, M, Greenfield, S & Paudyal, V 2021, 'Global prevalence and types of complementary and alternative medicines use amongst adults with diabetes: systematic review and meta-analysis', *European Journal of Clinical Pharmacology*. <https://doi.org/10.1007/s00228-021-03097-x>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Global prevalence and types of complementary and alternative medicines use amongst adults with diabetes: systematic review and meta-analysis

Abdulaziz S. Alzahrani¹ · Malcolm J. Price^{2,3,4} · Sheila M. Greenfield² · Vibhu Paudyal¹

Received: 7 September 2020 / Accepted: 21 January 2021

© The Author(s) 2021

Abstract

Aim This study aimed to undertake a systematic review and meta-analysis of global prevalence and types of complementary and alternative medicine (CAM) use amongst adults with diabetes.

Methods Nine databases, including MEDLINE and EMBASE, were searched for studies published between 2009 and 2019 which included extractable data for CAM use in adult patients with diabetes. Study characteristics, types of CAM, and overall and subgroup prevalence data in relation to CAM use were extracted. Meta-analysis of aggregate level data on prevalence and prevalence ratios (PRs) was performed using a random effects model.

Results From the 38 studies included in the review, a total of 37 types of CAM and 223 types of herbs were identified. Pooled prevalence of CAM use was 51%. A wide variation in prevalence rates (predictive interval 8–93%) was observed. In the context of high heterogeneity, we found no evidence that CAM use was associated with gender, chronicity or type of diabetes. Approximately one third of patients did not disclose their use of CAM to healthcare professionals (95% PrI 25%, 97%). Herbal medicines, acupuncture, homoeopathy and spiritual healing were the common CAM types reported.

Conclusions A wide variation in prevalence of CAM use by patients with diabetes was identified. Healthcare professionals should be aware of their patients' use of CAM to ensure treatment optimization, avoid herb–drug interactions and promote medication adherence in diabetes. Diabetic reviews and clinical guidelines should incorporate exploration of patient use of CAM as many patients do not proactively disclose the use of CAM to their healthcare professionals.

Registration The protocol for this study was registered with the Centre for Review and Dissemination (CRD). Protocol registration number CRD42019125036.

Keywords Prevalence · Complementary and alternative medicine · Diabetes · Systematic review

✉ Vibhu Paudyal
v.paudyal@bham.ac.uk

¹ School of Pharmacy, College of Medical and Dental Sciences, Sir Robert Aitken Institute for Medical Research, University of Birmingham, Birmingham, UK

² Institute of Applied Health Research, University of Birmingham, Birmingham, UK

³ NIHR Birmingham Biomedical Research Centre, University Hospitals Birmingham NHS Foundation Trust and University of Birmingham, Birmingham, UK

⁴ University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

Introduction

The World Health Organization (WHO) estimates that over 400 million people have diabetes worldwide, and this is projected to increase to reach 592 million by 2035 [1]. Poorly managed diabetes can lead to serious and possibly fatal complications such as cardiovascular disease, renal failure, nerve damage and blindness [2–4].

Diabetes mellitus (DM) is a chronic metabolic disorder in which blood glucose levels are higher than normal for a long period of time. These high blood glucose levels are attributed to abnormal disturbances of insulin production and/or function [5]. Diabetes is caused by either lack of insulin production by the pancreas (type 1 diabetes, T1D), when the amount of insulin produced by the pancreas is insufficient to carry out all blood glucose regulation processes, or by decreased insulin

sensitivity by the body cells (type 2 diabetes, T2D). Diabetes can also be caused by a combination of low insulin production as well as low insulin sensitivity or be due to hormonal dysregulation in pregnancy [5].

Self-care practices relevant to self-management of diabetes include adherence to prescribed treatment and clinical management plans, adopting a healthy lifestyle and having a balanced diet [6]. In addition, many patients also use complementary and alternative medicine (CAM) [7]. The WHO defines CAM as a ‘broad set of health care practices that are not part of that country’s own tradition or conventional medicine and are not fully integrated into the dominant health-care system’ [8].

CAM use is known to be prevalent in patients with diabetes as a supplement to their existing orthodox diabetes treatments, as a replacement, or for reasons that might not be directly related to diabetes such as using CAM for energy and general wellbeing [7]. Various factors may influence CAM use by patients with diabetes. A study of 3978 U.S. adults suggested that CAM use by patients who were diabetic for more than 10 years or patients who had a functional limitation caused by diabetes were more likely to use CAM compared to patients with less severe diabetes [9]. In addition, the study reported that 77% of patients who used CAM for the treatment of diabetes used CAM as a supplement to conventional treatment, while 23% used CAM as a replacement [9].

CAM users often perceive CAM to be an effective means of lowering blood glucose levels and treating side effects of prescribed diabetic medications [10–15]. However, adverse outcomes of CAM use have also been reported. For example, CAM can affect the management of diabetes by either direct herb–drug interaction with the use of herbal remedies or indirectly by affecting medication adherence when using herbal or any other CAM types [6].

There is a lack of an up-to-date systematic review that investigates the prevalence of CAM use by patients with diabetes. Patient sources of health-related information have changed immensely in the past decade [16]. In particular, increasing availability and use of web-based information sources, including social media and online health information in recent years, may encourage and inhibit CAM use in long term health conditions [17]. An up-to date systematic review on the prevalence of CAM use by patients with diabetes will help healthcare professionals to consider patient use of CAM when counselling patients, supporting adherence and identifying the risks of interactions and adverse effects when CAMs are used in conjunction with prescribed treatments.

The aim of this study was to systematically review the global prevalence of CAM use amongst adults with diabetes. Specific objectives were to identify the types of CAM that are used by the population with diabetes and to identify differences in CAM use amongst different populations with diabetes, including types of

diabetes, demographic characteristics, duration of diabetes and presence or absence of diabetic complications.

Methods

This systematic review was informed by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and checklist [18]. A protocol was developed as per the PRISMA protocol guideline (protocol ID CRD42019125036).

Data sources and searches

Cochrane Library, MEDLINE, Embase, CINAHL, AMED, Web of Science, Google Scholar and PROSPERO databases were searched for the past 10 years covering 2009 to June 2019. Open Grey was searched for grey literature. Search terms and an example search strategy are listed in Supplementary Table 1. The review was restricted to studies published in English. Studies that recruited participants who are adult patients with diabetes, 18 years of age and older and reported partially or exclusively the prevalence and use of CAM amongst patients with diabetes were included. Studies which either focused on CAM use in conjunction with conventional treatments or as a replacement were considered.

Study selection

Screening and selection were performed independently by two review authors (AA, VP) and were carried out in three phases. Titles and abstracts were screened for inclusion of possible relevant studies followed by assessment of full texts for eligibility. Reference lists of included studies were screened. If a title was considered relevant; the study was manually searched and the abstract examined.

Data extraction and quality assessment

Data on study characteristics, prevalence of CAM use as well as types of CAM used by patients with diabetes were extracted. Two review authors (AA, VP) independently assessed the quality of included studies using the critical appraisal tool from the Joanna Briggs Institute (JBI) checklist [19]. Studies were classified into high, moderate and low quality based on the results of the JBI checklist (Supplementary Table 1). The quality assessment in included studies was focused on three fields: clarity of inclusion criteria and study setting and sampling, appropriateness of approaches to data collection and analysis, and outcome measurement (i.e. use of CAM). Included studies were judged to be of ‘high quality’ if quality criteria were satisfied by at least 7 items, ‘moderate quality’ for scores

of 3–6 and ‘low quality’ for scores ≤ 2 [20]. All studies were included regardless of their quality.

Data synthesis and analysis

A quantitative synthesis of aggregate level data on prevalence was performed. Study specific results were reported as percentage prevalence with exact 95% confidence intervals (95%CI). When sufficient data were available for within-study comparisons of prevalence between dichotomous groups, e.g. sex, then relative prevalence ratios (PRs) together with 95% confidence intervals were calculated. Meta-analyses of proportions and PRs were performed using a random effects model fit using the method of Der Simonian & Laird [21]. Heterogeneity was assessed using the I^2 statistic, the between study standard deviation and calculation of 95% prediction intervals (95%PrI) for the prevalence in a new study [22, 23]. Data are presented in forest plots which include pooled estimates where appropriate. All analyses were performed using STATA version 15.

Results

A total of 2623 unique titles were screened of which 38 articles met the inclusion criteria (Fig. 1). After applying quality assessment, studies fell into these categories (8 high quality studies, 30 moderate quality studies and no low quality studies). Details of critical appraisal results are available in Supplementary Table 2.

Study characteristics

Included studies originated from 25 different countries. Participants were mostly recruited from diabetes clinics and healthcare centres (Table 1). Fifteen of the studies enrolled participants with either T1D or T2D, and 23 studies only included patients with T2D. Out of the included 38 studies, 37 were cross-sectional surveys and one analysed data from another cohort study (Table 1).

Fig. 1 Prisma flow diagram




Table 1 Study characteristics

Author and year	Country of study	Focus of the study	Study settings and recruitment of participants	Study design	Data collection method	Study participants
Alami et al. 2015 [24]	Morocco	Herbal supplements only	Mohammad VI university hospital, Oujda	Cross-sectional	Face-to-face interview using a semi-structured questionnaire	T1D and T2D patients
Al-Eidi et al. 2016 [25]	Saudi	Any CAM type	Diabetic Centre of King Salman bin Abdul-Aziz Hospital, in Riyadh city	Cross-sectional	Face-to-face interview using a structured questionnaire	T2D patients
Al-garni et al. 2017 [26]	Saudi	Herbal supplements only	Jeddah Diabetic Centre (JDC)	Cross-sectional	Interviewer-administered semi-structured questionnaire	T2D patients
Ali-Shtayeh et al. 2012 [14]	Palestine	Any CAM type	Patients attending outpatient departments at Governmental Hospitals in 7 towns in the Palestinian territories (Jenin, Nablus, Tulkarm, Qalqilia, Tubas, Ramalla, and Hebron)	Cross-sectional	Structured questionnaires	T1D and T2D patients
Amaeze et al. 2018 [27]	Nigeria	Herbal supplements only	5 secondary healthcare facilities across Lagos State	Cross-sectional	Interviewer-administered questionnaires	T2D patients
Andrews et al. 2018 [28]	Guatemala	Any CAM type	Interview three groups in the San Lucas Tolimán area	Cross-sectional	Semi-structured questionnaires	T2D patients Health promoters Traditional healers
Ashur et al. 2017 [29]	Libya	Any CAM type	National Centre for Diabetes and Endocrinology in Tripoli	Cross-sectional	Self-administered structured questionnaire	T2D patients
Avci et al. 2018 [30]	Turkey	Any CAM type	Van Yuzuncu Yil University, Van	Cross-sectional	Semi-structured questionnaires	T1D and T2D patients
Azizi-Fini et al. 2016 [11]	Iran	Herbal supplements only	Golabchi and Naqavi diabetes clinics in the Kashan city	Cross-sectional	Interviewer-administered structured questionnaires	T2D patients
Baharom et al. 2016 [31]	Malaysia	Any CAM type	45 government health clinics across Nigeria Sembilan	Cross-sectional	Interviewer-administered structured questionnaires	T2D patients
Bradley et al. 2012 [13]	USA	Any CAM type	Patients with moderately to poorly controlled T2D who receive care from Group Health Cooperative (GHC), a large non-profit, integrated health care system in Washington State	Cross-sectional	Telephone-administered questionnaires.	T2D patients
Candar et al. 2018 [32]	Turkey	Any CAM type	Patients registered with the Bursa Yuksek Ihtisas Training and Research Hospital Education Family Health Centre	Cross-sectional	Questionnaires	T1D and T2D patients
Chao et al. 2014 [33]	USA	Any CAM type	Patients who received primary care at one of four publicly funded clinics in the Community Health Network of San Francisco	Cross-sectional	Data collected for the Self-Management Automated and Real-Time Telephonic Support (SMART Steps) Study	T2D patients
Ching et al. 2013 [34]	Malaysia	Any CAM type	Primary health care clinic at Salak in Sepang	Cross-sectional	Face-to-face interview using a structured questionnaire	T2D patients
Damnjanovic et al. 2015 [35]	Serbia	Herbal supplements only	6 Remedia Pharmacy HealthFacilities in the territory of Nis	Cross-sectional	Structured questionnaires	T2D patients
	India	Any CAM type		Cross-sectional	Structured questionnaires	T2D patients

Table 1 (continued)

Author and year	Country of study	Focus of the study	Study settings and recruitment of participants	Study design	Data collection method	Study participants
Devi et al. 2015 [36]			Diabetes Health camp conducted by VS micro lab, Madurai			
Fabian et al. 2011 [37]	Austria	Herbal supplements only	Diabetes Centre of the Division of Endocrinology and Metabolism, Department of Internal Medicine, Medical University of Graz	Cross-sectional	Face-to-face interview using a structured questionnaire	T1D and T2D patients
Fan et al. 2013 [38]	Singapore	Any CAM type	Single centre study conducted in an outpatient diabetes Centre with an average load of 2500 patients a month	Cross-sectional	Self-administered questionnaires.	T2D patients
Hashempur et al. 2015 [39]	Iran	Any CAM type	Two outpatient diabetes clinics affiliated with the Shiraz University of Medical Sciences, Shiraz	Cross-sectional	Face-to-face interview using semi-structured questionnaire	T1D and T2D patients
Kamel et al. 2017 [40]	Saudi	Herbal supplements only	King Abdul-Aziz University and King Fahad General Hospitals in Jeddah	Cross-sectional	Interviewer-administered structured questionnaires	T1D and T2D patients
Karaman et al. 2018 [10]	Turkey	Herbal supplements only	Endocrinology clinics of two hospitals in Izmir	Cross-sectional	Face-to-face interview using a structured questionnaire	T1D and T2D patients
Khalaf and Whitford 2010 [41]	Bahrain	Any CAM type	Patients attending two hospital diabetes clinics	Cross-sectional	Questionnaires (administration not detailed)	T1D and T2D patients
Khalil et al. 2013 [42]	Egypt	Herbal supplements only	Outpatient clinics of Alexandria University Hospital, from seven health insurance centres, six MOH hospitals, and one private healthcare facility.	Cross-sectional	Questionnaires (administration method not reported)	T2D patients
Koren et al. 2015 [43]	Israel	Herbal supplements only	Internal medicine department at Assaf Harofeh Medical Centre, Zerifin	Cross-sectional	Interviewer-administered structured questionnaires	T2D patients
Lui et al. 2012 [44]	Australia	Any CAM type	Data reported here are taken from the Living with Diabetes Study (LWDS), a five-year, prospective cohort study being conducted in the State of Queensland	Data from cohort study	Questionnaires (administration not detailed)	T1D and T2D patients
Lunyera et al. 2016 [45]	Tanzania	Herbal supplements only	Kilimanjaro Region of Tanzania	Cross-sectional	Verbally administered structured questionnaire	T1D and T2D patients
Medagama et al. 2014 [46]	Sri Lanka	Herbal supplements only	Diabetes clinic at Teaching Hospital Peradeniya	Cross-sectional	Face-to-face interview using a structured questionnaire	T2D patients
Mekuria et al. 2018 [47]	Ethiopia	Herbal supplements only	Diabetes care clinic of University of Gondar comprehensive specialized hospital	Cross-sectional	Interviewer-administered questionnaires	T2D patients
Mohamed Ali and	Sudan	Herbal supplements only	125 primary health care centres in Khartoum	Cross-sectional	Interviewer-administered questionnaires	T2D patients

Table 1 (continued)

Author and year	Country of study	Focus of the study	Study settings and recruitment of participants	Study design	Data collection method	Study participants
Mahfouz 2014 [48] Naja et al. 2014 [49]	Lebanon	Any CAM type	Patients recruited from two major referral centres in Beirut- a public hospital and a private academic medical Centre	Cross-sectional	Face-to-face interview using a structured questionnaire	T2D patients
Nguyen et al. 2014 [50]	USA	Any CAM type	Patients recruited from seven primary care or endocrinology clinics affiliated with an academic medical centre in Southern California	Cross-sectional	Self-administered structured questionnaire	T2D patients
Putthapiban et al. 2017 [51]	Thailand	Herbal supplements only	At the Endocrine Clinic in Ramathibodi Hospital, Bangkok	Cross-sectional	Face-to-face interview using a structured questionnaire	T2D patients
Rhee et al. 2018 [52]	USA	Any CAM type	Non-institutionalized civilians in US	Cross-sectional	Data were from the 2012 NHIS, which was administrated by the National Centre for Health Statistics of the Centers for Disease Control and Prevention	T1D and T2D patients
Sethi et al. 2011 [12]	India	Any CAM type	Tertiary care Centre in Delhi	Cross-sectional	Face-to-face interview using a Semi-structured questionnaire	T1D and T2D patients
Vishnu et al. 2017 [53]	India	Any CAM type	Rural Kollam district of the Indian state of Kerala (community based)	Cross-sectional	Interviewer-administered structured questionnaires	T1D and T2D patients
Wanchai and Phrompayak 2016 [54]	Thailand	Any CAM type	Four primary healthcare units and two secondary hospitals in the north of Thailand	Cross-sectional	Semi-structured questionnaire	T2D patients
Wazaify et al. 2011 [15]	Jordan	Herbal supplements only	Outpatient departments at The National Centre for Diabetes, Endocrine and Genetics (NCDEG).	Cross-sectional	Face-to-face interview using a Semi-structured questionnaire	T1D and T2D patients
Yildirim and Marakoglu 2018 [55]	Turkey	Any CAM type	Outpatient diabetes from Selçuk University Family Medicine Diabetes Education Clinic	Cross-sectional	Face-to-face interview using a structured questionnaire	T2D patients

CAM Complementary and alternative medicine

Types of CAM

Sixteen studies focused exclusively on herbal and nutritional supplement use by patients with diabetes (Table 1). The remaining 22 studies discussed other CAM types. Fourteen of those 22 studies that investigated other CAM types also reported the use of herbal and nutritional supplements as a form of CAM [12–14, 25, 32, 34, 36, 38, 39, 41, 49, 52–54]. A total of 35 different CAM types were reported in at least one study. CAM types used by patients with diabetes and mentioned in the most studies were acupuncture ($n = 6$ studies), Mind–body

therapies ($n = 6$ studies) religious and spiritual healing ($n = 5$ studies) and homoeopathy ($n = 4$ studies) (Table 2).

Within the 31 studies which reported the use of herbal and nutritional supplements by patients with diabetes, a total of 223 different herbal and nutritional supplements were reported (Supplementary Table 3). The five herbs that were mentioned in the most studies were, cinnamon (*Cinnamomum verum*) and fenugreek (*Trigonella foenum-graecum*) each reported in 18 different studies, garlic (*Allium sativum*) reported in 17 studies, aloe vera (*Aloe Vera*) reported in 14 studies and black seed (*Nigella sativa*) reported in 12 studies.

Table 2 List of complementary and alternative medicine types as cited by included studies

CAM forms (other than herbal supplements)	Studies cited the CAM form	CAM forms (other than herbal supplements)	Studies cited the CAM form
Acupuncture	[25, 29, 36, 39, 50, 52]	Ruqyah (recitation) with the Quran	[25, 29]
Mind–body therapies	[32, 34, 36, 39, 41, 52]	Ruqyah water or oil	[25, 29]
Religious and spiritual healing	[29, 32, 49, 50, 54]	Balneotherapy	[32]
Homoeopathy	[12, 36, 52, 53]	Biofeedback	[52]
Meditation	[13, 36, 52, 54]	Chelation	[52]
Massage	[13, 25, 38, 49]	Chinese medicine	[49]
Ayurveda	[36, 52, 53]	Curandero	[50]
Chiropractic Massage	[13, 50, 52]	Daode Xinxi	[54]
Energy therapies	[34, 41, 52]	Deep breathing exercises	[13]
Specific diet	[13, 25, 36]	Leech (Hirudotherapy)	[32]
Yoga	[13, 36, 52]	Music therapy	[36]
Al-hijama (wet cupping)	[25, 29]	Prayer by religion person (imam)	[30]
Biologically based therapies	[36, 52]	Progressive muscle relaxation	[13]
Cupping	[32, 39]	Qi gong	[52]
Folk medicine	[13, 49]	Sugar therapy	[53]
Honey	[14, 25]	Tai chi	[52]
Movement therapies	[36, 52]	Traditional healers	[52]
Naturopathy	[50, 52]		

CAM Complementary and alternative medicine

Prevalence of CAM use

The highest prevalence of CAM (all types) use was reported at 89% by two studies, one each from India and Jordan followed by studies in Tanzania (78%), Sri Lanka (76%) and Iran (75%) (Table 3) [12, 35, 39, 45, 46]. The lowest prevalence of CAM use was 17% as reported by a study conducted in Jordan [15]. A study in Australia reported a prevalence of 8%, but the study gathered data from patients about their visits to CAM practitioners only and did not include data on CAM use in general by patients with diabetes [44]. Other studies reporting the lowest prevalence of CAM use included studies in Libya (29%), Saudi Arabia (26%), USA (26%), Israel (23%) and Jordan (17%) [15, 26, 29, 43, 52]. Pooled prevalence of CAM use was 51% (95%CI 43%, 59%). However, heterogeneity was very high ($I^2 = 99%$) with the predictive interval ranging from 8% to 93%. (Fig. 2).

Subgroup analysis

Study level factors

Meta-analysis was conducted for results stratified at the study level by continent. I^2 was 97.5% and predictive intervals were

found to be wide. The highest prevalence rates of 76% were observed in Europe (PrI inestimable), followed by Africa 55%, (95%PrI 0.17, 0.90) from seven studies. The lowest prevalence rates were observed in North America 45%, (95%PrI 0.04, 0.92) from five studies (Fig. 3).

Patient level factors

Subgroup analyses were conducted across ethnicity (reported in eight studies). All meta-analyses at subgroup level also showed high levels of heterogeneity. Results were as follows: for the ethnicity subgroup, no predictive interval could be estimated other than PrI for the group of ‘other ethnicities’ prevalence ratio 0.57 (95%CI 0.39–0.75); the estimated predictive intervals ranged between 0.00 and 1.00, $I^2 = 64.05%$ (Fig. 4).

For analysis stratified by binary subgroups within-study comparative data were extractable for sex (31 studies), type of diabetes (7 studies), duration of diabetes (15 studies) and presence or absence of diabetic complications (10 studies). Within study pooled estimates PRs for patients with no diabetic complications versus patients with diabetic complications gave a prevalence ratio (PR) 0.81 (95%CI 0.66, 0.99), (95%PrI 0.39–1.67) ($I^2 = 89%$) (Supplementary Fig. 1). For

Table 3 Included studies' overall and subgroups prevalence of CAM use


Country	Study	Sample size	Prevalence of CAM use	All participants		Female		Male		T1D		T2D		Had diabetes for ≤5y		Had diabetes for >5y	
				users	non users	users	non users	users	non users	users	non users	users	non users	users	non users	users	non users
India	Sethi et al. 2011 [12]	113	89.38%	101	12	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Serbia	Dannjanovic et al. 2015 [35]	519	88.82%	461	58	261	15	95%	200	43	82%	NR	NR	NR	NR	NR	NR
Tanzania	Lunyera et al. 2016 [45]	45	77.78%	35	10	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Sri Lanka	Medagama et al. 2014 [46]	252	76.19%	192	60	139	28	83%	53	32	62%	NR	NR	NR	NR	NR	NR
Iran	Hashempur et al. 2015 [39]	239	75.31%	180	59	124	37	77%	56	22	72%	10	7	59%	170	52	77%
Thailand	Wanchai and Phrompayak 2016 [54]	508	70.87%	360	148	282	102	73%	78	46	63%	NR	NR	NR	NR	NR	NR
Nigeria	Amaeze et al. 2018 [27]	453	67.33%	305	148	98	45	69%	207	103	67%	NR	NR	NR	96	66	59%
India	Devi et al. 2015 [36]	252	64.29%	162	90	98	41	71%	64	49	57%	NR	NR	NR	55	40	58%
Saudi	Kamel et al. 2017 [40]	214	64.02%	137	77	84	44	66%	53	33	62%	50	20	71%	87	57	60%
Guatemala	Andrews et al. 2018 [28]	55	63.64%	35	20	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Bahrain	Khalaf and Whitford 2010 [41]	402	62.69%	252	150	149	69	68%	103	81	56%	NR	NR	NR	48	51	48%
Malaysia	Ching et al. 2013 [34]	240	62.50%	150	90	96	49	66%	54	41	57%	NR	NR	NR	NR	NR	NR
Ethiopia	Mekuria et al. 2018 [47]	387	62.02%	240	147	149	73	67%	91	74	55%	NR	NR	NR	168	60	74%
Thailand	Putthapiban et al. 2017 [51]	200	61.00%	122	78	76	42	64%	46	36	56%	NR	NR	NR	NR	NR	NR
Sudan	Mohamed Ali and Mahfouz 2014 [48]	600	58.00%	348	252	206	167	55%	142	85	63%	NR	NR	NR	67	76	47%
Turkey	Karaman et al. 2018 [10]	455	57.58%	262	193	225	148	60%	37	45	45%	53	49	52%	209	114	65%
Iran	Azizi-Fini et al. 2016 [11]	500	56.20%	281	219	203	153	57%	78	66	54%	NR	NR	NR	NR	NR	NR
Morocco	Alami et al. 2015 [24]	279	54.84%	153	126	117	83	59%	36	43	46%	36	43	46%	117	83	59%
Palestine	Ali-Shtayeh et al. 2012 [14]	1883	51.89%	977	906	519	470	52%	458	436	51%	114	84	58%	863	822	51%
Malaysia	Baharom et al. 2016 [31]	680	49.41%	336	344	224	175	56%	112	169	40%	NR	NR	NR	NR	NR	NR
USA	Bradley et al. 2012 [13]	219	48.40%	106	113	47	50	48%	59	63	48%	NR	NR	NR	NR	NR	NR

Table 3 (continued)

Country	Study	Sample size	Prevalence of CAM use	All participants		Female		Male		T1D		T2D		Had diabetes for ≤5y		Had diabetes for >5y	
				users	non users	users	non users	users	non users	users	non users	users	non users	users	non users	users	non users
USA	Chao et al. 2014 [33]	278	47.48%	132	146	101	38	73%	31	108	22%	NR	NR	NR	NR	NR	NR
Turkey	Avci., 2018 [30]	386	46.37%	179	207	95	122	44%	84	85	50%	16	64%	150	191	44%	68
USA	Nguyen et al. 2014 [50]	410	45.85%	188	222	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Turkey	Candar et al. 2018 [32]	442	44.80%	198	244	137	134	51%	61	110	36%	NR	NR	NR	NR	NR	NR
Singapore	Fan et al., 2013 [38]	304	43.42%	132	172	67	69	49%	65	103	39%	NR	NR	NR	NR	NR	NR
Egypt	Khalil et al. 2013 [42]	1100	41.73%	459	641	252	359	41%	207	282	42%	NR	NR	87	202	30%	372
India	Vishnu et al. 2017 [53]	400	38.75%	155	245	73	109	40%	82	136	38%	NR	NR	95	142	40%	60
Lebanon	Naja et al. 2014 [56]	333	38.14%	127	206	51	98	34%	76	108	41%	NR	NR	34	77	31%	93
Turkey	Yildirim and Marakoglu 2018 [55]	400	36.75%	147	253	91	115	44%	56	138	29%	NR	NR	77	149	34%	70
Austria	Fabian et al. 2011 [37]	198	31.31%	62	136	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Saudi	Al-Eidi et al. 2016 [25]	302	30.46%	92	210	50	121	29%	42	89	32%	NR	NR	24	93	21%	68
Libya	Ashur et al. 2017 [29]	523	28.87%	151	372	102	206	33%	49	166	23%	NR	NR	86	226	28%	65
USA	Rhee et al. 2018 [52]	3386	26.17%	886	2500	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Saudi	Al-garni et al. 2017 [26]	310	25.81%	80	230	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Israel	Koren et al. 2015 [43]	111	23.42%	26	85	12	37	24%	14	48	23%	NR	NR	NR	NR	NR	NR
Jordan	Wazaify et al. 2011 [15]	1000	16.60%	166	834	99	432	19%	67	402	14%	8	15%	158	790	17%	NR
Australia	Lui et al. 2012 [44]	3337	7.73%	258	3079	157	1727	8%	101	1352	7%	NR	NR	NR	NR	NR	NR

CAM Complementary and alternative medicine, NR Not reported

Fig. 2 Forest plot showing pooled prevalence of complementary and alternative medicine in diabetes



patients who had had diabetes for more than 5 years versus less than 5 years pooled PR was 1.71 (95%CI 1.04, 1.32), (95%PrI 0.73, 1.88) (I^2 of 83%) (Supplementary Fig. 2). For male versus female participants, pooled PR was 0.86 (95%CI 0.81, 0.91), (95%PrI 0.64, 1.16) (I^2 of 72%) (Supplementary Fig. 3). Pooled PR for patients with T2D versus T1D patients was 1.00 (95%CI 0.83, 1.20), 95%PrI 0.56, 1.77) (I^2 = 75%) (Supplementary Fig. 4).

Additional outcomes

CAM as a complementary or alternative treatment Eight of the 38 included studies assessed whether CAM was used as an additional treatment or as an alternative treatment to conventional medicines. Prevalence of CAM use as an additional treatment to prescribed medicine was 78% (95%CI 56%, 94%) with 95% PrI (4%, 1.00%) (I^2 = 98%), and the

percentage of patients who used CAM as an alternative to their prescribed medicine was 21% (95%CI 12%, 31%) with 95% PrI (0%, 63%) (I^2 = 89%) (Supplementary Figs. 5, 6).

Patients' disclosure of CAM use to healthcare professionals

The percentage of patients who do not disclose their CAM use to healthcare professionals was 67% (95%CI 58%, 76%) with 95% PrI (25%, 97%) (I^2 = 98%) (Fig. 5).

Discussion

This study provides up-to-date data on the global prevalence of CAM use by patients with diabetes as reported in the peer reviewed research literature. The last literature review on CAM use of patients with diabetes was published in 2007 [7] which reviewed studies conducted in nine countries and




Fig. 3 Meta-analysis of study level factors in relation to CAM use (prevalence proportion by continent). CAM, complementary and alternative medicine

reported prevalence ranging from 17% to 73%. A similarly wide variation in prevalence rate of 8–89% was observed in our updated review that included studies from 25 countries.

According to the included studies, CAM use is common amongst patients with diabetes for the purpose of diabetes management. Most of the studies showed that the participants they recruited used CAM as an additional approach to conventional treatment, while in other studies, the reason for CAM use (additional or alternative) was not specified. Only seven studies reported that some patients with diabetes used

CAM as the sole means of managing their diabetes. Most of the included studies were conducted in healthcare settings. Therefore, patients who do not use conventional treatments for diabetes may not have been included. The prevalence of patients with diabetes who use CAM in the general population with diabetes is hence likely to be higher than the estimates provided by the included studies.

The meta-analysis of the prevalence data demonstrated extreme variation in prevalence of CAM use amongst patients with diabetes across studies.





Fig. 4 Ethnicity subgroup forest plot showing prevalence of CAM use. CAM, Complementary and alternative medicine

Strengths and limitations

This systematic review was a protocol driven review with a pre-specified aim, objectives and methodology. A range of relevant databases were used which covered the prevalence of CAM used by patients with diabetes globally. Data collection methods varied amongst studies. Some studies used

structured questionnaires while other studies used semi-structured questionnaires. In addition there was a wide variation in the nature of the study settings. The content of the questionnaires used to collect the data on prevalence and nature of CAMs are likely to influence patient response. Therefore, the included studies may have underreported the nature and extent of CAM use by study participants. In

Fig. 5 Patient who do not disclose the use of CAM to the healthcare professionals. CAM, complementary and alternative medicine



addition, our study only included studies published in the English language.

Implications for practice and research

This systematic review shows that CAM use amongst patients with diabetes is prevalent in many populations. This review suggests that healthcare professionals should consider use of CAM by patients with diabetes when advising them about using their prescribed treatments and monitor their medication adherence while using any forms of CAM. They should also be aware of patients' use of herbal supplements as some forms of herbal medicine can lead to herb–drug interactions [57]. For example, the most frequently mentioned herbal supplement used by patients with diabetes was cinnamon. It is reported that cinnamon has a potentiating effect on diabetic drugs increasing the risk of hypoglycaemia [58]. For example, cinnamon shows an inhibitory effect on the CYP3A4 enzyme in rabbits which potentiates the effect of pioglitazone if combined with cinnamon, leading to a hypoglycaemic effect [59]. *Aloe vera*, which was the most frequently reported CAM by the studies included in our review, has been linked to potential interaction with 45 different drugs, including diabetic drugs such as glimepiride [60]. Concomitant use of *Aloe vera* and glimepiride can produce hypoglycaemic effects as *Aloe vera* has an inhibiting effect on ATP sensitive potassium channels in pancreatic β cells leading to additional release of insulin [61].

Understanding CAM use patterns and considering any possible interactions between them and other potential medications could help healthcare professionals to appropriately

minimize drug-related problems or herb–drug interactions. It could help them to encourage their patient to discuss their CAM use and offer the opportunity to provide better advice for patients with diabetes.

The observed prevalence of CAM, and the many varieties of CAM that are used by patients with diabetes, call for revision of diabetes management guidelines. The National Institute for Health and Care Excellence (NICE) guideline on management of diabetes does not explicitly advise healthcare professionals to discuss patient use of herbal medicines or CAM in their consultation [6]. Guidelines should enable healthcare professionals to counsel patients with diabetes, their families and carers, who should all be educated about the safe use of CAM in conjunction with prescribed medicines.

Owing to the variable and often high prevalence of CAM use amongst patients with diabetes worldwide, research that generates evidence-based information about CAM is needed. This includes effectiveness and safety profiles of commonly used CAMs, including herbal medicines as identified in this systematic review.

This systematic review has identified that on average up to two-thirds of patients who use CAM do not disclose this to their healthcare professionals. Use of CAM such as herbal medicines could be incorporated as part of the comprehensive medication review services offered at community pharmacies and primary care [62].

Future studies need to consider the perspectives of patients with diabetes who do not visit conventional healthcare facilities for the management of diabetes to provide a better estimate of prevalence rates. In addition, there is a need to gather

evidence on the factors that affect use and non-use of CAM by patients with diabetes.

Conclusion

A wide variation in prevalence rate of CAM use in diabetes (8–89%) was observed, and pooled prevalence of CAM use was 51%. Our findings show that CAM use by patients with diabetes is common. Healthcare professionals should be aware of the use of CAM by patients with diabetes to ensure treatment optimization and medication adherence. Future studies should incorporate patient and healthcare professionals' perspectives of CAM use in diabetes, evaluate patient outcomes through the use of healthcare databases and carefully designed prospective studies, and identify opportunities to promote rational use of CAM through evidence-based guidelines and patient-centred approaches.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00228-021-03097-x>.

Acknowledgments This paper presents independent research supported by the NIHR Birmingham Biomedical Research Centre at the University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham in relation to MJP involvement. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. We would like to thank University of Birmingham Library Services for assisting with the literature search.

Authors contributions This study relates to AA's PhD. AA, SG and VP designed the study. VP and SG supervised AA's PhD. MJP led the statistical analysis and presentation of quantitative results. AA led the drafting of the manuscript to which all authors contributed through editing and revision. All authors had access to the data sets and agreed to the final version of the manuscript.

Funding This study was funded by University of Birmingham. AA was supported for his PhD studies by the Government of Saudi Arabia.

Data availability All data in relation to this study are presented in this manuscript.

Code availability Not applicable.

Declarations

Ethics approval statement Not applicable.

Patient consent statement Not applicable.

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. World Health Organization. WHO Global Report on Diabetes. <https://www.who.int/publications/i/item/9789241565257>. Accessed 26 Nov 2020
2. Moneta G (2011) Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Yearbook Vasc Surg* 17(10):49–51
3. Lim A (2014) Diabetic nephropathy: complications and treatment. *Int J Nephrol Renov Dis* 2014:361–381
4. Solomon S, Chew E, Duh E, Sobrin L, Sun J, VanderBeek B, Wykoff C, Gardner T (2017) Diabetic retinopathy: a position statement by the American Diabetes Association. *Diabetes Care* 40(3): 412–418
5. Gardner DG, Shoback DM (2017) Greenspan's basic and clinical endocrinology. McGraw-Hill Education, New York
6. National Institute of Clinical Excellence. NICE guideline: type 2 diabetes in adults: management NG28 2015. <https://www.nice.org.uk/guidance/ng28>. Accessed 26 Nov 2020
7. Chang H, Wallis M, Tiralongo E (2007) Use of complementary and alternative medicine among people living with diabetes: literature review. *J Adv Nurs* 58(4):307–319
8. World Health Organization (2018) Traditional, complementary and integrative medicine. https://www.who.int/health-topics/traditional-complementary-and-integrative-medicine#tab=tab_1. Accessed 26 Nov 2020
9. Nahin RL, Byrd-Clark D, Stussman BJ, Kalyanaraman N (2012) Disease severity is associated with the use of complementary medicine to treat or manage type-2 diabetes: data from the 2002 and 2007 National Health Interview Survey. *BMC Complement Altern Med* 12:193
10. Karaman E, Erkin O, Senman S, Yildirim Y (2018) The use of herbal supplements by individuals with diabetes mellitus. *J Pak Med Assoc* 68(4):587–594
11. Azizi-Fini I, Adib-Hajbaghery M, Gharehbaghloou Z (2016) Herbal medicine use among patients with type 2 diabetes in Kashan, Iran, 2015. *European J Integrat Med* 8(4):570–575
12. Sethi A, Srivastava S, Madhu SV (2011) Prevalence and pattern of use of indigenous medicines in diabetic patients attending a tertiary care Centre. *J Indian Med Assoc* 109(7):469–471
13. Bradley R, Sherman K, Catz S, Calabrese C, Oberg E, Cherkin D (2012) Self-care, use of CAM and satisfaction with health care in people with inadequately controlled type 2 diabetes. *BMC complementary and alternative medicine. International Research Congress on Integrative Medicine and Health* 12(SUPPL.1)
14. Ali-Shtayeh MS, Jamous RM, Jamous RM (2012) Complementary and alternative medicine use amongst Palestinian diabetic patients. *Complement Ther Clin Pract* 18(1):16–21
15. Wazaify M, Afifi FU, El-Khateeb M, Ajlouni K (2011) Complementary and alternative medicine use among Jordanian patients with diabetes. *Complement Ther Clin Pract* 17(2):71–75

16. Amante D, Hogan T, Pagoto S, English T, Lapan K (2015) Access to care and use of the internet to search for health information: results from the US National Health Interview Survey. *J Med Internet Res* 17(4):e106
17. Sharma V, Holmes JH (2016) And S. IN., identifying complementary and alternative medicine usage information from internet resources. A systematic review. *Methods Inf Med* 55(4):322–332
18. Moher D, Liberati A, Tetzlaff J, Altman DG (2009) The PRISMA group, preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 151(4)
19. Janna Briggs Institute (2019) Critical appraisal tools, The Joanna Briggs Institute The University of Adelaide, Australia. <https://joannabriggs.org/critical-appraisal-tools>. Accessed 26 Nov 2020
20. Janna Briggs Institute (2017) Joanna Briggs Institute reviewer's manual. The Joanna Briggs Institute. <https://nursing.lsuhs.edu/JBI/docs/ReviewersManuals/Scoping-pdf>. Accessed 26 Nov 2020
21. DerSimonian R, Laird N (1986) Meta-analysis in clinical trials. *Control Clin Trials* 7(3):177–188
22. Higgins J, Thompson S, Deeks J, Altman D (2003) Measuring inconsistency in meta-analyses. *BMJ* 327:557–560
23. Higgins J, Thompson S, Spiegelhalter D (2009) A re-evaluation of random-effects meta-analysis. *J R Stat Soc Ser A Stat Soc* 172:137–159
24. Alami Z, Aynaou H, Alami B, Hdidou Y, Latrech H (2015) Herbal medicines use among diabetic patients in oriental Morocco. *J Pharmacogn Phytother* 7(2):9–17
25. Al-Eidi S, Tayel S, Al-Slail F, Qureshi NA, Sohaibani I, Khalil M, Al-Bedah AM (2016) Knowledge, attitude and practice of patients with type 2 diabetes mellitus towards complementary and alternative medicine. *J Integrat Med* 14(3):187–196
26. Al-Garni AM, Al-Raddadi RM, Al-Amri TA (2017) Patterns and determinants of complementary and alternative medicine use among type 2 diabetic patients in a diabetic center in Saudi Arabia: herbal alternative use in type 2 diabetes. *J Fund Appl Sci* 9(1):1738–1748
27. Amaeze OU, Aderemi-Williams RI, Ayo-Vaughan MA, Ogundemuren DA, Ogunmola DS, Anyika EN (2018) Herbal medicine use among type 2 diabetes mellitus patients in Nigeria: understanding the magnitude and predictors of use. *Int J Clin Pharm* 40(3):580–588
28. Andrews CM, Wyne K, Svenson JE (2018) The use of traditional and complementary medicine for diabetes in rural Guatemala. *J Health Care Poor Underserved* 29(4):1188–1208
29. Ashur ST, Shah SA, Bosseri S, Shamsuddin K (2017) Use of traditional medicine among type 2 diabetic Libyans. *East Mediterr Health J* 23(5):375–382
30. Avci DK (2018) The use of traditional and complementary medicine among diabetes patients, and the awareness and attitudes of physicians. *J Pak Med Assoc* 68(11):1650–1654
31. Baharom N, Shah SA, Rotina AB (2016) Prevalence of complementary alternative medicine use among patients with type II diabetes in Negeri Sembilan, Malaysia. *Med Health (Universiti Kebangsaan Malaysia)* 11(2):257–266
32. Candar A, Demirci H, Baran AK, Akpınar Y (2018) The association between quality of life and complementary and alternative medicine use in patients with diabetes mellitus. *Complement Ther Clin Pract* 31:1–6
33. Chao MT, Handley M, Quan J, Sarkar U, Ratanawongsa N, Schillinger D (2014) Disclosure of complementary and alternative medicine use among diverse safety net patients with diabetes. *J Altern Complement Med* 20(5):A126
34. Ching SM, Zakaria ZA, Paimin F, Jalalian M (2013) Complementary alternative medicine use among patients with type 2 diabetes mellitus in the primary care setting: a cross-sectional study in Malaysia. *BMC Complement Altern Med* 13:148
35. Damjanovic I, Kitic D, Stefanovic N, Zlatkovic-Guberinic S, Catic-Djordjevic A, Velickovic-Radovanovic R (2015) Herbal self-medication use in patients with diabetes mellitus type 2. *Turkish J Med Sci* 45(4):964–971
36. Devi K, Santhini E, Manikandan R, Prabhu NM (2015) The prevalence, awareness and potential of complementary alternative medicine in type 2 diabetics living in Madurai, India. *European J Integrat Med* 7(5):469–473
37. Fabian E, Toscher S, Elmadfa I, Pieber TR (2011) Use of complementary and alternative medicine supplements in patients with diabetes mellitus. *Ann Nutr Metab* 58(2):101–108
38. Fan PE, Chan MF, Chan YL, Koh SL (2013) Patterns of complementary and alternative medicine use among a group of patients with type 2 diabetes receiving outpatient care in Singapore. *Int J Nurs Pract* 19(Suppl 3):44–55
39. Hashempur MH, Heydari M, Mosavat SH, Heydari ST, Shams M (2015) Complementary and alternative medicine use in Iranian patients with diabetes mellitus. *J Integrat Med* 13(5):319–325
40. Kamel FO, Magadmi RM, Hagraas MM, Magadmi B, AlAhmad RA (2017) Knowledge, attitude, and beliefs toward traditional herbal medicine use among diabetics in Jeddah Saudi Arabia. *Complement Ther Clin Pract* 29:207–212
41. Khalaf AJ, Whitford DL (2010) The use of complementary and alternative medicine by patients with diabetes mellitus in Bahrain: a cross-sectional study. *BMC Complement Altern Med* 10:35
42. Khalil SH, Zaki A, Ibrahim AM, El-Moughazi AM, Khater AM, Youssef AM, El-Sa'ed AT, Rashed EM (2013) Pattern of use of complementary and alternative medicine among type 2 diabetes mellitus patients in Alexandria, Egypt. *J Egypt Public Health Assoc* 88(3):137–142
43. Koren R, Lerner A, Tirosh A, Zaidenstein R, Ziv-Baran T, Golik A, Koren S (2015) The use of complementary and alternative medicine in hospitalized patients with type 2 diabetes mellitus in Israel. *J Altern Complement Med* 21(7):395–400
44. Lui C-W, Dower J, Donald M, Coll JR (2012) Patterns and determinants of complementary and alternative medicine practitioner use among adults with diabetes in Queensland, Australia. *Evid Based Complement Alternat Med* 2012:659419
45. Lunyera J, Wang D, Maro V, Karia F, Boyd D, Omolo J, Patel UD, Stanifer JW (2016) Traditional medicine practices among community members with diabetes mellitus in northern Tanzania: an ethnomedical survey. *BMC Complement Altern Med* 16:1–12
46. Medagama AB, Bandara R, Abeysekera RA, Imbulpitiya B, Pushpakumari T (2014) Use of complementary and alternative medicines (CAMs) among type 2 diabetes patients in Sri Lanka: a cross sectional survey. *BMC Complement Altern Med* 14:374
47. Mekuria AB, Belachew SA, Tegegn HG, Ali DS, Netere AK, Lemlemu E, Erku DA (2018) Prevalence and correlates of herbal medicine use among type 2 diabetic patients in teaching Hospital in Ethiopia: a cross-sectional study. *BMC Complement Altern Med* 18(1):85
48. Mohamed Ali BA, Mahfouz MS (2014) Herbal medicine use among patients with type 2 diabetes in North Sudan. *Annual Res Rev Biol* 4(11):1827–1838
49. Naja F, Alameddine M (2014) Prevalence and correlates of complementary and alternative medicine use among diabetic patients in Lebanon: a cross sectional study. *Clin Nutr* 1:S179–S180
50. Nguyen H, Sorkin DH, Billimek J, Kaplan SH, Greenfield S, Ngo-Metzger Q (2014) Complementary and alternative medicine (CAM) use among non-Hispanic white, Mexican American, and Vietnamese American patients with type 2 diabetes. *J Health Care Poor Underserved* 25(4):1941–1955
51. Putthapiban P, Sukhumthamarat W, Sriprapradang C (2017) Concealed use of herbal and dietary supplements among Thai patients with type 2 diabetes mellitus. *J Diab Metabol Dis* 16:36

52. Rhee TG, Westberg SM, Harris IM (2018) Complementary and alternative medicine in US adults with diabetes: reasons for use and perceived benefits. *J Diab* 10(4):310–319
53. Vishnu N, Mini GK, Thankappan KR (2017) Complementary and alternative medicine use by diabetes patients in Kerala, India. *Glob Health Epidemiol Genom* 2(e6):1–7
54. Wanchai A, Phrompayak D (2016) Use of complementary and alternative medicine among Thai patients with type 2 diabetes mellitus. *J Integrat Med* 14(4):297–305
55. Yildirim DI, Marakoglu K (2018) Complementary and alternative medicine use amongst Turkish type 2 diabetic patients: a cross-sectional study. *Complement Ther Med* 41:41–46
56. Naja F, Mousa D, Alameddine M, Shoaib H, Itani L, Mourad Y (2014) Prevalence and correlates of complementary and alternative medicine use among diabetic patients in Beirut, Lebanon: a cross-sectional study. *BMC Complement Altern Med* 14:185. <https://doi.org/10.1186/1472-6882-14-185>
57. Brantley SJ, Argikar AA, Lin YS, Nagar S, Paine MF (2014) Herb-drug interactions: challenges and opportunities for improved predictions. *Drug Metab Dispos* 42(3):301–317
58. Allen RW, Schwartzman E, Baker WL, Coleman CI, Phung OJ (2013) Cinnamon use in type 2 diabetes: an updated systematic review and meta-analysis. *Ann Fam Med* 11(4):452–459
59. Maminla S, Koganti VSRGP, Ravouru N, Koganti B (2017) Effect of Cinnamomum cassia on the pharmacokinetics and pharmacodynamics of pioglitazone. *Curr Clin Pharmacol* 12(1):41–49
60. Drugs.com (2020) *Aloe vera* drug interactions. 4 May 2020. <https://www.drugs.com/drug-interactions/aloe-vera-index.html>. Accessed 26 Nov 2020
61. Gupta RC, Chang D, Nammi S, Bensoussan A, Bilinski K, Roufogalis BD (2017) Interactions between antidiabetic drugs and herbs: an overview of mechanisms of action and clinical implications. *Diabetol Metab Syndr* 9:59
62. Rodgers RM, Gammie SM, Loo RL, Corlett SA, Krska J (2016) Comparison of pharmacist and public views and experiences of community pharmacy medicines-related services in England. *Patient Pref Adher* 10:1749–1758

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.