

## Microbe Profile

Lovering, Andrew L; Sockett, R Elizabeth

DOI:

[10.1099/mic.0.001043](https://doi.org/10.1099/mic.0.001043)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Lovering, AL & Sockett, RE 2021, 'Microbe Profile: Bdellovibrio bacteriovorus: a specialized bacterial predator of bacteria', *Microbiology*, vol. 167, no. 4, 001043. <https://doi.org/10.1099/mic.0.001043>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.


### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Microbe Profile: *Bdellovibrio bacteriovorus*: a specialized bacterial predator of bacteria

Andrew L. Lovering<sup>1,\*</sup> and R. Elizabeth Sockett<sup>2,\*</sup>



## Graphical abstract

Transmission electron micrograph of (curved) *B. bacteriovorus* HD100 invading and rounding *E. coli* S17-1, (Dr Carey Lambert, University of Nottingham).

## Abstract

*Bdellovibrio bacteriovorus* is an environmentally-ubiquitous bacterium that uses unique adaptations to kill other bacteria. The best-characterized strain, HD100, has a multistage lifestyle, with both a free-living attack phase and an intraperiplasmic growth and division phase inside the prey cell. Advances in understanding the basic biology and regulation of predation processes are paving the way for future potential therapeutic and bioremediation applications of this unusual bacterium.

## TAXONOMY

Domain Bacteria, phylum Proteobacteria, class Oligoflexia, order Bdellovibrionales, family Bdellovibrionaceae, genus *Bdellovibrio*, species *Bdellovibrio bacteriovorus*, strain HD100. Prior to the establishment of the Oligoflexia, *Bdellovibrio* was placed in the Deltaproteobacteria; a proposal has been made to reclassify into a distinct phylum, Bdellovibrionota.

## PROPERTIES

*Bdellovibrio bacteriovorus* is a predator of other Gram-negative bacteria. The type strain HD100 was isolated from soil, and utilizes whole cell invasion, with a staged lifecycle wherein it establishes itself in the periplasmic space of the host and kills it from within [1]. Related strains can be found in diverse environments (seawater, freshwater, digestive tracts) and adopt a similar endoperiplasmic method of predation, or the seemingly less complex method of epibiotic attachment and killing via external attachment only (*B. exovorus*). HD100 can be converted into a form competent for slower laboratory axenic growth via point mutations and culturing on amino-acid rich media. Some *B. bacteriovorus* strains such as *Tiberius* alternate naturally between slower axenic growth and predation. *Bdellovibrio* cells are small (0.2–0.5 by 0.5–2.5  $\mu\text{m}$ ), vibrioid with a monopolar, membrane-sheathed flagellum, relatively deformable during prey invasion and polymorphic in axenic growth, and they possess non-standard, mannose-sylated LPS.

## GENOME

The complete 3782950 bp genome of HD100 is relatively large and testament to the need for *Bdellovibrio bacteriovorus* to survive in a free-living stage in between prey killing events and for a percentage of its population to exist replicating slowly axenically as mutants [2]. As a specialized predator, there is some expected gene loss, largely in biosynthetic pathways including those of several amino acids. A significant proportion of novel hypotheticals (approximately a third of 3584 protein coding genes) encode proteins adapted for the unique predatory lifestyle e.g. a lysozyme variant that assists prey cell exit [3]. Likewise, a large fraction of the proteome is predicted to be secreted, consistent with deployment of enzymes into prey.

## PHYLOGENY

*Bdellovibrio* have both predatory (*Myxococcus xanthus*) and non-predatory (*Desulfovibrio*, *Geobacter*) relatives. There is some evidence to suggest that gene transfer to *Bdellovibrio* comes largely from non-prey bacteria including Firmicutes. It is currently unclear as to the exact relationships/evolutionary

path between endoperiplasmic and epibiotic strains, given that the latter can possess both similarly-sized genomes to HD100 (e.g. *B. qaytius*) or a relatively reduced genome (e.g. *B. exovorus*). Comparison between differing modes of predation, or to phylogenetically distant predators with a similar mode of killing (the Alphaproteobacterium *Micavibrio aeruginosavorus*) reveals that predator genome commonalities are mainly metabolic and that predation adaptations/mechanisms are likely to be predator-specific.

## KEY FEATURES AND DISCOVERIES

The staged lifecycle of predation is rich in cryptic, biologically interesting events, and begins with an attack phase in which the highly-motile *Bdellovibrio* collides with prey. An initially reversible attachment period progresses to dedicated invasion, in which the predator enters, through the prey outer membrane, sealing it neatly 'behind'. Consumption of mainly cytoplasmic, prey macromolecules then drives growth of the now-periplasmic predator (residing in a rounded, softened, dead host-cell, termed the bdelloplast). The predator elongates as the periplasm:cytoplasm ratio increases. Dependent on prey cell size, *Bdellovibrio* cells are able to divide into a variable number of progeny, odd or even, via synchronous filamentous septation. These progeny then mature, lyse the host cell and begin the cycle anew. Recognition between predator and prey is likely multifactorial, and unlike bacteriophage, susceptible prey populations do not develop genetic resistance to killing.

An initial burst of interest in *Bdellovibrio* was hampered by the available genetic tools and alternative of more tractable bacterial systems for study. In recent years this has changed, partly through the interest in the possibility of using *Bdellovibrio* and related predators as a 'living antibiotic' solution to problematic bacteria in both healthcare and agriculture. One large-scale initiative to study the feasibility of this idea was funded by the US research agency DARPA (<https://www.darpa.mil/program/pathogen-predators>). It is now apparent that *B. bacteriovorus* is able to kill prey regardless of antimicrobial resistance gene status [4]. Allied to this, injections of *B. bacteriovorus* were demonstrated to be effective in a zebrafish model of infection, working in tandem with the immune system to reduce pathogen numbers [5]. Pathogen killing has also been demonstrated inside rats, mice and chicks.

For the purposes of economy and exclusive access to the nutrients resulting from killing the prey cell, it is important that *B. bacteriovorus* crafts a stable intraperiplasmic niche and avoids premature lysis of the host cell. A swathe of prey-cell wall-manipulating enzymes assist in these tasks, relaxing host cell peptidoglycan to soften the invaded cell and signal occupancy to kin, reducing wasteful secondary invasion events.

Received 13 November 2020; Accepted 18 February 2021; Published 12 April 2021

Author affiliations: <sup>1</sup>Dept of Biosciences, University of Birmingham, Birmingham, UK; <sup>2</sup>University of Nottingham, Nottingham, UK.

\*Correspondence: Andrew L. Lovering, [a.lovering@bham.ac.uk](mailto:a.lovering@bham.ac.uk); R. Elizabeth Sockett, [liz.sockett@nottingham.ac.uk](mailto:liz.sockett@nottingham.ac.uk)

Keywords: predatory bacteria; bdellovibrio; molecular microbiology.

*B. bacteriovorus* itself also has peptidoglycan and must be careful not to act against self during predation. One of several mechanisms for this is to utilize an ankyrin-based immunity protein to block action of peptidoglycan endopeptidases; the importance of this is outlined in mutants that lack this protection and self-lyse upon invasion [6]. The use of fluorescent D-amino acid reporters to illuminate cell wall modifications revealed that a 'porthole' is built to support passage of *B. bacteriovorus* through the prey envelope. Two further modifications of the prey peptidoglycan serve to reseal this pore and provide general reinforcement of the invaded cell [7].

Use of fluorescence microscopy has also begun to provide details on the intracellular developmental stages of the predator. *B. bacteriovorus* cells place a block on replication, when external, between kills, and it was thought that a complete round of chromosome replication was required during a singular killing event. However, a detailed study of replication timing revealed that if the host cell provided insufficient resources for this, *B. bacteriovorus* could stall before a full replication cycle occurred and then complete this during killing of a second cell [8]. Other adaptations to maximize efficient use of the host cell material include cyclic nucleotide signalling that prey is sensed, inducing predatory processes, and uptake of nucleoside monophosphates without need for breakdown into their component units.

## OPEN QUESTIONS

- Which (surface?) features of prey determine susceptibility to *Bdellovibrio*?
- By what means is the decision to invade or not determined?
- What is the mechanistic basis for whole cell invasion, namely controlled entry through the outer membrane of prey?

- How is the remarkable variable number of progeny and resulting differential pattern of septation controlled?
- What are the key signalling events/molecules that inform on exhaustion of prey cell resources pre-exit?

### Funding information

R.E.S. and A.L.L. are currently supported by a Wellcome Trust Investigator Award 209437/Z/17/Z.

### Conflicts of interest

The authors declare that there are no conflicts of interest.

### References

1. Sockett RE. Predatory lifestyle of *Bdellovibrio bacteriovorus*. *Annu Rev Microbiol* 2009;63:523–539.
2. Rendulic S, Jagtap P, Rosinus A, Eppinger M, Baar C. A predator unmasked: life cycle of *Bdellovibrio bacteriovorus* from a genomic perspective. *Science* 2004;303:689–692.
3. Harding CJ, Huwiler SG, Somers H, Lambert C, Ray LJ *et al*. A lysozyme with altered substrate specificity facilitates prey cell exit by the periplasmic predator *Bdellovibrio bacteriovorus*. *Nat Commun* 2020;11:4817.
4. Dharani S, Kim DH, Shanks RMQ, Doi Y, Kadouri DE. Susceptibility of colistin-resistant pathogens to predatory bacteria. *Res Microbiol* 2018;169:52–55.
5. Willis AR, Moore C, Mazon-Moya M, Krokowski S, Lambert C *et al*. Injections of predatory bacteria work alongside host immune cells to treat *Shigella* infection in zebrafish larvae. *Current Biology* 2016;26:3343–3351.
6. Lambert C, Cadby IT, Till R, Bui NK, Lerner TR *et al*. Ankyrin-mediated self-protection during cell invasion by the bacterial predator *Bdellovibrio bacteriovorus*. *Nat Commun* 2015;6:8884.
7. Kuru E, Lambert C, Rittichier J, Till R, Ducret A *et al*. Fluorescent D-amino-acids reveal bi-cellular cell wall modifications important for *Bdellovibrio bacteriovorus* predation. *Nat Microbiol* 2017;2:1648–1657.
8. Makowski Łukasz, Trojanowski D, Till R, Lambert C, Lowry R *et al*. Dynamics of Chromosome Replication and Its Relationship to Predatory Attack Lifestyles in *Bdellovibrio bacteriovorus*. *Appl Environ Microbiol* 2019;85.

### Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at [microbiologyresearch.org](https://microbiologyresearch.org).