

Synchronization of Interhippocampal Ripple Events (80–200 Hz) by Long-projection Inhibitory Neurons

Tiesinga, PH; Li, Xiaoli; Sakatani, S; Boldogkoi, Z; Hirase, H; Sık, A

DOI:

[10.1186/1471-2202-8-S2-P45](https://doi.org/10.1186/1471-2202-8-S2-P45)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Tiesinga, PH, Li, X, Sakatani, S, Boldogkoi, Z, Hirase, H & Sık, A 2007, Synchronization of Interhippocampal Ripple Events (80–200 Hz) by Long-projection Inhibitory Neurons. in *BMC Neuroscience: Sixteenth Annual Computational Neuroscience Meeting: CNS*2007*. vol. 8, P45, BMC Neuroscience, no. (Suppl 2), vol. 8, Sixteenth Annual Computational Neuroscience Meeting, 2007, Toronto, Canada, 7/07/07. <https://doi.org/10.1186/1471-2202-8-S2-P45>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Poster presentation

Open Access

Synchronization of interhippocampal ripple events (80–200 Hz) by long-projection inhibitory neurons

Paul HE Tiesinga*¹, Xiaoli Li², Seiichi Sakatani³, Zsolt Boldogkői⁴, Hajime Hirase³ and Attila Sík*⁵

Address: ¹Department of Physics & Astronomy, University of North Carolina, Chapel Hill, NC, USA, ²CERCIA, School of Computer Science, The University of Birmingham, Birmingham, UK, ³RIKEN Brain Science Institute, Wako-shi, Saitama, Japan, ⁴Department of Biology, University of Szeged, Szeged, Hungary and ⁵Department of Dentistry, Laval University, Québec, Canada

Email: Paul HE Tiesinga* - tiesinga@physics.unc.edu; Attila Sík* - attila.sik@crulrg.ulaval.ca

* Corresponding authors

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, 8(Suppl 2):P45 doi:10.1186/1471-2202-8-S2-P45

© 2007 Tiesinga et al; licensee BioMed Central Ltd.

Network oscillations between the two hippocampi are highly synchronized. Synchronized theta is believed to be the result of the common input from the septal region, whereas the mechanism of the ripple synchronization is not well understood. It was previously demonstrated using partial coherence analysis that the "coupling" between the two CA1 regions of hippocampi during theta oscillations is stronger than that between the individual layers of the same hippocampus.

Hippocampal sharp wave-ripple complexes occur during slow-wave sleep and awake immobility and are thought to be important for memory consolidation. The delay between simultaneously recorded ripple events from the two hippocampi is remarkably short (1–2 ms). This observation suggests that some sort of fast communication mechanism should connect the two hippocampi. We demonstrate that the simultaneously occurring ripple events in the two hippocampi are highly coherent. This observation suggests an important role of the commissural projections in interhemispheric network synchronization. Using various anatomical methods we demonstrate that a subset of inhibitory neurons (NPY-expressing cells), located in the CA1, CA3 area and dentate gyrus, extensively project not only to the contralateral hippocampus, but also to the septal region. We use model simulations to determine to what extent and under which

conditions the highly synchronous ripple events can be produced by long-range intra- and interhippocampal inhibitory projections.