

# Peripheral causes of cognitive motor dissociation in patients with vegetative or minimally conscious state - Reply

Fernandez-Espejo, Davinia; Owen, Adrian M

DOI:

[10.1001/jamaneurol.2016.0143](https://doi.org/10.1001/jamaneurol.2016.0143)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Fernandez-Espejo, D & Owen, AM 2016, 'Peripheral causes of cognitive motor dissociation in patients with vegetative or minimally conscious state - Reply', *JAMA Neurology*, vol. 73, no. 5, pp. 608-609. <https://doi.org/10.1001/jamaneurol.2016.0143>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Published version available at: <http://dx.doi.org/10.1001/jamaneurol.2016.0143>

Validated Feb 2016

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 **Peripheral causes of cognitive motor dissociation in patients with vegetative or**  
2 **minimally conscious state – Reply**

3

4 Davinia Fernández-Espejo, PhD<sup>1\*</sup>, Adrian M. Owen, PhD<sup>2</sup>

5

6 <sup>1</sup> School of Psychology, University of Birmingham, Birmingham, B15 2TT, UK;

7 <sup>2</sup> The Brain and Mind Institute, The University of Western Ontario, London, Ontario, N6A  
8 5B7, Canada;

9

10

11 \*Correspondence:

12 Davinia Fernández-Espejo, School of Psychology, University of Birmingham, Birmingham,  
13 B15 2TT, UK; [D.Fernandez-Espejo@bham.ac.uk](mailto:D.Fernandez-Espejo@bham.ac.uk)

14

15 Word count: 303

16

17 We thank Latronico and colleagues for their comments regarding our article<sup>1</sup>. They proposed  
18 that peripheral nervous system and muscle pathology<sup>2</sup> may have contributed to the lack of  
19 behavioural responses exhibited by our patient. As mentioned in our Discussion section, Shea  
20 and Bayne<sup>3</sup> had previously argued a similar peripheral explanation for the absence of overt  
21 motor behaviour in patients with preserved covert motor behaviour<sup>4</sup>. In vegetative and  
22 minimally conscious patients, peripheral damage is most commonly related to motor axonal  
23 neuropathy<sup>5</sup>, which, as Latronico and colleagues point out, is a major cause of *paralysis*<sup>2</sup>.  
24 While we did not specifically test for electrophysiological signs of peripheral pathology, our  
25 patient showed no evidence of paralysis. In fact, he exhibited frequent spontaneous

26 movements of the limbs (more frequently upper limbs), head, and torso, as well as very  
27 consistent withdrawal to painful stimulation (see information about clinical assessments in  
28 the original supplementary information). In contrast, he was incapable of producing  
29 voluntarily motor responses to command. Therefore, the main deficit he exhibited, which our  
30 study aimed to explain, was not an absence of skeletal movement, but a lack of *voluntary*  
31 *control* of his motor responses, and thus the underlying mechanism is necessarily central <sup>6</sup>.  
32 Based on this, we disagree with Latronico and colleagues' suggestion for a role of peripheral  
33 pathology in explaining our patient's lack of overt command following capabilities.  
34 Nevertheless, as we mentioned in our Discussion, our patient exhibited other symptoms in  
35 addition to the lack of command following (e.g. lack of visual pursuit, or vocalizations) for  
36 which our results may not offer a complete explanation. In this context, we agree that the  
37 evaluation of the peripheral nervous system and muscles, in combination with neuroimaging  
38 and clinical assessments, may contribute to a more comprehensive understanding of the full  
39 clinical profile exhibited by each individual patient.

40

41

42

#### 43 **References**

44

- 45 1. Fernandez-Espejo D, Rossit S, Owen AM. A Thalamocortical Mechanism for the  
46 Absence of Overt Motor Behavior in Covertly Aware Patients. *JAMA Neurol.* October  
47 2015;1-9. doi:10.1001/jamaneurol.2015.2614.
- 48 2. Latronico N, Bolton CF. Critical illness polyneuropathy and myopathy: a major cause of  
49 muscle weakness and paralysis. *The Lancet Neurology.* 2011;10(10):931-941.  
50 doi:10.1016/S1474-4422(11)70178-8.

- 51 3. Shea N, Bayne T. The Vegetative State and the Science of Consciousness. *Br J Philos*  
52 *Sci.* 2010;61(3):459-484. doi:10.1093/bjps/axp046.
- 53 4. Fernandez-Espejo D, Owen AM. Detecting awareness after severe brain injury. *Nature*  
54 *Reviews Neuroscience.* October 2013. doi:10.1038/nrn3608.
- 55 5. Bagnato S, Boccagni C, Sant'angelo A, Prestandrea C, Romano MC, Galardi G.  
56 Neuromuscular involvement in vegetative and minimally conscious states following  
57 acute brain injury. *J Peripher Nerv Syst.* 2011;16(4):315-321. doi:10.1111/j.1529-  
58 8027.2011.00363.x.
- 59 6. Bracht T, Schnell S, Federspiel A, et al. Altered cortico-basal ganglia motor pathways  
60 reflect reduced volitional motor activity in schizophrenia. *Schizophrenia Research.*  
61 2013;143(2-3):269-276. doi:10.1016/j.schres.2012.12.004.

62