

Low nutrient intake and frailty among overweight and obese migrant women from ethnically diverse backgrounds aged 60+ years: a mixed-methods study

Castaneda Gamos, Diana; Redwood, Sabi; Thompson, Janice

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Castaneda Gamos, D, Redwood, S & Thompson, J 2017, 'Low nutrient intake and frailty among overweight and obese migrant women from ethnically diverse backgrounds aged 60+ years: a mixed-methods study', *Journal of Nutrition Education and Behavior*.

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Eligibility for repository: Checked on 16/9/2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **‘Low Nutrient Intake And Frailty Among Overweight And Obese Migrant**
2 **Women From Ethnically Diverse Backgrounds Aged 60+ Years: A Mixed-**
3 **Methods Study’**

4
5
6

7 **INTRODUCTION**

8
9

10 Frailty has become the focus of extensive research due to the ever-increasing aging of
11 the global population. Frailty is characterized as a disorder of multiple physiological
12 systems in which homeostatic mechanisms start failing, increasing the risk of declines
13 in cognitive and physical function.¹⁻³ Furthermore, longitudinal studies have
14 demonstrated a greater prevalence of cardiovascular disease and diabetes among frail
15 older people,⁴ and a greater frailty burden for women in comparison to men.^{5,6}
16 Therefore, identifying and treating individuals at risk of frailty may help delay its
17 negative consequences and reduce the financial, social, and personal burdens these
18 consequences place upon individuals, families and societies.^{7,8}

19

20 One of the most widely used definitions of frailty is the frailty phenotype proposed
21 and validated by Fried and colleagues.² This battery of tests identifies people as frail
22 when they meet three or more of five criteria: relatively weak grip strength,
23 unintentional weight loss, self-reported exhaustion, slow walking speed and low
24 levels of physical activity (PA). The inclusion of unintentional weight loss is used as a
25 proxy measure of dietary inadequacy, which is congruent with the conceptualization
26 of frailty as a wasting disorder.²⁻³ However, obesity can also be linked with frailty, as

27 indicated by the greater risk of physical function decline and pro-inflammatory state
28 commonly found among older adults who are obese.⁹

29

30 In older adults, the use of unintentional weight loss in the definition of frailty is
31 problematic as this measure may not be sensitive enough to reflect reduced energy
32 and nutrient intakes.⁸ Weight loss will not occur if energy intake matches energy
33 expenditure, however a diet that is adequate in energy can still be deficient in certain
34 nutrients, increasing a person's risk for frailty. Therefore, we hypothesize that a low
35 intake of energy and selected nutrients is a stronger predictor of frailty in
36 overweight/obese older women from diverse ethnic backgrounds than unintentional
37 weight loss. There is limited evidence examining the association between frailty as a
38 syndrome and nutrient intakes, and this has been conducted in predominantly White
39 older adults.⁸ Thus, very little is known about these associations in older adults from
40 diverse ethnic backgrounds. In addition, to develop interventions that can effectively
41 delay or prevent frailty in older women from diverse ethnic backgrounds, more
42 information is needed to explore if there are links between perceptions of body
43 weight, dietary intake and physical function in a population with disproportionately
44 higher rates of overweight and obesity.

45

46 Therefore, the aims of this study were to: 1) examine the associations between
47 dietary/nutrient intake and frailty in a sample of older women (≥ 60 years) from
48 diverse ethnic backgrounds living in the UK; 2) to gain a greater understanding of the
49 potential links between women's perceptions of body weight, dietary intake and
50 physical function.

51

52

METHODS

53

54

55 **Study Design**

56

57 A cross-sectional, mixed-methods design was employed, using 24-hr dietary recall
58 interviews that were enhanced with the addition of a qualitative semi-structured
59 interview. These methods allowed for the quantitative estimate of energy/nutrient
60 intake and its association with frailty, as well as providing insights into women's
61 perceptions of their body weight, dietary intake and physical function.

62

63 **Recruitment and Participants**

64

65 A convenience sample of first generation migrant women from Ireland, Jamaica,
66 Montserrat, St Kitts and Nevis, India, Pakistan, Bangladesh, Yemen, Sierra Leone,
67 Somalia, and Eritrea were recruited to participate in the study. Inclusion criteria
68 included being at least 60 years of age, with no medical conditions affecting memory
69 (e.g., dementia), and the ability to walk 15ft with no or minimal assistance (i.e., use of
70 a walking stick). Community-dwelling women living on their own or with family
71 members were recruited using maximum variation sampling¹⁰ to achieve our goal of
72 recruiting a sample across the ranges of age, migration backgrounds, socio-economic
73 status, and main ethnic groups living in the geographic region. This was achieved by
74 using the most recent Birmingham census data to identify the most representative
75 migrant groups.¹¹ Community centres serving specific migrant and older adult groups

76 were contacted and informed about the study. Those in leadership roles at these
77 centers facilitated access to potential participants so they could be approached and
78 informed of the purpose of the study. Participants were recruited via word-of-mouth
79 and snowballing.¹² Ethics approval was granted by The University of Birmingham
80 Ethics Committee (reference No. ERN_13-0557). All participants provided written
81 informed consent.

82

83 **Data Collection**

84

85 Data were collected at the participants' time and location of choice (e.g., homes or
86 community centers). For participants not fluent in English, trained interpreters fluent
87 in Punjabi, Bengali, Arabic and Somali provided simultaneous translation during
88 recruitment and data collection. Socio-demographic information was gathered via a
89 researcher-administrated questionnaire.

90 ***Dietary Intake.***

91 A multiple-pass 24-hr dietary recall interview was conducted to gather data on the
92 types and amounts of foods consumed on the previous day via a standard protocol.¹³
93 Information was also obtained on nutrient supplement use. A photographic food atlas
94 assisted with the estimation of portion sizes.¹⁴ The first author (DCG), a dietitian,
95 trained in dietary assessment conducted all 24-hr dietary recalls. Data coding and
96 processing was conducted by DCG, with oversight from JLT who has extensive
97 expertise in dietary assessment. These procedures enabled a standardized data entry
98 and analysis process. The dietary recall interview was audio-recorded to ensure
99 accuracy of quantitative data entry and to facilitate the collection of additional

100 qualitative information. When participants stated that the previous day did not reflect
101 their habitual diet (e.g., they had engaged in fasting practices), the 24-hr dietary recall
102 was repeated later in the same week on a day that was identified by participants as
103 being representative of their habitual intake. This occurred in 5 participants. Data
104 were not gathered during periods of major religious observances (e.g., Ramadan,
105 Diwali). All recalls were conducted during weekdays, excluding Monday. Nutrient
106 analysis was completed using DietPlan 6.0 software (Forestfield software Ltd 2006,
107 Horsham, UK), which included standard and supplemental food composition
108 databases that covered the range and ethnic diversity of foods consumed in the UK.
109
110 Similarly to methods reported by Bartali and colleagues,⁸ low intake was defined as
111 the lowest quintile of the distribution of energy (<13 kcal/kg) and specific nutrients:
112 protein <30 g, vitamin D <0.5 µg, vitamin E <2.5 mg, retinol <101 µg, vitamin C <32
113 mg, folate <127 µg, iron <5.6 mg, calcium <349 mg, and zinc <3.6 mg. A nutritional
114 score was obtained by summing the number of nutrients categorized as low intake.
115 This nutritional score was subsequently categorized into a low intake of 0, 1-3, or >3
116 nutrients. A low intake of >3 nutrients was classified as poor nutritional status.

117

118 *Anthropometric measures and assessment of frailty.*

119 Anthropometric measures included height measured to the nearest mm (SECA 213
120 portable stadiometer), weight to the nearest 0.1 kg (SECA 899 digital scale), and hip
121 and waist circumference (WC) measured to the nearest cm using an extractable tape
122 measure. All anthropometric measurements were taken with the participant wearing
123 light clothing and no shoes. Body mass index (BMI) was calculated as weight divided

124 by height squared (kg/m^2), and waist-to hip-ratio (WHR) as waist circumference
125 divided by hip circumference (cm).

126 Frailty status was assessed using a modified version of the original frailty definition
127 developed by Fried and colleagues.² This included: 1) Exhaustion, defined using self-
128 reported fatigue from two questions from the Center for Epidemiological Studies-
129 Depression (CES-D) depression scale (“I felt that everything I did was an effort,” and
130 “I could not get going.”) Participants who reported having these feelings for ≥ 3 days
131 over the previous week to either or both questions received positive scores for
132 exhaustion; 2) Slow walking speed, with the highest quintile of the time needed to
133 walk a distance of 15 feet, adjusted by height (>14.5 seconds for height ≤ 157.7 cm
134 and >9.7 seconds for height > 157.7 cm); 3) Weak grip strength was defined as the
135 lowest quintile for adjusted grip strength using a JAMAR hand-held dynamometer
136 (Sammons Preston Rolyan, Bolingbrook, Illinois, USA), adjusted by BMI.

137 Participants met the criteria for weak grip strength if their BMI and grip strength were
138 $\leq 25.8 \text{ kg}/\text{m}^2$ and $\leq 12 \text{ kg}$; $>25.9- 29.6 \text{ kg}/\text{m}^2$ and $\leq 11 \text{ kg}$; $>29.7- 31.6 \text{ kg}/\text{m}^2$ and ≤ 12
139 kg ; and $\geq 31.7 \text{ kg}/\text{m}^2$ and $\leq 14 \text{ kg}$. A low level of PA was defined as the lowest quintile
140 of caloric expenditure ($< 60 \text{ kcal}/\text{week}$) using the International Physical Activity
141 Questionnaire short-form modified for the elderly (IPAQ-E).¹⁵ This version of the
142 IPAQ provides examples of activities that are more common among older adults and
143 has shown a moderate correlation ($r=0.347$, $p<0.01$), and moderate agreement κ
144 ($95\% \text{CI}$)= 0.448 ($0.18-0.72$, $p < 0.001$) with accelerometry.¹⁶

145

146 Since the purpose of this study was to examine the association between dietary intake
147 and frailty, similar to Bartali’s study, unintentional weight loss (>10 pounds in the last
148 year) was excluded from the original frailty definition.⁸ Therefore, participants with

149 >2 positive criteria were categorized as frail, while those with ≤ 1 positive criteria
150 were categorized as not frail.

151

152 *Semi-structured interviews.*

153 A purposive sub-sample (n=46) across the range of age, ethnic groups and socio-
154 economic status was invited to participate in an interview that was guided by a list of
155 topics related to migration histories, dietary intake and eating behaviors, and
156 engagement in PA (migration histories and PA data not reported here). For the
157 purpose of this study, dietary topics centered on participants' perceptions of their diets
158 in relation to their body weight and frailty status (referred to as physical function
159 during the interviews). The interview guide was pilot-tested prior to the study and was
160 further revised via an iterative process throughout the data collection period. All
161 interviews were audio-taped and transcribed verbatim, with the interviews conducted
162 with participants who were not fluent in English being translated from their native
163 language into English by a trained interpreter during the interview process (n=16).

164

165 **Data Analysis**

166

167 *Quantitative data analysis.*

168 Descriptive characteristics (means, SDs, and percentages) were calculated for socio-
169 demographic variables. To identify potential confounding factors, independent t-tests
170 or Mann-Whitney U tests (for non-parametric data) were conducted to examine any
171 significant differences in continuous variables between those classified as frail or not
172 frail, with Chi-squared or Fisher's exact tests conducted for categorical variables.

173 Point-biserial correlations (r_{pb}) were used to determine the association between frailty
174 status (dichotomous variable), weight loss, and indices of overweight/obesity (e.g.,
175 BMI, WC and WHR). Multiple logistic regressions were used to evaluate the
176 association between frailty status and each of its components with low energy intake
177 and poor nutritional status. Separate models were conducted to test the association
178 between nutrient intakes with frailty adjusting for confounding factors and energy
179 intake. All statistical analyses were performed using SPSS version 21.0 (SPSS INC.,
180 Chicago, IL); alpha was set at $p < 0.05$.

181

182 *Qualitative data analysis.*

183 An inductive thematic analysis of the interview transcripts was conducted, allowing
184 for the identification of themes being driven by participants' perspectives of their
185 diets in the context of their body weight and physical function/frailty status rather
186 than fitting the data into a pre-existing theoretical framework.¹⁷ Initially, a subset of
187 transcripts were read several times by the first author and two independent researchers
188 to identify predominant topics across the data. An initial coding frame using
189 qualitative analysis software (QSR NVivo, version 10) was developed which formed
190 the basis of broad coding and analysis. All of the transcripts were then coded by the
191 first author. The coding frame was discussed and refined by all authors until
192 consensus was reached. Data saturation was considered to have been achieved when
193 no new or relevant information emerged from each of the various ethnic groups
194 included in the study.¹⁸

195

RESULTS

196

197

198

199 Table 1 includes the demographic characteristics of participants. On average,
200 participants (mean age= 70.5 ± 7.6 years) reported having 2.3 ± 1.5 diseases
201 previously diagnosed by a doctor, with hypertension, arthritis and type 2 diabetes the
202 most common. Over 88% of the sample was classified as overweight or obese. BMI
203 cut-points for overweight and obesity among the Arab, Indian, Pakistani and
204 Bangladeshi participants were those recommended by the World Health Organization
205 for Asian populations.¹⁹ Although participants came from all socioeconomic levels,
206 79% were categorized as being in the two most socio-economically deprived quintiles
207 based on the English indices of deprivation.²⁰ Seventeen participants (22.4%) were
208 classified as frail, while 23 (30.3%) and 36 (47.4%) were classified as pre-frail and
209 non-frail, respectively. Frail participants were older and had a higher number of
210 diagnosed diseases; these were the only demographic variables that were statistically
211 different between frail and non-frail participants.

212 **Frailty and low nutrient intake**

213

214 Among frail participants, 82.3% had a low nutrient intake of at least one selected
215 nutrient (Table 2). The percentage of women with frailty increased with the greater
216 number of nutrients classified as low intake. Logistic regression analyses indicated
217 that low energy intake was independently associated with frailty (odds ratio [OR]:
218 11.71, 95% confidence interval [CI]: 2.36-57.97). After adjusting for energy, age and
219 number of diseases, poor nutritional status (>3 low nutrient intakes) was significantly
220 associated with frailty (OR: 6.58, 95% CI: 1.01-43.08) in comparison to those women

221 who did not have a low intake of any nutrients. After adjusting for energy and other
222 confounding variables, only slow walking speed was significantly associated with
223 poor nutritional status (OR: 1.86, 95% CI: 1.31-3.07).

224

225 In addition, a low intake of retinol (OR: 10.33, 95% CI: 1.55- 68.94) and zinc (OR:
226 8.47, 95% CI: 1.04-68.80) were significantly associated with frailty after adjustment
227 for energy intake and other confounding variables (Table 3). Self-reported weight loss
228 ($p=0.3$ for Fisher's exact test), BMI ($r_{bp}= 0.09$, $p=0.4$), waist circumference ($r_{bp}= 0.2$,
229 $p=0.1$), and WHR ($r_{bp}= 0.03$, $p=0.8$) were not associated with frailty.

230

231 **Qualitative Interview Results**

232

233 Two main themes which linked women's perceptions of body weight, dietary intake
234 and physical function were identified. They were: 1) concerns about weight and body
235 image; and 2) perceptions about negative effects of unhealthy foods on physical
236 function and health. Specific quotes from participants have been used to demonstrate
237 the themes outlined above.

238

239 ***Weight and body image concerns.***

240 Weight and body image emerged as two issues that were particularly important to
241 participants. Data suggest that these women have become more aware of their weight
242 as they have aged. Furthermore, some participants emphasized that their weight status
243 worried them more than getting older or other health problems as the excerpts below
244 indicate:

245

246 *'I am very careful that I don't eat too much, though I am very hungry but I will leave*
247 *[the food uneaten]...I never say I want to eat more, no! ... I do not want to put on*
248 *weight, that is in the back of my mind, I never think of the heart [problem], I think of*
249 *my weight'* (Indian, 73y).

250

251 *'It doesn't bother me [the age], but when somebody says you are fat, then it hurts me!'*
252 *(Indian, 62y).*

253

254 Participants' narratives also highlighted a difficult relationship between their diets and
255 body weight, leading to feelings of frustration and shame:

256

257 *'My thinking was always eating healthy, but...I don't know how I put on so much*
258 *weight so quickly and I've been trying [to lose weight] for many years now, it's not*
259 *going down. I don't know what happened... I have gained so much I can't even get rid*
260 *of it... since I've put on weight and I am out of size as well, I think 'Oh God people,*
261 *don't see me!'* ...*That stops me from going out, dressing up as well, meeting people or*
262 *going into places'* (Pakistani, 62y)

263

264 Given pervasive concerns about weight gain, many participants described modifying
265 their diets in an effort to lose weight. However, adopting more restrictive diets have
266 led some women to link these changes with a negative impact on their strength:

267

268 *'When you are getting older is hard to lose weight ...well, I used to cut down my food*
269 *and then I think I was falling apart, I was getting weak... so I just said, "I'll just*
270 *continue [as normal]'"* (African-Caribbean, 79y)

271 Other participants who have also tried to reduce their food intake mentioned that they
272 occasionally complement their '*light diets*' with certain food items in order to meet
273 their perceived dietary requirements:

274

275 '*When I feel I haven't had enough protein... and need to rebuild some of the cells,*
276 *dying cells, ...then I would consciously have fish or chicken and try to eat a large*
277 *portion to try to convince myself that I'm eating enough protein...but no, I do a lot of*
278 *light days [of decreased consumption of fat and animal products]'* (African-
279 *Caribbean, 68y).*

280

281 ***Perceptions about negative effects of unhealthy foods on physical function and***
282 ***health.***

283 Participants' perceptions about the link between diet, physical function and general
284 health were mainly driven by their beliefs about the negative effects unhealthy foods
285 have on their mobility. For instance, some participants mentioned that eating
286 '*fattening food*' decreases their ability to be more active:

287

288 '*If I had fried food and I walk, I feel breathless yeah, so I keep in line what I am*
289 *eating*' (Indian, 71y)

290

291 '*Like...when you eat chips [French fries] you feel so heavy and you don't feel like*
292 *moving, you don't feel like running you know*' (Indian, 74y)

293

294 Overall, women felt that the quality of the food they eat is associated with their
295 general health, and that a healthy diet is an important component of healthy aging:

296 *'Health is related to what you put in your body, you are what you eat and if you put*
297 *healthy food in your body, you can expect to be healthy at this age' (African-*
298 *Caribbean, 69y)*

299

300

DISCUSSION

301

302

303 The present study examined the association between dietary intake and frailty in a
304 group of free-living first generation migrant older women using a mixed-methods
305 approach. Findings from this study indicated that having a low energy intake was
306 associated with frailty, and a poor nutritional status was significantly associated with
307 frailty after adjusting for energy and other confounding factors. Poor nutritional status
308 was also associated with slow walking speed, one of the criteria of the frailty
309 syndrome. The findings also provided rich insight into participants' perceptions about
310 the links between their body weight, dietary intake, and physical function.

311 Our findings support existing evidence associating frailty and its components to
312 nutrition at the nutrient level.⁸ Poor nutritional status and low serum levels of several
313 nutrient biomarkers (serum carotenoids, α -tocopherol, 25-hydroxyvitamin D, and
314 vitamin B6) have been found to be related to an increased risk of frailty among
315 predominantly White older adults.²¹⁻²³ These data, in addition to the findings from the
316 present study, suggest that an inadequate diet plays a crucial role in the physical
317 function of older adults. This is of particular importance due to the body composition
318 changes associated with old age leading to loss of muscle mass (sarcopenia) that can
319 contribute to morbidity and decreased quality of life.³

320 There are multiple pathways in which micronutrient deficiencies can increase the risk
321 of frailty in older adults by promoting conditions commonly associated with older age
322 such as oxidative stress, impaired immunity, muscle and bone metabolism, and
323 inflammation.²⁴ In our study, only retinol and zinc were independently associated with
324 frailty, suggesting that these two nutrients may be of particular concern in this sample.
325 Retinol is suggested to protect cell membranes from oxidative damage related to
326 aging,²⁵ while both retinol and zinc play an important role in maintaining the integrity
327 of the immune system.²⁶ Although malnourishment is typically associated with
328 underweight, this study confirms that overweight/obese individuals can also be
329 malnourished due to consuming a poor quality diet.²⁷ Thus, an individual can be frail
330 and not necessarily experience weight loss.

331 Among this sample, body weight concerns emerged as a key factor influencing energy
332 and nutrient intake. Therefore, the majority of participants were more conscious about
333 eating in moderation in order to lose weight, and did not identify being concerned
334 with how their dietary intake would affect nutrient adequacy. Although it is well
335 known that body dissatisfaction is highly associated with dietary intake in younger
336 adults, it is only recently that this has been reported in older adults, especially in
337 women.²⁸ Among women from minority ethnic groups, body weight perceptions have
338 been reported to be more positive and accepting of larger figures and a body weight
339 consistent with medically defined overweight or obesity.²⁹ However, our findings
340 indicate that the women in this ethnically diverse sample are concerned about their
341 body weight and the negative consequences associated with overweight/obesity.
342 These concerns may potentially lead them to adopt restrictive eating practices that
343 may cause more harm than good. Although body dissatisfaction has been previously
344 reported in younger migrant women,³⁰ to our knowledge, this is the first time that this

345 has been found in a sample of older migrant women with high rates of
346 overweight/obesity.

347 Regarding the negative effects of unhealthy foods on physical function and health, a
348 few studies have found that an unhealthy diet (i.e., poor consumption of fruits and
349 vegetables, low adherence to a Mediterranean-type diet) is associated with mobility
350 limitations and disability in older adults, particularly in women.³¹⁻³³ Although this
351 association has been found to be stronger in non-obese individuals,³² in our study
352 women felt that unhealthy foods, particularly fatty foods, were negatively related to
353 their mobility. Thus, in overweight/obese older women, healthier diets may be
354 perceived as a means of ameliorating mobility loss and further physical decline.

355 Given pervasive concerns about weight gain, findings from this study suggest that
356 older women from ethnically diverse backgrounds with a high prevalence of
357 overweight/obesity need dietary advice that promotes both the maintenance of a
358 healthy body weight and nutrient adequacy. Particularly, because both excess weight
359 and nutritionally inadequate diets are important determinants of morbidity and
360 premature mortality.³⁴

361 The major strength of the present study is the inclusion of a population commonly
362 under-represented in research,³⁵ and little is known about dietary intake, eating
363 behaviors, and frailty in older migrant women. The mixed-methods methodology is
364 also a strength, as it allowed for the examination of dietary intake and its association
365 with frailty as well as providing important insights into women's perceptions of their
366 dietary intake and its link with body weight and physical function. In addition, the
367 interview sample size was relatively large for a mixed-methods study, and data
368 saturation was reached in all participants across the range of age and ethnic groups.

369 Finally, some limitations of the study need to be considered. Due to the cross-
370 sectional study design and a relatively small sample size for the quantitative data,
371 causal inferences cannot be made and findings may not be generalizable to the wider
372 population of first generation older migrant women living in the UK. In addition,
373 almost 90% of the sample was overweight or obese. Although this could be
374 considered a strength as the sample reflects the higher prevalence of
375 overweight/obesity in ethnic groups in the UK,³⁶ the findings do not include data from
376 participants who were underweight. This could have limited the potential of finding
377 an association between frailty, protein and other micronutrients consistently found in
378 previous studies.^{8,23,37} In addition, BMI was used as a measure of weight status. This is
379 problematic as BMI does not distinguish between lean tissue and fat mass, and cannot
380 take into account the height loss that occurs with older age.³⁴ Studies including a
381 larger sample of older women from ethnically diverse backgrounds using an accurate
382 measure of body composition and nutritional biomarkers are needed to confirm our
383 findings. A larger sample will also allow for the examination of significant
384 differences between ethnic groups.

385 Another important limitation was the use of a single 24-hr dietary recall, a limitation
386 shared with other studies conducted with older adults and ‘hard to reach’
387 populations.³⁸ This method was considered the most appropriate as it minimized
388 participant burden and allowed participants with limited English literacy to fully
389 participate in both the quantitative and qualitative aspects of the study. Limitations in
390 willingness of participants to participate in a second 24-hr dietary recall interview, in
391 addition to budgetary constraints, prevented the use of repeated 24-hr dietary recalls.
392 In the present study, energy intake was relatively low and as such, under-reporting
393 cannot be ruled out. Under-reporting has been found to be associated with female

394 gender, higher age, lower socio-economic status, and overweight/obesity.³⁹ Because
395 of the day-to-day variability in dietary intake, the single 24-dietary recall provided
396 data for the sample rather than an estimate of an individual's dietary intake. The
397 interviews were conducted by a trained nutritionist, and when necessary with the aid
398 of interpreters with the same ethno-cultural background who were familiar with the
399 participants' dietary habits. In addition, we enhanced the 24-hr dietary recall with an
400 in-depth probing interview that allowed for a rich exploration of habitual dietary
401 behaviors not possible with a standard 24-hr dietary recall. Low dietary and nutrient
402 intakes in older adults are not uncommon given important changes in body
403 composition, intestinal absorption and decreased levels of PA.⁴⁰ In our study, women
404 were highly sedentary, which could have also influenced their energy intake.
405 Nevertheless, misreporting may have occurred and as such, our results should be
406 interpreted in light of this limitation.

407

408 **IMPLICATIONS FOR RESEARCH AND PRACTICE**

409

410

411 Findings from this study indicate that among a group of mainly overweight/obese
412 migrant women from ethnically diverse backgrounds, poor nutritional status is an
413 independent predictor of frailty. Given that weight loss may not necessarily be present
414 in community-dwelling older women, low energy and nutrient intakes make important
415 contributions to the development of frailty. Therefore, assessing dietary intake may
416 assist with screening for, and treating, frailty. Moreover, the mismatch found between
417 body weight and dietary inadequacy may potentially cause older women to engage in
418 self-imposed dietary restrictions that could cause further health problems. Future

419 strategies to prevent and detect frailty in this sub-group of the population should focus
420 on maintenance of a healthy body weight as well as in the overall nutritional quality
421 of the diet.

422

423

REFERENCES

424

425

426 1. Clegg A, Young J, Iliffe, S, et al. Frailty in elderly people. *Lancet*. 2013;381:752-
427 762. doi: 10.1016/S0140-6736(12)62167-9.

428

429 2. Fried LP, Tangen CM, Walston J., et al. Frailty in older adults: evidence for a
430 phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56:M146-56. doi:

431 10.1093/gerona/56.3.M146

432

433 3. Calvani R, Marini F, Cesari M, et al. Biomarkers for physical frailty and
434 sarcopenia: state of the science and future developments. *J Cachexia Sarcopenia*
435 *Muscle*. 2015;6:278-86. doi: 10.1002/jcsm.12051.

436

437 4. Gill TM, Gahbauer EA, Han L, et al. Trajectories of disability in the last year
438 of life. *N Engl J Med*. 2010;362:1173-80.

439

440 5. Collard RM, Boter H, Schoevers RA, et al. Prevalence of frailty in community-
441 dwelling older persons: a systematic review. *J Am Geriatr Soc*. 2012;60:1487-92. doi:

442 10.1111/j.1532-5415.2012.04054.x

443

444

- 445 6. Hubbard RE, Rockwood K. Frailty in older women. *Maturitas*. 2011;69:203-207.
446 doi: 10.1016/j.maturitas.2011.04.006.
447
- 448 7. Yang Z, Hall AG. The financial burden of overweight and obesity among elderly
449 Americans: the dynamics of weight, longevity, and health care cost. *Health Serv Res*.
450 2008;43: 849-868. doi: 10.1111/j.1475-6773.2007.00801.x.
451
- 452 8. Bartali B, Frongillo EA, Bandinelli S, et al. Low nutrient intake is an essential
453 component of frailty in older persons. *J Gerontol A Biol Sci Med Sci*. 2006;61:589-93.
454
- 455 9. Jensen GL, Hsiao PY. Obesity in older adults: relationship to functional
456 limitation. *Curr Opin Clin Nutr Metab Care*. 2010;13:46-51.
457
- 458 10. Teddlie C, Yu F. Mixed methods sampling a typology with examples.
459 *J Mix Methods Res*. 2007; 1:77-100.
460
- 461 11. Birmingham City Council. Population & Census.
462 <http://www.birmingham.gov.uk/census>. Accessed March 5, 2016.
463
- 464 12. Penrod J, Preston DB, Cain RE, et al. A discussion of chain referral as a method
465 of sampling hard-to-reach populations. *Journal Transcult Nur*. 2003;14:100-7. doi:
466 10.1177/1043659602250614
467
- 468 13. Adamson AJ, Collerton J, Davies K, et al. Nutrition in advanced age: dietary
469 assessment in the Newcastle 85+ study. *Eur J Clin Nutr*. 2009;63:S6-18.

470 doi:10.1038/ejcn.2008.60.

471

472 14. Nelson M, Atkinson M, Meyer J. *A photographic atlas of food portion sizes.*

473 MAFF publications.1997.

474

475 15. International Physical Activity Questionnaire website. <http://www.ipaq.ki.se/>

476 ipaq.htm. Accessed March, 2015.

477

478 16. Hurtig-Wennlöf A, Hagströmer M, Olsson LA. The nternational Physical

479 Activity Questionnaire modified for the elderly: aspects of validity and feasibility.

480 *Public Health Nutr.* 2010;13:1847-54.

481

482 17. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Health*

483 *Res.* 2006;3:77-101. doi:10.1191/1478088706qp063oa.

484

485 18. Bunce GG, Johnson L. How many interviews are enough? An experiment

486 with data saturation and variability. *Field method.* 2006;18:59-82. doi:

487 10.1177/1525822X05279903

488

489 19. WHO, Expert Consultation. 2004. Appropriate body-mass index for Asian

490 populations and its implications for policy and intervention strategies. *Lancet.*

491 2004;363:57.

492

493 20. Department for Communities and Local Government. 2011. English indices of

494 deprivation. <https://www.gov.uk/government/collections/english-indices-of>

495 deprivation. Accessed March, 2015.

496

497 21. Wilhelm-Leen ER, Hall YN, Deboer IH, et al. Vitamin D deficiency and frailty in

498 older Americans. *J Intern Med*. 2010;268:171-80. doi:10.1111/j.13652796.2010.

499 02248.x.

500

501 22. Michelson E, Blaum C, Semba RD, et al. Vitamin and carotenoid status in older

502 women: associations with the frailty syndrome. *J Gerontol A Biol Sci Med Sci*.

503 2006;61:600-7.

504

505 23. Semba RD, Bartali B, Zhou J, et al. Low serum micronutrient concentrations

506 predict frailty among older women living in the community. *J Gerontol A Biol Sci*

507 *Med Sci*. 2006;61:594-599.

508

509 24. Bonnefoy M, Berrut G, Lesourd B, et al. Frailty and nutrition: searching for

510 evidence. *J Nutr Health Aging*. 2015;19:250-7.

511

512 25. Denke MA. Dietary retinol--a double-edged sword. *JAMA*. 2002. 287(1):102-4.

513 doi:10.1001/jama.287.1.102.

514

515 26. Haase H, Rink L. The immune system and the impact of zinc during aging. *Immun*

516 *Ageing*. 2009;6:9. doi:10.1186/1742-4933-6-9.

517

518 27. Popkin BM. The nutrition transition and obesity in the developing world. *J Nutr*.

519 2001;131: 871S-873S.

- 520 28. Roy M, Shatenstein B, Gaudreau P, et al. Seniors' Body Weight Dissatisfaction
521 and Longitudinal Associations with weight changes, anorexia of aging, and obesity.
522 Results from the NuAge Study. *J Aging Health*. 2015;27:220-238. doi:
523 10.1177/0898264314546715.
- 524
- 525 29. Lynch EB, Kane J. Body size perception among African American women. *J Nutr*
526 *Educ Behav*. 2014;46:412-417. doi:10.1016/j.jneb.2014.03.002.
- 527
- 528 30. Greenhalgh T, Chowdhury MM, Wood GW. Big is beautiful? A survey of body
529 image perception and its relation to health in British Bangladeshis with diabetes.
530 *Psychol Health Med*. 2005;10:126-138.
- 531
- 532 31. Xu B, Houston DK, Locher JL, et al. Higher Healthy Eating Index-2005 scores
533 are associated with better physical performance. *J Gerontol A Biol Sci Med Sci*.
534 2012;67:93-99. doi: 10.1093/gerona/qlr159.
- 535
- 536 32. Koster A, Penninx BW, Newman AB, et al. Lifestyle Factors and Incident
537 Mobility Limitation in Obese and Non-obese Older Adults. *Obesity*. 2001;15:3122-
538 32. doi:10.1038/oby.2007.372.
- 539
- 540 33. Telegawkar SA, Bandinelli S, Bandeen-Roche, K, et al. A higher adherence to a
541 Mediterranean-style diet is inversely associated with the development of frailty in
542 community-dwelling elderly men and women. *J Nutr*. 2012;142:2161-2166. doi:
543 10.3945/jn.112.165498.
- 544

- 545 34. Villareal DT, Apovian CM, Kushner RF, et al. Obesity in older adults: technical
546 review and position statement of the American Society for Nutrition and NAASO,
547 The Obesity Society. *Am J Clin Nutr*. 2005;82:923-34. doi:10.1038/oby.2005.228.
548
- 549 35. Redwood S, Gill PS. Under-representation of minority ethnic groups in research—
550 call for action. *Br J Gen Pract*. 2013; 63:342-343. doi:10.1093/fampra/cms054.
551
- 552 36. Landman J, Cruickshank, JK. A review of ethnicity, health and nutrition-related
553 diseases in relation to migration in the United Kingdom. *Public Health Nutr*.
554 2001;4:647-657. doi:10.1079/PHN2001148.
555
- 556 37. Beasley JM, LaCroix AZ, Neuhaus ML, et al. Protein intake and incident frailty
557 in the Women's Health Initiative observational study. *J Gerontol A Biol Sci Med Sci*.
558 2010;58:1063-71. doi:10.1111/j.1532-5415.2010.02866.x.
559
- 560 38. Jonnalagadda SS, Diwan S. Nutrient intake of First Generation Gujarati Asian
561 Indian Immigrants in the U.S. *J Am Coll Nutr*. 2002; 21:372380. doi:10.1080/
562 07315724.2002.10719238
563
- 564 39. Poslusna K, Ruprich J, de Vries JH, Jakubikova M, van't Veer P. Misreporting of
565 energy and micronutrient intake estimated by food records and 24 hour recalls,
566 control and adjustment methods in practice. *Br J Nutr*. 2009;101:S73-85. doi:
567 10.1017/S0007114509990602.
568
- 569 40. Payette H, Shatenstein B. Determinants of healthy eating in community-dwelling

570 elderly people. *Can J Public Health*. 2005;1:S27-31.

571

572

573 **Table 1. Participant Demographic Characteristics**

Variable	Mean ± SD or %	Mean ± SD or %	Mean ± SD or %	p values
	Total n=76	Non-frail n=59	Frail n=17	
Age (y)	70.5 ± 7.6	69.9 ± 6.5	74.1 ± 9.3	0.04
Residency in the UK (y)	38.73 ± 17.1	37.2 ± 17.8	44.1 ± 13.5	0.10
No. of diseases	2.3 ± 1.5	2.1 ± 1.5	3.3 ± 1.2	<0.001
Ethnicity, %				0.06
African-Caribbean	21 (27.6)	14 (23.7)	7 (41.2)	
African	10 (13.2)	10 (10.6)	0 (0)	
Arab	8 (10.5)	5 (8.5)	3 (17.6)	
Indian	20 (26.3)	17 (28.8)	3 (17.6)	
Pakistani	7 (9.2)	6 (10.2)	1 (5.9)	
Bangladeshi	5 (6.6)	2 (3.4)	3 (17.6)	
Irish	5 (6.6)	5 (8.5)	0 (0)	
IMD quintile, %				0.20
1 (most deprived)	49 (64.5)	34 (57.6)	15 (88.2)	
2	11 (14.5)	10 (16.9)	1 (5.9)	
3	7 (9.2)	7 (11.9)	0 (0)	
4-5 (less deprived)	9 (11.8)	8 (13.6)	1 (5.9)	
Education, %				0.07
No qualifications	26 (34.2)	16 (27.1)	10 (58.8)	
Primary school	8 (10.5)	6 (10.2)	2 (11.8)	
Secondary school	18 (23.7)	15 (37.3)	3 (17.6)	
Tertiary	24 (31.6)	22 (37.3)	2 (11.8)	
Marital status, %				0.60
Married	34 (44.7)	31 (52.5)	3 (17.6)	

Widowed	30 (39.5)	21 (35.6)	9 (52.9)	
Single/ separated/divorced	12 (15.8)	7 (11.9)	5 (29.4)	
Living alone, %	26 (34.2)	20 (33.9)	6 (35.3)	0.60
BMI (kg/m ²)	29.3 ± 4.9	29.1 ± 4.8	30.2 ± 5.3	0.43
Normal, %	9 (11.8)	9 (15.3)	0 (0)	
Overweight, %	23 (30.3)	16 (27.1)	7 (41.2)	
Obese, %	44 (57.9)	34 (57.6)	10 (58.8)	
WC (cm) ^a	98.8 ± 10.8	97.8 ± 11.1	102.0 ± 9.3	0.15
WHR ^a	0.92 ± 0.8	0.92 ± 0.1	0.92 ± 0.6	0.70
Unintentional weight loss, %	9 (11.8)	6 (10.2)	6 (10.2)	0.41
Supplement use, %	30 (39.5)	24 (40.7)	6 (35.3)	0.46
Energy intake (Kcals)	1243.5 ± 524.4	1379.9 ± 507.9	819.7 ± 262.5	<0.01
Frailty score (No. of frailty components, %)				NA
0	36 (47.4)	36 (61)	0 (0)	
1	23 (30.3)	23 (39)	0 (0)	
≥2	17 (22.4)	0 (0)	17 (100)	

574 ^a n=68, BMI= Body Mass Index, IMD= Index of Multiple Deprivation, WC= waist
575 circumference, WHR= waist-to-hip ratio, NA=not applicable.
576
577

578 **Table 2. Association Between Frailty Syndrome and its Components According**
579 **to the Number of Nutrients with Low Intake (n=76)**

Number of nutrients with low intake				Adjusted Odds Ratios ^a	
Variables	0	1-3	>3	Low intake of 1-3 nutrients compared to 0	Low intake of >3 nutrients compared to 0
	%	%	%	OR (95% CI)	OR (95% CI)
Frailty	17.6	29.4	52.9	3.11 (0.56-17.35)	6.58 (1.01-43.08) ^b

syndrome						
Frailty components:						
Exhaustion	24.2	26.9	35.3	0.92 (0.26-3.17)	1.12 (0.17-7.20)	
Low PA	9.1	19.2	47.1	2.30 (0.46-11.33)	5.26 (0.72- 38.10)	
Weak grip strength	18.2	26.9	17.6	0.57 (0.15- 2.16)	1.23 (0.14-10.26)	
Slow walking speed	6.1	15.4	47.1	0.85 (0.11-6.79)	1.86 (1.13-3.07) ^b	

580 ^a Adjusted for low energy intake, age and number of diseases; ^b p< 0.05

581

582

583 **Table 3. Frailty Syndrome Associated with Low Intake of Specific Nutrients**

584 **(n=76)**

Frailty ^b	
Nutrient intake ^a	OR (95% CI)
Protein (g/day)	0.76 (0.09-5.99)
Retinol (µg/day)	10.33 (1.55- 68.94) ^c
Vitamin D (µg/day)	0.96 (0.18-5.19)
Vitamin E (mg/day)	0.98 (0.17-5.68)
Vitamin C (mg/day)	3.82 (0.67-21.64)
Folate (µg/day)	0.78 (0.12- 5.06)
Calcium (mg/day)	3.87 (0.65-22.85)
Iron (mg/day)	0.94 (0.17- 5.19)
Zinc (mg/day)	8.47 (1.04-68.8) ^c

585 ^a Defined as the lowest quintile of each selected nutrient, ^b Adjusted for low energy

586 intake, age and number of diagnosed diseases, ^c $p < 0.05$

587

588

589