

# Coping Strategies in Mothers of Children with Intellectual Disabilities Showing Multiple Forms of Challenging Behaviour

Adams, Dawn; Rose, J; Jackson, N; Karakatsani, E; Oliver, Christopher

DOI:

[10.1017/S1352465817000704](https://doi.org/10.1017/S1352465817000704)

License:

Other (please specify with Rights Statement)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Adams, D, Rose, J, Jackson, N, Karakatsani, E & Oliver, C 2017, 'Coping Strategies in Mothers of Children with Intellectual Disabilities Showing Multiple Forms of Challenging Behaviour: Associations with Maternal Mental Health', *Behavioural and Cognitive Psychotherapy*, pp. 1-19. <https://doi.org/10.1017/S1352465817000704>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

© British Association for Behavioural and Cognitive Psychotherapies 2017  
University of Birmingham, UK  
<https://doi.org/10.1017/S1352465817000704>  
Published online: 29 November 2017

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# **Coping Strategies in Mothers of Children with Intellectual Disabilities Showing Multiple Forms of Challenging Behaviour: Associations with Maternal Mental Health**

[D. Adams](#)

*Autism Centre of Excellence, School of Education and Professional Studies, Griffith University, Queensland, Australia and Cerebra Centre for Neurodevelopmental Disorders, School of Psychology, University of Birmingham, UK*

J. Rose

*School of Psychology, University of Birmingham, UK*

N. Jackson

*Cerebra Centre for Neurodevelopmental Disorders, School of Psychology, University of Birmingham, UK*

E. Karakatsani

*Cerebra Centre for Neurodevelopmental Disorders, School of Psychology, University of Birmingham, UK*

C. Oliver

*Cerebra Centre for Neurodevelopmental Disorders, School of Psychology, University of Birmingham, UK*

**Background:** It is well documented that mothers of children with intellectual disabilities experience elevated mental health difficulties and that these are exacerbated by the presence of challenging behaviour. However, comparatively little is known about the effect of specific coping strategies for managing such behaviours. **Aims:** This paper aims to document coping strategies used by mothers of children showing multiple forms of challenging behaviour and to explore how these relate to positive and negative maternal mental health. **Method:** Eighty-nine mothers of children with intellectual disabilities completed questionnaires assessing

maternal mental health (Hospital Anxiety and Depression Scale, Positive and Negative Affect Scale) and maternal coping strategies (Brief COPE). **Results:** Coping strategies were not associated with child age or ability, but were associated with maternal mental health. Higher levels of problem- and positive-coping strategies were associated with higher positive affect. Although active-avoidance coping was the least frequently reported, it was associated with higher levels of negative affect and increased anxiety and depression. Moderated mediation analyses identified that active-avoidance coping mediated the relationship between the number of forms of challenging behaviour and poor maternal mental health, but only in mothers with lower levels of problem-focused coping. **Conclusions:** Active-avoidance coping is associated with poorer negative mental health in mothers of children with intellectual disabilities who have average to low levels of problem-focused coping. This is reflective of that noted within a range of populations, highlighting it as a key area for intervention.

*Keywords:* mothers, parental cognitions, coping strategies, intellectual disabilities, challenging behaviour

## Introduction

It is widely accepted that caregivers of children with an intellectual disability (ID) are at increased risk of experiencing mental health difficulties compared with parents of typically developing children (Singer, 2006). However, as literature within this area has expanded and developed it has become increasingly apparent that this relationship between parenting a child with an intellectual disability and poorer mental health is more complex than initially described. Child factors (e.g. age, ability, challenging behaviour) have been explored with varying results.

A number of studies have focused upon the relationship between challenging behaviour and maternal mental health and have documented a strong association between the presence and the extent (e.g. frequency, severity) of challenging behaviour and elevated levels of maternal stress and mental health difficulties (e.g. Hastings, 2002; Mash and Johnston, 1990; Tomanik et al., 2004). However, the variability in how parents respond to the challenges of raising a child with an intellectual disability and challenging behaviour suggests that there may be a range of parental factors (e.g. socio-economic status; Giallo and Gavidia-Payne, 2006) and cognitive or psychological processes influencing the relationship (Horsley and Oliver, 2015), either by mediation or moderation. These have also begun to be explored in the positive psychology literature, exploring the underlying cognitive factors contributing to the variability in positive well-being in parents raising children with intellectual disabilities (Beighton and Wills, 2016). Models such as the Model of Parent–Child Interactive Stress (Mash and Johnston, 1990), aim to explain this variability through inter-linking factors that combine or interact to contribute to stress in parents and families.

Coping strategies have repeatedly been suggested as one of the key mechanisms by which parents adapt to the process and stresses associated with raising a child with a disability (Smith et al., 2008) and therefore influence their mental health and well-being. Hastings et al. (2005) reported the results of a multidimensional coping inventory, the Brief COPE, in mothers and fathers of children with autism, and explored the relationship between coping strategies and parental well-being. Factor analyses of the 28 items from the Brief COPE identified four factors; active-avoidance coping (nine items), problem-focused coping (seven items), positive coping (six items) and religious/denial coping (four items). There was no effect of child age on coping

strategies reported but there was an effect of parent gender, with mothers reporting more active-avoidance and problem-focused coping strategies than fathers. Mothers who showed more active-avoidance coping strategies reported more symptoms of anxiety and depression. Religious/denial coping was associated with depression in mothers and depression and anxiety in fathers. Finally, parents who adopted more positive coping strategies reported lower levels of depression. In a longitudinal study, Benson (2010) also used the Brief COPE in a sample of 113 mothers of children with autism. Although using a different factor structure, he found a similar profile of results; higher use of avoidant strategies (distraction and disengagement) increased parental distress but problem-focused coping strategies (engagement and cognitive reframing) had minimal impact on reducing maternal distress. Benson also noted that maternal well-being and positive health were associated with higher levels of cognitive reframing and lower levels of disengagement coping strategies. However, neither of these studies explored the impact of challenging behaviour on maternal coping strategies and how these may affect the relationship with maternal mental health.

Although there has been a considerable focus upon coping strategies in parents of children with autism, even within this population, the nature of parenting stress and coping remains inconclusive, especially when compared with parents of children without autism or ID (Lai and Oei, 2014; Lai et al., 2015; Sivberg, 2002). There has been comparatively less exploration of these factors within the ID literature. Consequently, although most studies agree on a relationship between challenging behaviour and elevated maternal distress, there is less consensus on the role of cognitions, with some research suggesting a mediating or moderating effect (e.g. Baker et al., 2005), whilst others (e.g. Hill and Rose, 2009) noting no effect. It may therefore be the case that such effects may not always be present, but may only be present in specific subgroups of individuals. As the coping styles literature has such mixed results, it would be reasonable to assume that perhaps coping styles may not have independent relationships with mental health, but may combine to make profiles of high/low coping styles that are at more or less risk for mental health difficulties. Such relationships have been noted in other areas of mental health and intervention research (e.g. Karakus, 2013; Wiedemann et al., 2009) but to date have not been explored within the intellectual disability literature. This, coupled with a very limited focus on specific types of cognitions or behaviours, such as coping strategies, in this field, highlight the need for further research in the area.

Rose et al. (2016) note that one limitation in the parental cognitions literature is the lack of discrimination between topographies of challenging behaviour. Given that meta-analyses have identified different risk markers and prevalence rates for different topographies of behaviour (e.g. McClintock et al., 2003; Petty and Oliver, 2005), that topographies of challenging different behaviour influence different referral patterns (Adams et al., 2008) and service use (Adams et al., 2016) and influence staff cognitions (e.g. Dilworth et al., 2011), it seems imperative that literature focusing upon parental well-being now begins to consider differences between different forms of challenging behaviours.

The aim of the present study is firstly to document the most frequently used or reported coping strategies within a sample of mothers with children with intellectual disabilities, and secondly to investigate whether these coping strategies mediate any relationship between child factors (age, challenging behaviour and adaptive functioning) and maternal psychological functioning. The final aim of this study is to explore whether each mediation within the Model of Parent–Child Interactive Stress is independent for each coping style, or whether coping styles may influence (or moderate) the mediation within other models. This will be explored through

moderated mediation, which explores whether the mediation model remains significant when another factor (the moderator) is changed.

Based upon the previous literature and the Model of Parent–Child Interactive Stress (Hassall *et al.*, 2005; Mash and Johnston, 1990; Rose *et al.*, 2016), the following hypotheses are made:

- (1) There will be a positive relationship between the number of forms of challenging behaviour and poorer maternal mental health (elevated maternal anxiety and depression, elevated negative affect and/or reduced positive affect).
- (2) There will be no association between child age or ability and measures of maternal mental health.
- (3) There will be a relationship between (a) increased active-avoidance coping strategies and poorer maternal mental health and (b) increased positive or problem-focused coping and better maternal mental health.
- (4) Maternal coping strategies will mediate the relationship between a number of forms of challenging behaviour and maternal mental health.
- (5) The mediating effect of maternal coping strategies will differ based on the topography of challenging behaviour.

As there is no previous work exploring moderated mediation in mothers of children with intellectual disabilities with or without challenging behaviour, no directional hypotheses are made.

## Method

### *Ethics*

Ethical approval was received from the University of Birmingham Research Ethics Committee.

### *Participants*

The sample consisted of a subgroup of parents/carers recruited to a larger study investigating the behavioural functioning of children and adults diagnosed with rare genetic syndromes associated with neurodevelopmental disabilities and intellectual disabilities (e.g. Moss *et al.*, 2009). This sample was deemed suitable for this study given that recruiting a sample of children with mixed aetiologies has been used in a range of previous studies exploring psychological factors in parents of children with ID (e.g. Lloyd and Hastings, 2009).

Three hundred and eighty-seven parents and carers of children aged 15 or under with one of eleven rare genetic syndromes (Angelman, Cri du Chat, Cornelia de Lange, Prader-Willi, Lowe, Smith-Magenis, Sotos, 1p36 deletion, 9q34 deletion (Kleefstra syndrome), 8p23 deletion and Phelan McDermid) were invited to complete an online questionnaire by letter or email. Recruitment adverts were also placed on the research group's Facebook page. A total of 113 (29.20%) participants agreed to take part. Twenty-four participants were excluded because they were not the child's mother or adoptive mother (i.e. were fathers, grandparents, foster carers or paid carers) or because they had not completed all of the measures in the online questionnaire.

Following the exclusions, the final sample consisted of 89 mothers of children with one of eleven neurodevelopmental disorders: Angelman syndrome (AS,  $n = 14$ ), Cornelia de Lange

**Table 1.** Demographic information for all participants ( $n = 89$ )

Demographic	$n$ (%)
Carer's age (years, mean, $SD$ )*	41.5 (6.5)
Carer's education (to degree level)*	39 (43.7%)
Child's age (years, mean, $SD$ )	10.2 (4.1)
Gender (male)	46 (51.6%)
Vision (normal)	57 (64%)
Speech (verbal/partly verbal)	60 (67.4%)
Mobility (mobile)	50 (56.2%)
Self-help category (partly able/able)	44 (49.4%)

\*Seven mothers chose not to provide this information.

syndrome (CdLS,  $n = 9$ ), Cri du Chat syndrome ( $n = 11$ ), Lowe syndrome ( $n = 2$ ), Phelan-McDermid syndrome (PMS,  $n = 5$ ), Prader-Willi syndrome (PWS,  $n = 15$ ), Soto syndrome ( $n = 6$ ), Smith-Magenis syndrome (SMS,  $n = 9$ ), 1p36 deletion syndrome ( $n = 12$ ), 9q34 syndrome ( $n = 1$ ) and 8p23 syndrome ( $n = 5$ ). All children lived at home full-time with their carers. Demographic information on mothers and children are presented in [Table 1](#).

### Measures

*Demographic information.* A background questionnaire was used to collect information about mothers' age and level of education, children's age, genetic or clinical diagnoses, gender, speech and mobility and living arrangements. Mothers completed the following three questionnaires addressing the dimensions of their well-being and coping strategies.

The *Brief COPE* is a 28-item shortened version of the original COPE scale developed by Carver et al. (1989). It consists of 14 subscales, each containing two items that represent a coping strategy. Participants rate the extent to which they use a coping strategy for their child by indicating either 'I haven't been doing this at all' (score of 1), 'I've been doing this a little bit' (score of 2), 'I've been doing this a medium amount' (score of 3) or 'I've been doing this a lot' (score of 4). At the point of development of the scale, Carver stated that all scales of the *Brief COPE* were at least minimally acceptable in terms of internal reliability with three of the scales having an  $\alpha$  value of 0.50–0.59, with all remaining scales having  $\alpha$  values of 0.60 upwards (Carver, 1997). The four-factor subscale structure proposed by Hastings et al. (2005) was chosen over the four-factor structure suggested by Benson, as the factors used by Hasting et al. are based on the individual items rather than subscales. In this sample, with the exception of the religious/denial subscale, the Cronbach's alpha statistics were similar to that reported by Hastings et al., being .74 for active-avoidance, .85 for problem-focused, .71 for positive coping and .54 for religious/denial coping. Results will be reported in two ways: coping strategies and factor (subscale) groupings suggested by Hastings et al.

The *Hospital Anxiety and Depression Scale* (HADS; Zigmond and Snaith, 1983) was used to examine self-reported levels of anxiety and depression. The items are rated on a scale of 0 to 3, and indicate the degree to which the participant agrees with a given item. This gives a maximum total of 21 for each of the subscales, with clinical cut-off points in terms of severity of presentation based upon the composite scores. Although this measure was published in

1983, it continues to have good test–retest reliability and concurrent validity data reported at subscale and total score levels for parents of children with neurodevelopmental disabilities (Bjelland et al., 2002; Jones et al., 2014). In this sample, Cronbach’s alpha was .72 for anxiety and .85 for depression.

The *Positive and Negative Affect Scale* (PANAS; Watson et al., 1988) consists of twenty descriptive words, ten reflecting positive affect and ten reflecting negative affect. Participants were asked to rate the extent to which they felt this way over the past week on a Likert-type scale. A higher score reflects a higher frequency of the emotions. The PANAS has been shown to have excellent internal reliability (Watson et al., 1988). In this sample, Cronbach’s alpha was .92 for the positive and the negative emotions subscales.

The *Wessex Scale* (Kushlick et al., 1973) measures both social and physical capacity of people with disabilities. Carers answer a number of questions that relate to physical capacity including vision, hearing and mobility, and social capacity such as communication, literacy and self-care. Since its development, the validity and reliability of the scale have been robustly supported (Oliver et al., 2012). Higher scores on this measure are an indication of greater level of ability.

The *Challenging Behaviour Questionnaire* (CBQ; Hyman et al., 2002) evaluates the presence or absence of self-injury, aggression, destruction of property and stereotyped behaviour in the last month. Previous examination of the psychometric properties of the questionnaire has demonstrated good inter-rater reliability with reliability coefficients ranging from 0.61 to 0.89 (Hyman et al., 2002). This study uses the data detailing the presence/absence of four behaviours that the questionnaire describes as challenging behaviours (self-injury, aggression, destruction of the environment and stereotyped behaviours) and the number of forms of challenging behaviour rated as present (0–4).

## Results

### *Number of challenging behaviours*

Only nine (10.1%) of children were reported to have shown no challenging behaviour in the previous month, 18 (20.2%) were reported to have shown one topography, 11 (12.4%) two topographies, 21 (23.6%) three topographies and 30 (33.7%) four topographies of challenging behaviour within the month preceding questionnaire completion.

### *Measures of maternal mental well-being*

The mean anxiety score was 10.8 ( $SD = 3.8$ ) with a minimum score of 2 and maximum of 17. The mean depression score was 6.9 ( $SD = 4.2$ ) with a minimum score of 0 and maximum of 18. Both the positive and negative affect scores ranged from the minimum to maximum (10 to 50) with a mean for positive affect of 30.1 ( $SD = 9.7$ ) and negative affect of 20.9 ( $SD = 8.9$ ).

### *Coping strategies used*

In order to complete the first aim of the study and document the most frequent coping strategies used by parents of children with intellectual disabilities, the mean score for each of the fourteen strategies and four subscales were calculated and are presented in Table 2. Due to differing

**Table 2.** Mean scores per item for each coping strategy (from most to least frequently used)

Subscale	Mean	SD
Acceptance	3.44	0.74
Planning	2.88	0.92
Active coping	2.86	0.90
Positive reframing	2.66	0.87
Instrumental support	2.58	0.82
Emotional support	2.27	0.89
Self-distraction	2.08	0.81
Self-blame	1.84	0.84
Humour	1.84	0.86
Venting	1.78	0.69
Substance use	1.50	0.83
Religion	1.45	0.84
Behavioural disengagement	1.27	0.50
Denial	1.14	0.44
<b>Factor</b>		
Positive coping	2.80	0.69
Problem-focused coping	2.70	0.70
Religious/denial	1.84	0.52
Active-avoidance coping	1.64	0.47

numbers of questions in each subscale, the mean score per question is reported for each of the four factors recommended by Hastings et al. (2005). For ease of comparison, the scores for each subscale are also presented as mean score per item. Therefore, for each row of data in Table 2, the minimum score is 1 and the maximum score is 4.

The most frequently used coping strategy was acceptance, followed by planning, active coping and positive reframing. The coping strategies that parents reported using the least were substance use, religion, behavioural disengagement and denial. This is also reflected in the four subscale scores, with positive coping being the most frequently reported coping method and the active-avoidance coping strategy being the least frequently reported.

#### *Association between child variables and maternal mental health*

In order to test the first and second hypotheses, that an increased number of forms of challenging behaviours will be associated with poorer maternal mental health but that there will be no relationship between forms of challenging behaviour and child variables (age and ability), a series of correlations were conducted. The results of these correlations are presented in Table 3. Alpha was set to .01 to correct for multiple correlations.

The results indicate a positive correlation between the number of forms of challenging behaviour and increased maternal negative affect, anxiety and depression, but no association with positive affect. Hypothesis 1 can be partially accepted. There was no association between child age or ability and maternal mental health. Hypothesis 2 can therefore be accepted.

**Table 3.** Pearson's correlation coefficients and *p* values between coping strategies, child variables, maternal age and maternal mental health

		1	2	3	4	5	6	7	8	9	10	11
Child factors	1. Child age	–										
	2. Wessex ability score	.33	–									
	3. Forms of challenging behaviour	.21	–.08	–								
	4. Maternal age	.38**	.11	–.02	–							
Maternal coping strategies	5. Problem-focused coping	–.06	.09	.02	.08	–						
	6. Positive coping	–.04	.10	–.02	.02	.60**	–					
	7. Religious/denial	–.03	.35	.08	–.12	.29*	.23	–				
	8. Active-avoidance coping	.03	.02	.39**	.03	.30*	.09	.38**	–			
Maternal mental health	9. Positive affect	–.04	–.02	–.23	.17	.33*	.45**	.13	–.27	–		
	10. Negative affect	–.14	–.04	.32*	–.06	.09	–.05	.18	.68**	–.36*	–	
	11. Anxiety	–.08	–.08	.33*	–.06	.16	–.07	.11	.42**	–.39**	.65**	–
	12. Depression	.01	–.04	.32*	–.03	–.05	–.25	–.01	.40**	–.58**	.55**	.68**

\**p* < .01; \*\**p* < .001.

### *Associations between coping strategies and maternal mental health*

To address hypothesis 3, the mean score per item was calculated for each of the four subscales and correlated with the four subscales assessing maternal mental health: positive affect, negative affect, anxiety and depression. As shown in Table 3, higher levels of problem-focused and positive coping strategies are associated with increased positive affect. Higher levels of active avoidance coping strategies are associated with higher levels of negative affect and increased levels of anxiety and depression. The religious/denial coping strategy was not associated with any of the maternal mental health variables measured. Hypothesis 3 can therefore be accepted.

### *The mediating effect of coping strategies between number of forms of challenging behaviour and maternal mental health*

The Parent–Child Interactive Stress model predicts that maternal cognitions mediate the relationship between challenging behaviour and maternal psychological distress. Table 3 shows that only the active-avoidance subscale was correlated with both the number of forms of challenging behaviour and three of the measures of maternal mental health (depression, anxiety and negative affect). Therefore, only this variable was entered into the mediation analyses which used the bootstrapping method with bias-corrected confidence estimates (MacKinnon et al., 2004; Preacher and Hayes, 2004). In the present study, the 95% confidence interval of the indirect effects was obtained with 5000 bootstrap resamples (Preacher and Hayes, 2008). The results are summarized in the upper section of Table 4.

#### *Maternal anxiety*

Results of the mediation analysis confirmed the mediating role of active-avoidance coping strategies in the relationship between challenging behaviour and maternal anxiety ( $b = .89$ ,  $SE = .28$ ,  $p = .002$ ). The overall model was significant ( $R^2 = .22$ ;  $p < .0001$ ) and accounted for 22% of the variance in maternal anxiety. The number of forms of challenging behaviour was no longer a significant predictor of maternal anxiety after controlling for the mediator ( $b = .55$ ,  $SE = .28$ ,  $p = .05$ ), consistent with full mediation.

#### *Maternal depression*

Results of the mediation analysis confirmed the mediating role of active-avoidance coping strategies in the relationship between challenging behaviour and maternal depression ( $b = .92$ ,  $SE = .32$ ,  $p = .005$ ). The overall model was significant ( $R^2 = .18$ ;  $p = .002$ ) and accounted for 18% of the variance in maternal depression. The number of forms of challenging behaviour was no longer a significant predictor of maternal depression after controlling for the mediator ( $b = .57$ ,  $SE = .32$ ,  $p = .08$ ), consistent with full mediation.

#### *Maternal negative affect*

Results of the mediation analysis confirmed the mediating role of active-avoidance coping strategies in the relationship between challenging behaviour and maternal negative affect ( $b = 2.06$ ,  $SE = .67$ ,  $p = .003$ ). The overall model was significant ( $R^2 = .47$ ;  $p < .0001$ )

**Table 4.** Mediation of the effect of challenging behaviour (number of forms and the specific topography of destruction of the environment) on maternal mental health through active-avoidance coping

			Point estimate	Bootstrap path estimate	Bias	Standard error	Lower BC 95% CI	Upper BC 95% CI
Number of forms of challenging behaviour	Maternal anxiety	Active-avoidance coping	.33	.32	-.008	.12	.14	.62
		Total	.33	.32	-.008	.12	.14	.62
	Maternal depression	Active-avoidance coping	.34	.35	.005	.14	.14	.70
		Total	.34	.35	.005	.14	.14	.70
	Maternal negative affect	Active-avoidance coping	1.4	1.4	.008	.44	.65	2.4
		Total	1.4	1.4	.008	.44	.65	2.4
Presence of destruction of the environment	Maternal anxiety	Active-avoidance coping	.72	.72	-.007	.33	.21	1.56
		Total	.72	.72	-.007	.33	.21	1.56
	Maternal depression	Active-avoidance coping	.84	.84	.006	.37	.26	1.73
		Total	.84	.84	.006	.37	.26	1.73

$n = 89$ ; bootstrap sample size = 5000; BC: bias-corrected.

and accounted for 47% of the variance in maternal negative affect. Forms of challenging behaviour were no longer a significant predictor of maternal negative affect after controlling for the mediator ( $b = .65, SE = .55, p = .24$ ), consistent with full mediation.

Hypothesis 4 can be partially accepted, as only the active-avoidance coping strategy mediated the relationship between the number of forms of challenging behaviour and maternal anxiety, depression and negative affect.

#### *Exploring a moderated mediation model of number of forms of challenging behaviour to maternal mental health*

Given the correlations between some of the coping styles reported in [Table 3](#), it is not unreasonable to assume some relationship between them or their influence on other factors. In order to identify whether any of the coping styles that were not entered into the above mediations have a statistical influence on the strength of the mediations between the number of forms of challenging behaviour to maternal mental health, moderated mediation analyses were undertaken. This aimed to explore whether the three coping styles that did not mediate the relationship between the number of forms of challenging behaviour and maternal mental health (problem-focused, positive and religious/denial coping) moderate the mediating effect of active-avoidant coping. A bootstrapped Model 8 of the PROCESS Macro (Hayes, 2012) identified two moderated mediation models (moderation of the direct effect only) that are summarized in [Fig. 1](#).


Further inspection of the data for both maternal anxiety and depression notes that at low (one standard deviation below the mean) or average levels of problem-focused coping, the indirect effect is significant, i.e. the mediation is present (anxiety low problem-focus coping CI .09 to .74; average problem-focus coping CI .07 to .57; depression low problem-focus coping CI .17 to .96; average problem-focus coping CI .13 to .73). However, when the levels of problem-focused coping are high (+1SD), both the direct and the indirect effect is non-significant and therefore the relationship between forms of challenging behaviour and maternal mental health is not mediated (indirect; anxiety CI  $-.01$  to  $.60$ , depression CI  $-.04$  to  $.74$ ; direct anxiety CI  $-.90$  to  $.57$ , depression high problem-focus coping CI  $-1.0$  to  $.70$ ).

#### *Exploring the mediating effect of coping strategies based upon topography of challenging behaviour*

Given that the topography of challenging behaviour has been shown to impact upon maternal stress and cognitions by Rose et al. (2016), the final stage of the analysis planned to repeat the mediation analyses with the dependent variable being the presence/absence of the specific topography of challenging behaviour. The active-avoidance subscale was the coping strategy subscale that correlated with both a specific topography of challenging behaviour (destruction of the environment;  $r = .29, p = .007$ ) and measures of maternal depression ( $r = .30, p = .004$ ) and anxiety ( $r = .32, p = .003$ ). Therefore, only these variables were entered into the mediation analyses. The results are summarized in the lower section of [Table 4](#).

#### *Maternal anxiety*

Results of the mediation analysis confirmed the mediating role of active-avoidance coping strategies in the relationship between the presence of destruction of the environment and



**Figure 1.** Moderated mediation model for the relationship between challenging behavior and maternal mental health. \* $p < .05$ ; \*\* $p < .01$ ; \*\*\* $p < .001$ .

maternal anxiety ( $b = 2.3$ ,  $SE = .80$ ,  $p = .004$ ). The overall model was significant ( $R^2 = .19$ ;  $p = .0001$ ) and accounted for 19% of the variance in maternal anxiety. The presence of destruction of the environment was no longer a significant predictor of maternal anxiety after controlling for the mediator ( $b = 1.6$ ,  $SE = .79$ ,  $p = .05$ ), consistent with full mediation.

#### *Maternal depression*

Results of the mediation analysis confirmed the mediating role of active-avoidance coping strategies in the relationship between the presence of destruction of the environment and

maternal depression ( $b = .92, SE = .32, p = .005$ ). The overall model was significant ( $R^2 = .19; p = .002$ ) and accounted for 19% of the variance in maternal depression. The presence of destruction of the environment was no longer a significant predictor of maternal depression after controlling for the mediator ( $b = .16, SE = .89, p = .07$ ), consistent with full mediation.

Hypothesis 5 can be partially accepted as only the active-avoidance coping strategy mediates the relationship between the presence of destruction of the environment and maternal anxiety and depression.

Follow-up exploration of potential moderated mediation pathways for the two mediation pathways described above were undertaken but no significant moderators of the mediation were found.

## Discussion

The results showed that although active-avoidance coping was the least frequently reported coping strategy, it is associated with higher levels of negative affect and increased levels of anxiety and depression. It mediates the relationship between the number of forms of challenging behaviour and maternal anxiety, depression and negative affect, but only when mothers have average or lower problem-focused coping. The mediation is not present in mothers with higher levels of problem-focused coping. Active-avoidance coping also mediates the relationship between the presence of destruction of the environment and maternal anxiety and depression. As per previous research into parents of children with autism, higher levels of problem-focused and positive coping strategies are associated with increased positive affect.

Active-avoidance coping, which includes behaviours such as self-blaming, self-criticizing, doing something to think about it less, and using drugs or alcohol to get through or make oneself feel better, have been associated with negative mental health in a range of samples, including cancer patients (e.g. Stanton and Snider 1993), caregivers for individuals with chronic disease (Billings et al., 2000), hospitalized burn patients (Fauerbach et al., 2002), and individuals coping with terrorist attacks (Silver et al., 2002). It is also associated with less positive outcomes for interventions (see Taylor and Stanton, 2007, for a thorough review). Given the robustness of its influence across a range of populations, it is unsurprising that it was the consistent mediating factor between forms of challenging behaviour and maternal anxiety, depression and negative affect, and the presence of destruction of the environment and maternal anxiety and depression in this study. The model accounted for 47% of the variability in negative affect scores, but only 22% of the anxiety and depression scores. This may be due to the different forms of measurement, with the negative affect requiring a Likert rating of the frequency of negative emotions and the anxiety and depression subscales requiring a Likert rating of specific behaviours, such as 'I get sudden feelings of panic' or 'I have lost interest in my appearance'. It may be that a rating scale such as the Hamilton Anxiety Rating Scale (HAM-A; Hamilton, 1959), which lists physical symptoms of anxiety rather than behaviours, would provide a different profile of results.

Longitudinal studies documenting coping strategies and mental health in parents of individuals with intellectual disabilities (e.g. Seltzer et al., 1995) suggest that certain coping strategies can 'buffer' the negative effects of caring for an individual with intellectual disabilities, therefore reducing the risk of later stress or mental health difficulties. The results of this study support this assertion, as the results of the moderated mediation model show that there was no significant direct or indirect relationship between the number of challenging

behaviours and maternal anxiety or depression in mothers who have strong problem-focused coping. The protective factor of problem-focused coping has been previously identified by Essex *et al.* (1999) who noted that problem-focused coping strategies at baseline were predictive of levels of depression 7.5 years later, but only for parents of children with more severe intellectual disabilities. Woodman and Hauser-Cram (2013) also noted that problem-focused coping strategies moderated the impact of caregiving demands on depressive symptoms three years later.

The recent focus upon parenting interventions for children with intellectual disabilities, most notably children with autism, has led to the recognition that parental factors, as well as child factors, may influence the effectiveness of this intervention and resulting child outcomes (see Hastings and Johnson, 2001). Given that parental mental health is associated with a child's progress and response to intervention (Robbins *et al.*, 1991), it could be posited that working with parental cognitions and behaviours in addition to a child's behaviour would be the most effective approach to reducing difficult or challenging behaviour within this population. The results within this research help to identify parents who may benefit most from interventions: those showing high active-avoidant coping styles with average or below problem-focused coping. Interventions would therefore encourage adjustments towards more helpful, problem-focused coping strategies (which could be achieved using a cognitive behavioural model) associated with improvements with parental mental health and reduce active-avoidant coping styles.

Although active-avoidant coping may be the least frequently reported coping style, its impact on the relationship between challenging behaviour and maternal anxiety, depression and negative affect is significant. Interventions specifically targeted at reducing the use of active-avoidance strategies and encouraging problem-focused and positive coping strategies have been shown to reduce perceived stress and depression and increase self-efficacy in populations, including patients who were HIV positive (e.g. Chesney *et al.*, 2003) and those with chronic pain (Keefe *et al.*, 2004; Rhee *et al.*, 2000). Given that active-avoidance coping appears to have a similar influence on a range of populations, it may be that these interventions could be easily adapted and trialled in parents of children with ID, perhaps initially targeting parents of children who have the mix of high active-avoidant coping styles with average or below problem-focused coping that strengthens the relationship between challenging behaviour and mental health.

The relationship between only one topography of challenging behaviour (destruction of the environment) and maternal anxiety and depression was mediated by the active-avoidance coping style, although the model only explained 19% of the variance. Given that the prevalence was similar for each of the four forms of challenging behaviour in the CBQ (60.7–67.4%), and that almost half (46%) of children who were showing one to three forms of challenging behaviour did not show destruction of the environment, the mediated relationship between numbers of topographies of challenging behaviour and maternal anxiety and depression cannot simply be explained by the presence or absence of destruction of the environment. Previous studies (e.g. Adams *et al.*, 2008, 2016) have discussed how behaviours that have more of an external impact (e.g. destruction of the environment, wandering) affect people differently to those that have more of an impact upon the person showing the behaviour (e.g. self-injury, withdrawal from social activities). Whilst this could be used to explain the relationship between destruction of the environment and maternal mental health (which is mediated by active-avoidance coping), it cannot explain why such a relationship is not present between aggression

and maternal mental health. It may be that aggression has a different impact than destruction of the environment, as one could argue that the locus of the impact differs (i.e. in one scenario, a person is at risk of being harmed whilst in the other, material goods are at risk). Further research, with larger samples and a more detailed measure of challenging behaviour, such as the Behavior Problem Inventory (Rojahn et al., 2001), would allow for greater insight into this model and explore such relationships with a range of topographies of challenging behaviour.

Whilst it is important to note the associations, it is also important to note the key non-significant associations, as these are also informative for researchers, yet are often over-looked. First, parental coping styles were not associated with child age or ability score on the Wessex scale. This is concordant with the finding that parental stress levels are influenced by some child factors, such as challenging behaviour, health problems and sleep difficulties, but not level of cognitive ability (Cahill and Glidden, 1996). Second, there was no association between any measures of parental mental well-being and the religious/denial coping subscale. However, it may be that these factors may have more of an individual influence rather than benefit measurable within a group design.

There are a number of limitations to this study. Whilst the Brief-COPE is a well-validated measure of coping strategies, it is limited in how much information it can accrue on the experience of parenting a child with an intellectual disability. The Cronbach's alpha for the religious/denial subscale was .54, suggesting that this subscale has unsatisfactory reliability within this sample. This subscale may therefore not reflect a unidimensional construct within this sample and may have reflected multiple constructs that had similar or differing relationships with the dependent or independent variables. Although there is a lot of research exploring mediator variables cross-sectionally, it is acknowledged that mediation models are stronger and more accurate when completed with longitudinal data. This study should therefore be repeated with a longitudinal design.

Whilst the heterogeneous sample in this study allowed exploration of coping strategies and well-being across mothers of children with an array of rare genetic syndromes, it may also bring increased variability into the sample. It may be that parents of children with rare genetic syndromes cope or adapt to behaviour problems differently than parents of idiopathic intellectual disabilities, perhaps by attributing the behaviour to the syndrome, although this is yet to be explored within the literature. Also, the data were collected via questionnaire rather than by direct assessment, limiting the detail with regard to behaviour and ability or IQ. Finally, it is important to reflect on the generalizability of the sample. Given that these mothers were originally recruited via parent support groups, and that they had the time to complete an online questionnaire, there is a potential for a recruitment bias.

### **Conclusions**

Understanding the factors that influence or explain increased levels of mental health difficulties, stress and elevated negative emotions in this population is important if effective treatments are to be implemented. This study highlights that, as reported in mothers of children with autism, specific coping strategies are associated with increased psychological well-being and distress. Higher levels of problem-focused and positive coping strategies are associated with higher levels of positive affect. Active-avoidance coping strategies, although the least commonly reported, have the strongest impact upon maternal mental health, being associated with higher levels of negative affect, increased levels of anxiety and depression and mediating

the relationship between challenging behaviour and maternal mental health, but only within those with average or below problem-focused coping skills. This highlights the importance of detailed research and analysis of data in order to inform interventions for these coping strategies when implementing both parental or behavioural interventions.

### Acknowledgements

We are grateful to all the parents and families for taking part in this research and for the syndrome support groups for facilitating the data collection.

*Conflicts of interest:* The authors have no conflicts of interest with respect to this publication.

*Ethical statement:* The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, and its most recent revision.

*Financial support:* This paper presents independent research partly funded by the National Institute for Health Research (NIHR) Post-Doctoral Fellowship Programme (D.A. grant number PDF-2010-03-02), Jerome Lejeune, and Cerebra. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, Cerebra or the Department of Health.

### References

- Adams, D., Handley, L., Simkiss, D., Walls, E., Jones, A., Knapp, M., Romeo, R. and Oliver, C. (2016). Service use and access in young children with developmental delay and challenging behavior. *Journal of Intellectual and Developmental Disability*. doi: [10.3109/13668250.2016.1238448](https://doi.org/10.3109/13668250.2016.1238448)
- Adams, D., Oliver, C., Kalsy, S., Peters, S., Broquard, M., Basra, T. et al. (2008). Behavioural characteristics associated with dementia assessment referrals in adults with Down syndrome. *Journal of Intellectual Disability Research*, 52, 358–368. doi: [10.1111/j.1365-2788.2007.01036.x](https://doi.org/10.1111/j.1365-2788.2007.01036.x)
- Baker, B. L., Blacher, J. and Olsson, M. B. (2005). Preschool children with and without developmental delay: behaviour problems, parents' optimism and well-being. *Journal of Intellectual Disability Research*, 49, 575–590. doi: [10.1111/j.1365-2788.2005.00691.x](https://doi.org/10.1111/j.1365-2788.2005.00691.x)
- Beighton, C. and Wills, J. (2016). Are parents identifying positive aspects to parenting their child with an intellectual disability or are they just coping? A qualitative exploration. *Journal of Intellectual Disabilities*. doi: [10.1177/1744629516656073](https://doi.org/10.1177/1744629516656073)
- Benson, P. R. (2010). Coping, distress, and well-being in mothers of children with autism. *Research in Autism Spectrum Disorders*, 4, 217–228. doi: [10.1016/j.rasd.2009.09.008](https://doi.org/10.1016/j.rasd.2009.09.008)
- Billings, D. W., Folkman, S., Acree, M. and Moskowitz, J. T. (2000). Coping and physical health during caregiving: the roles of positive and negative affect. *Journal of Personality and Social Psychology*, 79, 131–142. doi: [10.1037//0022-3514.79.1.131](https://doi.org/10.1037//0022-3514.79.1.131)
- Bjelland, I., Dahl, A. A., Haug, T. T. and Neckelmann, D. (2002). The validity of the Hospital Anxiety and Depression Scale – an updated literature review. *Journal of Psychosomatic Research*, 52, 69–77. doi: [S0022-3999\(01\)00296-3](https://doi.org/S0022-3999(01)00296-3)
- Cahill, B. M. and Glidden, L. M. (1996). Influence of child diagnosis on family and parental functioning: Down syndrome versus other disabilities. *American Journal on Mental Retardation*, 101, 149–160.
- Carver, C. S. (1997). You want to measure coping but your protocol's too long: consider the brief COPE. *International Journal of Behavioral Medicine*, 4, 92–100. doi: [10.1207/s15327558ijbm0401\\_6](https://doi.org/10.1207/s15327558ijbm0401_6)

- Carver, C. S., Scheier, M. F. and Weintraub, J. K.** (1989). Assessing coping strategies – a theoretically based approach. *Journal of Personality and Social Psychology*, *56*, 267–283. doi: [10.1037//0022-3514.56.2.267](https://doi.org/10.1037//0022-3514.56.2.267)
- Chesney, M. A., Chambers, D. B., Taylor, J. M., Johnson, L. M. and Folkman, S.** (2003). Coping effectiveness training for men living with HIV: results from a randomized clinical trial testing a group-based intervention. *Psychosomatic Medicine*, *65*, 1038–1046. doi: [10.1097/01.Psy.0000097344.78697.Ed](https://doi.org/10.1097/01.Psy.0000097344.78697.Ed)
- Dilworth, J. A., Phillips, N. and Rose, J.** (2011). Factors relating to staff attributions of control over challenging behaviour. *Journal of Applied Research in Intellectual Disabilities*, *24*, 29–38. doi: [10.1111/j.1468-3148.2010.00570.x](https://doi.org/10.1111/j.1468-3148.2010.00570.x)
- Essex, E. L., Seltzer, M. M. and Krauss, M. W.** (1999). Differences in coping effectiveness and well-being among aging mothers and fathers of adults with mental retardation. *American Journal on Mental Retardation*, *104*, 545–563. doi: [10.1352/0895-8017\(1999\)104](https://doi.org/10.1352/0895-8017(1999)104)
- Fauerbach, J. A., Heinberg, L. J., Lawrence, J. W., Bryant, A. G., Richter, L. and Spence, R. J.** (2002). Coping with body image changes following a disfiguring burn injury. *Health Psychology*, *21*, 115–121. doi: [10.1037//0278-6133.21.2.115](https://doi.org/10.1037//0278-6133.21.2.115)
- Giallo, R. and Gavidia-Payne, S.** (2006). Child, parent and family factors as predictors of adjustment for siblings of children with a disability. *Journal of Intellectual Disability Research*, *50*, 937–948.
- Hamilton, M.** (1959). The assessment of anxiety-states by rating. *British Journal of Medical Psychology*, *32*, 50–55.
- Hassall, R., Rose, J. and McDonald, J.** (2005). Parenting stress in mothers of children with an intellectual disability: the effects of parental cognitions in relation to child characteristics and family support. *Journal of Intellectual Disability Research*, *49*, 405–418. doi: [10.1111/j.1365-2788.2005.00673.x](https://doi.org/10.1111/j.1365-2788.2005.00673.x)
- Hastings, R. P.** (2002). Parental stress and behaviour problems of children with developmental disability. *Journal of Intellectual and Developmental Disability*, *27*, 149–160. doi: [10.1080/1366825021000008657](https://doi.org/10.1080/1366825021000008657)
- Hastings, R. P. and Johnson, E.** (2001). Stress in UK families conducting intensive home-based behavioral intervention for their young child with autism. *Journal of Autism and Developmental Disorders*, *31*, 327–336. doi: [10.1023/A:1010799320795](https://doi.org/10.1023/A:1010799320795)
- Hastings, R. P., Kovshoff, H., Brown, T., Ward, N. J., Degli Espinosa, F. and Remington, B.** (2005). Coping strategies in mothers and fathers of preschool and school-age children with autism. *Autism*, *9*, 377–391. doi: [10.1177/1362361305056078](https://doi.org/10.1177/1362361305056078)
- Hayes, A. F.** (2012). PROCESS: a versatile computational tool for observed variable mediation, moderation, and conditional process modeling [White Paper]. Retrieved from: <http://www.afhayes.com/public/process2012.pdf>
- Hill, C. and Rose, J.** (2009). Parenting stress in mothers of adults with an intellectual disability: parental cognitions in relation to child characteristics and family support. *Journal of Intellectual Disability Research*, *53*, 969–980. doi: [10.1111/j.1365-2788.2009.01207.x](https://doi.org/10.1111/j.1365-2788.2009.01207.x)
- Horsley, S. and Oliver, C.** (2015). Positive impact and its relationship to well-being in parents of children with intellectual disability: a literature review. *International Journal of Developmental Disabilities*, *61*, 1–19. doi: [10.1179/2047387713Y.0000000026](https://doi.org/10.1179/2047387713Y.0000000026)
- Hyman, P., Oliver, C. and Hall, S.** (2002). Self-injurious behavior, self-restraint, and compulsive behaviors in Cornelia de Lange syndrome. *American Journal on Mental Retardation*, *107*, 146–154. doi: [10.1352/0895-8017\(2002\)107](https://doi.org/10.1352/0895-8017(2002)107)
- Jones, L., Hastings, R. P., Totsika, V., Keane, L. and Rhule, N.** (2014). Child behavior problems and parental well-being in families of children with autism: the mediating role of mindfulness and acceptance. *American Journal on Intellectual and Developmental Disabilities*, *119*, 171–185. doi: [10.1352/1944-7558-119.2.171](https://doi.org/10.1352/1944-7558-119.2.171)
- Karakus, M.** (2013). Emotional intelligence and negative feelings: a gender specific moderated mediation model. *Educational Studies*, *39*, 1–15. doi: [10.1080/03055698.2012.671514](https://doi.org/10.1080/03055698.2012.671514)

- Keefe, F. J., Blumenthal, J., Baucom, D., Affleck, G., Waugh, R., Caldwell, D. S. et al.** (2004). Effects of spouse-assisted coping skills training and exercise training in patients with osteoarthritic knee pain: a randomized controlled study. *Pain*, *110*, 539–549. doi: [10.1016/j.pain.2004.03.022](https://doi.org/10.1016/j.pain.2004.03.022)
- Kushlick, A., Blunden, R. and Cox, G.** (1973). Method of rating behavior characteristics for use in large-scale surveys of mental handicap. *Psychological Medicine*, *3*, 466–478.
- Lai, W. W. and Oei, T. P. S.** (2014). Coping in parents and caregivers of children with autism spectrum disorders (ASD): a review. *Review Journal of Autism and Developmental Disorders*, *1*, 207–224. doi: [10.1007/s40489-014-0021-x](https://doi.org/10.1007/s40489-014-0021-x)
- Lai, W. W., Goh, T. J., Oei, T. P. S. and Sung, M.** (2015). Coping and well-being in parents of children with autism spectrum disorders (ASD). *Journal of Autism and Developmental Disorders*, *45*, 2582–2593. doi: [10.1007/s10803-015-2430-9](https://doi.org/10.1007/s10803-015-2430-9)
- Lloyd, T. J. and Hastings, R.** (2009). Hope as a psychological resilience factor in mothers and fathers of children with intellectual disabilities. *Journal of Intellectual Disability Research*, *53*, 957–968. doi: [10.1111/j.1365-2788.2009.01206.x](https://doi.org/10.1111/j.1365-2788.2009.01206.x)
- MacKinnon, D. P., Lockwood, C. M. and Williams, J.** (2004). Confidence limits for the indirect effect: distribution of the product and resampling methods. *Multivariate Behavioral Research*, *39*, 99–128. doi: [10.1207/s15327906mbr3901\\_4](https://doi.org/10.1207/s15327906mbr3901_4)
- Mash, E. J. and Johnston, C.** (1990). Determinants of parenting stress – illustrations from families of hyperactive children and families of physically abused children. *Journal of Clinical Child Psychology*, *19*, 313–328. doi: [10.1207/s15374424jccp1904\\_3](https://doi.org/10.1207/s15374424jccp1904_3)
- McClintock, K., Hall, S. and Oliver, C.** (2003). Risk markers associated with challenging behaviours in people with intellectual disabilities: a meta-analytic study. *Journal of Intellectual Disability Research*, *47*, 405–416. doi: [10.1046/j.1365-2788.2003.00517.x](https://doi.org/10.1046/j.1365-2788.2003.00517.x)
- Moss, J., Oliver, C., Aaron, K., Burbidge, C. and Berg, K.** (2009). The prevalence and phenomenology of repetitive behavior in genetic syndromes. *Journal of Autism and Developmental Disorders*, *39*, 572–588.
- Oliver, C., Petty, J., Ruddick, L. and Bacarese-Hamilton, M.** (2012). The association between repetitive, self-injurious and aggressive behavior in children with severe intellectual disability. *Journal of Autism and Developmental Disorders*, *42*, 910–919. doi: [10.1007/s10803-011-1320-z](https://doi.org/10.1007/s10803-011-1320-z)
- Petty, J. and Oliver, C.** (2005). Self-injurious behaviour in individuals with intellectual disabilities. *Current Opinion in Psychiatry*, *18*, 484–489. doi: [10.1097/01.yco.0000179484.62391.dc](https://doi.org/10.1097/01.yco.0000179484.62391.dc)
- Preacher, K. J. and Hayes, A. F.** (2004). SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods Instruments and Computers*, *36*, 717–731. doi: [10.3758/Bf03206553](https://doi.org/10.3758/Bf03206553)
- Preacher, K. J. and Hayes, A. F.** (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods*, *40*, 879–891. doi: [10.3758/Brm.40.3.879](https://doi.org/10.3758/Brm.40.3.879)
- Rhee, S. H., Parker, J. C., Smarr, K. L., Petroski, G. F., Johnson, J. C., Hewett, J. E. et al.** (2000). Stress management in rheumatoid arthritis: what is the underlying mechanism? *Arthritis Care and Research*, *13*, 435–442.
- Robbins, F. R., Dunlap, G. and Plien, A. J.** (1991). Family characteristics, family training, and the progress of young children with autism. *Journal of Early Intervention*, *15*, 173–184.
- Rojahn, J., Matson, J. L., Lott, D., Esbensen, A. J. and Smalls, Y.** (2001). The Behavior Problems Inventory: an instrument for the assessment of self-injury, stereotyped behavior, and aggression/destruction in individuals with developmental disabilities. *Journal of Autism and Developmental Disorders*, *31*, 577–588. doi: [10.1023/A:1013299028321](https://doi.org/10.1023/A:1013299028321)
- Rose, J., Nelson, L. and Hardiman, R.** (2016). The relationship between challenging behaviour, cognitions and stress in mothers of individuals with intellectual disabilities. *Behavioural and Cognitive Psychotherapy*, *44*, 691–704. doi: [10.1017/S1352465816000242](https://doi.org/10.1017/S1352465816000242)

- Seltzer, M. M., Greenberg, J. S. and Krauss, M. W.** (1995). A comparison of coping strategies of aging mothers of adults with mental illness or mental retardation. *Psychology and Aging*, *10*, 64–75. doi: [10.1037//0882-7974.10.1.64](https://doi.org/10.1037//0882-7974.10.1.64)
- Silver, R. C., Holman, E. A., McIntosh, D. N., Poulin, M. and Gil-Rivas, V.** (2002). Nationwide longitudinal study of psychological responses to September 11. *Journal of the American Medical Association*, *288*, 1235–1244. doi: [10.1001/jama.288.10.1235](https://doi.org/10.1001/jama.288.10.1235)
- Singer, G. H. S.** (2006). Meta-analysis of comparative studies of depression in mothers of children with and without developmental disabilities. *American Journal on Mental Retardation*, *111*, 155–169. doi: [10.1352/0895-8017\(2006\)111](https://doi.org/10.1352/0895-8017(2006)111)
- Sivberg, B.** (2002). Family system and coping behaviors – a comparison between parents of children with autistic spectrum disorders and parents with non-autistic children. *Autism*, *6*, 397–409. doi: [10.1177/1362361302006004006](https://doi.org/10.1177/1362361302006004006)
- Smith, L. E., Seltzer, M. M., Tager-Flusberg, H., Greenberg, J. S. and Carter, A. S.** (2008). A comparative analysis of well-being and coping among mothers of toddlers and mothers of adolescents with ASD. *Journal of Autism and Developmental Disorders*, *38*, 876–889. doi: [10.1007/s10803-007-0461-6](https://doi.org/10.1007/s10803-007-0461-6)
- Stanton, A. L. and Snider, P. R.** (1993). Coping with a breast cancer diagnosis – a prospective study. *Health Psychology*, *12*, 16–23. doi: [10.1037//0278-6133.12.1.16](https://doi.org/10.1037//0278-6133.12.1.16)
- Taylor, S. E. and Stanton, A. L.** (2007). Coping resources, coping processes, and mental health. *Annual Review of Clinical Psychology*, *3*, 377–401. doi: [10.1146/annurev.clinpsy.3.022806.091520](https://doi.org/10.1146/annurev.clinpsy.3.022806.091520)
- Tomanik, S., Harris, G. E. and Hawkins, J.** (2004). The relationship between behaviours exhibited by children with autism and maternal stress. *Journal of Intellectual and Developmental Disability*, *29*, 16–26. doi: [10.1080/13668250410001662892](https://doi.org/10.1080/13668250410001662892)
- Watson, D., Clark, L. A. and Tellegen, A.** (1988). Development and validation of brief measures of positive and negative affect – the PANAS Scales. *Journal of Personality and Social Psychology*, *54*, 1063–1070. doi: [10.1037/0022-3514.54.6.1063](https://doi.org/10.1037/0022-3514.54.6.1063)
- Wiedemann, A.U., Schuz, B., Sniehotta, F., Scholz, U. and Schwarzer, R.** (2009). Disentangling the relation between intentions, planning and behaviour: a moderated mediation analysis. *Psychology and Health*, *24*, 67–79.
- Woodman, A. C. and Hauser-Cram, P.** (2013). The role of coping strategies in predicting change in parenting efficacy and depressive symptoms among mothers of adolescents with developmental disabilities. *Journal of Intellectual Disability Research*, *57*, 513–530. doi: [10.1111/j.1365-2788.2012.01555.x](https://doi.org/10.1111/j.1365-2788.2012.01555.x)
- Zigmond, A. S. and Snaith, R. P.** (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, *67*, 361–370. doi: [10.1111/j.1600-0447.1983.tb09716.x](https://doi.org/10.1111/j.1600-0447.1983.tb09716.x)