

ERCC1 as predictive biomarker to platinum-based chemotherapy in adrenocortical carcinomas

Laufs, Valeria; Altieri, Barbara; Sbiera, Silviu; Kircher, Stefan; Steinhauer, Sonja; Beuschlein, Felix; Quinkler, Marcus; Willenberg, Holger S; Rosenwald, Andreas ; Fassnacht, Martin; Ronchi, Cristina

DOI:

[10.1530/EJE-17-0788](https://doi.org/10.1530/EJE-17-0788)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Laufs, V, Altieri, B, Sbiera, S, Kircher, S, Steinhauer, S, Beuschlein, F, Quinkler, M, Willenberg, HS, Rosenwald, A, Fassnacht, M & Ronchi, C 2018, 'ERCC1 as predictive biomarker to platinum-based chemotherapy in adrenocortical carcinomas', *European Journal of Endocrinology*, vol. 178, pp. 181-188.
<https://doi.org/10.1530/EJE-17-0788>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 19/04/2018

"Disclaimer: this is not the definitive version of record of this article. This manuscript has been accepted for publication in European Journal of Endocrinology, but the version presented here has not yet been copy-edited, formatted or proofed. Consequently, Bioscientifica accepts no responsibility for any errors or omissions it may contain. The definitive version is now freely available at 10.1530/EJE-17-0788 2017

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **ERCC1 as predictive biomarker to platinum-based chemotherapy in adrenocortical carcinomas.**

2 Valeria Laufs¹, Barbara Altieri^{1,2}, Silviu Sbiera¹, Stefan Kircher^{3,4}, Sonja Steinhauer¹, Felix
3 Beuschlein^{5,6}, Marcus Quinkler⁷, Holger S. Willenberg⁸, Andreas Rosenwald^{3,4}, Martin Fassnacht^{1,4},
4 Cristina L. Ronchi¹.

5

6 ¹Division of Endocrinology and Diabetes, Department of Internal Medicine I, University Hospital,
7 University of Wuerzburg, Wuerzburg, Germany.

8 ²Division of Endocrinology and Metabolic Diseases, Institute of Medical Pathology, Catholic
9 University of the Sacred Heart, Rome, Italy.

10 ³Institute of Pathology, University of Wuerzburg, Wuerzburg, Germany.

11 ⁴Comprehensive Cancer Center Mainfranken, University of Wuerzburg, Germany.

12 ⁵Medizinische Klinik and Poliklinik IV, Ludwig-Maximilians University, Munich, Germany.

13 ⁶Klinik für Endokrinologie, Diabetologie und Klinische Ernährung, Universitätsspital Zürich, Zürich,
14 Switzerland.

15 ⁷Endocrinology in Charlottenburg, Berlin, Germany.

16 ⁸Division of Endocrinology and Metabolism, Rostock University Medical Center, Germany.

17

18 **Short title:** ERCC1 in adrenocortical carcinoma

19 **Key words:** ERCC1, biomarker, adrenal cortex, carcinoma, chemotherapy.

20 **Word count:** 2723

21

22 **Corresponding author:**

23 Cristina L. Ronchi (MD, PhD)

24 Division of Endocrinology and Diabetes, Department of Internal Medicine I

25 Oberduerrbacher-Str 6

26 University Hospital of Wuerzburg

27 97080 Wuerzburg (Germany)

28 Ronchi_C@ukw.de

29 **Abstract**

30 **Objective:** Platinum-based chemotherapy (PBC) is the most effective cytotoxic treatment for
31 advanced adrenocortical carcinoma (ACC). Excision repair cross complementing group 1 (ERCC1)
32 plays a critical role in the repair of platinum-induced DNA damage. Two studies investigating the role
33 of ERCC1 immunostaining as a predictive marker for the response to PBC in ACC had reported
34 conflicting results. Both studies used the ERCC1-antibody clone 8F1 that later turned out to be not
35 specific. The aim of this study was to evaluate the predictive role of ERCC1 with the new specific
36 antibody in a larger series of ACC.

37 **Design and Methods:** 146 ACC patients with available FFPE slides were investigated. All patients
38 underwent PBC (median cycles=6), including cisplatin (n=131) or carboplatin (n=15), in most cases
39 combined with etoposide (n=144), doxorubicin (n=131) and mitotane (n=131). Immunostaining was
40 performed with the novel ERCC1-antibody clone 4F9. The relationship between ERCC1 expression
41 and clinico-pathological parameters, as well as best objective response to therapy and progression-free
42 survival (PFS) during PBC was evaluated.

43 **Results:** High ERCC1 expression was observed in 66% of ACC samples. During PBC, 43 patients
44 experienced objective response (29.5%), 49 stable disease (33.6%), 8 mixed response (5.5%) and 46
45 progressive disease (31,5%) without any relationship with the ERCC1 immunostaining. No significant
46 correlation was also found between ERCC1 expression and progression-free survival (median 6.5 vs 6
47 months, $P=0.33$, HR=1.23, 95%CI=0.82-2.0).

48 **Conclusion:** ERCC1 expression is not directly associated with sensitivity to PBC in ACC. Thus, other
49 predictive biomarkers are required to support treatment decisions in patients with ACC.

50
51
52
53
54
55
56

57 **Introduction**

58 Platinum-based chemotherapy (PBC) is the most effective cytotoxic treatment for advanced
59 adrenocortical carcinoma (ACC), mostly in combination with etoposide and doxorubicin plus mitotane
60 in the EDP-M regime¹. However, the best objective response rates remain below 30% and the impact
61 on overall survival is not satisfying as shown in the phase III clinical trial FIRM-ACT². Similarly,
62 other possible cytotoxic drugs such as streptozotocin² or gemcitabine did not show a better
63 effectiveness³ and no effective targeted therapies have emerged for ACC patients with advanced
64 disease⁴⁻⁶. Finally, PBC as other chemotherapeutic combinations is associated with relevant toxicity.
65 Thus, it is obvious that there is an urgent need of biomarkers that may serve to predict the response to
66 PBC.

67 Excision repair cross complementing group 1 (ERCC1) is an important member of the nucleoside
68 excision repair pathway, which plays a critical role in the DNA repair by removing DNA covalent
69 helix-distorting adducts caused by platinum compounds⁷. ERCC1 has been demonstrated to be a
70 predictive biomarker for platinum treatment in several cancers, such as non-small cell lung cancer,
71 testicular germ cell tumor, bladder cancer, pancreatic carcinoma and gastric cancer⁸⁻¹². Two previous
72 studies, one from our group¹³ and one from France¹⁴ investigated ERCC1 immunostaining in
73 relationship with the response to PBC in a relatively small series of ACC patients (n=45 and n=33,
74 respectively). These two studies described a similar overall response rate to PBC (25-30% of cases),
75 but reported conflicting results regarding the influence of ERCC1 on sensitivity to PBC, being
76 significant only in the first study. All the previous studies on ERCC1 immunostaining, including those
77 on ACC, have been performed by using the monoclonal anti-mouse antibody clone 8F1. However,
78 already some years ago, it had been suggested that this clone might be not specific, being ERCC1 not
79 the principal antigen recognized by the 8F1 antibody^{15, 16}. In fact, more recently, it has been
80 demonstrated that the clone 8F1 immunoglobulin recognizes also the choline phosphate
81 cytidyltransferase 1 alfa (PCYT1A), an unrelated nuclear membrane protein, involved in the
82 metabolism of phosphatidylcholine biosynthesis¹⁷. These findings raise doubts on previously
83 published data using the clone 8F1 to investigate ERCC1 as a predictive marker to PBC in several
84 solid tumors. Finally, a new highly specific clone 4F9 has been identified and then validated¹⁷⁻¹⁹.

85 Thus, the aim of the present study was to evaluate ERCC1 immunostaining with the new highly
86 specific clone 4F9^{17, 18} in a larger series of ACC and to correlate it with the response to PBC.

87

88 **Subjects and methods**

89 *Patients and treatment regimen*

90 Inclusion criteria were age of at least 18 years, histopathologic diagnosis of ACC, available formalin-
91 fixed paraffin embedded (FFPE) specimens and treatment with PBC. We identified a total of 153
92 patients that fulfilled these criteria and were treated with PBC in our centers between 2004 and 2015.
93 Seven of these patients received only one cycle of PBC and were then excluded from further analysis.
94 Thus, the final series included 146 patients with advanced ACC (F:M=90:56, median age 48 years).
95 None of these patients were already included in our previous paper on ERCC1¹³, while 49 participated
96 in the FIRM-ACT study². Specifically, 127 samples derived from primary surgery, 6 from local
97 recurrences, 4 from biopsies (patients not operable) and 9 from distant metastasis. The baseline
98 clinical parameters, such as sex, age at initial diagnosis, tumor size, biochemical evaluation, tumor
99 stage according to the European Network for the Study of Adrenal Tumors (ENSAT) classification²⁰,
100 Weiss score, Ki67 proliferation index, presence and number of distant metastases, and previous local
101 and/or pharmacological treatments are given in **Table 1**. All baseline data were collected through the
102 ENSAT Registry (www.ens@t.org/registry).

103 The treatment regimen included cisplatin (n=131) or carboplatin (n=15) and was in most cases
104 administered as combination therapy (see details **Table 1**). The median number of PBC cycles was 6
105 ranging from 2 to 15. Treatment was discontinued in cases of unacceptable toxicity, patient's refusal
106 or evidence of disease progression. A total of 131 patients (90% of cases) were treated with
107 concomitant mitotane (target plasma concentration: 14-20 mg/L). 114 patients received PBC as first-
108 line cytotoxic treatment (78% of cases), while the remaining 32 patients were treated with PBC as
109 second- or third- line therapy, with a history of failed streptozotocin² or gemcitabine + capecitabine³
110 (**Table 1**). All patients had undergone regular and standard follow-up visits with clinical, biochemical,
111 and radiological (abdominal and thoracic CT scan with contrast agent) evaluation with a staging
112 interval usually every 8 weeks. The sensitivity to PBC was evaluated as progression-free survival

113 during treatment and as best overall objective response. For this evaluation, according to our clinical
114 practice, all radiological images were reviewed by the local expert radiologists and discussed in our
115 multidisciplinary tumor board meetings to determine a final consensus response (progressive disease,
116 stable disease, partial or complete response). Clinical benefit was defined as stable disease or
117 treatment response for a minimum of 4 months.

118 The collection of the clinical data and the biomaterial for this retrospective study was approved by the
119 ethics committee of the University of Wuerzburg (No. 93/02 and 88/11) according to the Declaration
120 of Helsinki. Written informed consent was obtained from all patients.

121

122 ***Immunohistochemistry***

123 A total of 146 FFPE adrenocortical tissues on standard full slides were evaluated by
124 immunohistochemistry. In brief, sections were deparaffinized and immunohistochemical detection was
125 performed using an indirect immunoperoxidase technique after high temperature antigen retrieval in
126 10 mM citric acid monohydrate buffer (pH 6.5) in a pressure cooker for 13 min. Blocking of
127 unspecific protein-antibody interactions was performed with 20% human AB serum in PBS for 1h at
128 room temperature. Primary antibody for ERCC1 was the new highly specific monoclonal anti-mouse
129 antibody (mAb) clone 4F9 (UM500008, dilution 1:100) that was purchased from OriGene
130 Technologies, Inc (Rockville, USA). A mouse negative control was used (Dako North America Inc.,
131 Carpinteria, USA). The slices were incubated overnight at 4°C. Signal amplification was achieved
132 with En-Vision System Labeled Polymer-HRP Anti-Mouse (Dako) for 40 min and developed for 10
133 min with DAB Substrate Kit (Vector Laboratories, Burlingame, CA) according to the manufacturer's
134 instructions. Nuclei were counterstained with Mayer's hematoxin for 2 min. For positive controls,
135 sections of colon adenocarcinoma, renal cell carcinoma, breast cancer, hepatocellular carcinoma and
136 normal tonsil were chosen, while cells of the tumor stroma served as internal negative control.

137 All slides were analyzed independently by two investigators blinded to clinical information (V.L. and
138 S.S.) Nuclear staining intensity was graded as negative (0), low (1), medium (2), or strong (3). The
139 percentage of tumor cells with positive nuclei was calculated for each specimen and scored 0 if 0%
140 were positive, 0.1 if 1–9% were positive, 0.5 if 10–49% were positive and 1 if 50% or more were

141 positive. A semiquantitative H-score was then calculated by multiplying the staining intensity grading
142 score with the proportion score as described previously¹³. In case of discrepant results, staining
143 intensities were jointly assessed by both investigators, forming the final score by consensus. Inter-
144 observer agreement was investigated via Pearson's correlation coefficients 0.72 (95%CI: 0.63-0.79).

145

146 ***Comparison between anti-ERCC1 antibody clone 8F1 vs clone 4F9***

147 We also intended to re-evaluate our old results obtained with the mAb against ERCC1 clone 8F1 (old
148 batch)¹³ with the new high specific mAb clone 4F9. To this aim, we re-stained 38 ACC samples out of
149 the 45 previously published and re-investigated the relationship between ERCC1 expression and the
150 response to PBC in terms of both progression-free survival (PFS) and disease-specific survival (DSS)
151 after treatment. Moreover, the specificity of the currently available clone 8F1 has been shown to be
152 altered from the old clone 8F1^{21, 22}. In addition, we also evaluated a subgroup of 21 out of the 146
153 samples in our present series with the current clone 8F1 (new batch) in addition to the new clone 4F9.

154

155 ***Statistical analysis***

156 The Fisher's exact or the Chi-square tests were used to investigate dichotomic variables, while
157 continuous variables were investigated with a two-sided *t* test (or non-parametric test). A non-
158 parametric Kruskal-Wallis test, followed by Dunn's test, was used for comparison among several
159 groups for non-normal distributed variables. Correlations and 95% confidence intervals (95%CI)
160 between different parameters were evaluated by linear regression analysis. PFS was defined as the
161 time from the date of first administration of PBC to the first radiological evidence of disease
162 progression or death, as appropriate. DSS was defined as the time from the first administration of PBC
163 to disease-specific death or last follow-up. All survival curves were obtained with Kaplan-Meier
164 estimates, and the differences between survival curves were assessed by the log-rank (Mantel-Cox)
165 test. For the calculation of hazard ratios (HR), two ACC-groups with low or high protein expression
166 were considered (high expression: H-score ≥ 2). A multivariate regression analysis was performed via
167 a Cox proportional hazard regression model, aiming to identify factors that might independently
168 influence survival. Statistical analyses were made using GraphPad Prism (version 6.0, La Jolla, CA,

169 USA) and SPSS Software (PASW Version 21.0, SPSS Inc., Chicago, IL, USA). P values <0.05 were
170 considered as statistically significant.

171

172 **Results**

173 *Efficacy of platinum-based chemotherapy*

174 The data about efficacy of PBC in the current series of 146 patients with advanced ACC are
175 summarized in **Table 2**. Concerning the best objective response during PBC, one patient experienced
176 complete response (0.7%) and 42 patients partial remission (28.8%), 49 stable disease (33.5%), 8
177 mixed response (5.5%) and 46 progressive disease (31.5%), respectively. The median PFS during PBC
178 was 6 months, ranging from 2 to 18, while the median DSS was 17 months, ranging from 1.5 to 127.
179 Additionally, we observed a clinical benefit defined as at least a stable disease for a minimum of 4
180 months in 84 patients (58%) with a median PFS in this group of 6 months (range: 4-18). Only one
181 patient died unrelated to ACC during follow up. Thus, overall survival was more or less identical to
182 DSS (data not shown).

183

184 *ERCC1 expression and baseline clinical characteristics in ACC*

185 Nuclear ERCC1 immunostaining was homogeneous in individual ACC samples with a median
186 percentage of positive cells of 80% ($> 50\%$ in 135/146 samples, 92.5%). Tissue samples exemplifying
187 the range of staining intensity are shown in the **Figure 1**. ERCC1 expression was low (H-score 0-1) in
188 50 samples (34.2% of cases) and high (H-score 2-3) in 96 samples (65.7%). We did not observe any
189 significant differences in ERCC1 immunostaining among primary tumors, local recurrences and/or
190 distant metastasis. No significant correlation was also observed between the nuclear ERCC1
191 expression and the ENSAT tumor stage at the time of diagnosis, the Weiss score or the Ki67
192 proliferation index.

193

194 *Predictive role of ERCC1 expression on sensitivity to platinum-based chemotherapy*

195 Considering the potential predictive role of ERCC1 immunostaining on the objective response to PBC,
196 no significant differences were observed between the groups with high and low nuclear ERCC1

197 expression (*Table 2*). Similarly, no differences were found in terms of both PFS (median 6.5 vs 6
198 months, respectively, $P=0.33$, HR=1.23, 95%CI=0.82-2.0) and DSS (median 17 vs 16.5 months,
199 respectively, $P=0.87$, HR=1.03, 95%CI=0.70-1.53) (*Figure 2A-B*).

200

201 *Comparison between anti-ERCC1 antibody clone 8F1 vs clone 4F9*

202 We re-stained 38 out of 45 ACC samples of our previously published series (stained with the 8F1
203 clone old batch) with the new clone 4F9. Not unexpected, ERCC1 expression in terms of H-score
204 corresponded in only 49% of cases. As a consequence, ERCC1 nuclear expression did not longer
205 significantly correlate with response to PBC in terms of both PFS (data not shown) and DSS
206 (Supplementary Figure 1A and B).

207 Furthermore, we stained 21 out of the present 146 samples with the currently available clone 8F1 (new
208 batch) additionally to the clone 4F9. Two representative examples are shown in the Supplementary
209 Figure 2. Comparing the ERCC1 immunostaining results we observed here a correspondence between
210 the two antibodies in 81% of cases.

211

212 **Discussion**

213 We evaluated the potential role of ERCC1 nuclear expression as predictive biomarker to PBC in the
214 largest series of ACC patients up to date (n=146) by using for the first time the new high ERCC1-
215 specific monoclonal antibody clone 4F9. To note, ERCC1 has been previously demonstrated to be a
216 predictive biomarker for platinum treatment in several cancers, such as non-small cell lung cancer
217 (NSCLC), testicular germ cell tumors, bladder cancer, pancreatic carcinoma and gastric cancer⁸⁻¹². In
218 ACC, we previously demonstrated in a relatively small series of patients that ERCC1 immunostaining
219 was significantly correlated with overall survival during PBC¹³. Another study, however, did not
220 confirm this finding¹⁴. Nevertheless, several concerns about the reliability of the ERCC1
221 immunohistochemical analysis have been raised recently. First, it has been demonstrated that the clone
222 8F1 used in all the reported studies is not specific for ERCC1¹⁵⁻¹⁷. Specifically, the anti-ERCC1
223 antibody clone 8F1 has been identified to stain also the PCYT1A, a phospholipid synthesis enzyme
224 regulated by RAS^{17, 23} with no known clinical implication in platinum drug resistance. PCYT1A has

225 also been confirmed to play a role as prognostic biomarker in both lung and head and neck squamous
226 cell carcinomas²³.

227 Moreover, the batch of the clone 8F1 in use since 2011 seems not to be identical with the batch in use
228 in 2006¹⁹, thus rendering new data about NSCLC not comparable with previous ones²². According to
229 this new information, important previous results on the role of ERCC1 in the treatment of NSCLC⁸
230 have been revised by the same group²¹. Furthermore, this year the first randomized trial to evaluate
231 ERCC1 prospectively in 648 patients with NSCLC (ET trial) has been published definitively
232 demonstrating that selecting chemotherapy using the commercially available ERCC1 antibodies (clone
233 8F1) does not confer any additional survival benefit²⁴.

234 In parallel, a new highly ERCC1-specific clone 4F9 has been recently proposed and validated¹⁷⁻¹⁹. For
235 all these reasons, we decided to use the clone 4F9 to investigate a new large series of ACC samples in
236 order to re-evaluate our previous results on ERCC1 as predictive marker of sensitivity to PBC. Most
237 importantly, we could not confirm the previous results and our data now indicate that ERCC1 itself is
238 probably not the main factor involved in the response to PBC in ACC patients. In addition, we were
239 able to demonstrate that the current version of the clone 8F1 significantly differs from the old one that
240 we used for our pilot study¹³ and we were not able to reproduce the earlier results using now the same
241 tumor samples.

242 One reason that could explain the lack of correlation between ERCC1 and PBC, independently from
243 the issues with immunohistochemistry, is that ERCC1 works together with the XPF protein, codified
244 by *ERCC4*. ERCC1–XPF complex is a two subunit structure-specific endonuclease that plays a key
245 role during the nucleotide excision repair (NER) process^{7, 25}. Thus, XPF itself might be involved in the
246 sensitivity to the response to PBC^{26, 27}. However, the ET trial demonstrated that XPF expression is not
247 predictive for response to 648 patients with NSCLC²⁴. Moreover, the ERCC1–XPF complex makes
248 incisions on the damaged DNA strand on the 5' side and acts in cooperation with several other
249 proteins, like XPC–RAD23B, XPA, RPA, TFIIH and XPG, during the NER process^{28, 29}. Thus,
250 although ERCC1 plays a major role in the NER, several other proteins and mechanisms could
251 influence the response to PBC.

252 Another explanation, why ERCC1 expression and clinical outcome in our and other series did not
253 correlate could be the fact that virtually all patients have received in parallel to the platinum derivate
254 1-3 other additional cytotoxic drugs (mostly doxorubicin, etoposide and mitotane) diluting the
255 hypothesized correlation. Other potential biomarker could for instance be involved in the prediction of
256 response to these concomitant treatments (i.e. TOP2A³⁰). Finally, one potential limitation in our study
257 as well as in several others might be that ERCC1 was assessed on tumor specimens obtained months
258 or even years before the start of chemotherapy. Nevertheless, we did not observe any significant
259 differences in ERCC1 immunostaining among primary tumors, local recurrences and/or distant
260 metastasis, thus suggesting that the ERCC1 levels remain quite stable over the time and tumor
261 progression.

262 More generally, the search for predictive biomarkers to conventional cytotoxic chemotherapy has been
263 proven challenging due to frequent discrepant and non-replicable findings. And this is true not only for
264 protein expression where issues with antibodies and immunohistochemical analysis are common, but
265 also for gene expression. Thus, if a plethora of biomarkers predicting chemotherapy efficacy have
266 been evaluated also in the clinical setting, none of them is ready for clinical implementation yet³¹.
267 Considering that most mechanisms of resistance or sensitivity to chemotherapy are multifactorial, a
268 combinatorial approach and further efforts are required³².

269 Concerning the response rate to PBC in general, we observed an objective partial response in 29.5% of
270 cases and a stable disease in further 33.5%, thus confirming that PBC is the currently most effective
271 cytotoxic therapy for advanced ACC. These data are generally superimposable to those reported in the
272 FIRM-ACT study on EDP-M².

273 In conclusion, ERCC1 expression as detected by immunostaining is not directly associated with
274 sensitivity to PBC in ACC. Thus, the search for predictive biomarkers in this devastating disease with
275 poor response to medical therapy has to continue.

276

277 **Declaration of interest**

278 All authors declare that there is no conflict of interest that could be perceived as prejudicing the
279 impartiality of the research reported.

280 **Funding**

281 This work has been supported by the Deutsche Forschungsgemeinschaft (DFG) within the
282 CRC/Transregio 205/1 “The Adrenal: Central Relay in Health and Disease” to M.F. and by an
283 individual grant (FA466/4-1 and 4-2 to M.F.).

284

285 **Acknowledgements**

286 The authors are grateful to Martina Zink for excellent technical support and to Michaela Haaf for
287 coordinating the ENSAT Registry in Wuerzburg.

288

289 **References**

- 290 1. Berruti A, Terzolo M, Sperone P, Pia A, Della Casa S, Gross DJ, Carnaghi C, Casali
291 P, Porpiglia F, Mantero F, Reimondo G, Angeli A & Dogliotti L. Etoposide,
292 doxorubicin and cisplatin plus mitotane in the treatment of advanced adrenocortical
293 carcinoma: a large prospective phase II trial. *Endocr Relat Cancer* 2005 **12** 657-666.
- 294 2. Fassnacht M, Terzolo M, Allolio B, Baudin E, Haak H, Berruti A, Welin S, Schade-
295 Brittinger C, Lacroix A, Jarzab B, Sorbye H, Torpy DJ, Stepan V, Schteingart DE,
296 Arlt W, Kroiss M, Leboulleux S, Sperone P, Sundin A, Hermsen I, Hahner S,
297 Willenberg HS, Tabarin A, Quinkler M, de la Fouchardiere C, Schlumberger M,
298 Mantero F, Weismann D, Beuschlein F, Gelderblom H, Wilmink H, Sender M,
299 Edgerly M, Kenn W, Fojo T, Muller HH, Skogseid B & Group F-AS. Combination
300 chemotherapy in advanced adrenocortical carcinoma. *N Engl J Med* 2012 **366** 2189-
301 2197.
- 302 3. Sperone P, Ferrero A, Daffara F, Priola A, Zaggia B, Volante M, Santini D, Vincenzi
303 B, Badalamenti G, Intrivici C, Del Buono S, De Francia S, Kalomirakis E, Ratti R,
304 Angeli A, Dogliotti L, Papotti M, Terzolo M & Berruti A. Gemcitabine plus
305 metronomic 5-fluorouracil or capecitabine as a second-/third-line chemotherapy in
306 advanced adrenocortical carcinoma: a multicenter phase II study. *Endocr Relat Cancer*
307 2010 **17** 445-453.
- 308 4. Ronchi CL, Kroiss M, Sbiera S, Deutschbein T & Fassnacht M. EJE prize 2014:
309 current and evolving treatment options in adrenocortical carcinoma: where do we
310 stand and where do we want to go? *Eur J Endocrinol* 2014 **171** R1-R11.
- 311 5. Else T, Kim AC, Sabolch A, Raymond VM, Kandathil A, Caoili EM, Jolly S, Miller
312 BS, Giordano TJ & Hammer GD. Adrenocortical carcinoma. *Endocr Rev* 2014 **35**
313 282-326.
- 314 6. Fassnacht M, Kroiss M & Allolio B. Update in adrenocortical carcinoma. *J Clin*
315 *Endocrinol Metab* 2013 **98** 4551-4564.
- 316 7. Manandhar M, Boulware KS & Wood RD. The ERCC1 and ERCC4 (XPF) genes and
317 gene products. *Gene* 2015 **569** 153-161.
- 318 8. Olaussen KA, Dunant A, Fouret P, Brambilla E, Andre F, Haddad V, Taranchon E,
319 Filipits M, Pirker R, Popper HH, Stahel R, Sabatier L, Pignon JP, Tursz T, Le
320 Chevalier T, Soria JC & Investigators IB. DNA repair by ERCC1 in non-small-cell

- 321 lung cancer and cisplatin-based adjuvant chemotherapy. *N Engl J Med* 2006 **355** 983-
322 991.
- 323 9. Mendoza J, Martinez J, Hernandez C, Perez-Montiel D, Castro C, Fabian-Morales E,
324 Santibanez M, Gonzalez-Barrios R, Diaz-Chavez J, Andonegui MA, Reynoso N,
325 Onate LF, Jimenez MA, Nunez M, Dyer R & Herrera LA. Association between
326 ERCC1 and XPA expression and polymorphisms and the response to cisplatin in
327 testicular germ cell tumours. *Br J Cancer* 2013 **109** 68-75.
- 328 10. Bellmunt J, Paz-Ares L, Cuello M, Cecere FL, Albiol S, Guillem V, Gallardo E,
329 Carles J, Mendez P, de la Cruz JJ, Taron M, Rosell R, Baselga J & Spanish Oncology
330 Genitourinary G. Gene expression of ERCC1 as a novel prognostic marker in
331 advanced bladder cancer patients receiving cisplatin-based chemotherapy. *Ann Oncol*
332 2007 **18** 522-528.
- 333 11. Strippoli A, Rossi S, Martini M, Basso M, D'Argento E, Schinzari G, Barile R,
334 Cassano A & Barone C. ERCC1 expression affects outcome in metastatic pancreatic
335 carcinoma treated with FOLFIRINOX: A single institution analysis. *Oncotarget* 2016.
- 336 12. De Dosso S, Zanellato E, Nucifora M, Boldorini R, Sonzogni A, Biffi R, Fazio N,
337 Bucci E, Beretta O, Crippa S, Saletti P & Frattini M. ERCC1 predicts outcome in
338 patients with gastric cancer treated with adjuvant cisplatin-based chemotherapy.
339 *Cancer Chemother Pharmacol* 2013 **72** 159-165.
- 340 13. Ronchi CL, Sbiera S, Kraus L, Wortmann S, Johanssen S, Adam P, Willenberg HS,
341 Hahner S, Allolio B & Fassnacht M. Expression of excision repair cross
342 complementing group 1 and prognosis in adrenocortical carcinoma patients treated
343 with platinum-based chemotherapy. *Endocr Relat Cancer* 2009 **16** 907-918.
- 344 14. Malandrino P, Al Ghuzlan A, Castaing M, Young J, Caillou B, Travagli JP, Elias D,
345 de Baere T, Dromain C, Paci A, Chanson P, Schlumberger M, Leboulleux S & Baudin
346 E. Prognostic markers of survival after combined mitotane- and platinum-based
347 chemotherapy in metastatic adrenocortical carcinoma. *Endocr Relat Cancer* 2010 **17**
348 797-807.
- 349 15. Niedernhofer LJ, Bhagwat N & Wood RD. ERCC1 and non-small-cell lung cancer. *N*
350 *Engl J Med* 2007 **356** 2538-2540; author reply 2540-2531.
- 351 16. Bhagwat NR, Roginskaya VY, Acquafondata MB, Dhir R, Wood RD & Niedernhofer
352 LJ. Immunodetection of DNA repair endonuclease ERCC1-XPF in human tissue.
353 *Cancer Res* 2009 **69** 6831-6838.
- 354 17. Ma D, Baruch D, Shu Y, Yuan K, Sun Z, Ma K, Hoang T, Fu W, Min L, Lan ZS,
355 Wang F, Mull L & He WW. Using protein microarray technology to screen anti-
356 ERCC1 monoclonal antibodies for specificity and applications in pathology. *BMC*
357 *Biotechnol* 2012 **12** 88.
- 358 18. Bahamon BN, Gao F & Danaee H. Development and Validation of an ERCC1
359 Immunohistochemistry Assay for Solid Tumors. *Arch Pathol Lab Med* 2016 **140**
360 1397-1403.
- 361 19. Smith DH, Fiehn AM, Fogh L, Christensen IJ, Hansen TP, Stenvang J, Nielsen HJ,
362 Nielsen KV, Hasselby JP, Brunner N & Jensen SS. Measuring ERCC1 protein
363 expression in cancer specimens: validation of a novel antibody. *Sci Rep* 2014 **4** 4313.
- 364 20. Fassnacht M, Johanssen S, Quinkler M, Bucszy P, Willenberg HS, Beuschlein F,
365 Terzolo M, Mueller HH, Hahner S, Allolio B, German Adrenocortical Carcinoma
366 Registry G & European Network for the Study of Adrenal T. Limited prognostic value
367 of the 2004 International Union Against Cancer staging classification for
368 adrenocortical carcinoma: proposal for a Revised TNM Classification. *Cancer* 2009
369 **115** 243-250.
- 370 21. Friboulet L, Olausson KA, Pignon JP, Shepherd FA, Tsao MS, Graziano S, Kratzke R,
371 Douillard JY, Seymour L, Pirker R, Filipits M, Andre F, Solary E, Ponsonnailles F,

- 372 Robin A, Stoclin A, Dorvault N, Commo F, Adam J, Vanhecke E, Saulnier P,
373 Thomale J, Le Chevalier T, Dunant A, Rousseau V, Le Teuff G, Brambilla E & Soria
374 JC. ERCC1 isoform expression and DNA repair in non-small-cell lung cancer. *N Engl*
375 *J Med* 2013 **368** 1101-1110.
- 376 22. Wislez M, Barlesi F, Besse B, Mazieres J, Merle P, Cadranet J, Audigier-Valette C,
377 Moro-Sibilot D, Gautier-Felizot L, Goupil F, Renault A, Quoix E, Souquet PJ,
378 Madroszyck A, Corre R, Perol D, Morin F, Zalcman G & Soria JC. Customized
379 adjuvant phase II trial in patients with non-small-cell lung cancer: IFCT-0801 TASTE.
380 *J Clin Oncol* 2014 **32** 1256-1261.
- 381 23. Vaezi AE, Bepler G, Bhagwat NR, Malysa A, Rubatt JM, Chen W, Hood BL, Conrads
382 TP, Wang L, Kemp CE & Niedernhofer LJ. Choline phosphate cytidyltransferase-
383 alpha is a novel antigen detected by the anti-ERCC1 antibody 8F1 with biomarker
384 value in patients with lung and head and neck squamous cell carcinomas. *Cancer* 2014
385 **120** 1898-1907.
- 386 24. Lee SM, Falzon M, Blackhall F, Spicer J, Nicolson M, Chaudhuri A, Middleton G,
387 Ahmed S, Hicks J, Crosse B, Napier M, Singer JM, Ferry D, Lewanski C, Forster M,
388 Rolls SA, Capitanio A, Rudd R, Iles N, Ngai Y, Gandy M, Lillywhite R & Hackshaw
389 A. Randomized Prospective Biomarker Trial of ERCC1 for Comparing Platinum and
390 Nonplatinum Therapy in Advanced Non-Small-Cell Lung Cancer: ERCC1 Trial (ET).
391 *J Clin Oncol* 2017 **35** 402-411.
- 392 25. Scharer OD. Nucleotide excision repair in eukaryotes. *Cold Spring Harb Perspect Biol*
393 2013 **5** a012609.
- 394 26. Olausson KA & Soria JC. Validation of ERCC1-XPF immunodetection--letter.
395 *Cancer Res* 2010 **70** 3851-3852; author reply 3852.
- 396 27. Kirschner K & Melton DW. Multiple roles of the ERCC1-XPF endonuclease in DNA
397 repair and resistance to anticancer drugs. *Anticancer Res* 2010 **30** 3223-3232.
- 398 28. Evans E, Moggs JG, Hwang JR, Egly JM & Wood RD. Mechanism of open complex
399 and dual incision formation by human nucleotide excision repair factors. *EMBO J*
400 1997 **16** 6559-6573.
- 401 29. Aboussekhra A, Biggerstaff M, Shivji MK, Vilpo JA, Moncollin V, Podust VN, Protic
402 M, Hubscher U, Egly JM & Wood RD. Mammalian DNA nucleotide excision repair
403 reconstituted with purified protein components. *Cell* 1995 **80** 859-868.
- 404 30. Roca E, Berruti A, Sbiera S, Rapa I, Oneda E, Sperone P, Ronchi CL, Ferrari L,
405 Grisanti S, Germano A, Zaggia B, Scagliotti GV, Fassnacht M, Volante M, Terzolo M
406 & Papotti M. Topoisomerase 2alpha and thymidylate synthase expression in
407 adrenocortical cancer. *Endocr Relat Cancer* 2017 **24** 299-307.
- 408 31. Bergot E, Levallet G, Campbell K, Dubois F, Lechapt E & Zalcman G. Predictive
409 biomarkers in patients with resected non-small cell lung cancer treated with
410 perioperative chemotherapy. *Eur Respir Rev* 2013 **22** 565-576.
- 411 32. Olausson KA & Postel-Vinay S. Predictors of chemotherapy efficacy in non-small-cell
412 lung cancer: a challenging landscape. *Ann Oncol* 2016 **27** 2004-2016.
- 413
- 414
- 415
- 416
- 417
- 418

419 **Figure legends**

420

421 **Figure 1. Representative examples of nuclear ERCC1 immunostaining in adrenocortical tissue**
422 **samples using the monoclonal ERCC1 antibody clone 4F9.** A) Normal adrenal gland; B)
423 Adrenocortical carcinoma with high intensity and high percentage of positive cells (H-score 3). C)
424 Adrenocortical carcinoma with intermediate intensity and high percentage of positive cells (H-score
425 2). D) Adrenocortical carcinoma with low intensity and low percentage of positive cells (H-score 0,5).
426 Magnification 1x10.

427

428 **Figure 2. Relationship between ERCC1 expression and response to platinum-based**
429 **chemotherapy in 146 patients with adrenocortical carcinoma (ACC).** Progression-free survival
430 (A) and overall survival (B) during treatment (Kaplan-Meyer curves and log-rank test) in ACC
431 patients with high (H-score ≥ 2) and low staining (H-score ≤ 1) of ERCC1.

432

433 **Supplementary data**

434

435 **Supplementary Figure 1. Re-evaluation of the overall survival in the old series of 38 patients**
436 **with adrenocortical carcinoma treated with platinum-based chemotherapy** ¹³. (A) ERCC1
437 immunostaining with the 8F1 clone (old batch) (B) ERCC1 immunostaining with the new specific 4F9
438 clone.

439

440 **Supplementary Figure 2. Direct comparison between ERCC1 antibodies 4F9 (A) and C) and 8F1**
441 **clone (new batch) (B) and D))** in one normal adrenal gland (A) and B)) and in one adrenocortical
442 carcinoma (C) and D)). Magnification 1x20.

443

444