

## Study of states in $^{14}\text{C}$ via the $^{10}\text{Be}(4\text{He},4\text{He})^{10}\text{Be}$ reaction

Malcolm, J.D.; Freer, M.; Ashwood, N.I.; Curtis, N.; Munoz-Britton, T.; Wheldon, C.

DOI:

[10.1088/1742-6596/381/1/012077](https://doi.org/10.1088/1742-6596/381/1/012077)

### Document Version

Early version, also known as pre-print

### Citation for published version (Harvard):

Malcolm, J.D., Freer, M., Ashwood, N.I., Curtis, N., Munoz-Britton, T & Wheldon, C 2012, 'Study of states in  $^{14}\text{C}$  via the  $^{10}\text{Be}(4\text{He},4\text{He})^{10}\text{Be}$  reaction', *Journal of Physics: Conference Series*, vol. 381, no. 1.  
<https://doi.org/10.1088/1742-6596/381/1/012077>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Study of states in $^{14}\text{C}$ via the $^{10}\text{Be}(^4\text{He}, ^4\text{He})^{10}\text{Be}$ reaction

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2012 J. Phys.: Conf. Ser. 381 012077

(<http://iopscience.iop.org/1742-6596/381/1/012077>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 147.188.40.113

The article was downloaded on 28/09/2012 at 10:47

Please note that [terms and conditions apply](#).

# Study of states in $^{14}\text{C}$ via the $^{10}\text{Be}(^4\text{He},^4\text{He})^{10}\text{Be}$ reaction

J D Malcolm<sup>1</sup>, M Freer<sup>1</sup>, N I Ashwood<sup>1</sup>, N Curtis<sup>1</sup>,  
T Munoz-Britton<sup>1</sup>, C Wheldon<sup>1</sup>, V A Ziman<sup>1</sup>, W N Catford<sup>2</sup>,  
S Brown<sup>2</sup>, G Wilson<sup>2</sup>, N Soic<sup>3</sup>, D Bardayan<sup>4</sup>, S D Pain<sup>4</sup>,  
N L Achouri<sup>5</sup>, K Chipps<sup>6</sup> and K Crzywacz-Jones<sup>7</sup>

<sup>1</sup> School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

<sup>2</sup> School of Electronics and Physical Sciences, University of Surrey, Guildford, Surrey, GU2 7XH, UK

<sup>3</sup> Rudjer Boskovic Institute, Department of Experimental Physics, Bijenicka 54, HR-10000 Zagreb, Croatia

<sup>4</sup> Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

<sup>5</sup> Laboratoire de Physique Corpusculaire, ISMRA and Universite de Caen, IN2P3-CNRS, 14050 Caen Cedex, France

<sup>6</sup> Rutgers University, New Jersey, USA

<sup>7</sup> School of Physics and Astronomy, University of Tennessee, Knoxville, TN 37996, USA

E-mail: jxm354@bham.ac.uk

E-mail: M.Freer@bham.ac.uk

**Abstract.** A study of the  $^{10}\text{Be}(^4\text{He},^4\text{He})^{10}\text{Be}$  reaction has been performed at  $^{10}\text{Be}$  beam energies of 25.0, 27.0, 29.0, 32.0, 34.0, 38.0, 40.0, 42.0, 44.0 and 46.0 MeV. The measurements were to explore possible molecular rotational bands in  $^{14}\text{C}$ . Three states at excitation energies of  $E_x = 18.8, 19.76$  and  $20.66$  MeV have been measured and their spins have been determined to be  $5^-, 5^-$  and  $6^+$ , respectively.

## 1. Introduction

From previous work [1, 2] it is well known that  $^8\text{Be}$  has a well defined  $\alpha$ - $\alpha$  cluster structure. The ground state of  $^8\text{Be}$  is, however, unbound as the  $\alpha$ - $\alpha$  interaction is weak and the Pauli principle causes a repulsive core. The introduction of one more neutron into the system, to produce  $^9\text{Be}$ , has the effect of adding enough binding energy to stabilise the system. In this case the neutron orbits the  $\alpha$  cores in a delocalised region. Extending this system to add another neutron to create  $^{10}\text{Be}$  produces a system where there are  $\pi$  and  $\sigma$  molecular bonds binding the core [3]. There is now strong evidence supporting the molecular  $\alpha$  cluster structure in these isotopes [4, 5].


The same principles can be applied to carbon where the nucleus can be thought of as a three  $\alpha$  particle configuration. In neutron-rich carbon nuclei, such as  $^{13}\text{C}$  and  $^{14}\text{C}$ , the valence neutrons reside in delocalised molecular orbits around the core. The extra neutrons again have the effect of increasing the binding energy and creating a more stable system. It is expected that there are three possible configurations of the nucleus in  $^{14}\text{C}$ : a triangular configuration

with the valence neutrons forming  $\sigma$  bonds between two of the  $\alpha$  particles, a symmetrical linear configuration where the neutrons are spaced between the  $\alpha$  particles ( $\alpha$ - $n$ - $\alpha$ - $n$ - $\alpha$ ) and an asymmetric configuration where both neutrons are between two  $\alpha$  particles ( $\alpha$ - $2n$ - $\alpha$ - $\alpha$ ). The final configuration should give rise to two rotational bands with opposite parity.

In an attempt to study the potential prolate and oblate molecular band nature of  $^{14}\text{C}$  the spins and partial decay widths of resonances in  $^{14}\text{C}$  were measured.

## 2. Experimental Details

The resonant scattering reaction  $^{10}\text{Be}(^4\text{He},^4\text{He})^{10}\text{Be}$  was studied at beam energies of 25.0, 27.0, 29.0, 32.0, 34.0, 38.0, 40.0, 42.0, 44.0 and 46.0 MeV. This was performed at the Holyfield Radioactive Ion Beam Facility (HRIBF) at the Oak Ridge National Laboratory (ORNL), USA. In the experiment the  $^{10}\text{Be}$  beam was incident on gaseous  $^4\text{He}$ , which filled the reaction chamber at a pressure of 720 torr, providing a range of interaction energies as the beam was slowed by the gas. The gas was contained in the chamber by a 5 micron Havar window. The Q-value for the break-up of  $^{14}\text{C}$  into the constituent isotopes,  $^{10}\text{Be}$  and  $^4\text{He}$ , is  $-12.011$  MeV.




**Figure 1.** Schematic of the two ‘lampshade’ detectors.

Three sets of detectors were utilised in the experiment, two ‘lampshade’ detectors (shown in Fig. 1), each made from six silicon strip detectors, and a zero degree monitor. The lampshades were positioned 175 and 434 mm (to the back of the detectors) from the window with the silicon strips tilted at an angle of  $46^\circ$  to the beam axis. The twelve silicon strip detectors were 500 microns thick and segmented into 16 strips each, creating 96 channels per lampshade. For beam energies below 38 MeV the zero degree monitor consisted of two silicon surface barrier detectors (160 microns each) behind a 46 micron Mylar foil absorber. At 38 MeV and above the zero degree monitor was made from three surface barrier detectors (160 microns each) behind a 144 micron Mylar absorber. These were positioned 336 mm from the window.

The detectors were calibrated using  $\alpha$  particles from  $^{241}\text{Am}$  and  $^{244}\text{Cm}$ , and the elastic scattering of the  $^{10}\text{Be}$  beam from a gold target.


### 3. Analysis and Results

Calculations were made to correct for energy loss in the Havar window and energy loss of the beam as it traversed through the helium gas. The zero degree monitor was used for particle identification using E- $\Delta$ E. The energy deposited in the front detector is compared to that in the back detector, shown in Fig. 2, allowing the recoil  $^4\text{He}$  to be identified.




**Figure 2.** Particle identification plot for the zero degree monitors. Units are arbitrary

Using the monitor and correlating the data from the multiple beam energies a spectrum (Fig. 3) showing the states in  $^{14}\text{C}$  can be constructed. The lampshade detectors were used to determine the spins of the states identified by the zero degree monitors. Both the  $^4\text{He}$  and the  $^{10}\text{Be}$  particles were detected by the lampshades in opposite silicon detectors.




**Figure 3.** Normalised excitation energy spectrum of  $^{14}\text{C}$  from the zero degree monitors.


The energy deposited in a detector was plotted against the angle to produce a two dimensional plot shown in Fig. 4. Resonances appear on the plot periodically along a line as shown in the figure. Once the resonance had been identified its spin was determined by comparing the periodicity of the data with simulated data. A range of spins were compared with the resonances shown in the two dimensional plots until a match was acquired.



**Figure 4.** Real data from the lampshade detectors. Units are arbitrary



**Figure 5.** Simulated data for the lampshades. Units are arbitrary.



**Figure 6.** Comparison of real data with simulated data of different spins.

Fig. 6 is an example of the comparison process that each of the resonances underwent. As spin is increased the periodicity of the angular distribution is also increased. By matching the periodicity of the real data to that of the simulated data allowed for identification of the spin of the states. Via this method three of the resonances determined in the spectrum of  $^{14}\text{C}$  could have their spins identified. These are the resonances at excitation energies,  $E = 18.8, 19.76$  and  $20.66$  MeV which have spins of  $5^-$ ,  $5^-$  and  $6^+$  respectively.

#### 4. Summary and Future Work

The  $^{10}\text{Be}(^4\text{He}, ^4\text{He})^{10}\text{Be}$  reaction has been studied at various  $^{10}\text{Be}$  beam energies ranging from 25 to 46 MeV. A spectrum showing clear energy states in  $^{14}\text{C}$  has been created and the spins of some of those states has been determined. The unknown spins of the other resonances shown in Fig. 3 will be calculated using R-Matrix analysis. Once the spins of the states have been deduced possible rotational bands may become evident.

#### References

- [1] Buck B, Friedrich H and Wheatley C 1977 *Nucl. Phys. A* **275** 246
- [2] Hiura J and Tamagaki R 1972 *Suppl. Prog. Theor. Phys.* **52** 25
- [3] Kanada-En'yo Y, Horiuchi H and Dote A 1999 *Phys. Rev. C* **60** 064304
- [4] von Oertzen W 1997 *Z. Phys. A* **357** 355
- [5] Freer M *et al* 1999 *Phys. Rev. Lett.* **82** 1383